

# Quantum Mechanics

Stephen P. Martin

[spmartin@niu.edu](mailto:spmartin@niu.edu)

August 12, 2025



# Preface

This book is intended as a text for a core graduate course on quantum physics. Quantum mechanics started to assume its modern form in a brilliant flash of activity from 1924-1930. In the subsequent century, there have been many excellent textbooks written on the topic, and one might well wonder why someone might decide to create another. The short answer is this: I've attempted to imagine, and then write, the book that would have made me happiest to read, as a graduate student. Whether my selection of topics, explanations, and derivations works well for someone else is an entirely different and personal question, of course!

Quantum mechanics is a big subject, and it is not possible to cover everything. So I didn't. Even so, there is more material in this book than can be reasonably covered in a typical academic year of lectures. This is intentional, because every course and every student will have different priorities. I did consider it essential to include certain modern subjects that are not always covered in older books at this level, notably the hidden variables alternative, Bell inequalities, entangled subsystems and open systems, generalized measurements, decoherence, and quantum information. The instructor will likely have to make some hard choices of what to leave out, informed by the interests and preparation of the students.

Although this book aims to be a self-contained discussion, starting from the basics, it will probably be more readily accessible for those who have already taken undergraduate courses in classical mechanics, electromagnetism, and quantum mechanics, with at least some exposure to the concepts of wavefunctions, operators, commutators, Schrödinger's equation, and the uncertainty principle. It is also assumed that the reader is already familiar with matrices and linear algebra, multivariable calculus, simple differential equations, the algebra of complex numbers, and basic concepts of probability.

The exercises at the end of each chapter vary in length and difficulty. For this I make no apology, since at the wild frontiers of research, problems do not present themselves with convenient labels saying "I am easy, you can solve me in one line" or "I am challenging, good luck". With that said, none of the exercises given here are remotely close to research level, and each has been chosen to reinforce some concept or method.

I am grateful to the many students who have provided feedback, with special thanks to those who pointed out corrections to earlier drafts: Sameneh Ahmadijad, Jared Coles, Matthew Dudak, William Emark, David Iglesias Tinoco, Abdulrahman Kauther, Spencer Kelham, Mark Mekosh, Dillon Merenich, Olabisi Olayinka, Zhenghao Pan, Cassandra Phillips, Deeksha Sinha, Vladimir Tsitrin, and Tasfia Yeashna.

Stephen P. Martin  
Northern Illinois University, DeKalb, Illinois, USA

# Contents

<b>1</b>	<b>Introduction: the quantum revolution</b>	<b>8</b>
1.1	Classical instability of charged matter . . . . .	8
1.2	Black-body radiation and Planck's constant . . . . .	10
1.3	Photo-electric effect and particle-like features of light . . . . .	12
1.4	Electron diffraction and wave-like features of matter . . . . .	15
1.5	Spin and the Stern–Gerlach experiment . . . . .	16
1.6	Exercises . . . . .	21
<b>2</b>	<b>Math tools: states, operators, and representations</b>	<b>23</b>
2.1	Complex linear vector spaces . . . . .	23
2.2	Inner products, Hilbert spaces, and orthonormal bases . . . . .	26
2.3	Dual vector spaces . . . . .	32
2.4	Operators . . . . .	33
2.5	Matrix representations of operators . . . . .	39
2.6	Eigenvalues and eigenvectors . . . . .	44
2.7	Observables . . . . .	53
2.8	Wavefunctions . . . . .	56
2.9	Tensor product Hilbert spaces . . . . .	66
2.10	Exercises . . . . .	68
<b>3</b>	<b>The core principles of quantum mechanics</b>	<b>71</b>
3.1	The basic postulates . . . . .	71
3.2	Valid and invalid questions . . . . .	73
3.3	Expectation values and uncertainties . . . . .	76
3.4	How states change . . . . .	78
3.5	Mixed ensembles and the density matrix operator . . . . .	85
3.6	Exercises . . . . .	95
<b>4</b>	<b>Canonical variables and the Hamiltonian</b>	<b>98</b>
4.1	Quantum observables from classical dynamics . . . . .	98
4.2	The two-body problem . . . . .	100
4.3	Charged particle in external electromagnetic fields . . . . .	102
4.4	Exercises . . . . .	107
<b>5</b>	<b>Transformations, symmetries, and conservation laws</b>	<b>108</b>
5.1	Continuous unitary transformations, symmetries, and Noether's principle . . . . .	108
5.2	Translations . . . . .	111
5.3	Rotations . . . . .	115
5.4	Parity . . . . .	121
5.5	Gauge transformations . . . . .	123
5.6	Currents and local conservation of probability . . . . .	128
5.7	Exercises . . . . .	131
<b>6</b>	<b>Particle moving in one dimension</b>	<b>132</b>
6.1	Gaussian wavefunctions . . . . .	132
6.2	Time evolution of free-particle state in one dimension . . . . .	134
6.3	Properties of stationary states in one-dimensional potentials . . . . .	137
6.4	Particle in a one-dimensional box . . . . .	140

6.5	Bound states for the one-dimensional square well . . . . .	142
6.6	Scattering problems in one dimension . . . . .	146
6.7	Particle acted on by a constant force . . . . .	153
6.8	Exercises . . . . .	156
<b>7</b>	<b>The harmonic oscillator</b>	<b>159</b>
7.1	The unreasonable effectiveness of the harmonic oscillator . . . . .	159
7.2	Position and momentum representations: the differential equations approach . . . . .	162
7.3	Energy representation: the algebraic approach . . . . .	169
7.4	Coherent states of the harmonic oscillator . . . . .	174
7.5	Three-dimensional harmonic oscillator . . . . .	180
7.6	Exercises . . . . .	182
<b>8</b>	<b>Angular momentum and its representations</b>	<b>186</b>
8.1	The eigenvalue problem for angular momentum . . . . .	186
8.2	The $j = 1/2$ representation: Pauli matrices and spin . . . . .	190
8.3	Matrix representation for $j = 1$ . . . . .	193
8.4	Matrix representation for arbitrary $j$ . . . . .	194
8.5	Matrix representations for unitary rotation operators . . . . .	195
8.6	Spherical harmonic representation of orbital angular momentum . . . . .	199
8.7	Parity of angular momentum eigenstates . . . . .	208
8.8	Exercises . . . . .	210
<b>9</b>	<b>Charged particle in a magnetic field</b>	<b>212</b>
9.1	Spin precession in a constant magnetic field . . . . .	212
9.2	Magnetic spin resonance and Rabi oscillations . . . . .	215
9.3	Landau levels for a charged particle moving in a constant uniform magnetic field . . . . .	218
9.4	Exercises . . . . .	226
<b>10</b>	<b>Examples with spherical symmetry</b>	<b>228</b>
10.1	Stationary states with spherical symmetry . . . . .	228
10.2	Free particle in spherical coordinates . . . . .	229
10.3	Particle confined to a sphere . . . . .	236
10.4	Particle in a spherical potential well . . . . .	239
10.5	Isotropic three-dimensional harmonic oscillator . . . . .	240
10.6	Exercises . . . . .	245
<b>11</b>	<b>Coulomb potential and hydrogen-like atoms</b>	<b>248</b>
11.1	Bound states for hydrogen atom . . . . .	248
11.2	Unbound states of Coulomb potentials . . . . .	261
11.3	Exercises . . . . .	264
<b>12</b>	<b>Addition of angular momenta</b>	<b>266</b>
12.1	Statement of the problem . . . . .	266
12.2	Addition of two spins . . . . .	268
12.3	Addition of orbital angular momentum and spin . . . . .	272
12.4	The general case and Clebsch–Gordan coefficients . . . . .	275
12.5	How spherical harmonics combine . . . . .	285
12.6	Exercises . . . . .	288

<b>13 Tensor operators and useful rules they obey</b>	<b>291</b>
13.1 Irreducible tensor operators . . . . .	291
13.2 Selection rules for scalar and vector operators and the Landé projection formula . . . .	295
13.3 The Wigner–Eckart Theorem and selection rules for tensor operators . . . . .	299
13.4 Exercises . . . . .	302
<b>14 Entanglement, the EPR problem, hidden variables, and Bell inequalities</b>	<b>304</b>
14.1 The Einstein–Podolsky–Rosen problem in Bohm’s formulation . . . . .	304
14.2 Hidden variables and Bell’s inequality . . . . .	309
14.3 Quantum mechanics vs. hidden variables, without inequalities . . . . .	313
14.4 Aspect’s experiments and the demise of local hidden variables . . . . .	315
14.5 Exercises . . . . .	319
<b>15 Stationary-state perturbation theory</b>	<b>321</b>
15.1 Perturbative expansion for energy eigenstates . . . . .	321
15.2 Simple examples of perturbation theory . . . . .	326
15.3 Helium atom ground state from first-order perturbation theory . . . . .	332
15.4 Brillouin–Wigner perturbation theory . . . . .	334
15.5 Dalgarno–Lewis method for simplifying perturbation theory . . . . .	335
15.6 Degenerate perturbation theory . . . . .	337
15.7 Hydrogen atom in electric field (Stark effect) . . . . .	343
15.8 Almost-degenerate perturbation theory . . . . .	348
15.9 Exercises . . . . .	352
<b>16 The variational method</b>	<b>355</b>
16.1 Estimate and upper bound on the ground state energy . . . . .	355
16.2 Variational method for excited states . . . . .	356
16.3 Examples of the variational method . . . . .	358
16.4 Helium atom ground state from the variational method . . . . .	361
16.5 Exercises . . . . .	363
<b>17 Fine, hyperfine, and magnetic effects for the hydrogen atom</b>	<b>365</b>
17.1 Relativistic kinetic, spin-orbit, and Darwin corrections . . . . .	365
17.2 Hyperfine structure of hydrogen . . . . .	369
17.3 Hydrogen atom in external magnetic field (Zeeman and Paschen–Back effects) . . . . .	375
17.4 Exercises . . . . .	381
<b>18 Identical particles and multi-electron atoms</b>	<b>383</b>
18.1 Intrinsic indistinguishability of identical particles . . . . .	383
18.2 Wavefunctions and spin for two identical particles . . . . .	391
18.3 Excited states of the helium atom . . . . .	395
18.4 Multi-electron atoms . . . . .	398
18.5 Exercises . . . . .	412
<b>19 Simple molecules</b>	<b>415</b>
19.1 Hierarchies of scales and the Born–Oppenheimer approximation . . . . .	415
19.2 The simplest molecule: the $\text{H}_2^+$ ion . . . . .	419
19.3 Diatomic bonds and vibrations from the Morse potential . . . . .	422
19.4 Rigid bodies and molecular rotational states . . . . .	426
19.5 Vibrational and rotational spectra for diatomic molecules . . . . .	429

19.6	The most important molecule: $\text{H}_2\text{O}$	435
19.7	Exercises	437
<b>20</b>	<b>Heisenberg and interaction representations</b>	<b>438</b>
20.1	The Heisenberg picture and equations of motion	438
20.2	The interaction picture and transition amplitudes	441
20.3	Exercises	445
<b>21</b>	<b>Time-dependent perturbation theory</b>	<b>446</b>
21.1	The short-time and sudden approximations	446
21.2	Transition amplitudes and probabilities in perturbation theory	448
21.3	Applying first-order time-dependent perturbation theory, and Fermi's golden rule	452
21.4	Harmonic time-dependent perturbations	457
21.5	Exercises	461
<b>22</b>	<b>Absorption and emission of light</b>	<b>463</b>
22.1	Electrons in the presence of electromagnetic waves	463
22.2	Absorption of electromagnetic waves	464
22.3	Induced and spontaneous emission of light	468
22.4	Electric dipole approximation	471
22.5	Magnetic dipole, electric quadrupole, and higher orders	475
22.6	Photo-electric effect for atoms	479
22.7	Exercises	484
<b>23</b>	<b>Scattering in three dimensions</b>	<b>486</b>
23.1	Cross-sections and scattering amplitudes	486
23.2	The scattering matrix ( $S$ -matrix) operator and the Lippmann–Schwinger equation	490
23.3	The Optical Theorem	501
23.4	Born approximation	502
23.5	Spherical potential scattering and the partial wave expansion	507
23.6	Bound states, resonances, and poles in scattering amplitudes	514
23.7	Examples of scattering from spherical potentials	519
23.8	Neutron-proton scattering and the deuteron	523
23.9	Scattering of identical particles	526
23.10	Exercises	529
<b>24</b>	<b>Subsystems: entanglement, evolution, and generalized measurements</b>	<b>532</b>
24.1	Open and bipartite systems and entanglement	532
24.2	Dynamical maps and evolution of open systems	537
24.3	Generalized measurements	545
24.4	Summary of rules of quantum mechanics for open systems	551
24.5	Local time evolution approximation and the Lindblad equation	553
24.6	Exercises	556
<b>25</b>	<b>Decoherence</b>	<b>557</b>
25.1	Emergence of classical behavior	557
25.2	Loss of coherence for a damped harmonic oscillator	561
25.3	Measurement and decoherence: phase damping of a spin	564
25.4	Decoherence for a spin in an exactly solvable model	567
25.5	Exercises	570

<b>26 Invitation to quantum information</b>	<b>571</b>
26.1 Qubits and the Bloch ball . . . . .	571
26.2 Quantum registers and parallelism . . . . .	573
26.3 Sending information using entanglement: teleportation and dense coding . . . . .	579
26.4 Copying quantum information: the no-cloning theorem and redundant encoding . . . . .	584
26.5 Keeping secrets with quantum private key generation . . . . .	586
26.6 Fighting decoherence and other faults with error correction . . . . .	588
26.7 Exercises . . . . .	596
<b>27 Relativistic quantum mechanics</b>	<b>597</b>
27.1 Special relativity, four-vectors, and Lorentz transformations . . . . .	597
27.2 Klein-Gordon wave equation . . . . .	601
27.3 Dirac equation . . . . .	603
27.4 Electromagnetic fields and the Dirac equation . . . . .	609
27.5 Dirac equation solutions for the hydrogen atom . . . . .	613
27.6 Exercises . . . . .	618
<b>28 Feynman path integral approach</b>	<b>620</b>
28.1 Propagators . . . . .	620
28.2 Summing over paths . . . . .	623
28.3 Evaluation of the path integral for the harmonic oscillator . . . . .	628
28.4 The Ehrenberg–Siday and Aharonov–Bohm effect . . . . .	633
28.5 Dirac quantization condition for magnetic monopoles . . . . .	636
28.6 Classical limit from the sum over paths . . . . .	640
28.7 Exercises . . . . .	642
<b>Index</b>	<b>643</b>



# 1 Introduction: the quantum revolution

## 1.1 Classical instability of charged matter

Quantum mechanics is often portrayed as nonintuitive, weird, or even paradoxical. We begin with a retort: it is actually classical physics that is conspicuously incompatible with basic features of the world we live in. Most strikingly, if classical mechanics governed the universe, then matter made up of charged particles would necessarily be unstable against rapid and catastrophic collapse.

Consider, for example, a classical model of the hydrogen atom, consisting of a point-like<sup>†</sup> electron and much heavier proton, separated by a distance  $r$ . The electric potential energy is

$$V(r) = -\frac{Q_e^2}{4\pi\epsilon_0 r} = -\frac{e^2}{r}, \quad (1.1.1)$$

where in SI metric system units,  $Q_e = -1.60218 \times 10^{-19}$  C is the electronic charge and  $\epsilon_0 = 8.85419 \times 10^{-12}$  C<sup>2</sup>/N · m<sup>2</sup> is the permittivity of free space. The positive quantity  $e$ , which will appear often in this book, is equal to the proton's charge in the Gaussian cgs metric system units. Its square is given numerically by

$$e^2 = 1.43996 \times 10^{-9} \text{ eV} \cdot \text{m} = 2.30708 \times 10^{-28} \text{ J} \cdot \text{m} = 2.30708 \times 10^{-19} \text{ erg} \cdot \text{cm}. \quad (1.1.2)$$

Clearly, classical physics has a serious problem: the potential  $V(r)$  is unbounded from below as  $r$  approaches 0, implying that a classical atom should release an arbitrarily large amount of energy as it shrinks to zero size.

While this may seem dangerous (or possibly useful, depending on your imagination!), it is certainly not what is observed. One might suppose that a way to achieve safety is to somehow force the electron to travel in a fixed orbit about the much heavier proton. However, this cannot work in a classical theory, because of energy conservation. Maxwell's equations imply that the classical electron will continuously lose energy in the form of electromagnetic radiation, due to its centripetal acceleration. For a circular orbit, the acceleration is

$$a = \frac{v^2}{r} = \frac{F}{m_e} = \frac{e^2}{m_e r^2}, \quad (1.1.3)$$

---

<sup>†</sup>The small size of the electron compared to atoms was understood in 1897 by Joseph J. Thomson from observations of cathode rays. All experiments to the present day are consistent with the electron having no substructure. Protons and nuclei are certainly not point-like, as they are now known to be composed of quarks and gluons. However, the important thing for the following discussion is just that they are tiny ( $\sim 10^{-15}$  meters) compared to atoms ( $\sim 10^{-10}$  meters). This had become apparent by 1911, before the development of the quantum theory, from the results of Ernest Rutherford's experiments with Hans Geiger and Ernest Marsden in which alpha particles were observed scattering at large angles off of gold nuclei.



where  $m_e = 9.109390 \times 10^{-31} \text{ kg} = 0.510999 \text{ MeV}/c^2$  is the electron's mass, with  $c = 2.99792458 \times 10^8 \text{ m/sec}$ , the speed of light in vacuum. The Larmor formula for the radiated power of an accelerating charge,

$$P = -\frac{dE}{dt} = \frac{Q_e^2 a^2}{6\pi\epsilon_0 c^3} = \frac{2e^2 a^2}{3c^3}, \quad (1.1.4)$$

therefore tells us that the classical electron must lose energy at a rate proportional to  $1/r^4$ , which in turn will decrease its orbit size at an ever-increasing rate.

Just for fun, let us estimate the tragic fate of the electronic orbit in this classical model, making some simplifying assumptions. If the orbit stays nearly circular, and nonrelativistic, as it decays, then the energy will be

$$E = \frac{1}{2}m_e v^2 - \frac{e^2}{r} = -\frac{e^2}{2r}. \quad (1.1.5)$$

Combining eqs. (1.1.4) and (1.1.5) gives

$$3r^2 \frac{dr}{dt} = -K, \quad (1.1.6)$$

where  $K \equiv 4e^4/m_e^2 c^3$  is a constant. If  $r = r_0$  at  $t = 0$ , this integrates to  $r^3 - r_0^3 = -Kt$ , so

$$r = r_0 \left(1 - Kt/r_0^3\right)^{1/3}. \quad (1.1.7)$$

This shows that the decay of the classical electron's orbit is even worse than asymptotic; it collapses all the way to  $r = 0$  in a finite time  $r_0^3/K$ , which turns out to be very short (see Exercise 1.1 to find out just how short). Larger atoms, and crystal structures of electrons and nuclei, would have similar instabilities if classical physics governed them.

Quantum mechanics addresses this catastrophe, and allows matter composed of charged particle constituents to be stable, by changing the rules. In the quantum theory, there is no counterpart to the decaying classical orbit with unbounded negative energy. In particular, there are no physical quantum states of the hydrogen atom with arbitrarily low energy, corresponding to a classical electron localized arbitrarily near  $r = 0$ . Instead, as we will see, there is a single state with the lowest possible energy (about 13.6 eV below a state of ionization), which therefore is stable since there is no lower energy state into which it could decay by emitting electromagnetic radiation. In this way, quantum mechanics saves the universe. More generally, in quantum mechanics the energies of bound states turn out to be quantized (discrete). There are also unbound (ionized) states with a continuum of allowed energies, but those energies are bounded from below.

## 1.2 Black-body radiation and Planck's constant

In the late 19th century, it became apparent that classical physics theory makes a similarly catastrophic prediction for electromagnetic radiation. Consider a cavity whose walls are as close as possible to idealized black (a perfect absorber and emitter of electromagnetic radiation), heated as an oven so that it is kept in thermal equilibrium at temperature  $T$  with the electromagnetic radiation inside it. The spectrum of the electromagnetic radiation is characterized by the energy density  $\rho_E$ , per unit volume  $V$  and per unit frequency  $\nu$ , in terms of which the total energy inside the cavity is

$$\text{Total energy} = V \int_0^\infty d\nu \rho_E(\nu, T). \quad (1.2.1)$$

The function  $\rho_E(\nu, T)$  can be determined experimentally by making a small hole in the walls of the cavity and measuring the radiation that escapes, analyzed for different frequencies using diffraction gratings, for example.

To obtain a theoretical prediction for  $\rho_E(\nu, T)$ , we first need to quantify the density of electromagnetic modes per unit frequency. For simplicity, assume that the cavity is a cubic box of side  $L$ , and that the electromagnetic radiation modes satisfy periodic boundary conditions, with fields proportional to  $\cos(\vec{k} \cdot \vec{r})$  and  $\sin(\vec{k} \cdot \vec{r})$ , with allowed wavevectors  $\vec{k} = (2\pi/L)\vec{n}$ , where  $\vec{n} = \hat{x}n_x + \hat{y}n_y + \hat{z}n_z$  for integers  $n_x, n_y, n_z$ . The corresponding wavelengths and frequencies are

$$\lambda = L/n, \quad \nu = cn/L, \quad (1.2.2)$$

where  $n = |\vec{n}| = \sqrt{n_x^2 + n_y^2 + n_z^2}$ . Because the allowed  $n_x, n_y$ , and  $n_z$  are integers, the number of modes with frequency between  $\nu$  and  $\nu + d\nu$  can be enumerated by integrating the volume in  $\vec{n}$  space, using radial coordinate  $n$ , with

$$d^3\vec{n} \rightarrow 4\pi n^2 dn = \frac{4\pi L^3}{c^3} \nu^2 d\nu. \quad (1.2.3)$$

So, if  $\overline{E}$  is the average energy of a mode with frequency  $\nu$  when in thermal equilibrium with the walls at temperature  $T$ , then the total energy for frequencies between  $\nu$  and  $\nu + d\nu$  is

$$2 \overline{E} \frac{4\pi L^3}{c^3} \nu^2 d\nu, \quad (1.2.4)$$

where the first factor of 2 accounts for the fact that each electromagnetic mode can have two transverse polarizations. Setting eq. (1.2.4) equal to  $L^3 \rho_E d\nu$  in accord with eq. (1.2.1), we find

$$\rho_E(\nu, T) = \frac{8\pi}{c^3} \nu^2 \overline{E}. \quad (1.2.5)$$

We next need to evaluate  $\overline{E}$  as a function of temperature.

A fundamental prediction of statistical mechanics is that for an ensemble of identical systems in thermal equilibrium, the probability for one of the systems to have energy  $E$  is proportional to the **Boltzmann factor**, named after Ludwig Boltzmann,

$$P(E) \propto e^{-E/k_B T}, \quad (1.2.6)$$

where  $T$  is the temperature in Kelvin, and Boltzmann's constant

$$k_B = 1.380649 \times 10^{-23} \text{ J/K} = 1.380649 \times 10^{-16} \text{ erg/K} = 8.617333 \times 10^{-5} \text{ eV/K} \quad (1.2.7)$$

is the conversion factor between units of temperature and energy. If the allowed energies of electromagnetic modes are continuous and unrestricted, we therefore have the simple classical prediction for the average energy,

$$\overline{E} = \left( \int_0^\infty dE e^{-E/k_B T} E \right) / \int_0^\infty dE e^{-E/k_B T} = k_B T, \quad (1.2.8)$$

independent of  $\nu$ . Plugging this into eq. (1.2.5) gives the result

$$\rho_E(\nu, T) = 8\pi k_B T \nu^2 / c^3. \quad (1.2.9)$$

This is the Rayleigh–Jeans prediction, developed by John W. Strutt, 3rd Baron Rayleigh, and James Jeans. Even without consulting experimental data, this formula is clearly problematic, since it claims that the energy density grows quadratically with frequency. After integrating  $\int_0^\infty d\nu \rho_E(\nu, T)$ , we would find an infinite total energy density per unit volume in black-body radiation. This impossible prediction of classical physics is called the ultraviolet catastrophe.

In order to explain the existing data and avoid the ultraviolet catastrophe, Max Planck proposed in 1900 that the black-body walls can only emit modes with frequency  $\nu$  if the energy is equal to an integer multiple of  $h\nu$ , where, using modern data,

$$h = 4.13567 \times 10^{-15} \text{ eV}\cdot\text{s} = 6.62607 \times 10^{-34} \text{ J}\cdot\text{s} = 6.62607 \times 10^{-27} \text{ erg}\cdot\text{s} \quad (1.2.10)$$

is known as **Planck's constant**. In that case, the integrals in eq. (1.2.8) are replaced by sums over only the discrete allowed energies,

$$\overline{E} = \left( \sum_{n=0}^{\infty} n h \nu e^{-n h \nu / k_B T} \right) / \sum_{n=0}^{\infty} e^{-n h \nu / k_B T} = \frac{h \nu}{e^{h \nu / k_B T} - 1}. \quad (1.2.11)$$




Figure 1.2.1: Comparison of the classical Rayleigh–Jeans (dashed line) and Planck (solid curve) predictions for the electromagnetic radiation energy per unit volume and unit frequency,  $\rho_E$ , in a black-body cavity, as a function of the frequency  $\nu$ , for temperature  $T = 1000\text{K}$ .

For very small  $h\nu/k_B T$ , this agrees with eq. (1.2.8). However, unlike that formula, it depends on the frequency, and is exponentially suppressed in the ultraviolet limit of large  $h\nu/k_B T$ . Putting eq. (1.2.11) into eq. (1.2.5) yields<sup>†</sup>

$$\rho_E(\nu, T) = \frac{8\pi h}{c^3} \frac{\nu^3}{e^{h\nu/k_B T} - 1} \quad (1.2.12)$$

Figure 1.2.1 compares Planck’s formula eq. (1.2.12) with the Rayleigh–Jeans prediction eq. (1.2.9), for black-body radiation with  $T = 1000\text{K}$ . In the far infrared, the two results agree, but they differ significantly in the near infrared, and the Planck prediction for  $\rho_E$  decreases with frequency above  $\nu = 5.9 \times 10^{13} \text{ Hz}$ . In the visible range of  $\nu$  between about  $4 \times 10^{14}$  and  $8 \times 10^{14} \text{ Hz}$ , the Planck formula is many orders of magnitude smaller, in accord with observation. Furthermore, the Planck prediction for the total energy density per unit volume

$$\int_0^\infty d\nu \rho_E(\nu, T) = \frac{8\pi^5 (k_B T)^4}{15h^3 c^3}, \quad (1.2.13)$$

is finite, resolving the ultraviolet catastrophe. The agreement of this prediction with observation established that the electromagnetic radiation modes inside the cavity are quantized in energy units equal to Planck’s constant multiplied by the frequency.

### 1.3 Photo-electric effect and particle-like features of light

The photo-electric effect is the ejection of electrons from the surface of a material when light shines on it. This is observed to occur if the angular frequency  $\omega$  of the light is sufficiently high.

---

<sup>†</sup>Historically, Planck’s original proposal of this formula was simply an inspired guess to fit the data, and his subsequent derivation of it was not particularly compelling by modern standards, but the result was correct. The origin of eq. (1.2.11) from deeper principles is derived in section 3.5, see eq. (3.5.47).




Figure 1.3.1: Sketch of results for the photoelectric effect. Shining electromagnetic radiation on the surface of a metal produces a current of ejected electrons, but to a good approximation this occurs only if the angular frequency of the light exceeds a minimum value  $\omega_{\min}$ . The maximum kinetic energy of the ejected electrons then rises like  $\hbar(\omega - \omega_{\min})$ .

For  $\omega$  less than a certain threshold value characteristic of the materials used in the experiment, almost no electrons are ejected, even as the intensity of the light is increased. The threshold angular frequency  $\omega_{\min}$  typically corresponds to visible or ultraviolet light. For  $\omega > \omega_{\min}$ , the maximum energy of ejected electrons is observed to rise linearly, as sketched in Figure 1.3.1.

This behavior was unexpected when it was first found, because in classical electrodynamics, the frequencies and energies of electromagnetic waves are continuous and independent of each other. One might have supposed that increasing the intensity of the light would result in electrons being ejected for any  $\omega$ , no matter how small. To explain the observations, Albert Einstein proposed in 1905 that light of a given angular frequency always occurs in chunks, or quanta, which are now called photons. For each photon, the energy is related to the angular frequency by the same formula proposed by Planck, which can be rewritten, using  $\omega = 2\pi\nu$ , as

$$E = \hbar\omega, \quad (1.3.1)$$

where the **reduced Planck's constant** (or just “h bar”, when speaking) is defined to be related to the ordinary Planck's constant by

$$\hbar = \frac{h}{2\pi} = 6.58212 \times 10^{-16} \text{ eV}\cdot\text{s} = 1.05457 \times 10^{-34} \text{ J}\cdot\text{s} = 1.05457 \times 10^{-27} \text{ erg}\cdot\text{s}. \quad (1.3.2)$$

Einstein's quantization condition conceptually generalized Planck's proposal, which was only intended to apply to electromagnetic modes absorbed and emitted by the black-body cavity.

It follows from special relativity and wave kinematics that the momentum  $p$  and the wavelength  $\lambda$  of each photon are related by

$$p = E/c = \hbar\omega/c = 2\pi\hbar/\lambda. \quad (1.3.3)$$

In the photo-electric effect, the discrete particle-like nature of light explains the existence of  $\omega_{\min}$ , because  $\hbar\omega_{\min}$  is the minimum energy jump needed for the electron to escape the metal




Figure 1.3.2: The wave nature of electromagnetic radiation causes interference and diffraction phenomena when light from a coherent source passes through holes or slits in a screen.

and be collected on another plate when it is struck by a single photon.<sup>†</sup> Experiments reported by Robert A. Millikan in 1916 later verified, using sodium and lithium as the targets, that the maximum kinetic energy of the ejected electron indeed behaves like

$$E = \hbar(\omega - \omega_{\min}). \quad (1.3.4)$$

The value of  $\omega_{\min}$  depends on the collector plate material as well as the material being illuminated, but the most important result of the experiment is the slope of the line in Figure 1.3.1, which is  $\hbar$ . Millikan's measured value for  $\hbar$  was consistent with Planck's result from black-body radiation, but was significantly more accurate and precise.

Although light behaves like a particle in the photo-electric effect, it also has interference properties governed by the wavelength. These effects appear in interference and diffraction experiments like the double-slit experiment illustrated in Figure 1.3.2. Light from a coherent source passing through holes or slits in a screen yields a pattern of intensity maxima and minima, where the interference of amplitudes is constructive and destructive, respectively. However, the interpretation of this effect is slightly different in quantum mechanics than in the corresponding classical theory. Classically, the intensity is proportional to the magnitude of the time-averaged Poynting vector,

$$\text{Classical intensity} \propto |\vec{E} \times \vec{B}|, \quad (1.3.5)$$

where  $\vec{E}$  and  $\vec{B}$  are the electric and magnetic fields that each obey superposition and interference due to the linearity of Maxwell's equations. In the quantum theory, the *real* quantity  $\vec{E} \times \vec{B}$  is replaced by the squared magnitude of a *complex* quantity  $\psi$ ,

$$\text{Quantum intensity} \propto |\psi|^2. \quad (1.3.6)$$

---

<sup>†</sup>Actually, two photons can team up to eject a single electron, as has been observed in experiments with high-power lasers, but the rate for this is very small except when the intensity is extremely large.

Here  $\psi(x, y, z, t)$  is called a **wavefunction**, and it is a probability density amplitude. This means that  $|\psi(x, y, z, t)|^2$  is proportional to the probability to detect a photon at position  $(x, y, z)$  at time  $t$ . It is a fundamental feature of quantum mechanics that we must deal with probabilities, not definite outcomes for individual events, even if the initial conditions of a physical situation are known as perfectly as possible.

## 1.4 Electron diffraction and wave-like features of matter

Inspired by the dual wave-particle nature of light exhibited in the photoelectric effect, Louis de Broglie in his 1924 PhD thesis suggested that matter particles, such as electrons, will also have wavefunctions  $\psi(x, y, z, t)$  that are subject to interference. He proposed that what is now called the **de Broglie wavelength**  $\lambda$  (and the corresponding wavenumber  $k$ ) of the wavefunction should be related to the momentum of a particle in the same way as for light, eq. (1.3.3). That is, for a particle with mass  $m$  and energy  $E = \sqrt{p^2 c^2 + m^2 c^4}$ ,

$$\lambda = 2\pi/k = 2\pi\hbar/p. \quad (1.4.1)$$

De Broglie's wave hypothesis for matter was verified by the discovery of electron diffraction in a series of experiments by Clinton Davisson and Lester Germer from 1923-1928.

In the Davisson–Germer experiments, electrons with controlled energies (of order a few hundred eV) were made to hit a target made of a crystal of nickel,<sup>†</sup> and a detector was used to observe the intensity of electrons scattered at various fixed angles, as shown schematically in the first panel of Figure 1.4.1. The results as a function of varying incident electron momentum (proportional to the inverse de Broglie wavelength) are shown in the second panel. They feature maxima and minima of constructive and destructive interference, depending on the differences in path lengths of the electrons scattering from the regularly spaced crystal sites, with a distance between nuclei of order  $3.5 \times 10^{-10}$  meters. This interference effect was similar to the previously known phenomenon of Bragg peaks in x-ray scattering. The electrons' wavelength corresponded to the prediction of the de Broglie relation to momentum in eq. (1.4.1). This provided the first direct evidence that matter particles are also described by a wavefunction which can be interpreted as a probability amplitude, with the crucial feature that this amplitude is subject to superposition and interference. It also provided another example of the central role of Planck's constant in the quantum theory, through the connection between momentum, wavelength, and  $\hbar$  in eq. (1.4.1).

---

<sup>†</sup>Their original motivation was to study the surface of nickel, not to check de Broglie's idea, which came later.






Figure 1.4.1: Schematic setup for the Davisson–Germer electron diffraction experiments (left), and some of their data for intensity of electrons scattered at fixed angle, as a function of the reciprocal of the electron’s de Broglie wavelength (right). Source: C.J. Davisson and L.H. Germer, “Reflection and refraction of electrons by a crystal of nickel”, Proceedings of the National Academy of Sciences of the USA, vol. 14, no. 8, p. 619 (1928).

## 1.5 Spin and the Stern–Gerlach experiment

In classical mechanics, angular momentum takes on continuous values, but in quantum mechanics it always occurs in multiples of a fundamental unit. For angular momentum associated with the motion of particles, called **orbital angular momentum**, the fundamental quantum unit is  $\hbar$ , as we will show in section 5.3. However, quantum mechanics also allows for **intrinsic angular momentum**, or **spin**, which has no classical counterpart. The spin of a particle is quantized in units of  $\hbar/2$ , as we will prove in section 8.1, and has a fixed magnitude that depends only on the identity of the particle. Electrons, protons, neutrons, muons, tau leptons, neutrinos, and quarks all carry this type of intrinsic angular momentum. Since their spin angular momentum is half of the quantized unit for orbital angular momentum, they are called spin-1/2 particles. Of these, all but the proton and neutron are currently believed to be fundamental (not composite collections of other particles). But even composite particles like atoms, atomic nuclei, and mesons and baryons bound together by the strong nuclear force, always have spin that comes in integer multiples of  $\hbar/2$ .

The first experimental hints of the existence of spin came from the otherwise mysterious doubling of certain spectral lines from atomic transitions. Wolfgang Pauli suggested in 1924 that this was due to the presence of some extra quantum number that could only take on two values. The following year, Samuel Goudsmit and George Uhlenbeck proposed the intrinsic angular momentum interpretation that we now know as spin. This idea took some time to gain




Figure 1.5.1: The Stern–Gerlach experiment. Silver atoms are heated in a furnace, and passed through an inhomogeneous magnetic field, which exerts a force on them proportional to the  $\hat{z}$  component of the atomic magnetic dipole moment. The observed result is that the silver beam splits into two “spatially quantized” components on the detection screen. In contrast, the (incorrect) classical prediction is a continuum of deflection magnitudes.

acceptance, and the many doubters famously included Pauli. Part of the problem was that, at first, the spin was erroneously thought to be due to some kind of internal rotational motion of the structure of the electron, but nobody could get this idea to work in detail, and it turns out to be incorrect. Spins of fundamental particles like the electron are now understood to be a form of angular momentum that is distinct from the mechanical type.

In 1922, Otto Stern and Walter Gerlach reported an experiment which eventually provided profound insights into spin and the emerging quantum theory. They heated silver atoms in a furnace to vaporize them. The atoms escaped through a narrow collimating structure to form a beam, which then moved through a region where they were deflected by an inhomogeneous magnetic field, and finally were collected on a measurement screen, as shown in Figure 1.5.1.

The total magnetic dipole moment for the silver atom is a vector  $\vec{\mu}$  whose magnitude  $\mu$  is a constant, very nearly the same as that of a single electron,<sup>†</sup> which in turn is proportional to the electron’s spin, although Stern and Gerlach did not know this at the time. The furnace thoroughly randomizes the directions of the magnetic moments. As a result of the inhomogeneous magnetic field  $\vec{B}$ , there is a classical force on the atoms,

$$\vec{F} = \vec{\nabla}(\vec{\mu} \cdot \vec{B}), \quad (1.5.1)$$

causing them to deflect. For simplicity, suppose the magnetic field in the deflection region has

---

<sup>†</sup>The explanation for this is as follows. Silver atoms have 47 electrons, each of them carrying a magnetic dipole moment along that electron’s spin direction. However, 46 of the electrons pair up in such a way that their spins are opposite, and cancel. Furthermore, there is no net orbital angular momentum of the electrons, and the contribution of the heavy nucleus to the atomic magnetic moment is relatively small.

a cylindrically symmetric form<sup>‡</sup>

$$\vec{B}(x, y, z) = B_0 \left[ \hat{x} \frac{x}{2a} + \hat{y} \frac{y}{2a} + \hat{z} \left( 1 - \frac{z}{a} \right) \right] \quad (1.5.2)$$

where the coordinate system has unit vector  $\hat{z}$  (up in Figure 1.5.1) perpendicular to the beam, and  $a$  is a length that is large compared to the size of the region where the magnetic field acts on the atoms in the beam. The atomic dipole moment experiences a torque, and since  $\vec{\mu}$  is proportional to the angular momentum, it obeys an equation of motion of the form

$$\frac{d\vec{\mu}}{dt} \propto \vec{\mu} \times \vec{B}. \quad (1.5.3)$$

The dominant magnetic field component  $B_0 \hat{z}$  therefore causes  $\vec{\mu}$  to rotate rapidly about the  $\hat{z}$  direction, keeping the magnitude of  $\mu_z$  nearly constant but causing the oscillating  $\mu_x$  and  $\mu_y$  components to average to 0 over the time scale in which the atom is moving through the macroscopic magnetic field region. This implies that  $\mu_x$  and  $\mu_y$  can be neglected when computing the deflection of the atom. Since only  $\mu_z$  contributes, eq. (1.5.1) becomes

$$\vec{F} = \mu_z \vec{\nabla} B_z = \hat{z} \mu_z \frac{\partial B_z}{\partial z} = -\hat{z} (B_0/a) \mu_z. \quad (1.5.4)$$

Because the gradient of the vertical magnetic field  $-B_0/a$  is known and fixed, measuring the deflection of the atom is equivalent to measuring the  $\hat{z}$  component of its magnetic dipole moment.

Note that in the limit of a homogeneous field ( $a \rightarrow \infty$ ) there would be no deflection at all. Thus, the role of the large homogeneous part of the magnetic field  $\hat{z} B_0$  is to determine which component of  $\vec{\mu}$  will be measured, by washing out the effects of the other components, while the smaller inhomogeneous part  $-\hat{z} B_0/a$  provides the force needed to produce the deflection and actually make the measurement.

Since the magnetic dipole moments of silver atoms emerging from the furnace are random in direction, classical physics reasoning suggests that the measured values of  $\mu_z$  should have equal likelihood to be anything between  $-|\mu|$  and  $|\mu|$ . This in turn would imply that their deflections as observed on the screen should form a continuum between two extremes. Instead, they form two spots with equal numbers of atoms deposited, with a gap in between. The important and surprising conclusion is that the result of measuring  $\mu_z$  for silver atoms can only give two discrete, quantized values, with equal probabilities.

The amount of deflection can be related to the spin carried by each atom, which in the case of silver comes mostly from a single unpaired electron. The proportionality between magnetic

---

<sup>‡</sup>The  $\hat{x}$  and  $\hat{y}$  components of  $\vec{B}$  play no essential role here, except being necessary for consistency in order to satisfy the magnetostatic field equations  $\vec{\nabla} \cdot \vec{B} = 0$  and  $\vec{\nabla} \times \vec{B} = 0$ . More generally, the field need not be cylindrically symmetric, but at least one of  $B_x$  or  $B_y$  must be non-zero. The  $\vec{B}$  given here has the form that would result from a dipole magnet fixed at a distance  $3a$  below the beam, to linear order in  $1/a$ .

moment and spin for an electron is very close to

$$\vec{\mu} = -\frac{e}{m_e c} \vec{S}. \quad (1.5.5)$$

The numerical magnitudes of the deflections observed in the Stern–Gerlach experiment imply that the measurement of  $S_z$  for an electron can only give the values

$$\text{measured } S_z = \pm \hbar/2. \quad (1.5.6)$$

There is nothing special about the  $\hat{z}$  direction, so this also applies to  $\hat{n} \cdot \vec{S}$  for any unit vector  $\hat{n}$ .

The Stern–Gerlach experiment has been performed for other types of atoms and nuclei,<sup>§</sup> in which the angular momentum (and its relationship to the magnetic moment) can be different, resulting in more than two spots on the screen. This can be used to reveal the possible angular momentum properties (“quantum numbers”) of the atom in question. A Stern–Gerlach apparatus can even be used to isolate samples with particular desired angular momentum quantum numbers. The results of such experiments are always consistent with quantization of the components of any angular momentum vector in integer multiples of  $\hbar/2$ . The quantum theory must account for this property, and we will see how in Chapter 8.

One can have more general Stern–Gerlach analyzers (called  $\text{SG}\hat{n}$  in the following) with the inhomogeneous magnetic field element oriented in any chosen unit vector  $\hat{n}$  direction. The result of analyzing silver atoms fresh from the furnace with  $\text{SG}\hat{n}$  is that half of them will be found to have  $\hat{n} \cdot \vec{S} = +\hbar/2$  and the other half will have  $-\hbar/2$ , for any  $\hat{n}$ . Stern–Gerlach analyzers play a dual role in further efforts to understand quantum mechanics. First, they are measuring devices, if the output beams are sent directly to a detection screen. Second, because the outgoing beams are separated (sometimes called “spatial quantization”),  $\text{SG}\hat{n}$  provides a way of preparing a sample of atoms in which the spin component along  $\hat{n}$  is known to be either  $+\hbar/2$  or  $-\hbar/2$ . Instead of impacting a detection screen, one or both of the output beams can be sent off to some other component of the experiment, which could be another analyzer. Idealized versions of the Stern–Gerlach deflection analyzers are commonly used as modular components in experiments, real or imagined, to probe the implications of quantum theory.

For example, consider the sequence of two ideal Stern–Gerlach analyzers shown in Figure 1.5.2. The atoms in the experiment start in a furnace, which is assumed to produce completely randomized spins. After passing through a first analyzer  $\text{SG}\hat{z}$ , the atoms that had the result  $S_z = +\hbar/2$  are sent into a second analyzer  $\text{SG}\hat{z}$ , while those that had  $S_z = -\hbar/2$  are thrown

---

<sup>§</sup>However, the Stern–Gerlach setup does not work directly for free electrons, because the necessarily non-zero  $B_x$  and/or  $B_y$  cause a Lorentz force  $-e\vec{v} \times \vec{B}$  on the electron. This produces a large deflection due to the electron’s small mass, washing out the magnetic moment effect. The Stern–Gerlach setup relies on the deflected particles either being electrically neutral, or heavy, or both as in the case of silver atoms.




Figure 1.5.2: An experiment with two sequential Stern–Gerlach analyzers. The first analyzer prepares a sample of atoms with  $S_z = +\hbar/2$ , which are then fed into the second analyzer, which again measures  $S_z = +\hbar/2$ , with probability  $P = 1$  in the idealized case.




Figure 1.5.3: An experiment with three sequential Stern–Gerlach analyzers. The first analyzer prepares a sample of atoms with  $S_z = +\hbar/2$ , which are then fed into the second analyzer, which measures  $S_x = +\hbar/2$  and  $-\hbar/2$ , each with probability 0.5. The atoms with  $S_x = +\hbar/2$  are then sent to the third analyzer, which measures  $S_z = +\hbar/2$  and  $-\hbar/2$ , each with probability 0.5. The act of measuring  $S_x$  restores the possibility of measuring  $S_z = -\hbar/2$  at the end, even though the first analyzer had been used to select only atoms with  $S_z = +\hbar/2$ .

away. In this case, the prediction of quantum mechanics for the output of the second analyzer is unlikely to surprise anyone. All of the output atoms on the far right again have  $S_z = +\hbar/2$ ; the second analyzer simply confirms the measurement made by the first.

A more interesting setup is shown in Figure 1.5.3, which differs only by placing a  $\text{SG}\hat{x}$  analyzer between the two  $\text{SG}\hat{z}$  analyzers. As before, the experiment uses the first analyzer to select a pure sample of atoms with  $S_z = +\hbar/2$ . However, now the second analyzer separates the sample by measuring  $S_x$ . Since the  $\hat{z}$  direction has no way of preferring one of  $\pm\hat{x}$  over the other, it is no surprise that the output of  $\text{SG}\hat{x}$  is 50% for each of  $S_x = +\hbar/2$  and  $-\hbar/2$ .

The experiment in Figure 1.5.3 then throws away the atoms with  $S_x = -\hbar/2$ , and feeds those with  $S_x = +\hbar/2$  into a third analyzer  $\text{SG}\hat{z}$ . The final results for  $S_z$  can then be determined with a detection screen (not shown). One might perhaps suppose that we should find that the final atoms will all have  $S_z = +\hbar/2$ , since the first analyzer already selected only atoms with that property. However, this is wrong. In reality, an equal number are measured to have  $S_z = +\hbar/2$  and  $-\hbar/2$ . Inserting the  $\text{SG}\hat{x}$  analyzer in the middle of the chain affects the atoms in such a way as to restore the possibility of obtaining  $S_z = -\hbar/2$ .

Another way of thinking about the experiment shows that the restoration of the  $S_z = -\hbar/2$  outcomes is a logical necessity, assuming only that the spin is the only thing that makes a

difference. (In particular, this assumes that the velocity direction of the atoms has no impact on the spin measurement, and furthermore that there are no “hidden variables” associated with the atom that are involved in the measurements in some mysterious way that we do not know how to take into account.) To see this, cover up everything in Figure 1.5.3 except the last analyzer. Feeding directly into it are atoms with spin known to be aligned in the  $+\hat{x}$  direction. By assumption, the  $+\hat{x}$  direction has no reason to prefer  $+\hat{z}$  over  $-\hat{z}$ , or vice versa. So, no matter what may have occurred earlier, the only possibility is that the final probabilities for  $S_z = +\hbar/2$  and  $-\hbar/2$  are equal.

It is important that this result does *not* have anything to do with the fact that we threw away the atoms that were measured to have  $S_x = -\hbar/2$  coming out of the second analyzer. We could feed those atoms into the final analyzer as well, and they would also be measured to have probability  $P = 0.5$  for each of  $S_z = +\hbar/2$  and  $-\hbar/2$ , by the same argument. Evidently, it is the mere act of measuring  $S_x$  that causes the restoration of the  $S_z = -\hbar/2$  outcomes. When we give the postulates of quantum mechanics in Chapter 3.1, the act of measurement will play a special role in one of them (Postulate 5), consistent with the preceding discussion. This also carries with it the implication that quantum mechanics, as defined by these postulates, is an inherently probabilistic, rather than deterministic, theory.

If you are familiar with the behavior of polarizing filters for light, you may recognize that the preceding example is closely analogous to a similar experiment that is common and easy to do in optics. Two linear polarization filters arranged with axes of polarization at right angles will not allow any light to pass through, but a third polarizer inserted between the two, with axis at a  $45^\circ$  angle with respect to the axes of each of the others, will restore the transmission of a fraction of the light. This analogy carries over into the quantum regime. In fact, the most sensitive experiments testing quantum mechanics are often done with the polarization of photons replacing the role of the spin in Stern–Gerlach type experiments. One famous example will be discussed in detail in section 14.4.

## 1.6 Exercises

**Exercise 1.1.** Estimate the time in seconds needed for a classical hydrogen atom to reach zero size and infinite binding energy, under the assumptions used to obtain eq. (1.1.7), and taking  $r_0 = 5 \times 10^{-11}$  meters as the initial condition.


**Exercise 1.2.** The cosmic background radiation is nearly blackbody radiation with a present temperature of 2.73 Kelvin. Compute the numerical energy per cubic meter in this radiation:

(a) within the visible frequency range, defined here as  $4 \times 10^{14} \text{ Hz} < \nu < 8 \times 10^{14} \text{ Hz}$ , assuming eq. (1.2.9), the classical Rayleigh-Jeans formula. (For comparison, the energy density in starlight

is roughly  $10^{-15}$  J/m<sup>3</sup>.)

(b) integrated over all frequencies, as given by eq. (1.2.12), the quantum Planck formula. (Most of this is in the microwave spectrum; convince yourself that the visible range contribution is tiny.)

**Exercise 1.3.** (a) What velocity should an electron have so that its de Broglie wavelength will be 1 meter? What if its de Broglie wavelength is  $4.5 \times 10^{-7}$  meters (the same as for blue light)? (b) Electrons are accelerated from rest by a potential difference of 1 volt, and then pass through a screen with two very narrow long parallel slits. What is the de Broglie wavelength of the electrons? How far apart must the slits be in order for the first minimum of the interference pattern at a distant detector to be at an angle  $\theta = 0.2^\circ$  away from the central maximum?



**Exercise 1.4.** A coin is a thin disk of 30 grams of pure silver, with radius 2 centimeters. Each silver atom has one unpaired electron spin with magnitude  $\hbar/2$  and a nucleus that also has spin magnitude  $\hbar/2$ . In the (extraordinarily unlikely) case that all of these spins were lined up in the same direction, what would be the total intrinsic angular momentum of the (nonrotating) coin? How does this compare to the ordinary angular momentum that the coin would have if it rotated once per second about the symmetry axis perpendicular to the disk?



## 2 Math tools: states, operators, and representations

From the results of many experiments, physicists have arrived at a set of postulates that govern quantum theory. These will be given in the next chapter, but first we must introduce the mathematical language necessary to frame these ideas. This chapter therefore contains many definitions of key concepts, and some useful theorems. In doing so, we employ Paul A.M. Dirac's **bra-ket** notation, which is the modern standard in quantum mechanics.

### 2.1 Complex linear vector spaces

Quantum mechanics is based on the algebra of complex numbers. For any complex number

$$c = a + ib, \quad (2.1.1)$$

where  $a$  and  $b$  are real and  $i = \sqrt{-1}$ , the real and imaginary parts are denoted in this book by

$$\text{Re}[c] = a, \quad \text{Im}[c] = b, \quad (2.1.2)$$

and the **complex conjugate** is denoted using an asterisk,

$$c^* = (a + ib)^* = a - ib. \quad (2.1.3)$$

The **magnitude** (also known as the **modulus**) is defined by

$$|c| = |a + ib| = \sqrt{cc^*} = \sqrt{a^2 + b^2}. \quad (2.1.4)$$

It follows that  $1/c = c^*/|c|^2$  for every nonzero complex number  $c$ . Any complex number can be written as a product of its magnitude and a phase factor,

$$a + ib = |c|e^{i\phi}, \quad \text{with } \phi = \arctan(b/a), \quad (2.1.5)$$

where Euler's formula says that the unit-magnitude phase factor is

$$e^{i\phi} = \cos \phi + i \sin \phi. \quad (2.1.6)$$

We say that  $\phi$  is the phase of the complex number  $c$ .

A **complex linear vector space** is a set of vectors, also known as **kets**, denoted for example as  $|v\rangle, |w\rangle, \dots$ , such that the following properties hold:

- Additive closure: if  $|v\rangle$  and  $|w\rangle$  are kets, then so is their sum, or **superposition**,  $|v\rangle + |w\rangle$ .
- Multiplicative closure: if  $|v\rangle$  is a ket, then so is the product  $c|v\rangle$ , where  $c$  is any complex number.

- There exists a **null ket**,  $|\text{Null}\rangle$ , with the properties that  $|v\rangle + |\text{Null}\rangle = |v\rangle$  for every ket  $|v\rangle$ , and  $c|\text{Null}\rangle = |\text{Null}\rangle$  for every complex number  $c$ . Multiplying any ket by the complex number 0 results in the null ket:  $0|v\rangle = |\text{Null}\rangle$ .
- Addition of kets and multiplication by complex numbers satisfy the usual commutative, associative, and distributive properties. This means that we have  $|v\rangle + |w\rangle = |w\rangle + |v\rangle$ , and  $(|v\rangle + |w\rangle) + |x\rangle = |v\rangle + (|w\rangle + |x\rangle)$ , and  $c_1(c_2|v\rangle) = (c_1c_2)|v\rangle$ , and  $(c_1 + c_2)|v\rangle = c_1|v\rangle + c_2|v\rangle$ , and  $c(|v\rangle + |w\rangle) = c|v\rangle + c|w\rangle$ .

In quantum mechanics, the physical state of a system is completely described by a non-null ket, known as the **state ket** or **state vector**, which we will often denote by  $|\psi\rangle$ . However, one of the rules is that the ket  $c|\psi\rangle$  represents the same physical state as  $|\psi\rangle$ , provided that  $c$  is a non-zero complex number. This is true even though  $c|\psi\rangle$  and  $|\psi\rangle$  are mathematically distinct members of the vector space; they are identified with each other physically.

The null ket, although it is part of the vector space, cannot describe any physical state. There is also a convenient and obvious notational shortcut: if we want to write down that some expression is equal to the null ket, we will just write “= 0” instead of “=  $|\text{Null}\rangle$ ”. Therefore, the null ket will not appear explicitly any more, but one should always consider the possibility that a ket arising in some calculation might actually be the null ket. A warning: it is common to use the notation  $|0\rangle$  for some specific ket that does represent a physical state and must not be confused with the null ket. For example, depending on the system under consideration,  $|0\rangle$  might represent the lowest energy state of a system, also known as the **ground state**.

Let us list some examples of complex linear vector spaces, which you can check satisfy the properties in the definition.

Example 1: The set of complex numbers  $z$ .

Example 2: The set of all ordered triples  $(z_1, z_2, z_3)$ , where  $z_1$ ,  $z_2$ , and  $z_3$  are complex numbers.

Example 3: The set of all complex linear combinations  $c_1|\uparrow\rangle + c_2|\downarrow\rangle$  of two basic kets  $|\uparrow\rangle$  and  $|\downarrow\rangle$ . (This turns out to be the state space for a single spin-1/2 quantum system.)

Example 4: The set of all complex linear combinations of an infinite number of basic kets  $|0\rangle$ ,  $|1\rangle$ ,  $\dots$ ,  $|n\rangle$ ,  $\dots$ , in one-to-one correspondence with the non-negative integers. (This turns out to be a natural notation for the states of fixed energy for a harmonic oscillator in one dimension.)

Example 5: The set of all complex functions of a real variable,  $f(x)$ , defined on the domain  $-\infty < x < \infty$ . One can choose to add extra conditions on the functions in a variety of ways, for example requiring that they are continuous, or that they vanish at certain points or on specified intervals, or that they satisfy certain integrability conditions.

Intuitively, these vector spaces have different sizes. To make this precise, we define the

notions of linear independence and dependence, dimension, and basis. A set of kets  $|\psi_j\rangle$  are **linearly independent** if the equation  $\sum_j c_j |\psi_j\rangle = 0$  can only be satisfied by taking all  $c_j = 0$ . In other words,  $|\psi_j\rangle$  are linearly independent if we cannot write any of them as a complex linear combination of the others. Otherwise, the kets are **linearly dependent**.

A vector space is said to have **dimension**  $d$  if one can choose a set of  $d$ , but not more, linearly independent vectors. The  $d$  linearly independent vectors are then said to form a **basis** for the vector space. The choice of basis is certainly not unique for  $d > 1$ ; it is a common problem that one wants to change the choice of basis, either to make some calculation easier or to make some result simpler to interpret.

You can now check that for our five examples, the dimensions are:

Example 1 has dimension  $d = 1$ .

Example 2 has dimension  $d = 3$ .

Example 3 has dimension  $d = 2$ .

Example 4 has dimension  $d = \infty$  (countable, discrete basis).

Example 5 has dimension  $d = \infty$  (uncountable, continuous basis).

The cardinality (countable vs. uncountable) of the basis for infinite-dimensional vector spaces involves mathematical issues that we will find it convenient to mostly ignore. In physics, we are interested in kets that are members of a vector space with the additional structure of a Hilbert space with an inner product, as discussed in the next section. The requirement that all physical state vectors must have a finite inner product turns out to imply that the basis is always countable. Despite this, we will see that it is often a very convenient fiction to include continuous and uncountable sets of vectors that are not members of the physical state space but are nevertheless extremely useful, both as idealizations and in practical calculations. The most common examples are the kets that describe idealized states in which either a particle's position or its momentum (but not both!) is perfectly known. These continuous sets of kets can be chosen to obey the other properties of a basis, so we will often simply call them basis vectors, as a slight abuse of terminology.

Given a specific basis set  $\{|\beta_j\rangle\}$ , any vector  $|v\rangle$  can be expressed as

$$|v\rangle = \sum_{j=1}^d v_j |\beta_j\rangle \quad (2.1.7)$$

where the  $d$  complex numbers  $v_j$  are called the **components** of  $|v\rangle$  in that basis. Using the preceding definitions, one can show that, for a given  $|v\rangle$  and a given choice of basis  $\{|\beta_j\rangle\}$ , the components  $v_j$  are unique. In writing eq. (2.1.7) as a sum, we have implicitly assumed that the basis vectors are discrete and countable. If the basis kets are instead continuous,<sup>†</sup> then the sum

---

<sup>†</sup>Notice that we are already engaging in the slight abuse of terminology mentioned in the previous paragraph.

must be replaced by an integral. For example, if the basis set is  $\{|\beta_q\rangle\}$  where  $q$  is a continuous real variable with domain  $a < q < b$ , then we can write any ket  $|v\rangle$  as

$$|v\rangle = \int_a^b dq v(q) |\beta_q\rangle. \quad (2.1.8)$$

Here  $v(q)$  are the components, which in this case form a function of  $q$ . The symbol  $q$  might represent a coordinate on ordinary space, but it could also be a momentum, or an energy, or some other continuous quantity of interest.

A **subspace** of a vector space is a subset of the elements that also form a vector space by themselves. Below, we will often work with subspaces that consist of states with some feature in common, such as a fixed energy or fixed angular momentum magnitude.

## 2.2 Inner products, Hilbert spaces, and orthonormal bases

In quantum mechanics, the vector space of states has the additional structure of a **Hilbert space**, which implies that the complex linear vector space is endowed with an inner product. Given a ket  $|v\rangle$  and a second ket  $|w\rangle$ , the **inner product** (also known as a **scalar product**) returns a complex number, denoted  $\langle w|v\rangle$ , which must satisfy the following rules:

- The order matters, in such a way that exchanging the two kets gives the complex conjugate,

$$\langle v|w\rangle = (\langle w|v\rangle)^*. \quad (2.2.1)$$

It immediately follows that  $\langle v|v\rangle$  is always real. But also...

- If  $|v\rangle$  is not the null ket, then  $\langle v|v\rangle$  is positive.
- If either  $|v\rangle$  or  $|w\rangle$  is the null ket, then  $\langle v|w\rangle = \langle w|v\rangle = 0$ .
- Linearity is satisfied. The inner product of  $c_1 |v\rangle + c_2 |w\rangle$  and  $|x\rangle$ , in that order, is

$$\langle x| (c_1 |v\rangle + c_2 |w\rangle) = c_1 \langle x|v\rangle + c_2 \langle x|w\rangle. \quad (2.2.2)$$

It follows from eqs. (2.2.1) and (2.2.2) that the inner product of those same kets, but in the opposite order, must be

$$c_1^* \langle v|x\rangle + c_2^* \langle w|x\rangle = (c_1^* \langle v| + c_2^* \langle w|) |x\rangle. \quad (2.2.3)$$

The inner product should be thought of as similar to the dot product in the familiar three-dimensional real vector space. However, because quantum mechanics uses complex linear vector spaces, the inner product treats the two input vectors asymmetrically, and interchanging them is the same as taking the complex conjugate.

The **norm** of a ket  $|v\rangle$  is defined by  $\sqrt{\langle v|v\rangle}$ . It follows from the preceding that the norm of a non-null ket is real and positive,<sup>†</sup> and it is 0 if  $|v\rangle$  is the null ket. It is sometimes written as  $\| |v\rangle \| \equiv \sqrt{\langle v|v\rangle}$ .

Two useful inequalities that govern the inner product follow.

**Theorem 2.2.1. (Cauchy–Schwarz inequality)** *For any two kets  $|v\rangle$  and  $|w\rangle$ ,*

$$| \langle v|w\rangle |^2 \leq \langle v|v\rangle \langle w|w\rangle, \quad (2.2.4)$$

*Also, equality holds if and only if  $|w\rangle$  and  $|v\rangle$  are proportional to each other or one of them is the null ket.*

Proof: If either  $|v\rangle$  or  $|w\rangle$  is the null ket, then eq. (2.2.4) is trivially satisfied with equality. Therefore, we can assume for the remainder of the proof that neither of them is null. Consider the ket  $|z\rangle = c_1 |v\rangle - c_2 |w\rangle$ , where  $c_1$  and  $c_2$  are complex numbers. Since  $|z\rangle$  is a ket by the additive closure property,  $\langle z|z\rangle$  must be non-negative, which gives

$$\langle z|z\rangle = |c_1|^2 \langle v|v\rangle + |c_2|^2 \langle w|w\rangle - c_1 c_2^* \langle w|v\rangle - c_1^* c_2 \langle v|w\rangle \geq 0. \quad (2.2.5)$$

Choosing  $c_1 = \langle w|w\rangle$  and  $c_2 = \langle w|v\rangle$ , and using  $\langle v|w\rangle^* = \langle w|v\rangle$ , eq. (2.2.5) becomes

$$(\langle w|w\rangle)^2 \langle v|v\rangle - \langle w|w\rangle | \langle v|w\rangle |^2 \geq 0 \quad (2.2.6)$$

Now, since  $|w\rangle$  is not null, we can divide by  $\langle w|w\rangle$  to get eq. (2.2.4). If the equality condition holds, then it follows that  $|z\rangle$  is the null ket, which implies that  $|v\rangle$  and  $|w\rangle$  are proportional.  $\square$

**Theorem 2.2.2. (Triangle inequality)** *The norms of the kets  $|v\rangle$  and  $|w\rangle$  and their superposition  $|v\rangle + |w\rangle$  must obey*

$$\| (|v\rangle + |w\rangle) \| \leq \| |v\rangle \| + \| |w\rangle \|, \quad (2.2.7)$$

*with equality if and only if  $|w\rangle$  and  $|v\rangle$  are proportional or one of them is the null ket.*

The proof can be obtained from the Cauchy–Schwarz inequality, and is left to Exercise 2.1. The triangle inequality is similar to the statement in ordinary plane geometry that the sum of the lengths of two sides of a triangle must exceed that of the third side.

Given a non-null ket  $|v\rangle$ , one can define a new ket by dividing it by its norm. The result

$$|v\rangle / \sqrt{\langle v|v\rangle} \quad (2.2.8)$$

---

<sup>†</sup>In quantum field theories with gauge invariance, it is sometimes useful, as a book-keeping trick, to modify the rules by allowing some kets to satisfy  $\langle v|v\rangle \leq 0$ . However, these kets represent fictitious (unphysical) states, which must decouple from the true physical states. We will not encounter this issue in the present book.

then has norm 1, and in quantum mechanics it is physically equivalent to the original ket, in the sense that it represents the same physical state. We say that the ket has been normalized to unity. This still leaves the freedom to multiply the ket by a complex phase, because

$$e^{i\theta} |v\rangle \quad (2.2.9)$$

has the same norm as  $|v\rangle$ , for any real number  $\theta$ . This freedom will appear very often as an ambiguity in the determination of a ket that has to satisfy some other specified properties. The resolution of such ambiguities is arbitrary, and equivalent to a choice of convention.

Although the phase of a single ket is not a physically measurable quantity, the relative phases between different kets can be physically meaningful if one considers their sum, also known as their superposition. This is because if we multiply two non-null kets by different phases,

$$|v\rangle \rightarrow e^{i\theta_v} |v\rangle, \quad |w\rangle \rightarrow e^{i\theta_w} |w\rangle, \quad (2.2.10)$$

then their sum is replaced by

$$|v\rangle + |w\rangle \rightarrow e^{i\theta_v} (|v\rangle + e^{i(\theta_w - \theta_v)} |w\rangle), \quad (2.2.11)$$

which, by removing the overall phase  $e^{i\theta_v}$ , is physically equivalent to

$$|v\rangle + e^{i(\theta_w - \theta_v)} |w\rangle. \quad (2.2.12)$$

In general, this is not proportional to, and therefore not physically equivalent to,  $|v\rangle + |w\rangle$ , unless  $|v\rangle$  and  $|w\rangle$  are proportional, or  $\theta_w$  and  $\theta_v$  differ by an integer multiple of  $2\pi$ .

Two kets  $|v\rangle$  and  $|w\rangle$  are **orthogonal** (also known as **perpendicular**) if  $\langle w|v\rangle = 0$ . An **orthonormal basis**, which we will call an **orthobasis** for short, is a basis of kets  $\{|\varphi_j\rangle\}$  with  $j = 1, 2, \dots$  that satisfies the additional property

$$\langle \varphi_j | \varphi_k \rangle = \delta_{jk}. \quad (2.2.13)$$

Here  $\delta_{jk}$  is the **Kronecker delta symbol**,

$$\delta_{jk} = \begin{cases} 1 & (\text{for } j = k), \\ 0 & (\text{for } j \neq k), \end{cases} \quad (2.2.14)$$

and is defined only when  $j$  and  $k$  are labels that take on discrete values.

As noted in the previous section, we will also often want to deal with basis kets labeled by one or more continuous parameters. In that case, we need a different orthonormality condition, called **Dirac orthonormality**, in which the Kronecker delta symbol is replaced by a **Dirac**

**delta function.**<sup>‡</sup> The Dirac orthonormality condition for kets  $|\varphi_q\rangle$  labeled by a single continuous real variable  $q$  can be taken to be

$$\langle\varphi_q|\varphi_{q'}\rangle = \delta(q - q'). \quad (2.2.15)$$

Here  $q$  could be, for example, a position coordinate of a particle. The delta function  $\delta(x)$  has the properties that

$$\delta(x) = \begin{cases} \infty & (\text{for } x = 0), \\ 0 & (\text{for } x \neq 0), \end{cases} \quad (2.2.16)$$

and

$$\int_{-\infty}^{\infty} dx \delta(x) = 1, \quad (2.2.17)$$

or, more generally, for sufficiently well-behaved functions  $f(x)$ ,

$$\int_{-\infty}^{\infty} dx \delta(x - c) f(x) = f(c). \quad (2.2.18)$$

For most purposes, eq. (2.2.18) can be taken as the practical definition of the delta function.

An important technical note: as we have already warned in the previous section, kets satisfying the Dirac orthonormality condition eq. (2.2.15) cannot be physical states, strictly speaking. This is because they do not have finite norm, due to  $\delta(0) = \infty$ . Such kets with continuous labels are still very useful in practical calculations, where they often represent idealizations with perfectly known position, or momentum, or some other continuous quantity. These are sometimes known as **generalized kets** or **non-normalizable kets**, and they act as a basis for the Hilbert space of physical states, even though they are not themselves part of the Hilbert space. They are so useful as idealizations that mere non-normalizability is not a good enough reason for us to eliminate them from our toolbox.

It is often useful to think of  $\delta(x)$  as the “limit” (in a sense that we will not bother to try to make mathematically precise) of a sequence of increasingly narrow and sharply peaked functions that have unit area, in various different ways. For example, the sequence of functions can be taken to be rectangular functions with width  $\Delta$  and height  $1/\Delta$ ,

$$\delta(x) = \lim_{\Delta \rightarrow 0} \begin{cases} 1/\Delta & \text{for } |x| < \Delta/2, \\ 0 & \text{for } |x| > \Delta/2. \end{cases} \quad (2.2.19)$$

Another useful representation of  $\delta(x)$  is

$$\delta(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} dk e^{ikx}. \quad (2.2.20)$$

---

<sup>‡</sup>Technically, the Dirac delta function is not a function, but a distribution in mathematical language. But, this is not a math textbook, so we will not belabor the distinction.






Figure 2.2.1: The Dirac delta function  $\delta(x)$  can be thought of as the  $\Delta \rightarrow 0$  “limit” of functions peaked near  $x = 0$  with width of support proportional to  $\Delta$  and unit integrated area, as in the rectangular functions in eq. (2.2.19) [left], the Gaussian functions in eq. (2.2.21) [center], or the normalized sine functions in eq. (2.2.22) [right].

To make better sense of this technically ill-defined expression, we can “cut off” the integration over  $k$ . One way to do this is to insert a convergence factor  $e^{-k^2\Delta^2/2}$  into the integrand, and then take  $\Delta \rightarrow 0$ ,

$$\delta(x) = \lim_{\Delta \rightarrow 0} \frac{1}{2\pi} \int_{-\infty}^{\infty} dk e^{-k^2\Delta^2/2} e^{ikx} = \lim_{\Delta \rightarrow 0} \frac{\exp(-x^2/2\Delta^2)}{\sqrt{2\pi}\Delta}. \quad (2.2.21)$$

This interprets  $\delta(x)$  as the  $\Delta \rightarrow 0$  limit of Gaussian functions with height  $1/\sqrt{2\pi}\Delta$  and full width at half maximum (FWHM) equal to  $2\sqrt{2\ln 2}\Delta \approx 2.3548\Delta$ . Another way to make sense of eq. (2.2.20) is to limit the  $k$  integration to a large but finite range  $-1/\Delta < k < 1/\Delta$ ,

$$\delta(x) = \lim_{\Delta \rightarrow 0} \frac{1}{2\pi} \int_{-1/\Delta}^{1/\Delta} dk e^{ikx} = \lim_{\Delta \rightarrow 0} \frac{\sin(x/\Delta)}{\pi x}. \quad (2.2.22)$$

Each of the expressions in eqs. (2.2.19), (2.2.21), and (2.2.22) has the crucial property of unit integrated area for all  $\Delta$ , even before taking the limit  $\Delta \rightarrow 0$ . This justifies the normalization factor of  $1/2\pi$  in eq. (2.2.20). They are illustrated in Figure 2.2.1. Each of them, as well as the formal integral representation of eq. (2.2.20), can be useful, depending on the situation.

The representation of the delta function in eq. (2.2.20) is related to the theory of Fourier transforms. To see how this works, we can use it to write, for any sufficiently well-behaved function  $f(x)$ ,

$$f(x) = \int_{-\infty}^{\infty} dx' f(x') \delta(x - x') = \int_{-\infty}^{\infty} dx' f(x') \left( \frac{1}{2\pi} \int_{-\infty}^{\infty} dk e^{ik(x-x')} \right). \quad (2.2.23)$$

By rearranging the order of integrations, this becomes

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} dk e^{ikx} \int_{-\infty}^{\infty} dx' e^{-ikx'} f(x'). \quad (2.2.24)$$

Now, we define the Fourier transform of  $f(x)$  by the second integral,

$$F(k) \equiv \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dx' e^{-ikx'} f(x'), \quad (2.2.25)$$

for  $-\infty < k < \infty$ . Then eq. (2.2.24) becomes

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dk e^{ikx} F(k), \quad (2.2.26)$$

which is the inverse Fourier transform relation.

For a finite-dimensional<sup>§</sup> Hilbert space, there is a systematic way to find an orthobasis:

**Theorem 2.2.3.** *Given a finite-dimensional Hilbert space with an inner product and a known basis  $\{|\beta_j\rangle\}$  that is not necessarily orthonormal, one can always construct an orthobasis  $\{|\varphi_j\rangle\}$  by a systematic procedure known as the **Gram–Schmidt process**.*

Proof: We prove this constructively, by giving the steps of the Gram–Schmidt process:

Step 1. Choose any ket in the original basis; call it  $|\beta_1\rangle$ . The first member of the orthobasis will be the re-scaled version of this ket with unit norm,  $|\varphi_1\rangle = |\beta_1\rangle / \sqrt{\langle\beta_1|\beta_1\rangle}$ .

Step 2. Choose a second ket  $|\beta_2\rangle$  from the original basis set. From it, define a new vector  $|\tilde{\varphi}_2\rangle = |\beta_2\rangle - |\varphi_1\rangle \langle\varphi_1|\beta_2\rangle$ , which we can describe in words as subtracting off the projection along the vector  $|\varphi_1\rangle$ . Thus  $|\tilde{\varphi}_2\rangle$  is orthogonal to  $|\varphi_1\rangle$  by construction. Now take  $|\varphi_2\rangle = |\tilde{\varphi}_2\rangle / \sqrt{\langle\tilde{\varphi}_2|\tilde{\varphi}_2\rangle}$ , which has norm 1. This is the second member of the orthobasis we are constructing.

Step 3. Choose a third ket  $|\beta_3\rangle$  from the original basis set. From it, define a new vector  $|\tilde{\varphi}_3\rangle = |\beta_3\rangle - |\varphi_1\rangle \langle\varphi_1|\beta_3\rangle - |\varphi_2\rangle \langle\varphi_2|\beta_3\rangle$ . In words, we are subtracting off the projections along both of the vectors  $|\varphi_1\rangle$  and  $|\varphi_2\rangle$ . Since  $|\tilde{\varphi}_3\rangle$  is orthogonal to both  $|\varphi_1\rangle$  and  $|\varphi_2\rangle$  by construction, again we just need to re-scale it to have unit norm. Therefore, define  $|\varphi_3\rangle = |\tilde{\varphi}_3\rangle / \sqrt{\langle\tilde{\varphi}_3|\tilde{\varphi}_3\rangle}$  as the third member of the orthobasis.

...

Step  $n$ . Take the  $n$ th ket  $|\beta_n\rangle$  from the original basis set, and subtract off the projections along all of the previously found orthobasis kets, by defining

$$|\tilde{\varphi}_n\rangle = |\beta_n\rangle - \sum_{j=1}^{n-1} |\varphi_j\rangle \langle\varphi_j|\beta_n\rangle. \quad (2.2.27)$$

By construction, this is orthogonal to all of the previously found kets  $|\varphi_j\rangle$  with  $j = 1, \dots, n-1$ . It is not null, because of the linear independence of the original basis. So, to include it in the orthobasis, we only need to re-scale it to also have unit norm,

$$|\varphi_n\rangle = |\tilde{\varphi}_n\rangle / \sqrt{\langle\tilde{\varphi}_n|\tilde{\varphi}_n\rangle}. \quad (2.2.28)$$

---

<sup>§</sup>In the infinite-dimensional case, the Gram–Schmidt algorithm cannot end in a finite number of steps, but in cases relevant to quantum mechanics a suitable orthobasis is often easy to identify anyway. For example, the Hilbert space may naturally split into an infinite number of mutually orthogonal finite-dimensional subspaces.

Continuing in this way, when we are finished with Step  $d$ , where  $d$  is the dimension of the Hilbert space, we will have constructed the full orthobasis  $\{|\varphi_n\rangle\}$ . This concludes the proof.  $\square$

Suppose that we have chosen an orthobasis  $\{|\varphi_j\rangle\}$ . To obtain the components  $v_j$  of an arbitrary ket

$$|v\rangle = \sum_j v_j |\varphi_j\rangle, \quad (2.2.29)$$

we can take the inner product with  $|\varphi_k\rangle$ , resulting in

$$\langle\varphi_k|v\rangle = \sum_j v_j \langle\varphi_k|\varphi_j\rangle = v_k, \quad (2.2.30)$$

where the last equality relies on the orthonormality property eq. (2.2.13). Thus, the components of the ket are simply given by its inner products with the orthobasis vectors.

## 2.3 Dual vector spaces

Given a Hilbert space of kets with an inner product, it is useful to construct another complex linear vector space, called the **dual vector space**. Dual vectors are defined mathematically as linear maps from the space of kets to the complex numbers. For each ket  $|v\rangle$ , there is a corresponding dual vector denoted  $\langle v|$ , and we write the association as

$$\begin{array}{cc} \text{vector} & \text{dual vector} \\ |v\rangle & \rightarrow \langle v|. \end{array} \quad (2.3.1)$$

Specifically, the dual vector  $\langle v|$  is defined to map each ket  $|w\rangle$  to the complex number given by the inner product of the ket  $|w\rangle$  with the corresponding ket  $|v\rangle$ ,

$$\langle v|(|w\rangle) \equiv \langle v|w\rangle. \quad (2.3.2)$$

A dual vector defined in this way is also called a **bra**, a silly bit of terminology devised long ago so that the inner product is a “bra-ket”, or bracket. From the properties of the inner product, specifically by comparing eqs. (2.2.2) and (2.2.3), one finds that the bra associated with a linear combination of kets is

$$\begin{array}{cc} \text{vector} & \text{dual vector} \\ c_1 |v\rangle + c_2 |w\rangle & \rightarrow c_1^* \langle v| + c_2^* \langle w|. \end{array} \quad (2.3.3)$$

One must remember to take the complex conjugates of the coefficients.

The bra  $\langle v|$  is also known as the **Hermitian adjoint** of the corresponding ket  $|v\rangle$ , and vice versa. To understand the relation better, it is useful to consider the following linear algebra

analogy: kets are like complex  $d$ -dimensional column vectors, and bras are like complex  $d$ -dimensional row vectors, with

$$\vec{v} = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_d \end{pmatrix}, \quad \vec{w}^\dagger = (w_1^* \quad w_2^* \quad \cdots \quad w_d^*). \quad (2.3.4)$$

The fact that a dual vector maps vectors to complex numbers is just expressed as

$$\vec{w}^\dagger \vec{v} = \sum_k w_k^* v_k. \quad (2.3.5)$$

As we will discuss more fully in section 2.5, this is not just an analogy; if one has chosen an orthobasis  $\{|\varphi_k\rangle\}$ , then  $v_k = \langle\varphi_k|v\rangle$  are the components of the ket  $|v\rangle$ , while  $w_k^* = \langle w|\varphi_k\rangle$  are the components of the bra  $\langle w|$ . However, the bra-ket notation has the great virtue of being independent of any particular choice of orthobasis.

According to the dual vector definition, for each ket  $|v\rangle$ , there is always a unique corresponding bra  $\langle v|$ . If the Hilbert space has a finite dimension, then there is also a unique ket for every bra, and the correspondence is one-to-one. However, in the infinite-dimensional case it is possible to construct bras that have no associated ket within the Hilbert space, although this fact is of little practical consequence. For example, consider the generalized kets like those that satisfy the Dirac orthonormality condition eq. (2.2.15); as we have noted, these have infinite norm and are therefore not part of the physical Hilbert space. The corresponding bras are nevertheless well-defined maps from the physical Hilbert space to the complex numbers, and so are perfectly respectable members of the dual vector space.

You may find it useful to think of kets (vectors) as representing possible states of a system, while bras (dual vectors) represent possible questions that one may ask about the state. For example, we can interrogate the actual state of a system  $|\psi\rangle$  about its overlap with another possible state  $|\chi\rangle$ . Then the question that we are asking is associated with the dual vector  $\langle\chi|$ . As we will discuss further near the end of section 3.2, the answer that one receives is that the probability of finding the system in the state  $|\chi\rangle$  is nothing other than  $|\langle\chi|\psi\rangle|^2$ , assuming that both  $|\psi\rangle$  and  $|\chi\rangle$  were normalized to 1.

## 2.4 Operators

An **operator**  $A$  is a map from the space of kets to itself. This means that the result of acting with  $A$  on any ket  $|v\rangle$  must be another ket in the Hilbert space, which we can call either  $|Av\rangle$ , or equivalently,  $A|v\rangle$ . As a notational convention, we will usually use capital letters to represent operators, except when established tradition says otherwise.

In quantum mechanics, we are almost always interested in **linear operators**, which obey

$$A(c_1 |v\rangle + c_2 |w\rangle) = c_1 A|v\rangle + c_2 A|w\rangle. \quad (2.4.1)$$

Because there is a bra associated to each ket, a linear operator also maps the dual space to itself. Specifically, for each bra  $\langle w|$ , the bra  $\langle w|A$  resulting from the operation of  $A$  is defined by the relation

$$(\langle w|A) |v\rangle = \langle w| (A|v\rangle). \quad (2.4.2)$$

Adopting this definition, it follows that the operation of  $A$  on the dual space also obeys linearity,

$$(c_1 \langle v| + c_2 \langle w|)A = c_1 \langle v|A + c_2 \langle w|A. \quad (2.4.3)$$

Because of the equality of the two sides of eq. (2.4.2), we can define the **matrix element** of the operator  $A$  between  $\langle w|$  and  $|v\rangle$  as their common value  $\langle w|A|v\rangle$ , without parentheses.

Addition and subtraction of linear operators and multiplication by complex numbers are defined in the obvious ways, such that

$$(c_1 A + c_2 B) |v\rangle = A(c_1 |v\rangle) + B(c_2 |v\rangle). \quad (2.4.4)$$

The product of two operators  $A$  and  $B$  is defined by

$$(AB) |v\rangle \equiv A(B|v\rangle) = AB|v\rangle. \quad (2.4.5)$$

As before, the parentheses make no difference and can be omitted, as indicated in the last equality. Operators also obey **associativity**; for any three operators  $A$ ,  $B$ , and  $C$ ,

$$(AB)C = A(BC). \quad (2.4.6)$$

However, the order of operators matters, so that  $BA$  and  $AB$  are different, in general. One therefore defines the **commutator** of  $A$  and  $B$  as

$$[A, B] = AB - BA, \quad (2.4.7)$$

and the **anticommutator** by

$$\{A, B\} = AB + BA. \quad (2.4.8)$$

The simplest example of a linear operator is the **identity operator**  $I$ , defined by

$$I|v\rangle = |v\rangle, \quad \langle v|I = \langle v| \quad (2.4.9)$$

for every  $|v\rangle$ . The **inverse of an operator**  $A$ , if it exists, is denoted  $A^{-1}$ , and is defined by the requirements<sup>†</sup>

$$A^{-1}A = I = AA^{-1}. \quad (2.4.10)$$

However, it is important to recognize that not all operators have an inverse.

There is an enormously useful way of writing the identity operator, given an orthobasis. Equations (2.2.29) and (2.2.30) can be combined to write

$$|v\rangle = \sum_j |\varphi_j\rangle \langle \varphi_j | v \rangle. \quad (2.4.11)$$

Since this is true for all  $|v\rangle$ , one has simply

$$I = \sum_j |\varphi_j\rangle \langle \varphi_j|. \quad (2.4.12)$$

This extraordinarily important identity is known as the **completeness relation**, or the **closure relation**. We will use it very often.

If  $|v\rangle$  and  $|w\rangle$  are any two kets, then

$$A = |v\rangle \langle w| \quad (2.4.13)$$

is a linear operator, sometimes called the **outer product** of  $|v\rangle$  and  $\langle w|$ . It is defined by its action on any other ket  $|x\rangle$ ,

$$A|x\rangle = (|v\rangle \langle w|)|x\rangle = |v\rangle (\langle w|x\rangle). \quad (2.4.14)$$

As a useful special case, we can form the **projection operator**  $P_v$  onto a ket  $|v\rangle$ , defined by

$$P_v = \frac{|v\rangle \langle v|}{\langle v|v\rangle}. \quad (2.4.15)$$

If  $|v\rangle$  has norm 1, then one can simply write

$$P_v = |v\rangle \langle v|. \quad (2.4.16)$$

Projection operators have the property

$$P_v^2 = P_v. \quad (2.4.17)$$

---

<sup>†</sup>For a finite dimensional vector space, the last equality in eq. (2.4.10) is redundant, because  $BA = I$  can be shown to imply  $AB = I$  as well. However, in an infinite dimensional vector space this is not true. For example, consider the vector space with general element  $v = (v_1, v_2, v_3, \dots)$ , and define the right-shift and left-shift operators by  $Rv = (0, v_1, v_2, v_3, \dots)$  and  $Lv = (v_2, v_3, v_4, \dots)$ . Then  $LR = I$ , but  $RL \neq I$ .

Intuitively, the projection operator  $P_v$  acts on a ket by throwing away the part orthogonal to  $|v\rangle$ , and keeping the rest intact; doing this twice has the same effect as doing it once. As an aside, this is a good example of an operator that has no inverse (with the trivial exception of the case that the state space is one-dimensional). The reason is that when  $P_v$  acts on any ket  $|w\rangle$  that is orthogonal to  $|v\rangle$ , it yields 0 (the null ket), and then there is no way to resurrect  $|w\rangle$  by acting with another linear operator, the purported inverse.

The completeness relation eq. (2.4.12) can now be equivalently expressed as the statement that the identity operator is equal to

$$I = \sum_j P_{\varphi_j}, \quad (2.4.18)$$

which is the sum over the projection operators for all of the members of an orthobasis, spanning the whole Hilbert space. Similarly, one can project onto a subspace of the Hilbert space, by summing over projection operators for only a subset of the orthobasis vectors.

For any linear operator  $A$ , the **Hermitian adjoint** (or just adjoint)  $A^\dagger$  can be defined by the way that it acts on dual vectors,

$$\langle v| A^\dagger = \langle Av|, \quad (2.4.19)$$

where  $\langle Av|$  is the bra corresponding to the ket  $|Av\rangle = A|v\rangle$ . Consider a generic matrix element involving  $A^\dagger$ ,

$$\langle w| A^\dagger |v\rangle = \langle Aw|v\rangle = (\langle v|Aw\rangle)^*, \quad (2.4.20)$$

where the second equality has made use of eq. (2.2.1). This we can restate as the useful relation

$$\langle w| A^\dagger |v\rangle = (\langle v| A |w\rangle)^*. \quad (2.4.21)$$

It is not hard to use the definition to show the following facts. The adjoint of the operator  $A = |w\rangle\langle v|$  is

$$(|w\rangle\langle v|)^\dagger = |v\rangle\langle w|. \quad (2.4.22)$$

The adjoint of the operator  $cI$ , where  $c$  is a complex number, is  $(cI)^\dagger = c^*I$ . The adjoint of a product of operators is

$$(AB)^\dagger = B^\dagger A^\dagger, \quad (2.4.23)$$

where the order matters.

It is useful to generalize the concept of taking the adjoint to whole expressions and equalities involving kets, bras, and operators. Given any expression, define the adjoint of it according to the following rules:



- Substitute  $A \rightarrow A^\dagger$  for all operators.
- Substitute  $c \rightarrow c^*$  for all complex numbers.
- Substitute  $|v\rangle \leftrightarrow \langle v|$  for all kets and bras.
- Reverse the order of kets, bras, and operators within each term.

(Of course, the complex number factors of any term can be written in any desired order.) Then one can show that the adjoint of any valid equality will also be a valid equality. Equations (2.2.1), (2.4.21), (2.4.22) and (2.4.23) are examples of this. As a more random example, if one has

$$ABC |v\rangle \langle v|w\rangle + B |x\rangle = c |z\rangle, \quad (2.4.24)$$

where  $A$ ,  $B$ , and  $C$  are operators and  $c$  is a complex number, then one must also have

$$\langle w|v\rangle \langle v|C^\dagger B^\dagger A^\dagger + \langle x|B^\dagger = c^* \langle z|. \quad (2.4.25)$$

An operator  $A$  is called **Hermitian** (or **self-adjoint**) if it is the same as its adjoint,  $A^\dagger = A$ . Hermitian operators are particularly important in quantum mechanics, where they are associated with physically measurable quantities. An **anti-Hermitian** operator is one that satisfies  $A^\dagger = -A$ . Note that any operator can be written as the sum of a Hermitian part and an anti-Hermitian part,

$$A = \left( \frac{A + A^\dagger}{2} \right) + \left( \frac{A - A^\dagger}{2} \right). \quad (2.4.26)$$

The product of two Hermitian operators is Hermitian if, and only if, they commute. It is also not hard to show that for two Hermitian operators  $A$  and  $B$ , the commutator is anti-Hermitian, so that  $i[A, B]$  is Hermitian.

An operator  $U$  is **unitary** if its adjoint is equal to its inverse, so  $U^\dagger = U^{-1}$ . The product of two unitary operators is always unitary. In quantum mechanics, unitary operators are associated with a change of orthobasis, as we will discuss in the next section. They often appear in the context of defining or exploiting the symmetries of the physical system. The time evolution of a system will also be associated with a unitary operator.

It is possible to define operators that are functions of other operators. The most common example of this that we will encounter in quantum mechanics is the exponentiation of an operator. If  $A$  is an operator, then the operator  $\exp(A)$  can be defined in two different but equivalent ways, which are useful in different circumstances. Both treat  $A$  just as if it were an ordinary

number, exploiting the facts that it obviously commutes with itself, and no other operators are involved. First, one can define it as the usual limit,

$$e^A = \lim_{N \rightarrow \infty} \left( I + \frac{A}{N} \right)^N. \quad (2.4.27)$$

Alternatively, one can define it as a power series expansion:

$$e^A = \sum_{n=0}^{\infty} \frac{A^n}{n!}, \quad (2.4.28)$$

where of course  $A^0 = I$ . It follows that  $e^A$  is always invertible, and the inverse is  $e^{-A}$ . Taking  $A = iB$ , where  $B$  is Hermitian, we then find:

**Theorem 2.4.1.** *If  $B$  is a Hermitian operator, then the operator  $\exp(iB)$  is unitary.*

Some other useful results involving exponentials of operators follow.

**Theorem 2.4.2.** *For any two operators  $A$  and  $B$ ,*

$$e^A B e^{-A} = B + [A, B] + \frac{1}{2!}[A, [A, B]] + \frac{1}{3!}[A, [A, [A, B]]] + \cdots. \quad (2.4.29)$$

Proof: define the operator  $F(\lambda) = e^{\lambda A} B e^{-\lambda A}$ , where  $\lambda$  is a variable. Now

$$\frac{dF}{d\lambda} = AF - FA = [A, F(\lambda)], \quad (2.4.30)$$

where we have used the fact that  $\frac{d}{d\lambda} e^{\lambda A} = A e^{\lambda A} = e^{\lambda A} A$ . Repeating this gives

$$\frac{d^2 F}{d\lambda^2} = [A, [A, F(\lambda)]], \quad \frac{d^3 F}{d\lambda^3} = [A, [A, [A, F(\lambda)]]], \quad (2.4.31)$$

etc. Using these to compute the Taylor series expansion for  $F(\lambda)$  about the point  $\lambda = 0$ , and using  $F(0) = B$ , we have

$$F(\lambda) = B + \lambda[A, B] + \frac{\lambda^2}{2!}[A, [A, B]] + \frac{\lambda^3}{3!}[A, [A, [A, B]]] + \cdots, \quad (2.4.32)$$

and eq. (2.4.29) now follows by taking  $\lambda = 1$ .  $\square$

As a special case application of Theorem 2.4.2, we have:

**Theorem 2.4.3.** *If  $A$  and  $B$  are operators such that  $[A, B] = cB$  where  $c$  is a number, then*

$$e^A B e^{-A} = e^c B. \quad (2.4.33)$$

Another useful special case of Theorem 2.4.2 occurs if instead  $[A, B]$  commutes with  $A$ :

**Theorem 2.4.4.** *For any two operators  $A$  and  $B$  such that  $[A, [A, B]] = 0$ ,*

$$[e^A, B] = [A, B]e^A. \quad (2.4.34)$$

Finally, the following formula is often useful:

**Theorem 2.4.5. (Baker–Campbell–Hausdorff, special case)** *For any two operators  $A$  and  $B$  such that  $[A, B]$  commutes with both  $A$  and  $B$ , then*

$$e^A e^B = e^{A+B} e^{\frac{1}{2}[A, B]}. \quad (2.4.35)$$

The proof is left as Exercise 2.6. The previous two theorems hold in particular if the commutator of  $A$  and  $B$  is proportional to the identity operator.

## 2.5 Matrix representations of operators

Suppose we have selected an orthobasis  $\{|\varphi_j\rangle\}$ , and consider two kets  $|v\rangle, |w\rangle$  given as

$$|v\rangle = \sum_j v_j |\varphi_j\rangle, \quad |w\rangle = \sum_j w_j |\varphi_j\rangle, \quad (2.5.1)$$

where  $v_j$  and  $w_j$  are their components in the chosen orthobasis. The inner product can be written as

$$\langle w|v\rangle = \sum_j \sum_k w_j^* v_k \langle \varphi_j | \varphi_k \rangle = \sum_j w_j^* v_j = (w_1^* \cdots w_d^*) \begin{pmatrix} v_1 \\ \vdots \\ v_d \end{pmatrix}, \quad (2.5.2)$$

where the second equality was obtained using the orthonormality property eq. (2.2.13). In the case of a Hilbert space with finite dimension  $d$ , we can therefore consider the bra  $\langle w|$  as associated with a row vector of complex numbers, and the ket  $|v\rangle$  with a column vector,

$$\langle w| \leftrightarrow (w_1^* \cdots w_d^*), \quad |v\rangle \leftrightarrow \begin{pmatrix} v_1 \\ \vdots \\ v_d \end{pmatrix}. \quad (2.5.3)$$

The double-arrow notation is used to indicate that the bra and ket are represented by the corresponding row and column vectors. However, it is important to remember that while this representation depends on the choice of basis, the bra and ket themselves exist as abstract objects independent of the choice of basis.

Given an operator  $A$ , one can construct a matrix representation for it corresponding to the orthobasis  $\{|\varphi_j\rangle\}$ . We start with

$$A|v\rangle = A \sum_k v_k |\varphi_k\rangle = \sum_k v_k A |\varphi_k\rangle. \quad (2.5.4)$$

Now taking the inner product with a basis ket  $|\varphi_j\rangle$  gives

$$\langle\varphi_j|A|v\rangle = \sum_k \langle\varphi_j|A|\varphi_k\rangle v_k. \quad (2.5.5)$$

We therefore define the  $d \times d$  complex matrix with elements

$$A_{jk} = \langle\varphi_j|A|\varphi_k\rangle, \quad (2.5.6)$$

so that the components of

$$|v'\rangle = A|v\rangle \quad (2.5.7)$$

are

$$v'_j = \langle\varphi_j|v'\rangle = \sum_k A_{jk}v_k, \quad (2.5.8)$$

or, in matrix form,

$$\begin{pmatrix} v'_1 \\ \vdots \\ v'_d \end{pmatrix} = \begin{pmatrix} A_{11} & \cdots & A_{1d} \\ \vdots & \ddots & \vdots \\ A_{d1} & \cdots & A_{dd} \end{pmatrix} \begin{pmatrix} v_1 \\ \vdots \\ v_d \end{pmatrix}. \quad (2.5.9)$$

Thus, we complete the correspondence of eq. (2.5.3) by writing, for any operator  $A$ , the matrix representation

$$A \leftrightarrow \begin{pmatrix} A_{11} & \cdots & A_{1d} \\ \vdots & \ddots & \vdots \\ A_{d1} & \cdots & A_{dd} \end{pmatrix}, \quad (2.5.10)$$

with the orthobasis matrix elements given by eq. (2.5.6).

It is often convenient to dispense with formalities by replacing the  $\leftrightarrow$  symbol in equations like (2.5.3) and (2.5.10) with an  $=$  symbol. However, again we emphasize that the kets  $|v\rangle$ , bras  $\langle w|$ , and operators  $A$  are objects that are conceptually independent of the choice of orthobasis, while their components  $v_j$ ,  $w_j^*$ , and  $A_{jk}$  do depend on the choice of orthobasis, and will be different if a different orthobasis is used.

The matrix representation for a product of operators  $AB$  is given by

$$(AB)_{jk} = \sum_n A_{jn}B_{nk}. \quad (2.5.11)$$

The proof of this is an almost trivial consequence of the completeness relation eq. (2.4.12), because it can also be read as  $\langle\varphi_j|AB|\varphi_k\rangle = \sum_n \langle\varphi_j|A|\varphi_n\rangle \langle\varphi_n|B|\varphi_k\rangle$ . It is also easy to show that the orthobasis matrix elements of an operator  $A$  and its adjoint  $A^\dagger$  are related by

$$(A^\dagger)_{jk} = (A_{kj})^*. \quad (2.5.12)$$

Many operator relations and properties are easiest to see by using the matrix representation associated with an appropriately chosen orthobasis, using the tools of linear algebra. For instance, a Hermitian (or unitary) operator  $A$  is represented by a Hermitian (or unitary) matrix with elements  $A_{jk}$ . The matrix representation of  $A^{-1}$ , if it exists, is equal to the inverse of the matrix representation of  $A$ . It is a standard result in linear algebra that  $A^{-1}$  exists if and only if the determinant of the matrix  $A$  is non-zero,

$$\text{Det}(A) \neq 0. \quad (2.5.13)$$

In that case, the inverse matrix for  $A$  is

$$A^{-1} = \frac{1}{\text{Det}(A)} \text{cof}[A]^T \quad (2.5.14)$$

where  $\text{cof}[A]$  is the cofactor matrix of  $A$ , and  $T$  denotes the transpose. For a general  $d \times d$  matrix  $A$ , the cofactor matrix is defined by

$$(\text{cof}[A])_{jk} = (-1)^{j+k} M_{jk} \quad (2.5.15)$$

where the minor  $M_{jk}$  is equal to the determinant of the  $(d-1) \times (d-1)$  submatrix obtained from  $A$  by deleting row  $j$  and column  $k$ .

A unitary operator  $U$ , as defined in the previous section, satisfies

$$U^\dagger U = U U^\dagger = I. \quad (2.5.16)$$

The components of the matrix representation of  $U$  in an orthobasis  $\{|\varphi_k\rangle\}$ ,

$$U = \begin{pmatrix} U_{11} & U_{12} & \cdots & U_{1d} \\ U_{21} & U_{22} & \cdots & U_{2d} \\ \vdots & \vdots & \ddots & \vdots \\ U_{d1} & U_{d2} & \cdots & U_{dd} \end{pmatrix}, \quad (2.5.17)$$

obey the rule

$$U_{jk} = j\text{th component of } |\varphi'_k\rangle, \quad (2.5.18)$$

where

$$|\varphi'_k\rangle = U |\varphi_k\rangle. \quad (2.5.19)$$

Now, using eq. (2.5.16), it is not too difficult to show that the set  $\{|\varphi'_k\rangle\}$  defined in this way is *another* orthobasis. From eqs. (2.5.17) and (2.5.18), we see that the components of the new orthobasis element  $|\varphi'_k\rangle$  are equal to the  $k$ th column of the matrix representation of  $U$  in the

original orthobasis. In this sense, a unitary operator  $U$  performs a change of orthobasis, which can be thought of as a complex rotation in the Hilbert space. The operator  $U^\dagger$  performs the inverse change of basis,

$$|\varphi_k\rangle = U^\dagger |\varphi'_k\rangle, \quad (2.5.20)$$

and it follows that the bras for the two orthobases are related by

$$\langle\varphi'_k| = \langle\varphi_k| U^\dagger, \quad \langle\varphi_k| = \langle\varphi'_k| U. \quad (2.5.21)$$

Conversely, given any two orthonormal bases  $\{|\varphi_k\rangle\}$  and  $\{|\varphi'_k\rangle\}$ , the inner products are the elements of a unitary matrix,

$$\langle\varphi_j|\varphi'_k\rangle = \langle\varphi_j|U|\varphi_k\rangle = U_{jk}. \quad (2.5.22)$$

To verify directly that  $U$  as defined by eq. (2.5.22) is indeed unitary, one can use the completeness relation eq. (2.4.12) to show

$$\sum_n U_{nj}^* U_{nk} = \delta_{jk}, \quad (2.5.23)$$

or, rewriting,

$$\sum_n (U^\dagger)_{jn} U_{nk} = \delta_{jk}. \quad (2.5.24)$$

This expresses the unitarity of the matrix representation of the operator  $U$ , and is equivalent to the operator equation (2.5.16). Also,

$$|\text{Det}(U)| = 1, \quad (2.5.25)$$

follows from the general linear algebra facts that  $\text{Det}(AB) = (\text{Det}A)(\text{Det}B)$  and  $\text{Det}(A^\dagger) = (\text{Det}A)^*$ , for matrices  $A, B$ .

An important feature of such a unitary change of basis is that it preserves inner products. To see this, consider the transformation<sup>†</sup>

$$|v\rangle \rightarrow |v'\rangle = U |v\rangle, \quad (2.5.26)$$

$$|w\rangle \rightarrow |w'\rangle = U |w\rangle, \quad (2.5.27)$$

so that we have performed the same complex rotation on both  $|v\rangle$  and  $|w\rangle$ . Then

$$\langle w|v\rangle \rightarrow \langle w'|v'\rangle = \langle w|U^\dagger U|v\rangle = \langle w|v\rangle, \quad (2.5.28)$$

---

<sup>†</sup>Here, the arrow “ $\rightarrow$ ” can be read as “transforms to”.

so that inner products are unaffected by a change of orthobasis.

We can also see how the unitary change of basis is realized on the matrix elements of any operator  $A$ . For any two kets  $|v\rangle$  and  $|w\rangle$ ,

$$\langle w'|A|v'\rangle = \langle w|U^\dagger A U|v\rangle. \quad (2.5.29)$$

If we now define a transformation of the operator  $A$  according to

$$A \rightarrow A' = U A U^\dagger, \quad (2.5.30)$$

to go along with eqs. (2.5.26) and (2.5.27), then

$$\langle w'|A'|v'\rangle = (\langle w|U^\dagger)(U A U^\dagger)(U|v\rangle) = \langle w|(U^\dagger U)A(U^\dagger U)|v\rangle = \langle w|A|v\rangle. \quad (2.5.31)$$

In words, the matrix element for the transformed states of the transformed operator  $A'$  is the same as the original matrix element of  $A$ .

Functions of operators are often easiest to deal with using their matrix representations. If the operator  $A$  has a matrix representation that is diagonal in some appropriate orthobasis,

$$A \leftrightarrow \text{diag}(a_1, a_2, \dots, a_d), \quad (2.5.32)$$

then immediately from the series definition of eq. (2.4.28), with  $A$  replaced by  $iA$ , one finds

$$\exp(iA) \leftrightarrow \text{diag}(e^{ia_1}, e^{ia_2}, \dots, e^{ia_d}). \quad (2.5.33)$$

Also, if the powers of  $A$  obey some recurrence relation, then the series for  $\exp(iA)$  can often be resummed in the matrix representation, even if it is not diagonal. As an example, suppose  $A$  has matrix representation

$$A \leftrightarrow a \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (2.5.34)$$

Then we recognize that  $A^2 = a^2 I$ , and so the unitary operator  $e^{iA}$  has matrix representation, obtained by separating the even powers of  $A$  from the odd powers of  $A$ ,

$$\sum_{n=0}^{\infty} \frac{(ia)^{2n}}{(2n)!} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \sum_{n=0}^{\infty} \frac{(ia)^{2n+1}}{(2n+1)!} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (2.5.35)$$

or, summing each series,

$$\exp(iA) \leftrightarrow \begin{pmatrix} \cos(a) & i \sin(a) \\ i \sin(a) & \cos(a) \end{pmatrix}. \quad (2.5.36)$$

## 2.6 Eigenvalues and eigenvectors

In quantum mechanics, we will often need to solve eigenvalue problems. If, for some operator  $A$ , we can find a complex number  $\alpha$  and a non-null ket  $|v\rangle$  such that

$$A|v\rangle = \alpha|v\rangle, \quad (2.6.1)$$

then  $\alpha$  is called the **eigenvalue** and  $|v\rangle$  is the corresponding **eigenvector** or **eigenket**, associated to the **eigenstate** of  $A$ . When eq. (2.6.1) holds, then it follows immediately that

$$\langle v|A^\dagger = \alpha^*\langle v|, \quad (2.6.2)$$

or, in words,  $\langle v|$  is an eigenbra of  $A^\dagger$ , with eigenvalue  $\alpha^*$ .

The **eigenvalue problem** is to find all solutions for both  $\alpha$  and  $|v\rangle$  as a pair, given  $A$ . Rewriting eq. (2.6.1) as

$$(A - \alpha I)|v\rangle = 0, \quad (2.6.3)$$

we see that for a solution with a particular  $\alpha$  to exist, the operator  $(A - \alpha I)^{-1}$  must not exist; otherwise, we could act with it on both sides to discover that  $|v\rangle$  could only be the null ket, in contradiction of the assumption.

If the Hilbert space is finite dimensional, we can go to the matrix representation to find that a solution for  $\alpha$  must satisfy

$$\text{Det}(A - \alpha I) = 0. \quad (2.6.4)$$

This is called the characteristic equation, and the left side is a polynomial of degree  $d$ , the same as the dimension of the Hilbert space. The fundamental theorem of algebra states that this always has exactly  $d$  complex solutions,  $\alpha_1, \dots, \alpha_d$ , allowing for the possibility that some of them may be repeated. The integer number of times a particular eigenvalue  $\alpha_n$  is repeated in the list is called its **degeneracy**, and we will denote it  $g_{\alpha_n}$ .

After the eigenvalues  $\alpha_n$  have been found, for each of them we can then solve

$$(A - \alpha_n I)|v_n\rangle = 0 \quad (2.6.5)$$

for the corresponding eigenket  $|v_n\rangle$ . If a particular eigenvalue  $\alpha_n$  has degeneracy  $g_{\alpha_n}$ , then there will be a set of  $g_{\alpha_n}$  linearly independent eigenkets  $|v_n\rangle$  with that eigenvalue. Sometimes the solution of eq. (2.6.5) can be done by inspection, sometimes it will require linear algebra, and in still other cases it will involve solving some differential equation(s). It is often not feasible to solve it exactly, which makes life interesting and necessitates approximation methods. In any



case, note that if  $|v_n\rangle$  satisfies the equation, then so does  $c|v_n\rangle$  for every complex constant  $c$ . That constant can always be chosen, for each eigenket, to normalize it as desired. Even after doing so, the complex phase of each eigenket remains an arbitrary choice.

The most trivial example is the case that  $A = I$ , the identity operator. Since  $I|v\rangle = |v\rangle$ , every non-null ket  $|v\rangle$  is an eigenket, and the corresponding eigenvalue is 1.

Another simple case is the projection operator for a ket  $|w\rangle$ , as introduced in eq. (2.4.15):

$$P_w = \frac{|w\rangle\langle w|}{\langle w|w\rangle}. \quad (2.6.6)$$

Then the equation  $P_w|v\rangle = \alpha|v\rangle$  becomes

$$\frac{\langle w|v\rangle}{\langle w|w\rangle} |w\rangle = \alpha |v\rangle. \quad (2.6.7)$$

There are two ways that this eigenvalue equation can be satisfied. If  $\langle w|v\rangle \neq 0$ , then  $|v\rangle$  must be proportional to  $|w\rangle$ . Indeed, any  $|v\rangle = c|w\rangle$ , where  $c$  is any non-zero complex number, is a solution, and the corresponding eigenvalue is  $\alpha = 1$ . The second way to satisfy the equation is if  $\langle w|v\rangle = 0$ , which then implies  $\alpha = 0$ . So, any ket  $|v\rangle$  orthogonal to  $|w\rangle$  is an eigenket, with  $\alpha = 0$  as the eigenvalue. These are the only solutions to the eigenvalue problem for  $P_w$ .

As another example, take a Hilbert space that is spanned by an orthobasis of three kets, which we will call  $|a\rangle$ ,  $|b\rangle$ , and  $|c\rangle$ . Consider an operator  $R_\theta$ , which depends on a continuous parameter  $\theta$ , and is defined by

$$R_\theta|a\rangle = \cos\theta|a\rangle + \sin\theta|b\rangle, \quad (2.6.8)$$

$$R_\theta|b\rangle = -\sin\theta|a\rangle + \cos\theta|b\rangle, \quad (2.6.9)$$

$$R_\theta|c\rangle = |c\rangle. \quad (2.6.10)$$

Because we have specified how  $R_\theta$  acts on each member of the orthobasis, it is completely defined. We can now adopt a representation associated with this orthobasis, so that

$$|a\rangle \leftrightarrow \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad |b\rangle \leftrightarrow \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad |c\rangle \leftrightarrow \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad (2.6.11)$$

and

$$\langle a| \leftrightarrow (1 \ 0 \ 0), \quad \langle b| \leftrightarrow (0 \ 1 \ 0), \quad \langle c| \leftrightarrow (0 \ 0 \ 1). \quad (2.6.12)$$

The matrix representation of  $R_\theta$  in this orthobasis is,<sup>†</sup> by applying eqs. (2.5.6) and (2.5.10),

$$R_\theta \leftrightarrow \begin{pmatrix} \cos\theta & -\sin\theta & 0 \\ \sin\theta & \cos\theta & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (2.6.13)$$

---

<sup>†</sup>Beware of a common mistake: it is tempting to scan the form of eqs. (2.6.8)–(2.6.10) and incorrectly write down the *transpose* of the matrix representation eq. (2.6.13). For similar examples, see eqs. (2.6.33) and (2.6.56).

This is a unitary matrix, and  $R_\theta^{-1} = R_\theta^\dagger = R_{-\theta} = R_\theta^T$ , where the last equality follows because it is a real matrix. The change of basis produced by  $R_\theta$  is a rotation in the  $|a\rangle, |b\rangle$  subspace.

To find the eigenvalues and eigenkets, first note that the characteristic equation is

$$\text{Det}(R_\theta - \alpha I) = (1 - 2\alpha \cos \theta + \alpha^2)(1 - \alpha) = 0, \quad (2.6.14)$$

which has eigenvalue solutions

$$\alpha = e^{i\theta}, e^{-i\theta}, 1. \quad (2.6.15)$$

These are non-degenerate, and it is not hard to solve  $(R_\theta - I\alpha)|V\rangle = 0$  in matrix form for the corresponding eigenvectors. The results are

$$\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -i \\ 0 \end{pmatrix} \leftrightarrow \frac{1}{\sqrt{2}} (|a\rangle - i|b\rangle) \quad \text{for } \alpha = e^{i\theta}, \quad (2.6.16)$$

$$\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ i \\ 0 \end{pmatrix} \leftrightarrow \frac{1}{\sqrt{2}} (|a\rangle + i|b\rangle) \quad \text{for } \alpha = e^{-i\theta}, \quad (2.6.17)$$

$$\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \leftrightarrow |c\rangle \quad \text{for } \alpha = 1. \quad (2.6.18)$$

We have chosen the multiplicative constants in front of these so that each eigenket has norm 1. There remains a freedom to choose each of the phases of the eigenkets; this can only be resolved by arbitrary choice.

We will now discuss several theorems regarding eigenvalues and eigenkets that are important for quantum mechanics.

**Theorem 2.6.1. (Superposition principle)** *If a linear operator  $A$  has some eigenkets  $|v_n\rangle$  with a common eigenvalue  $\alpha$ , then any complex linear combination of them,  $\sum_n c_n |v_n\rangle$ , is also an eigenket of  $A$ , with the same eigenvalue  $\alpha$ , unless it is the null ket.*

The proof is an immediate consequence of the definitions, including the assumed linearity of  $A$ .

**Theorem 2.6.2. (Common eigenvectors for commuting operators)** *If operators  $A$  and  $B$  commute, and  $|v\rangle$  is an eigenket of  $A$  with eigenvalue  $\alpha$ , then (i)  $B|v\rangle$  is also an eigenket of  $A$  with the same eigenvalue  $\alpha$ , and (ii) if  $\alpha$  is non-degenerate, then  $|v\rangle$  is also an eigenket of  $B$ .*

Proof: To establish (i) takes only one line,

$$A(B|v\rangle) = AB|v\rangle = BA|v\rangle = B\alpha|v\rangle = \alpha(B|v\rangle). \quad (2.6.19)$$

The second equality is where we used that  $A$  and  $B$  commute. To prove (ii), note that the assumption that  $\alpha$  is non-degenerate simply means that the corresponding eigenket is unique up to a multiplicative constant. So, from (i),  $B|v\rangle = \beta|v\rangle$  for some constant  $\beta$ , which is therefore the eigenvalue of  $B$  for  $|v\rangle$ .  $\square$

**Theorem 2.6.3.** *All eigenvalues  $\alpha$  of a Hermitian operator  $A$  are real.*

Proof: Let  $A|v\rangle = \alpha|v\rangle$ . Taking the inner product with  $|v\rangle$ , we have  $\langle v|A|v\rangle = \alpha\langle v|v\rangle$ . However, we also have  $\langle v|A|v\rangle = \langle v|A^\dagger|v\rangle = (\langle v|A|v\rangle)^* = \alpha^*\langle v|v\rangle$ , where the first equality made use of the assumption that  $A$  is Hermitian, and the second employs eq. (2.4.21). Combining these expressions gives

$$(\alpha - \alpha^*)\langle v|v\rangle = 0, \quad (2.6.20)$$

which establishes that  $\alpha = \alpha^*$ , because  $|v\rangle$ , being an eigenket, is not null.  $\square$

In general, a **selection rule** is a statement that some matrix element vanishes under certain specified conditions. The following theorem is an example.

**Theorem 2.6.4. (Matrix element selection rule)** *If  $A$  is a Hermitian operator and  $B$  is an operator that commutes with  $A$ , and  $A$  has eigenkets  $|v_1\rangle$  and  $|v_2\rangle$  with respective eigenvalues  $\alpha_1 \neq \alpha_2$ , then  $\langle v_1|B|v_2\rangle = 0$ .*

Proof: Since  $A$  and  $B$  commute,  $AB$  and  $BA$  are the same operator, so

$$\langle v_1|AB|v_2\rangle - \langle v_1|BA|v_2\rangle = 0. \quad (2.6.21)$$

Now we can use  $A|v_2\rangle = \alpha_2|v_2\rangle$  on the second term, and use eq. (2.6.2), which says  $\langle v_1|A = \alpha_1^*\langle v_1|$ , on the first term, to get

$$(\alpha_1 - \alpha_2)\langle v_1|B|v_2\rangle = 0, \quad (2.6.22)$$

where we have also used Theorem 2.6.3 which implies  $\alpha_1^* = \alpha_1$ , since  $A$  is Hermitian. Now, since  $\alpha_1 \neq \alpha_2$  by assumption, the matrix element  $\langle v_1|B|v_2\rangle$  must vanish.  $\square$

**Theorem 2.6.5. (Inner product selection rule)** *If  $A$  is a Hermitian operator with eigenkets  $|v_1\rangle$  and  $|v_2\rangle$  with different eigenvalues  $\alpha_1 \neq \alpha_2$ , then the eigenkets are orthogonal,  $\langle v_1|v_2\rangle = 0$ .*

The proof follows immediately as a special case of Theorem 2.6.4, by taking  $B = I$ .

The next result establishes a useful connection between a Hermitian operator and a corresponding orthobasis for the Hilbert space:

**Theorem 2.6.6. (Orthobasis of eigenkets of a Hermitian operator)** *If  $A$  is a Hermitian operator on a Hilbert space with finite dimension  $d$ , then one can find a set of its eigenkets that form an orthobasis. The resulting orthobasis elements corresponding to non-degenerate eigenvalues of  $A$  are unique (up to multiplication by complex phases).*

Proof: The characteristic equation for the eigenvalue problem for  $A$  (using its matrix representation in any orthobasis) has  $d$  solutions, according to the fundamental theorem of algebra. The eigenkets corresponding to non-degenerate eigenvalues are orthogonal (according to Theorem 2.6.5) and non-null (from the definition of an eigenket), and so can be normalized to be orthonormal. For each eigenvalue  $\alpha$  with degeneracy  $g_\alpha > 1$ , one can use the Gram–Schmidt process to construct (non-uniquely) a set of orthonormal kets which have the same eigenvalue  $\alpha$  due to the linearity of the eigenvalue problem, and which are all orthogonal to the kets corresponding to the other eigenvalues (again using Theorem 2.6.5). The union of all of the kets found in this way are orthonormal with each other, and there are  $d$  of them, so they form an orthobasis for the whole Hilbert space.  $\square$

Let us make some important comments on Theorem 2.6.6. First, it is often convenient to adopt a notation such that the orthobasis eigenkets of  $A$  are named by using the corresponding eigenvalue as a label. However, if the eigenvalue  $\alpha$  is degenerate, then we need to introduce another label  $u_\alpha$ , which we refer to as a **degeneracy label**, to distinguish the orthobasis kets that have the same  $\alpha$ . Thus, the eigenkets can be called  $|\alpha, u_\alpha\rangle$ , and the eigenvalue equation is written as

$$(A - \alpha I) |\alpha, u_\alpha\rangle = 0, \quad (2.6.23)$$

while the orthonormality condition reads

$$\langle \alpha', u'_{\alpha'} | \alpha, u_\alpha \rangle = \delta_{\alpha\alpha'} \delta_{u_\alpha u'_{\alpha'}}. \quad (2.6.24)$$

In the orthobasis of eigenkets of  $A$ , the matrix representation of  $A$  will be diagonal, as follows immediately from eq. (2.6.23), so that

$$A \leftrightarrow A_{\text{diag}} = \begin{pmatrix} \alpha_1 & 0 & \cdots & 0 \\ 0 & \alpha_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \alpha_d \end{pmatrix} \quad (2.6.25)$$

in that basis.

Another useful comment on Theorem 2.6.6 is that if we form the matrix whose columns are the components of the orthobasis eigenkets of  $A$ , then the result is a unitary matrix  $U$ . This follows from the general discussion surrounding eqs. (2.5.17)–(2.5.24), if in that discussion we

let  $\{|\varphi_k\rangle\}$  be the “original” basis we are working in, and let the set  $\{|\varphi'_k\rangle\} = \{|\alpha, u_\alpha\rangle\}$  be the orthobasis of eigenkets of  $A$ . Thus, we define the components of the matrix  $U$  according to the rule that, in the original basis,

$$\begin{aligned} U_{jk} &= \langle \varphi_j | U | \varphi_k \rangle = \langle \varphi_j | \varphi'_k \rangle \\ &= \text{jth component of (kth orthonormal eigenvector of } A). \end{aligned} \quad (2.6.26)$$

The eigenvalue equation (2.6.23) can then be written as, by taking the inner product with  $\langle \varphi_n |$ ,

$$\sum_j A_{nj} U_{jk} = \alpha_k U_{nk} = \sum_l U_{nl} (\delta_{lk} \alpha_l), \quad (2.6.27)$$

or, in index-free matrix notation,

$$AU = UA_{\text{diag}}. \quad (2.6.28)$$

Now, multiplying from the left by  $U^\dagger$ , we obtain

**Theorem 2.6.7. (Diagonalization of Hermitian operators)** *The matrix representation of a Hermitian operator  $A$  in an arbitrary orthobasis  $\{|\varphi_k\rangle\}$  is diagonalized by the unitary matrix  $U$  specified in eq. (2.6.26), according to*

$$U^\dagger AU = A_{\text{diag}}, \quad (2.6.29)$$

where  $A_{\text{diag}}$  is the matrix representation of the operator  $A$  in the new orthobasis  $\{|\varphi'_k\rangle\}$  consisting of its eigenvectors.

Let us do an extended example to illustrate some of the preceding results. Consider a Hilbert space of dimension 3 with an operator  $A$  defined by its action on an orthobasis  $\{|\varphi_1\rangle, |\varphi_2\rangle, |\varphi_3\rangle\}$ ,

$$A|\varphi_1\rangle = 3|\varphi_1\rangle, \quad (2.6.30)$$

$$A|\varphi_2\rangle = 4|\varphi_2\rangle - i|\varphi_3\rangle, \quad (2.6.31)$$

$$A|\varphi_3\rangle = i|\varphi_2\rangle + 4|\varphi_3\rangle. \quad (2.6.32)$$

The matrix representation of  $A$  in this original orthobasis is, using eqs. (2.5.6) and (2.5.10),

$$A \leftrightarrow \begin{pmatrix} 3 & 0 & 0 \\ 0 & 4 & i \\ 0 & -i & 4 \end{pmatrix}. \quad (2.6.33)$$

Since the matrix representation is Hermitian,  $A$  is a Hermitian operator. The characteristic equation for its eigenvalues is  $\text{Det}(A - \alpha I) = (3 - \alpha)(15 - 8\alpha + \alpha^2) = 0$ , which yields

$$\alpha = 3, 3, 5. \quad (2.6.34)$$

These eigenvalues are real, as guaranteed by Theorem 2.6.3. For the non-degenerate eigenvalue  $\alpha = 5$ , the solution of the matrix equation for the corresponding eigenvector gives, after normalization and choice of an arbitrary phase, and conversion to ket language,

$$|5\rangle = \frac{1}{\sqrt{2}} (|\varphi_2\rangle - i|\varphi_3\rangle). \quad (2.6.35)$$

Note that we have named the eigenket after its eigenvalue. For the eigenvalue  $\alpha = 3$ , the degeneracy is 2, so there are two linearly independent eigenkets, which can be chosen to be

$$|3, 1\rangle = \frac{1}{\sqrt{2}} (|\varphi_2\rangle + i|\varphi_3\rangle), \quad (2.6.36)$$

$$|3, 2\rangle = |\varphi_1\rangle. \quad (2.6.37)$$

The second entry (1 or 2) on each  $\alpha = 3$  ket is a degeneracy label. These kets are automatically orthogonal to  $|5\rangle$ , as promised by Theorem 2.6.5, but we had to make a choice of linear combination to make them orthonormal, since arbitrary linear combinations of  $|3, 1\rangle$  and  $|3, 2\rangle$  would not be orthogonal to each other and would not have norm 1. In this example, the linear combination that makes an orthobasis is not hard to find, but given any other (non-orthonormal) basis for the degenerate eigenvalue subspace, the Gram–Schmidt procedure of Theorem 2.2.3 can be used to construct an orthobasis.

Our orthobasis of eigenkets of  $A$  is related to the original orthobasis by a unitary transformation. Let us define an operator  $U$  by

$$U|\varphi_1\rangle = |5\rangle, \quad (2.6.38)$$

$$U|\varphi_2\rangle = |3, 1\rangle, \quad (2.6.39)$$

$$U|\varphi_3\rangle = |3, 2\rangle. \quad (2.6.40)$$

Now using eqs. (2.6.35)–(2.6.37), and applying the operator-matrix correspondence rules of eqs. (2.5.6) and (2.5.10) again, one finds the matrix representation of  $U$  in the  $\{|\varphi_1\rangle, |\varphi_2\rangle, |\varphi_3\rangle\}$  basis,

$$U \leftrightarrow \begin{pmatrix} 0 & 0 & 1 \\ 1/\sqrt{2} & 1/\sqrt{2} & 0 \\ -i/\sqrt{2} & i/\sqrt{2} & 0 \end{pmatrix}, \quad (2.6.41)$$

which is indeed a unitary matrix.

Acting on eqs. (2.6.38)–(2.6.40) with  $U^\dagger$ , and using  $U^\dagger U = I$ , we get

$$U^\dagger |5\rangle = |\varphi_1\rangle = |3, 2\rangle, \quad (2.6.42)$$

$$U^\dagger |3, 1\rangle = |\varphi_2\rangle = (|5\rangle + |3, 1\rangle)/\sqrt{2}, \quad (2.6.43)$$

$$U^\dagger |3, 2\rangle = |\varphi_3\rangle = i(|5\rangle - |3, 1\rangle)/\sqrt{2}. \quad (2.6.44)$$

The second equality in each of these equations follows by inverting eqs. (2.6.35)–(2.6.37) to solve for  $|\varphi_1\rangle$ ,  $|\varphi_2\rangle$ , and  $|\varphi_3\rangle$ . Equations (2.6.42)–(2.6.44) provide the matrix representation of  $U^\dagger$  in the orthobasis  $\{|5\rangle, |3, 1\rangle, |3, 2\rangle\}$ ,

$$U^\dagger \leftrightarrow \begin{pmatrix} 0 & 1/\sqrt{2} & i/\sqrt{2} \\ 0 & 1/\sqrt{2} & -i/\sqrt{2} \\ 1 & 0 & 0 \end{pmatrix}. \quad (2.6.45)$$

As a check, this matrix is indeed equal to the transpose conjugate of eq. (2.6.41). As promised by Theorem 2.6.7 on diagonalization of Hermitian operators,

$$U^\dagger A U = A_{\text{diag}} = \begin{pmatrix} 5 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{pmatrix}, \quad (2.6.46)$$

where  $A$ ,  $U$ , and  $U^\dagger$  on the left side are taken to be the matrices given by eqs. (2.6.33), (2.6.41), and (2.6.45), respectively. Note that  $A_{\text{diag}}$  on the right side is the matrix representation of the operator  $A$  in the orthobasis  $\{|5\rangle, |3, 1\rangle, |3, 2\rangle\}$ .

The choice of orthobasis made in eqs. (2.6.35)–(2.6.37) is not the unique one composed of eigenkets of  $A$ , due to the superposition principle of Theorem 2.6.1. The most general orthonormal pair of basis kets with  $A$  eigenvalue  $\alpha = 3$  is obtained using the freedom to do unitary transformations within the degenerate eigenvalue subspace, and so is parameterized by two complex numbers  $c, s$ , subject to the constraint  $|c|^2 + |s|^2 = 1$ ,

$$|3, 1'\rangle = c|3, 1\rangle + s|3, 2\rangle, \quad (2.6.47)$$

$$|3, 2'\rangle = -s^*|3, 1\rangle + c^*|3, 2\rangle. \quad (2.6.48)$$

These kets have new degeneracy labels  $(1', 2')$  to distinguish this orthobasis from the unprimed one with  $c = 1, s = 0$ . The change in orthobasis is brought about by another unitary operator  $V$ , defined by

$$V|5\rangle = |5\rangle, \quad (2.6.49)$$

$$V|3, 1\rangle = |3, 1'\rangle, \quad (2.6.50)$$

$$V|3, 2\rangle = |3, 2'\rangle, \quad (2.6.51)$$

or, in a matrix representation in the basis  $\{|5\rangle, |3, 1\rangle, |3, 2\rangle\}$ ,

$$V \leftrightarrow \begin{pmatrix} 1 & 0 & 0 \\ 0 & c & -s^* \\ 0 & s & c^* \end{pmatrix}, \quad (|5\rangle, |3, 1\rangle, |3, 2\rangle \text{ basis}). \quad (2.6.52)$$

By combining the information in eqs. (2.6.35)–(2.6.37) and (2.6.47)–(2.6.51), one also finds

$$V |\varphi_1\rangle = c^* |\varphi_1\rangle - \frac{s^*}{\sqrt{2}} |\varphi_2\rangle - i \frac{s^*}{\sqrt{2}} |\varphi_3\rangle, \quad (2.6.53)$$

$$V |\varphi_2\rangle = \frac{s}{\sqrt{2}} |\varphi_1\rangle + \frac{1}{2}(1+c) |\varphi_2\rangle + \frac{i}{2}(c-1) |\varphi_3\rangle, \quad (2.6.54)$$

$$V |\varphi_3\rangle = -i \frac{s}{\sqrt{2}} |\varphi_1\rangle + \frac{i}{2}(1-c) |\varphi_2\rangle + \frac{1}{2}(1+c) |\varphi_3\rangle, \quad (2.6.55)$$

so that the matrix representation of  $V$  in the original orthobasis is

$$V \leftrightarrow \begin{pmatrix} c^* & s/\sqrt{2} & -is/\sqrt{2} \\ -s^*/\sqrt{2} & (1+c)/2 & i(1-c)/2 \\ -is^*/\sqrt{2} & i(c-1)/2 & (1+c)/2 \end{pmatrix}, \quad (|\varphi_1\rangle, |\varphi_2\rangle, |\varphi_3\rangle \text{ basis}). \quad (2.6.56)$$

The comparison of eqs. (2.6.52) and (2.6.56) illustrates the point that even though  $V$  is unambiguous as a Hilbert space operator, its matrix representation looks very different depending on the orthobasis used.

The **trace of an operator** in a finite-dimensional Hilbert space can be defined to be equal to the trace of its matrix representation. For an operator  $A$  and any orthobasis  $\{|\varphi_k\rangle\}$ ,

$$\text{Tr}(A) = \sum_k \langle \varphi_k | A | \varphi_k \rangle = \sum_k A_{kk}. \quad (2.6.57)$$

It is left as Exercise 2.4 to show that the trace of an operator is independent of the choice of orthobasis, and in particular is equal to the sum of its eigenvalues, and that

$$\text{Tr}(AB) = \text{Tr}(BA), \quad (2.6.58)$$

$$\text{Tr}(|v\rangle \langle w|) = \langle w | v \rangle, \quad (2.6.59)$$

which can be proved very quickly using completeness of the orthobasis.

We conclude this section by stating two useful theorems involving unitary operators, omitting the proofs (which are not too difficult).

**Theorem 2.6.8.** *The eigenvalues of a unitary operator  $U$  have magnitude 1.*

Note that eq. (2.6.15) illustrates this.

**Theorem 2.6.9.** *If an operator  $A$  has eigenvectors  $|\alpha_n\rangle$  with eigenvalues  $\alpha_n$ , and  $U$  is a unitary operator, then  $UAU^\dagger$  has eigenvectors  $U|\alpha_n\rangle$  with the same eigenvalues  $\alpha_n$ .*

This implies the important result that the eigenvalues of an operator do not depend on the choice of basis we use to calculate them.



## 2.7 Observables

An operator  $A$  is an **observable** if it is Hermitian and its eigenkets form an orthobasis  $|\alpha, u_\alpha\rangle$ , which satisfies the orthonormality relation eq. (2.6.24) and the completeness relation

$$\sum_{\alpha} \sum_{u_\alpha=1}^{g_\alpha} |\alpha, u_\alpha\rangle \langle \alpha, u_\alpha| = I. \quad (2.7.1)$$

Here,  $\alpha$  are the eigenvalues with degeneracies  $g_\alpha$ , and  $u_\alpha$  are the degeneracy labels. For a Hilbert space of finite dimension, the completeness part of this definition is redundant, following automatically from Theorem 2.6.6. However it is not automatic for a Hermitian operator in an infinite-dimensional Hilbert space; that is one reason why it must be included in the definition here. The other reason is that we will later find it convenient to slightly generalize our definition of an observable to include ordered sets of Hermitian operators, so that its eigenvalues can be not just numbers, but ordered sets of numbers. (An example is the vector position operator in three-dimensional space, which is really an ordered triple of three Hermitian operators  $X$ ,  $Y$ , and  $Z$ .) In such cases, eq. (2.7.1) is again certainly not automatic. From a physical point of view, the crucial requirement we are ensuring with eq. (2.7.1) is that if  $A$  is really supposed to be an observable, then every state in the Hilbert space can always be expressed as a linear combination of its eigenstates.

The way that eq. (2.7.1) is written assumes that the eigenvalues of  $A$  are countable and discrete. If instead they are uncountable and continuous, then the orthonormality and completeness conditions are of the Dirac type [compare to eq. (2.2.15)],

$$\langle \alpha', u'_{\alpha'} | \alpha, u_\alpha \rangle = \delta(\alpha - \alpha') \delta_{u_\alpha u'_{\alpha'}}, \quad (2.7.2)$$

$$\int d\alpha \sum_{u_\alpha=1}^{g_\alpha} |\alpha, u_\alpha\rangle \langle \alpha, u_\alpha| = I, \quad (2.7.3)$$

where the integral is over the range or ranges of  $\alpha$  that occur as eigenvalues. Similarly, it is also possible for the degeneracy labels  $u_\alpha$  to be continuous, in which case the Kronecker  $\delta_{u_\alpha u'_{\alpha'}}$  is replaced by a Dirac delta function  $\delta(u_\alpha - u'_{\alpha'})$  and  $\sum_{u_\alpha=1}^{g_\alpha}$  is replaced by  $\int du_\alpha$ .

It is even possible for the eigenvalues  $\alpha$  or the degeneracy labels  $u_\alpha$  to have a spectrum that includes both discrete (countable) and continuous (uncountable) components. In that case, the orthonormality relations will include both Kronecker and Dirac deltas, and the completeness relation will include both a sum and an integral. We will see an example of this when we study the bound and unbound states of the hydrogen atom, in eqs. (11.2.11)-(11.2.13) and (11.2.22).

A particularly useful consequence of completeness follows from acting with  $A$  on both sides of eq. (2.7.1). Then, evaluating  $A = \alpha$  when acting on its eigenkets, we obtain the **spectral**

**decomposition** of an observable operator  $A$ ,

$$A = \sum_{\alpha} \sum_{u_{\alpha}=1}^{g_{\alpha}} \alpha |\alpha, u_{\alpha}\rangle \langle \alpha, u_{\alpha}|, \quad (2.7.4)$$

with an obvious counterpart for continuous eigenvalues by replacing summation with integration. Solving problems in quantum mechanics is often the art of turning expressions involving operators into numbers. The spectral decomposition idea allows us to do this in a systematic way. When we see an observable operator  $A$  appearing in a matrix element, a standard trick is to insert a completeness sum directly before or after it, so as to replace  $A$  by a sum (or integral) over its eigenvalues, as in eq. (2.7.4). Of course, a prerequisite for using this trick is to solve the eigenvalue problem for  $A$ .

At the end of section 2.4, we mentioned that it is possible to define functions of operators, and gave the most common example, that of an exponential of an operator. If we have solved the eigenvalue problem for an observable  $A$ , completeness now gives us yet another way to define a completely general function of it,  $F(A)$ . For the case of discrete eigenvalues  $\alpha$ , the spectral decomposition of  $F(A)$  is

$$F(A) = \sum_{\alpha} \sum_{u_{\alpha}=1}^{g_{\alpha}} F(\alpha) |\alpha, u_{\alpha}\rangle \langle \alpha, u_{\alpha}|, \quad (2.7.5)$$

again with an obvious counterpart for the continuous case involving integration rather than summation.

We next state a very useful theorem about commuting observables, which we will prove only in the case of a Hilbert space of finite dimension.

**Theorem 2.7.1. (Compatible Observables)** *If  $A$  and  $B$  are observables, and  $[A, B] = 0$ , then there exists an orthobasis of kets that are eigenkets of both  $A$  and  $B$ . In other words, for commuting observables  $A$  and  $B$ , there is an orthobasis in which their matrix representations are both diagonal.*

Proof: From the definition of an observable, we already know that we can find an orthobasis of kets  $|\alpha, u_{\alpha}\rangle$  that are eigenkets of  $A$ . Furthermore, since  $[A, B] = 0$ , we know from Theorem 2.6.4 that whenever  $\alpha \neq \alpha'$ ,

$$\langle \alpha', u'_{\alpha'} | B | \alpha, u_{\alpha} \rangle = 0. \quad (2.7.6)$$

Therefore, the matrix representation of  $B$  has a block-diagonal form,

$$\begin{pmatrix} B_{\alpha_1} & 0 & 0 & \cdots & 0 \\ 0 & B_{\alpha_2} & 0 & \cdots & 0 \\ 0 & 0 & B_{\alpha_3} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & B_{\alpha_n} \end{pmatrix}, \quad (2.7.7)$$

where  $n \leq d$  is the number of distinct eigenvalues  $\alpha_j$  of  $A$ , and the blocks  $B_{\alpha_j}$  along the diagonal represent the possibly non-zero entries, which are confined to the subspaces of the Hilbert space that have the same eigenvalues  $\alpha_j$ . Each  $B_{\alpha_j}$  is a Hermitian  $g_{\alpha_j} \times g_{\alpha_j}$  matrix, where  $g_{\alpha_j}$  is the degeneracy<sup>†</sup> of  $\alpha_j$ , and has entries

$$\langle \alpha_j, u'_{\alpha_j} | B | \alpha_j, u_{\alpha_j} \rangle, \quad (u_{\alpha_j}, u'_{\alpha_j} = 1, \dots, g_{\alpha_j}). \quad (2.7.8)$$

Therefore, according to Theorem 2.6.7, each of the matrices  $B_{\alpha_j}$  can be diagonalized by a change of basis corresponding to a unitary operator. Each of these unitary operators acts non-trivially only within the corresponding  $B_{\alpha_j}$  block, and is the identity operator outside of that block. After doing these basis transformations, we will be left in an orthobasis in which the operators  $A$  and  $B$  are represented by diagonal matrices of the form

$$A = \text{diag}(\underbrace{\alpha_1, \dots, \alpha_1}_{g_{\alpha_1} \text{ times}}, \underbrace{\alpha_2, \dots, \alpha_2}_{g_{\alpha_2} \text{ times}}, \dots, \underbrace{\alpha_n, \dots, \alpha_n}_{g_{\alpha_n} \text{ times}}) \quad (2.7.9)$$

and

$$B = \text{diag}(\beta_{\alpha_1,1}, \dots, \beta_{\alpha_1,g_{\alpha_1}}, \beta_{\alpha_2,1}, \dots, \beta_{\alpha_2,g_{\alpha_2}}, \dots, \beta_{\alpha_n,1}, \dots, \beta_{\alpha_n,g_{\alpha_n}}), \quad (2.7.10)$$

where the  $\beta_{\alpha_j,k}$  with  $k = 1, \dots, g_{\alpha_j}$  are the eigenvalues of the  $B_{\alpha_j}$  sub-matrix. This is the desired orthobasis that achieves the requirements of the theorem.  $\square$

Even with two observables  $A$  and  $B$ , there can be unresolved degeneracies, by which we mean that there can be more than one orthobasis ket with the same eigenvalues  $\alpha$  and  $\beta$ . This raises the question of how to tell apart the corresponding degenerate states. From a physics perspective, if two states are genuinely distinct, then there must be some observable that distinguishes them. This leads to the important concept of a **Complete Set of Commuting Observables**, called a **CSCO** from here on. For any Hilbert space of states, a CSCO is a set of observables  $A, B, C, \dots$  that all commute with each other, and whose common eigenkets form an orthobasis with no degeneracies. Thus, we can write

$$A |\alpha, \beta, \gamma, \dots\rangle = \alpha |\alpha, \beta, \gamma, \dots\rangle, \quad (2.7.11)$$

$$B |\alpha, \beta, \gamma, \dots\rangle = \beta |\alpha, \beta, \gamma, \dots\rangle, \quad (2.7.12)$$

$$C |\alpha, \beta, \gamma, \dots\rangle = \gamma |\alpha, \beta, \gamma, \dots\rangle, \quad (2.7.13)$$

etc., where each of the eigenkets  $|\alpha, \beta, \gamma, \dots\rangle$  is *uniquely* determined by specifying its CSCO eigenvalues. The eigenvalues of a CSCO are sometimes known as **good quantum numbers**,

---

<sup>†</sup>Note that  $j = 1, \dots, n$ , and  $\sum_{j=1}^n g_{\alpha_j} = d$  is the dimension of the Hilbert space.

especially if one of them is the energy. For each one of the eigenvalue labels  $\alpha$  or  $\beta$  or  $\gamma$  etc., the others can be viewed as degeneracy labels.

If we have a set of commuting observables, but the degeneracy in the eigenkets is not fully eliminated, it means that we have not actually found a CSCO yet, and at least one more observable (commuting with all of the others) can be found to add to the list. Conversely, if the degeneracy has been fully eliminated by a list of commuting observables, then we have a CSCO; there is no need to add more observables to the list, even if it is easy to find more observables that commute with all of the others. As we will see in many examples, the number of observables needed to form a CSCO is always finite and typically not enormous for quantum systems with a fixed number of particles, even if the Hilbert space is infinite dimensional.

For a given quantum system, the choice of which operators to include in a CSCO is not unique. For example, even in the case of a spinless free particle moving in 3 dimensions, there are an infinite number of different choices we can make for the CSCO, all of which have 3 members. One CSCO choice turns out to be just the three spatial coordinates that specify the position of the particle. Another CSCO choice consists of the three momentum components of the particle. In some cases (those with rotational symmetry about some choice of origin), still another CSCO choice is the Hamiltonian (total energy) of the particle together with two operators associated with its angular momentum with respect to the origin. In general, the best choice of CSCO depends on what physics questions we would like to answer. Furthermore, it is often useful to be able to translate between the orthobases defined by different CSCOs.

Note that the word “complete” in CSCO does not have the same meaning as in the completeness relation. For a CSCO it is the operators that are complete, while for the completeness relation it is the orthobasis that is complete. Also, the second C in CSCO can equivalently be interpreted as standing for the word “compatible”. Observables are called **compatible** if they commute with each other, and are called **incompatible** if they have a non-zero commutator.

## 2.8 Wavefunctions

In the preceding, we have been mainly concerned with matrix representations in a finite dimensional Hilbert space. Let us now consider in more detail the case of an observable with a continuous spectrum of eigenvalues. A quintessential example is the position coordinate operator  $X$  for a particle moving in one dimension. The eigenvalues and eigenkets of  $X$  will be labeled as  $x$  and  $|x\rangle$ , respectively, so that the eigenvalue equation is

$$X|x\rangle = x|x\rangle. \quad (2.8.1)$$

The physical interpretation is that if the state is  $|x\rangle$ , then the position of the particle is known to be  $x$ , possibly because it has just been measured to be there. The allowed values of  $x$  form an

uncountable, continuous set, perhaps  $-\infty < x < \infty$ , or  $a < x < b$  if the particle is confined to that range. In the following, we will assume the latter, in order to be more general, and reserve the option to take  $a \rightarrow -\infty$  and  $b \rightarrow \infty$ .

Since a particle measured to be at position  $x$  is not at any other position  $x'$  at that precise moment, the states  $|x\rangle$  and  $|x'\rangle$  are orthogonal. Furthermore, a measurement of the particle's position must return one of the allowed values. These two statements are made precise as a special case of eqs. (2.7.2) and (2.7.3),

$$\langle x|x'\rangle = \delta(x - x'), \quad (2.8.2)$$

$$\int_a^b dx |x\rangle\langle x| = I. \quad (2.8.3)$$

To check that this is consistent, and that  $\delta(x - x')$  in eq. (2.8.2) should really be the Dirac delta function, consider an arbitrary ket  $|\psi\rangle$  in the Hilbert space. Then we define

$$\psi(x) \equiv \langle x|\psi\rangle \quad (2.8.4)$$

to be the **position wavefunction**. The wavefunction  $\psi(x)$  can also be viewed as comprising the components of the ket  $|\psi\rangle$  in the **position representation**. Using completeness, we have

$$\psi(x) = \langle x|\psi\rangle = \int_a^b dx' \langle x|x'\rangle\langle x'|\psi\rangle = \int_a^b dx' \delta(x - x') \psi(x'). \quad (2.8.5)$$

The fact that this should hold for every function  $\psi(x)$  is just the definition of the Dirac delta function  $\delta(x - x')$ ; see eq. (2.2.18).

The kets  $|x\rangle$  do not have finite norm, since  $\langle x|x\rangle = \delta(0)$  is not finite. This means that they are really generalized kets in the sense defined earlier [see the paragraph after eq. (2.2.18)], and are not associated with physical states in the Hilbert space, strictly speaking. However, eqs. (2.8.3)–(2.8.5) show that they do satisfy the requirements of orthonormality and completeness in the Dirac sense. Therefore, we will refer to the uncountable continuous set  $\{|x\rangle\}$  as the **position eigenstate orthobasis**, despite the fact that they are only idealized limits of physical states.

The inner product of two kets can be written in terms of an integral of the wavefunctions, using the completeness relation, as

$$\langle\phi|\psi\rangle = \int_a^b dx \langle\phi|x\rangle\langle x|\psi\rangle = \int_a^b dx \phi^*(x) \psi(x). \quad (2.8.6)$$

In particular, the squared norm of a state can be written in the manifestly non-negative form

$$\langle\psi|\psi\rangle = \int_a^b dx |\psi(x)|^2. \quad (2.8.7)$$

For valid physical states, both of eqs. (2.8.6) and (2.8.7) should be finite, and the latter must be non-zero. In other words, physical states correspond to wavefunctions that are **normalizable**. This set of functions forms a Hilbert space. However, for physics purposes, there is a further constraint that the wavefunctions must also be continuous. The reason for this is as follows. The norm of a ket is not physically significant, but it is often most convenient to choose the normalization of it so that it has norm 1. In particular, if the wavefunction for a single particle is chosen to have unit norm,  $\int_a^b dx |\psi(x)|^2 = 1$ , then  $|\psi(x)|^2$  can be interpreted as the probability density, in the sense that the probability to find the particle between  $x$  and  $x + dx$  is

$$d\mathcal{P}(x) = |\psi(x)|^2 dx. \quad (2.8.8)$$

This is known as the **Born rule** for the probabilistic interpretation of the wavefunction, after Max Born. On physical grounds, the probability density  $|\psi(x)|^2$  can in principle be measured, and therefore must be unambiguous, so it must be the same for any limiting approach to a given point  $x$ , implying continuity.

There is a one-to-one correspondence between kets  $|\psi\rangle$  and their wavefunctions  $\psi(x) = \langle x|\psi\rangle$ . Now consider the ket corresponding to the derivative of the wavefunction,  $d\psi/dx$ . Let us write

$$|d\psi/dx\rangle = D|\psi\rangle, \quad (2.8.9)$$

which defines a linear operator  $D$  on the space of kets. We then have

$$\langle x|D|\psi\rangle = \langle x|d\psi/dx\rangle = d\psi/dx. \quad (2.8.10)$$

Using completeness gives a condition satisfied by the position-eigenket matrix elements of  $D$ ,

$$d\psi/dx = \int_a^b dx' \langle x|D|x'\rangle \langle x'|\psi\rangle = \int_a^b dx' \langle x|D|x'\rangle \psi(x'). \quad (2.8.11)$$

We also have

$$d\psi/dx = \frac{d}{dx} \left( \int_a^b dx' \psi(x') \delta(x - x') \right) = \int_a^b dx' \psi(x') \frac{d}{dx} \delta(x - x'). \quad (2.8.12)$$

Comparison of eqs. (2.8.11) and (2.8.12) implies

$$\langle x|D|x'\rangle = \frac{d}{dx} \delta(x - x'). \quad (2.8.13)$$

To see this, plug in  $\psi(x') = \delta(x' - x'')$ , do the integrations, and then relabel  $x'' \rightarrow x'$ .

We now see that  $D$  cannot be an observable, because it is certainly not Hermitian, since

$$\langle x'|D^\dagger|x\rangle = \left( \langle x|D|x'\rangle \right)^* = \frac{d}{dx} \delta(x - x') = -\frac{d}{dx'} \delta(x' - x) = -\langle x'|D|x\rangle. \quad (2.8.14)$$

Although  $D$  is not Hermitian, the minus sign at the end of eq. (2.8.14) suggests that it is anti-Hermitian, and we can simply multiply by a factor of  $i$  or  $-i$  to make an observable. We therefore define<sup>†</sup> the **momentum operator**,

$$P = -i\hbar D, \quad (2.8.15)$$

which could be Hermitian and thus a candidate to be an observable. However, since this is an infinite-dimensional Hilbert space, and eq. (2.8.14) was derived using the technically ill-defined derivatives of the Dirac delta function, it is prudent to be suspicious about the Hermiticity of  $P$ . So, let us check.

From eq. (2.8.9),

$$P|\psi\rangle = -i\hbar|d\psi/dx\rangle. \quad (2.8.16)$$

Therefore, given any two kets  $|f\rangle$  and  $|g\rangle$ , with wavefunctions  $\langle x|f\rangle = f(x)$  and  $\langle x|g\rangle = g(x)$ ,

$$\langle g|P|f\rangle = \int_a^b dx \langle g|x\rangle \langle x|P|f\rangle = -i\hbar \int_a^b dx g(x)^* \frac{d}{dx} f(x), \quad (2.8.17)$$

where the first equality used completeness. Interchanging the roles of  $f$  and  $g$ , we also have

$$\langle f|P|g\rangle = -i\hbar \int_a^b dx f(x)^* \frac{d}{dx} g(x). \quad (2.8.18)$$

Therefore, the difference between  $P^\dagger$  and  $P$ , in an arbitrary matrix element, is

$$\langle f|P^\dagger|g\rangle - \langle f|P|g\rangle = (\langle g|P|f\rangle)^* - \langle f|P|g\rangle \quad (2.8.19)$$

$$= i\hbar \int_a^b dx g(x) \frac{d}{dx} f(x)^* + i\hbar \int_a^b dx f(x)^* \frac{d}{dx} g(x) \quad (2.8.20)$$

$$= i\hbar \int_a^b dx \frac{d}{dx} [f(x)^* g(x)] \quad (2.8.21)$$

$$= i\hbar [f(b)^* g(b) - f(a)^* g(a)]. \quad (2.8.22)$$

So, for  $P$  to be a Hermitian operator, we must have

$$f(b)^* g(b) = f(a)^* g(a). \quad (2.8.23)$$

This can be considered a condition on allowed wavefunctions (and their corresponding kets) in the physical Hilbert space of states if  $P$  is to be an observable.

For example, for a particle confined to a box with impenetrable walls at  $x = a$  and  $x = b$ , eq. (2.8.23) is satisfied because all wavefunctions are required to obey  $\psi(a) = \psi(b) = 0$ . It is also

---

<sup>†</sup>The sign and the  $\hbar$  normalization in the definition of  $P$  are mysteriously arbitrary at this point, but will be justified in what follows; see the discussions following eq. (2.8.39) and surrounding eqs. (3.4.17) and (3.4.18). For now, note that the presence of  $\hbar$  at least has the correct units to make  $P$  a momentum.

satisfied if all wavefunctions satisfy the weaker condition  $\psi(b) = e^{i\varphi}\psi(a)$ , where  $\varphi$  is any fixed constant. This can occur if  $X$  is interpreted to be some kind of periodic or angular coordinate  $\theta$ , defined for example on the interval  $0 \leq \theta \leq 2\pi$ ; then  $\varphi = 0$  implies that wavefunctions are single-valued at the point  $\theta = 0$ , which is identified with  $\theta = 2\pi$ .

If the domain of  $x$  is infinite, with  $a = -\infty$  and  $b = \infty$ , then things are trickier. In practical solutions for wavefunctions, one sometimes finds that the condition  $\psi(x) \rightarrow 0$  at large spatial distances is naturally satisfied. States with this property are called **bound states**. If either  $|f\rangle$  or  $|g\rangle$  is one of these, then eq. (2.8.23) is satisfied. However, there may also be other solutions that do not obey this property, called **unbound states** or **scattering states**. These are very useful idealized solutions to simple problems, but strictly speaking they are generalized ket states rather than physical ones, since they cannot have finite norm. In fact, finiteness of the norms for true physical states  $|f\rangle$  and  $|g\rangle$  ensures that  $f(x)$  and  $g(x)$  must approach 0 for large  $|x|$ . One resolution is that in a more precise formulation that maintains Hermiticity of  $P$ , the idealized unbound or scattering states should be replaced by more complicated states whose wavefunctions are required to vanish only at very large distances, say, outside of a box with sides of length  $L = \text{several light-years}$ , to be safe. Another possibility is to impose periodic boundary conditions on the wavefunctions in the large box, so that they need not vanish even at large distances, but rather obey  $\psi(x - L/2) = \psi(x + L/2)$ . For  $|x| \ll L/2$ , the unbound state wavefunctions of interest will be closely approximated by a linear combination of these periodic wavefunctions, for which eq. (2.8.23) holds. These modifications will have a completely negligible effect on physical questions about phenomena localized far from the edges of the box, and in practice one can use the idealized unbound states with impunity in most cases.

Next, consider the eigenvalue problem for the momentum operator. The eigenkets with definite and constant momentum  $p$ , corresponding to a **free particle**, satisfy

$$P|p\rangle = p|p\rangle. \quad (2.8.24)$$

Similarly to the case of position eigenstates, these are taken to obey Dirac orthonormality and completeness conditions, again as a special case of eqs. (2.7.2) and (2.7.3),

$$\langle p|p'\rangle = \delta(p - p'), \quad (2.8.25)$$

$$\int_{-\infty}^{\infty} dp |p\rangle\langle p| = I. \quad (2.8.26)$$

Here the allowed range of continuous  $p$  is taken to be from  $-\infty$  to  $\infty$ . As should now be familiar, the generalized kets  $|p\rangle$  are, strictly speaking, not associated with physical states, because they do not have finite norm, but are still very useful as idealizations, because they satisfy the properties of an orthobasis in the Dirac sense. For any ket  $|\psi\rangle$ , we define the **momentum**



## wavefunction

$$\tilde{\psi}(p) \equiv \langle p | \psi \rangle, \quad (2.8.27)$$

which can be viewed as the components of the ket  $|\psi\rangle$  in the **momentum representation**. Using the completeness relation eq. (2.8.26), the inner product of any two states can be written

$$\langle \phi | \psi \rangle = \int_{-\infty}^{\infty} dp \tilde{\phi}^*(p) \tilde{\psi}(p). \quad (2.8.28)$$

If one has chosen the ket  $|\psi\rangle$  to have norm 1, then

$$\int_{-\infty}^{\infty} dp |\tilde{\psi}(p)|^2 = 1, \quad (2.8.29)$$

and we can interpret

$$d\mathcal{P}(p) = |\tilde{\psi}(p)|^2 dp \quad (2.8.30)$$

as the probability for the particle in the state  $|\psi\rangle$  to have momentum between  $p$  and  $p + dp$ .

Consider the position wavefunctions of the free-particle momentum eigenstates, defined by

$$\phi_p(x) \equiv \langle x | p \rangle. \quad (2.8.31)$$

Now we have

$$\langle x | P | p \rangle = p \langle x | p \rangle = p \phi_p(x), \quad (2.8.32)$$

but also, using the completeness relation for position eigenkets,

$$\langle x | P | p \rangle = \int_{-\infty}^{\infty} dx' \langle x | P | x' \rangle \langle x' | p \rangle = \int_{-\infty}^{\infty} dx' \left( -i\hbar \frac{d}{dx} \delta(x - x') \right) \phi_p(x') \quad (2.8.33)$$

$$= -i\hbar \frac{d}{dx} \left( \int_{-\infty}^{\infty} dx' \delta(x - x') \phi_p(x') \right) = -i\hbar \frac{d}{dx} \phi_p(x). \quad (2.8.34)$$

Comparing this to eq. (2.8.32), we see that the momentum eigenvalue equation in the position representation is a linear first-order differential equation,

$$\frac{d}{dx} \phi_p(x) = i \frac{p}{\hbar} \phi_p(x), \quad (2.8.35)$$

with the solution

$$\phi_p(x) = c_p e^{ipx/\hbar}, \quad (2.8.36)$$

where  $c_p$  is a non-zero complex number. To fix the magnitude of  $c_p$ , we use completeness in  $x$ ,

$$\langle p' | p \rangle = \int_{-\infty}^{\infty} dx \langle p' | x \rangle \langle x | p \rangle = \int_{-\infty}^{\infty} dx \phi_{p'}(x)^* \phi_p(x) = c_{p'}^* c_p \int_{-\infty}^{\infty} dx e^{ix(p-p')/\hbar} \quad (2.8.37)$$

$$= |c_p|^2 2\pi\hbar \delta(p - p'), \quad (2.8.38)$$

where the last equality used the delta function expression eq. (2.2.20). So, in order to enforce the Dirac orthonormality condition of eq. (2.8.25), we choose  $c_p = 1/\sqrt{2\pi\hbar}$ , and so

$$\langle x|p\rangle = \phi_p(x) = \frac{1}{\sqrt{2\pi\hbar}} e^{ipx/\hbar}, \quad (2.8.39)$$

up to an unavoidably arbitrary choice of constant phase. From the physical consideration that the wavefunctions  $\phi_p(x)$  should not blow up at large  $|x|$ ,  $p$  must be real, as appropriate for the eigenvalue of a Hermitian operator  $P$ . The position dependence of the momentum eigenstate is seen to have the form of a plane wave, with wavenumber  $p/\hbar$ , and therefore wavelength  $2\pi\hbar/p$ . Comparison with de Broglie's eq. (1.4.1) confirms that we made the correct choice of magnitude of normalization of the momentum operator  $P$  in eq. (2.8.15).

As a further check of the completeness condition for momentum eigenkets in eq. (2.8.26), take the matrix element of it between two arbitrary position eigenkets,

$$\langle x'|x\rangle = \int_{-\infty}^{\infty} dp \langle x'|p\rangle \langle p|x\rangle = \frac{1}{2\pi\hbar} \int_{-\infty}^{\infty} dp e^{ip(x'-x)/\hbar} = \delta(x-x'). \quad (2.8.40)$$

In particular, this confirms that we must integrate over the whole range  $-\infty < p < \infty$  in the momentum completeness relation eq. (2.8.26).

We can also use completeness to find formulas that convert between the momentum wavefunction and the position wavefunction. Using completeness in  $x$ ,

$$\tilde{\psi}(p) = \langle p|\psi\rangle = \int_{-\infty}^{\infty} dx \langle p|x\rangle \langle x|\psi\rangle = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} dx e^{-ipx/\hbar} \psi(x). \quad (2.8.41)$$

Conversely, given  $\tilde{\psi}(p)$ , we can reconstruct  $\psi(x)$  using completeness in  $p$ ,

$$\psi(x) = \langle x|\psi\rangle = \int_{-\infty}^{\infty} dp \langle x|p\rangle \langle p|\psi\rangle = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} dp e^{ipx/\hbar} \tilde{\psi}(p). \quad (2.8.42)$$

In words, the position and momentum wavefunctions are Fourier transforms of each other, with some  $\hbar$ 's included. It is also easy to check that

$$\langle x|X|\psi\rangle = x\psi(x), \quad \langle x|P|\psi\rangle = -i\hbar \frac{d}{dx} \psi(x), \quad (2.8.43)$$

$$\langle p|P|\psi\rangle = p\tilde{\psi}(p), \quad \langle p|X|\psi\rangle = i\hbar \frac{d}{dp} \tilde{\psi}(p). \quad (2.8.44)$$

These are analogous to matrix representations of observables in a finite dimensional Hilbert space. If one works in the position representation, with wavefunctions of  $x$ , then the observable  $X$  is represented by  $x$  and the observable  $P$  is represented by  $-i\hbar d/dx$ . Inner products are accomplished by integration over all  $x$ , with complex conjugation for the wavefunction of the bra vector. If, instead, one works in the momentum representation with wavefunctions  $\tilde{\psi}(p)$ ,

Representation	$ \psi\rangle$	$\langle\psi $	$X$	$P$	$\langle\psi_1 \psi_2\rangle$
position	$\psi(x)$	$\psi(x)^*$	$x$	$-i\hbar\frac{d}{dx}$	$\int dx \psi_1(x)^* \psi_2(x)$
momentum	$\tilde{\psi}(p)$	$\tilde{\psi}(p)^*$	$i\hbar\frac{d}{dp}$	$p$	$\int dp \tilde{\psi}_1(p)^* \tilde{\psi}_2(p)$

Table 2.8.1: Summary of the position and momentum representations for a particle moving in one dimension. The position and momentum wavefunctions are related to each other as in eqs. (2.8.41) and (2.8.42).

then  $P$  is represented by  $p$  and  $X$  is represented by  $i\hbar d/dp$ , and inner products are done by integration over all  $p$ . These results are summarized in Table 2.8.1. For example,

$$\langle\psi_1|X|\psi_2\rangle = \int_{-\infty}^{\infty} dx \psi_1(x)^* x \psi_2(x) = \int_{-\infty}^{\infty} dp \tilde{\psi}_1(p)^* \left(i\hbar\frac{d}{dp}\right) \tilde{\psi}_2(p), \quad (2.8.45)$$

$$\langle\psi_1|P|\psi_2\rangle = \int_{-\infty}^{\infty} dx \psi_1(x)^* \left(-i\hbar\frac{d}{dx}\right) \psi_2(x) = \int_{-\infty}^{\infty} dp \tilde{\psi}_1(p)^* p \tilde{\psi}_2(p). \quad (2.8.46)$$

The position representation provides a convenient way to find the commutator of  $X$  and  $P$ . First we evaluate

$$\langle x|XP|\psi\rangle = x\left(-i\hbar\frac{d}{dx}\right)\psi(x) = -i\hbar x\frac{d\psi}{dx}, \quad (2.8.47)$$

$$\langle x|PX|\psi\rangle = -i\hbar\frac{d}{dx}(x\psi(x)) = -i\hbar x\frac{d\psi}{dx} - i\hbar\psi(x). \quad (2.8.48)$$

Therefore, for every ket  $|\psi\rangle$ ,

$$\langle x|[X, P]|\psi\rangle = i\hbar\langle x|\psi\rangle, \quad (2.8.49)$$

so we can conclude that

$$[X, P] = i\hbar, \quad (2.8.50)$$

where the identity operator on the right side is understood. This position-momentum commutation relation was derived after defining the operator  $P$  by its action on the orthobasis of position eigenkets, but one could just as easily have worked in the other direction, taking eq. (2.8.50) to be the fundamental definition, and then deriving the operation of  $P$  on the kets.

The preceding can all be generalized in a straightforward way to three position and three momentum observables. We define position and momentum operators that are vectors in real space (not the Hilbert vector space, in which they are Hermitian operators), by

$$\vec{R} = \hat{x}X + \hat{y}Y + \hat{z}Z, \quad (2.8.51)$$

$$\vec{P} = \hat{x}P_x + \hat{y}P_y + \hat{z}P_z. \quad (2.8.52)$$

They satisfy commutation relations:

$$[X, P_x] = i\hbar, \quad [Y, P_y] = i\hbar, \quad [Z, P_z] = i\hbar, \quad (2.8.53)$$

with all other relevant commutators vanishing. It is convenient to adopt a notation in which the components of  $\vec{R}$  and  $\vec{P}$  are given an index  $a = 1, 2, 3$  corresponding to  $x, y, z$  respectively, so that  $R_1 = X$ ,  $R_2 = Y$ ,  $R_3 = Z$ , and  $P_1 = P_x$ ,  $P_2 = P_y$ ,  $P_3 = P_z$ . Then the commutation relations take the form

$$[R_a, P_b] = i\hbar\delta_{ab}, \quad [R_a, R_b] = 0, \quad [P_a, P_b] = 0. \quad (2.8.54)$$

The operators  $\vec{R}$  and  $\vec{P}$  have eigenkets  $|\vec{r}\rangle$  and  $|\vec{p}\rangle$  with<sup>†</sup> eigenvalues  $\vec{r} = \hat{x}x + \hat{y}y + \hat{z}z$  and  $\vec{p} = \hat{x}p_x + \hat{y}p_y + \hat{z}p_z$ , so that

$$\vec{R}|\vec{r}\rangle = \vec{r}|\vec{r}\rangle, \quad \vec{P}|\vec{p}\rangle = \vec{p}|\vec{p}\rangle. \quad (2.8.55)$$

These eigenkets satisfy completeness relations

$$I = \int d^3\vec{r} |\vec{r}\rangle\langle\vec{r}|, \quad (2.8.56)$$

$$I = \int d^3\vec{p} |\vec{p}\rangle\langle\vec{p}|, \quad (2.8.57)$$

and have inner products realizing Dirac orthonormality,

$$\langle\vec{r}'|\vec{r}\rangle = \delta^{(3)}(\vec{r} - \vec{r}'), \quad \langle\vec{p}'|\vec{p}\rangle = \delta^{(3)}(\vec{p} - \vec{p}'). \quad (2.8.58)$$

Here the three-dimensional delta function is given in rectangular coordinates by

$$\delta^{(3)}(\vec{r} - \vec{r}') = \delta(x - x')\delta(y - y')\delta(z - z'), \quad (2.8.59)$$

and in spherical coordinates by

$$\delta^{(3)}(\vec{r} - \vec{r}') = \frac{1}{r^2} \delta(r - r') \delta(\phi - \phi') \delta(\cos\theta - \cos\theta'), \quad (2.8.60)$$

with a practical definition that, when integrating over a volume  $V$ ,

$$\int_V d^3\vec{r}' f(\vec{r}') \delta^{(3)}(\vec{r} - \vec{r}') = \begin{cases} f(\vec{r}) & \text{if } \vec{r} \text{ is inside the volume } V, \\ 0 & \text{if } \vec{r} \text{ is outside the volume } V. \end{cases} \quad (2.8.61)$$

The wavefunction for a free particle with momentum eigenvalue  $\vec{p}$  is a plane wave,

$$\langle\vec{r}|\vec{p}\rangle = \frac{1}{(2\pi\hbar)^{3/2}} e^{i\vec{p}\cdot\vec{r}/\hbar}. \quad (2.8.62)$$

---

<sup>†</sup>Here we have stretched the previous notion of eigenvalue slightly, as foreshadowed after eq. (2.7.1), because our eigenvalues here are actually not just numbers, but vectors in real space, or equivalently ordered triples of numbers  $(x, y, z)$  or  $(p_x, p_y, p_z)$ . This was a sneaky thing to do, but it is convenient, and is perfectly valid *if* the components are compatible observables, as here. This will be discussed further in section 3.2.

The position and momentum wavefunctions for a general state  $|\psi\rangle$  are

$$\psi(\vec{r}) = \langle \vec{r} | \psi \rangle, \quad \tilde{\psi}(\vec{p}) = \langle \vec{p} | \psi \rangle, \quad (2.8.63)$$

and they are related by Fourier transforms,

$$\tilde{\psi}(\vec{p}) = \frac{1}{(2\pi\hbar)^{3/2}} \int d^3\vec{r} e^{-i\vec{p}\cdot\vec{r}/\hbar} \psi(\vec{r}), \quad (2.8.64)$$

$$\psi(\vec{r}) = \frac{1}{(2\pi\hbar)^{3/2}} \int d^3\vec{p} e^{i\vec{p}\cdot\vec{r}/\hbar} \tilde{\psi}(\vec{p}), \quad (2.8.65)$$

which follows from eq. (2.8.62) and the completeness relations (2.8.56) and (2.8.57). The squared norm of a state can be written in terms of either the position or momentum wavefunction, as

$$\langle \psi | \psi \rangle = \int d^3\vec{r} |\psi(\vec{r})|^2 = \int d^3\vec{p} |\tilde{\psi}(\vec{p})|^2. \quad (2.8.66)$$

The interpretation of the integrands is that the probability for the particle described by the state  $|\psi\rangle$  to be found in an infinitesimal volume  $d^3\vec{r}$  near  $\vec{r}$  is given by the Born rule,

$$d\mathcal{P}(\vec{r}) = d^3\vec{r} |\psi(\vec{r})|^2, \quad (2.8.67)$$

while the probability for it to have momentum in a volume  $d^3\vec{p}$  in momentum space near  $\vec{p}$  is

$$d\mathcal{P}(\vec{p}) = d^3\vec{p} |\tilde{\psi}(\vec{p})|^2. \quad (2.8.68)$$

The last claims are based on postulates that will be stated more generally in section 3.1.

We also note, for future reference, that in the position representation,  $\vec{P}$  is represented by  $-i\hbar\vec{\nabla}$ , where  $\vec{\nabla}$  is the gradient, and the momentum squared operator  $P^2 = \vec{P}\cdot\vec{P}$  is represented by  $-\hbar^2\nabla^2$ , where  $\nabla^2$  is the Laplacian. The rules for the position and momentum representations for a particle in three dimensions are summarized in Table 2.8.2.

Representation	$ \psi\rangle$	$\langle\psi $	$\vec{R}$	$\vec{P}$	$P^2$	$\langle\psi_1 \psi_2\rangle$
Position	$\psi(\vec{r})$	$\psi(\vec{r})^*$	$\vec{r}$	$-i\hbar\vec{\nabla}$	$-\hbar^2\nabla^2$	$\int d^3\vec{r} \psi_1(\vec{r})^* \psi_2(\vec{r})$
Momentum	$\tilde{\psi}(\vec{p})$	$\tilde{\psi}(\vec{p})^*$	$i\hbar\vec{\nabla}_p$	$\vec{p}$	$p^2$	$\int d^3\vec{p} \tilde{\psi}_1(\vec{p})^* \tilde{\psi}_2(\vec{p})$

Table 2.8.2: Summary of the position and momentum representations for a particle moving in three dimensions. The position and momentum wavefunctions are related as in eqs. (2.8.64) and (2.8.65). In rectangular coordinates,  $\vec{\nabla}_p = \hat{x}\partial/\partial p_x + \hat{y}\partial/\partial p_y + \hat{z}\partial/\partial p_z$ .

## 2.9 Tensor product Hilbert spaces

We will often want to consider state spaces that are combinations of simpler state spaces. For example, consider a particle moving in three dimensions. In the previous subsection, we worked with a Hilbert space spanned by an orthobasis of kets that are eigenkets of all three position coordinates. An equivalent way to proceed is to first define three separate Hilbert spaces that only describe the  $x$ ,  $y$ , and  $z$  degrees of freedom, and then combine them to form the Hilbert space that describes all three degrees of freedom. As another example, a Hilbert space for a system of two particles can be constructed from the Hilbert spaces of the two particles separately.

The formal construction of Hilbert spaces from simpler components is called a **tensor product** Hilbert space. Consider two distinct Hilbert spaces  $\mathcal{H}_1$  and  $\mathcal{H}_2$ , spanned respectively by orthobases  $\{|\varphi_j\rangle\}$  with dimension  $d_1$  and  $\{|v_k\rangle\}$  with dimension  $d_2$ . Then the tensor product Hilbert space  $\mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2$  is defined to be a complex linear vector space with inner product with the following properties:

- There is an orthobasis for  $\mathcal{H}$  denoted  $|\varphi_j\rangle \otimes |v_k\rangle$  for  $j = 1, \dots, d_1$  and  $k = 1, \dots, d_2$ . Thus,  $\mathcal{H}$  has dimension  $d = d_1 d_2$ , and any ket in  $\mathcal{H}$  can be written in a unique way as a linear combination of the  $d$  kets  $|\varphi_j\rangle \otimes |v_k\rangle$ .
- Tensor product kets satisfy linearity and distributive properties. This means that if  $|\Phi\rangle$  and  $|\Psi\rangle$  are any kets in  $\mathcal{H}_1$ , and  $|V\rangle$  and  $|W\rangle$  are kets in  $\mathcal{H}_2$ , and  $a, b, c, d$  are complex numbers, then

$$\begin{aligned} (a|\Phi\rangle + b|\Psi\rangle) \otimes (c|V\rangle + d|W\rangle) &= ac|\Phi\rangle \otimes |V\rangle + ad|\Phi\rangle \otimes |W\rangle \\ &\quad + bc|\Psi\rangle \otimes |V\rangle + bd|\Psi\rangle \otimes |W\rangle. \end{aligned} \quad (2.9.1)$$

- The dual tensor product Hilbert space has an orthobasis  $\langle\varphi_j| \otimes \langle v_k|$ .
- The inner product of orthobasis kets in  $\mathcal{H}$  is inherited from the inner products of the  $\mathcal{H}_1$  and  $\mathcal{H}_2$  orthobasis kets, according to

$$(\langle\varphi_j| \otimes \langle v_k|)(|\varphi_l\rangle \otimes |v_m\rangle) = \delta_{jl} \delta_{km}. \quad (2.9.2)$$

- Given an operator  $A$  that acts on  $\mathcal{H}_1$  and an operator  $B$  that acts on  $\mathcal{H}_2$ , the tensor product operator  $A \otimes B$  is defined to act on  $\mathcal{H}$  according to

$$(A \otimes B)(|\Psi\rangle \otimes |V\rangle) = (A|\Psi\rangle) \otimes (B|V\rangle), \quad (2.9.3)$$

for any kets  $|\Psi\rangle$  in  $\mathcal{H}_1$  and  $|V\rangle$  in  $\mathcal{H}_2$ . In the very common case that  $B$  is the identity operator, we simply write  $A$  instead of  $A \otimes I$ . Thus, if it is understood that the operator

$A$  acts non-trivially only on  $\mathcal{H}_1$ , then we write  $A(|\Psi\rangle \otimes |V\rangle) = (A|\Psi\rangle) \otimes |V\rangle$ . Similarly, if  $A$  is the identity operator, then we simply write  $B$  instead of  $I \otimes B$ .

Even though an orthobasis of kets for  $\mathcal{H}$  can be chosen so that their elements are tensor products  $|\varphi_j\rangle \otimes |v_k\rangle$ , not all kets in  $\mathcal{H}$  are tensor products. Only special kets, called **separable**, can be written as a tensor product of kets,

$$|\Phi\rangle \otimes |W\rangle. \quad (2.9.4)$$

A ket in a tensor product Hilbert space that cannot be written in this form, for any choice of  $|\Phi\rangle$  and  $|W\rangle$ , is said to be an **entangled state**, a concept first highlighted by Erwin Schrödinger in 1935. Because  $\mathcal{H}$  is defined to be a vector space, all linear combinations of its elements must also be included in it. Therefore, for example,

$$|\Psi_1\rangle \otimes |V_1\rangle + |\Psi_2\rangle \otimes |V_2\rangle \quad (2.9.5)$$

is certainly an element of  $\mathcal{H}$ . However, it cannot be written as a tensor product of kets  $|\Phi\rangle \otimes |W\rangle$  unless either  $|\Psi_1\rangle = c|\Psi_2\rangle$  or  $|V_1\rangle = c|V_2\rangle$  for some complex constant  $c$ . We will have much more to say about the properties of entangled states in Chapters 14, 24, 25, and 26.

One of the common uses of the tensor product formalism is as a divide-and-conquer strategy for solving problems. This includes situations where we can make use of an already-solved problem for one or more components of the tensor product. As a simple example, the Hilbert space  $\mathcal{H}$  for a free particle moving in three dimensions can be written as a tensor product of Hilbert spaces that would describe a particle moving in the three rectangular coordinates separately,

$$\mathcal{H} = \mathcal{H}_x \otimes \mathcal{H}_y \otimes \mathcal{H}_z. \quad (2.9.6)$$

An orthobasis for this Hilbert space is

$$|\vec{r}\rangle = |x\rangle \otimes |y\rangle \otimes |z\rangle, \quad (2.9.7)$$

where  $|x\rangle$  describes a state in which a particle is known to have  $X$  eigenvalue  $x$ . In this example, as in many other cases, the tensor product notation  $|x\rangle \otimes |y\rangle \otimes |z\rangle$  has no real advantage over just writing the typographically cleaner

$$|\vec{r}\rangle = |x, y, z\rangle, \quad (2.9.8)$$

so that is what we will do from now on. Then,  $X|x, y, z\rangle = x|x, y, z\rangle$  and  $Y|x, y, z\rangle = y|x, y, z\rangle$ , and  $Z|x, y, z\rangle = z|x, y, z\rangle$ . This is similar to the separation-of-variables strategy for solving partial differential equations problems with several independent variables.

In the following, we will study the orbital angular momentum eigenvalue problem by isolating only the relevant angular  $(\theta, \phi)$  component of the three-dimensional particle Hilbert space, and then apply the results more generally to the whole Hilbert space. Schematically, this relies on using an orthobasis that is a tensor product of the form

$$|\text{radial part}\rangle \otimes |\text{angular momentum part}\rangle, \quad (2.9.9)$$

although it is not necessarily notationally convenient to write it that way explicitly. This will be made more precise in section 8.6 after we develop the theory of angular momentum.

Another important use of tensor product spaces is to describe systems of more than one particle. For example, we can construct a Hilbert space that describes two distinct spinless particles labeled 1 and 2, with an orthobasis of kets

$$|\vec{r}_1, \vec{r}_2\rangle = |\vec{r}_1\rangle \otimes |\vec{r}_2\rangle. \quad (2.9.10)$$

Here, each  $|\vec{r}_n\rangle$  is the Hilbert space for just one particle  $n = 1, 2$  moving in three dimensions. This construction can naturally be generalized to a tensor product Hilbert space for  $N$  spinless particles with orthobasis kets

$$|\vec{r}_1, \dots, \vec{r}_N\rangle = |\vec{r}_1\rangle \otimes \dots \otimes |\vec{r}_N\rangle. \quad (2.9.11)$$

However, when the particles have spin (intrinsic angular momentum) or are identical, things are more complicated, as we will discuss in Chapter 18.

## 2.10 Exercises

**Exercise 2.1.** Use the Cauchy–Schwarz inequality (2.2.4) to prove the triangle inequality (2.2.7), and show that equality holds if and only if  $|w\rangle = n|v\rangle$  where  $n$  is a non-negative real number. (Hint: start with  $\|(|v\rangle + |w\rangle)\|^2 = |(\langle v| + \langle w|)(|v\rangle + |w\rangle)|$ , and use the property of complex numbers  $\text{Re}(z) \leq |z|$ .)

**Exercise 2.2.** Use the definition of the adjoint to show that  $(|w\rangle\langle v|)^\dagger = |v\rangle\langle w|$  and that  $(AB)^\dagger = B^\dagger A^\dagger$ .

**Exercise 2.3.** Suppose that  $U$  is a unitary operator.

- (a) Show that if  $A$  is a Hermitian operator, then  $U^\dagger A U$  is also Hermitian.
- (b) Show that if  $V$  is another unitary operator, then  $VU$  and  $U^\dagger V U$  are each also unitary.

**Exercise 2.4.** Consider the trace of an operator  $A$  defined in terms of an orthobasis  $|\varphi_k\rangle$  by eq. (2.6.57).



- (a) Show that  $\text{Tr}(A)$  does not depend on the choice of orthonormal basis. To do this, consider any other orthobasis of kets  $|\chi_q\rangle$ , and show that  $\text{Tr}(A)$  is also equal to  $\sum_q \langle \chi_q | A | \chi_q \rangle$ .
- (b) Show that  $\text{Tr}(AB) = \text{Tr}(BA)$ .
- (c) Show that the trace of the outer product is the inner product,  $\text{Tr}(|v\rangle \langle w|) = \langle w | v \rangle$ .

**Exercise 2.5.** Suppose  $A$ ,  $B$ , and  $C$  are operators.

- (a) Show that  $[A, BC] = [A, B]C + B[A, C]$ .
- (b) Prove the **Jacobi identity**,  $[A, [B, C]] + [B, [C, A]] + [C, [A, B]] = 0$ .
- (c) If  $[[A, B], B] = 0$ , show that  $[A, B^n] = n[A, B]B^{n-1}$ . (Hint: do this by mathematical induction, by showing that if it is true for some  $n$ , then it is also true for  $n + 1$ .)

**Exercise 2.6.** Prove Theorem 2.4.5. (Hints: define  $F(\lambda) = e^{\lambda(A+B)}e^{-\lambda B}e^{-\lambda A}$ . Then show that  $dF/d\lambda = c\lambda[A, B]F(\lambda)$  where  $c$  is a number that you will find. You may use the fact that  $\frac{d}{d\lambda}e^{\lambda X} = Xe^{\lambda X} = e^{\lambda X}X$  for any operator  $X$ , and the power series expansion for the exponential function, and the result of Exercise 2.5(c). Then solve the differential equation for  $F(\lambda)$  using the boundary condition at  $\lambda = 0$ , and plug in the value  $\lambda = 1$  to get the desired result.)

**Exercise 2.7.** Consider a vector space spanned by three orthobasis kets  $|1\rangle$ ,  $|2\rangle$ , and  $|3\rangle$ , and a Hermitian operator  $A$  defined by

$$A|1\rangle = 5|1\rangle + i|2\rangle, \quad A|2\rangle = -i|1\rangle + 5|2\rangle, \quad A|3\rangle = 4|3\rangle. \quad (2.10.1)$$

- (a) Write down the  $3 \times 3$  matrix representation of  $A$  in this basis.
- (b) Find the eigenvalues of  $A$ . Two of them should be equal (degenerate).
- (c) For the non-degenerate eigenvalue, show that the most general form of the eigenvector is, in ket form,  $c(|1\rangle + i|2\rangle)$ , where  $c$  is an arbitrary complex number.
- (d) For the degenerate eigenvalue, find the most general form for a single eigenvector; you should write it in terms of two arbitrary complex numbers. Write your answer both in three-component column vector and ket forms. Check that this vector is orthogonal to the result in part (c).

**Exercise 2.8.** Consider a state space spanned by an orthobasis consisting of three kets  $|1\rangle$ ,  $|2\rangle$ , and  $|3\rangle$ . Let two Hermitian operators  $A$  and  $B$  be defined by

$$A|1\rangle = 2a|1\rangle + a|3\rangle, \quad A|2\rangle = a|2\rangle, \quad A|3\rangle = a|1\rangle + 2a|3\rangle, \quad (2.10.2)$$

$$B|1\rangle = b|1\rangle + 2b|3\rangle, \quad B|2\rangle = 0, \quad B|3\rangle = 2b|1\rangle + b|3\rangle, \quad (2.10.3)$$

where  $a$  and  $b$  are constant real numbers. The orthobasis kets can be represented by vectors

$$|1\rangle \leftrightarrow v_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad |2\rangle \leftrightarrow v_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad |3\rangle \leftrightarrow v_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}. \quad (2.10.4)$$

- (a) What are the  $3 \times 3$  matrix representations of the operators  $A$  and  $B$  in this basis?
- (b) Show that  $[A, B] = 0$ , which implies that  $A$  and  $B$  can be simultaneously diagonalized.
- (c) Find the eigenvalues of  $A$  and the eigenvalues of  $B$ . (One of them has non-degenerate eigenvalues; you would be wise to use this to plan your strategy for the next part.)
- (d) Find an orthonormal basis  $v'_1, v'_2$ , and  $v'_3$  of vectors that are eigenvectors of both  $A$  and  $B$ . For each of them, give their eigenvalues with respect to each of  $A$  and  $B$ .
- (e) Write the matrix  $U$  that transforms the original orthobasis into the one you found in the previous part. This means that  $v'_1 = Uv_1$  and  $v'_2 = Uv_2$  and  $v'_3 = Uv_3$ . Check by direct computation that  $U$  is unitary.
- (f) Check that  $U$  diagonalizes the matrices  $A$  and  $B$ , by computing  $U^\dagger AU$  and  $U^\dagger BU$ . Note that this is an illustration of Theorem 2.6.7.

**Exercise 2.9.** Find expressions for the following matrix elements, where  $X$  and  $P$  are the position and momentum operators for a particle moving in 1 dimension, with eigenkets  $|x\rangle$  and  $|p\rangle$  respectively, with the Dirac normalizations  $\langle x'|x\rangle = \delta(x - x')$  and  $\langle p'|p\rangle = \delta(p - p')$ .

- (a)  $\langle x|X|p\rangle$ ,      (b)  $\langle p|P|x\rangle$ ,      (c)  $\langle x|PX|p\rangle$ ,      (d)  $\langle p|PX|x\rangle$ ,      (e)  $\langle x|P^2X^2|p\rangle$ .

## 3 The core principles of quantum mechanics

### 3.1 The basic postulates

We are now ready to present the six basic postulates that define quantum mechanics. Although they are not postulates in the traditional formal and rigorous mathematical sense, they constitute the key principles that should govern all quantum systems and can be used to make physical predictions. Some sources give versions that differ from the ones presented here, by combining two or all three of Postulates 2, 3, and 4, or by leaving out the controversial Postulate 5, or by including a requirement that the state space of a composite system is always the tensor product of its subsystems. Others include a postulate governing identical particles, which we do not include for reasons discussed in Chapter 18.

**Postulate 1: States.** The state of a quantum system is specified by a non-null ket  $|\psi\rangle$ , a vector in a Hilbert space (a complex linear vector space with an inner product). For a physical state, the norm  $\langle\psi|\psi\rangle$  is real and positive and finite, but is not otherwise physically significant; two state kets  $|\psi\rangle$  and  $c|\psi\rangle$  are physically equivalent if  $c$  is any non-zero complex number.

**Postulate 2: Observables.** Physically measurable quantities, called observables, correspond to Hermitian operators whose eigenstates can be chosen to be a complete orthonormal basis (orthobasis) of the state space. This means that for an observable  $A$  there is a basis  $|\alpha, u_\alpha\rangle$  satisfying  $A|\alpha, u_\alpha\rangle = \alpha|\alpha, u_\alpha\rangle$ , where  $\alpha$  are the eigenvalues, and  $u_\alpha$  are the degeneracy labels for each  $\alpha$ , with

$$\langle\alpha', u'_{\alpha'}|\alpha, u_\alpha\rangle = \delta_{\alpha\alpha'} \delta_{u_\alpha u'_{\alpha'}}, \quad (3.1.1)$$

and the completeness relation

$$I = \sum_{\alpha} \sum_{u_\alpha} |\alpha, u_\alpha\rangle \langle\alpha, u_\alpha|. \quad (3.1.2)$$

The preceding assumes that  $\alpha$  and  $u_\alpha$  have discrete values. If instead  $\alpha$  has continuous values, then Dirac orthonormality and completeness are used:  $\delta_{\alpha\alpha'}$  is replaced by  $\delta(\alpha - \alpha')$  and  $\sum_{\alpha}$  is replaced by a definite integral  $\int d\alpha$  over the range of allowed  $\alpha$ . Similarly, if  $u_\alpha$  is continuous, then  $\delta_{u_\alpha u'_{\alpha'}}$  is replaced by  $\delta(u_\alpha - u'_{\alpha'})$ , and  $\sum_{u_\alpha}$  is replaced by a definite integral  $\int du_\alpha$ . In some cases  $\alpha$  and/or  $u_\alpha$  can have both some discrete and some continuous values, which are then summed and integrated over the possible values.

**Postulate 3: Allowed results of measurements.** The result of the measurement of an observable  $A$  is always one of its eigenvalues,  $\alpha$ . This rule is sensible and consistent because the eigenvalues of a Hermitian operator are always real, and do not depend on the arbitrary choice of orthobasis used to calculate them, as we observed following Theorem 2.6.9.

**Postulate 4: Probabilities of results of measurements.** If a system is in a state given by a ket  $|\psi\rangle$  which has been normalized so that  $\langle\psi|\psi\rangle = 1$ , and an observable  $A$  is measured, then the probability of getting a particular discrete result  $\alpha$  is

$$\mathcal{P}(\alpha) = \sum_{u_\alpha} |\langle\alpha, u_\alpha|\psi\rangle|^2 = \langle\psi|P_\alpha|\psi\rangle, \quad (3.1.3)$$

where  $|\alpha, u_\alpha\rangle$  are normalized as in eq. (3.1.1), and

$$P_\alpha = \sum_{u_\alpha} |\alpha, u_\alpha\rangle\langle\alpha, u_\alpha| \quad (3.1.4)$$

is the projection operator to the subspace of states with eigenvalue  $\alpha$  for  $A$ . If instead  $\alpha$  is continuous, then the probability of getting a result between  $\alpha$  and  $\alpha + d\alpha$  is

$$d\mathcal{P}(\alpha) = d\alpha \sum_{u_\alpha} |\langle\alpha, u_\alpha|\psi\rangle|^2 = d\alpha \langle\psi|P_\alpha|\psi\rangle. \quad (3.1.5)$$

This postulate generalizes the Born rule for the probabilistic interpretation of the position wavefunction, eq. (2.8.67), and the corresponding rule for momentum, eq. (2.8.68). Note that eq. (3.1.3) ensures that the probabilities are always positive,

$$\mathcal{P}(\alpha) > 0, \quad (3.1.6)$$

and the completeness relation (3.1.2) ensures that the sum of the probabilities for all possible outcomes for a measurement must be unity,

$$\sum_{\alpha} \mathcal{P}(\alpha) = 1. \quad (3.1.7)$$

Thus, these important consistency requirements demanded by any sensible theory of probability are built into the postulates of quantum mechanics.

**Postulate 5: Collapse of the state due to measurement.** If a system is in a state  $|\psi\rangle$  and an observable  $A$  is measured and found to have the value  $\alpha$ , then the state immediately after the measurement will be an eigenstate of  $A$  with eigenvalue  $\alpha$ , specifically,

$$P_\alpha |\psi\rangle, \quad (3.1.8)$$

where  $P_\alpha$  is the projection operator given by eq. (3.1.4). This ket has a norm that is typically less than 1; in fact, from eq. (3.1.3) and the projection operator rules  $P_\alpha^\dagger = P_\alpha$  and  $P_\alpha^2 = P_\alpha$ , we see that the squared norm is simply equal to the probability  $\mathcal{P}(\alpha)$ . Therefore, one can divide the ket (3.1.8) by its norm, to obtain the normalized state after the measurement,

$$\frac{P_\alpha |\psi\rangle}{\sqrt{\langle\psi|P_\alpha|\psi\rangle}} = \frac{P_\alpha |\psi\rangle}{\sqrt{\mathcal{P}(\alpha)}}. \quad (3.1.9)$$

There is a nice shortcut in the special case that the eigenvalue  $\alpha$  is not degenerate, because then this post-measurement state is simply the corresponding eigenstate  $|\alpha\rangle$ , up to an unobservable phase. More generally, it is always a linear combination of the states  $|\alpha, u_\alpha\rangle$ .

**Postulate 6: Time evolution of the quantum state.** Between measurements described by Postulate 5, the time dependence of a state obeys the Schrödinger differential equation,

$$i\hbar \frac{d}{dt} |\psi(t)\rangle = H |\psi(t)\rangle, \quad (3.1.10)$$

where  $H$  is a Hermitian observable operator, called the **Hamiltonian**, whose eigenvalues are the allowed energies of the system.

## 3.2 Valid and invalid questions

Scientific advances often reveal not just how to answer certain difficult questions, but that other seemingly sensible questions do not need to be answered or even considered. From the theory of special relativity, we learn that it makes no sense to ask questions about collisions of particles with relative speed larger than the speed of light in vacuum  $c$ ; these are not valid questions because the very structure of the theory implies that such collisions do not occur. One also learns not to ask questions concerning spatially separated events that are supposedly simultaneous in two different reference frames that are moving with respect to each other, because this also is not meaningful in special relativity. Even though our experience with nonrelativistic systems might make such questions seem legitimate, they are in fact invalid.

Similarly, in quantum mechanics, there are questions that have no good answer because they are not valid to begin with. A prominent example is “what are the position and momentum of this particle at time  $t$ ?”. In classical mechanics this question makes perfect sense, and we learn to calculate the answer given some initial conditions and the equations of motion. However, in quantum mechanics, even in the most idealized case, we can only ask for the probability that the measurement of an observable has a specific result from among the allowed list of eigenvalues. In any particular measurement, that observable could be the position of a particle, or it could be the momentum, but it cannot be both.

To see why, suppose we attempt to define a clever new multi-component operator

$$\Omega = (\vec{R}, \vec{P}), \quad (3.2.1)$$

which is the ordered pair whose components are the position and momentum vectors of a particle. If one could measure  $\Omega$ , the result would be the answer to the simultaneous position and momentum of a particle. Since  $\vec{R}$  and  $\vec{P}$  are each Hermitian,  $\Omega$  may indeed be defined as a Hermitian operator. However, it is not an observable, because it fails the part of the definition

that calls for the eigenstates to form an orthobasis that satisfies the completeness relation. In fact,  $\Omega$  has no eigenstates at all; this is directly related to the fact that  $\vec{R}$  and  $\vec{P}$  do not commute. Notice that if they had a vanishing commutator, then Theorem 2.7.1 says that they would have had a whole orthobasis of simultaneous eigenstates, and so would  $\Omega$ , which could therefore have been an acceptable observable. But, as things are,  $\Omega$  is not an observable, and the postulates of quantum mechanics do not allow for its measurement.

More generally, we can ask whether a set of observables can be measured simultaneously (as a multi-component operator), and whether it makes sense to ask what the results of such a measurement are. This depends on the commutation relations of the observable operators.

First, suppose that the operators  $A, B, C, \dots$  all commute with each other (are compatible). Then the **simultaneous measurement** can be performed, and can be defined as a sequence of consecutive measurements, performed immediately after one another so that there is no time for the system to evolve between the measurements. It is left to Exercise 3.1 to show, using Postulates 4 and 5, that for compatible observables the final results for the probabilities of the different outcomes  $(\alpha, \beta, \gamma, \dots)$  and for the corresponding final state do not depend on which order one performs the measurements, as long as they are all performed with no intervening time delay, so that Postulate 6 does not come into play.

Next, suppose that two of the operators are incompatible, with a commutator  $[A, B]$  that is an operator with no vanishing eigenvalues. This includes the case that  $[A, B]$  is a non-zero multiple of the identity operator, notably if  $A$  and  $B$  are a position operator and the corresponding momentum. In this case, Postulate 5 tells us that after measuring  $B$  the system will be left in some state  $|\beta\rangle$  that is an eigenstate of  $B$ , but it is definitely not an eigenstate of  $A$ . (Otherwise, it would be an eigenstate of  $[A, B]$  with eigenvalue 0, which we are assuming does not exist.) Similarly, if  $A$  is measured, the system will be left in a state that is definitely not an eigenstate of  $B$ . Therefore, the order of making the measurements certainly makes a difference, and one cannot define their simultaneous measurement.

A third possibility is that  $[A, B]$  is an operator that has some vanishing but also some non-vanishing eigenvalues. In this case, one might measure  $A$ , and find a result that leaves the state in an eigenstate of  $B$ , but this will not always happen. The same is true if  $B$  is measured first. An evaluation using Postulates 4 and 5 will be necessary on a case-by-case basis to decide what the outcomes are that might leave  $A$  and  $B$  simultaneously determined in the final state.

Consider a classical observable, for example  $f(a, b, c, \dots)$  where  $a, b, c, \dots$  are quantities that have quantum observable counterparts  $A, B, C, \dots$ . Then, there is always at least one quantum operator  $F(A, B, C, \dots)$  which is also an observable. However, one must be careful in defining it if  $A, B, C, \dots$  do not all commute, due to quantum ordering ambiguities. For example, if  $f(x, p) =$

$xp$  where  $x$  and  $p$  are the position and momentum of a particle moving in one dimension, we could try quantum operator versions  $F(X, P) = XP$  or  $F(X, P) = PX$  or  $F(X, P) = (XP + PX)/2$ . The first two of these are not Hermitian. However, the last is Hermitian, and is an observable. More generally, by completely symmetrizing each term of  $F(A, B, C, \dots)$ , we can always systematically construct a Hermitian version of it, as can easily be proved using eq. (2.4.23).

For the converse, there can be quantum observables that have no classical counterpart; the quintessential example of this is spin, or intrinsic angular momentum. The magnitude of the spin of a particle is a fixed multiple of  $\hbar/2$ . In particular, unlike ordinary angular momentum, there are no states in which it can take on classical values arbitrarily large compared to  $\hbar$ .

A perhaps unexpected example of a valid question in quantum mechanics is: “Given a system in a state  $|\psi\rangle$ , what is the probability of finding it in another state  $|\chi\rangle$ ?” (Such a question has a valid counterpart in classical physics, but there it is somewhat trivial since all information about the state of a classical system is, in principle, more directly accessible.) The observable corresponding to this question is the projection operator

$$P_\chi = |\chi\rangle\langle\chi|. \quad (3.2.2)$$

It is Hermitian, and has eigenvalues 1 (with eigenstate  $|\chi\rangle$ ) and 0 (with eigenstates consisting of all states orthogonal to  $|\chi\rangle$ ). The result of the measurement of  $P_\chi$  will therefore always be either 1 (“yes, we are in the state  $|\chi\rangle$ ”) or 0 (“no, we are not in the state  $|\chi\rangle$ ”), even though the state  $|\psi\rangle$  before the measurement need not have had either of these definite attributes. The probability of obtaining the result 1 is  $|\langle\chi|\psi\rangle|^2$ , assuming both kets are normalized. If we do measure  $P_\chi$  and obtain the result 1, then the state after the measurement will be  $P_\chi|\psi\rangle$ , which is simply the same as  $|\chi\rangle$ , up to normalization. If we obtain the result 0, then the state after the measurement will be  $(I - P_\chi)|\psi\rangle$ , which is orthogonal to  $|\chi\rangle$ . A crucial feature of quantum mechanics, as embodied in the collapse Postulate 5, is that making the measurement changes the state; it is not the same after the measurement as it was before, unless the system was already in an eigenstate of  $P_\chi$ .

One can also construct an observable that generalizes eq. (3.2.2) in a natural way to a weighted sum of projection operators for any orthobasis  $\{|\varphi_n\rangle\}$  with  $n = 1, \dots, d$ , where  $d$  is the dimension of the Hilbert space. Such an observable is associated with the valid question “Given a state  $|\psi\rangle$ , what are the probabilities of finding it in each of these orthobasis states?” For example, consider the Hermitian operator

$$Q = \sum_{n=1}^d n P_n, \quad (3.2.3)$$




Figure 3.2.1: A meter for the  $Q$  observable of eq. (3.2.3), for a system with  $d = 6$ , showing the result after measurement of  $Q$  has resolved the quantum system with initial state  $|\psi\rangle$  into the orthobasis state  $|\varphi_3\rangle$  from among the possibilities  $|\varphi_n\rangle$  with  $n = 1, 2, 3, 4, 5, 6$ . The probability of this outcome is  $|\langle\varphi_3|\psi\rangle|^2$ .

where  $P_n = |\varphi_n\rangle\langle\varphi_n|$  are the orthobasis state projection operators. The eigenstates of  $Q$  are the orthobasis states  $|\varphi_n\rangle$ , and the corresponding eigenvalues are the coefficients of the orthogonal projection operators, namely the integers  $n = 1, \dots, d$ . Therefore, the possible outcomes of measuring  $Q$  are  $n = 1, \dots, d$ , as visualized in Figure 3.2.1. The probabilities for these outcomes are  $\mathcal{P}(n) = |\langle\varphi_n|\psi\rangle|^2$ , assuming that the ket  $|\psi\rangle$  has norm 1. After the measurement outcome  $n$ , the system will be left in the state  $|\varphi_n\rangle$ . Thus, measuring the observable  $Q$  changes the state by resolving it into one of the given orthobasis states, and the outcome  $n$  tells you which one.

### 3.3 Expectation values and uncertainties

In addition to the quantities directly associated to single measurements of an observable, there are statistical quantities that result from making many measurements. Consider an idealized situation in which we have access to an arbitrarily large number  $N$  of copies of a system in the same state  $|\psi\rangle$ . This mythical group of identical and independent quantum systems is called a **pure ensemble**. The **expectation value** of an operator  $A$  in the state  $|\psi\rangle$  is defined to be the average value obtained by measuring  $A$  in these independent experiments, as  $N \rightarrow \infty$ . According to the frequentist interpretation of probabilities, this is the same as the sum of the possible outcomes for each experiment multiplied by their respective probabilities, which can be evaluated using Postulate 4 as

$$\sum_{\alpha} \alpha \mathcal{P}(\alpha) = \sum_{\alpha} \sum_{u_{\alpha}} \alpha \langle\psi|\alpha, u_{\alpha}\rangle\langle\alpha, u_{\alpha}|\psi\rangle = \sum_{\alpha} \sum_{u_{\alpha}} \langle\psi|A|\alpha, u_{\alpha}\rangle\langle\alpha, u_{\alpha}|\psi\rangle = \langle\psi|A|\psi\rangle. \quad (3.3.1)$$

Here, the first equality used eq. (3.1.3) and assumed that  $|\psi\rangle$  is normalized to unity, the second equality used the fact that  $|\alpha, u_{\alpha}\rangle$  are eigenstates of  $A$  with eigenvalue  $\alpha$ , and the completeness relation was used to get the last equality. In cases where the state  $|\psi\rangle$  is understood by context, it is customary to denote the expectation value by

$$\langle A \rangle \equiv \langle\psi|A|\psi\rangle, \quad (3.3.2)$$

still assuming that  $|\psi\rangle$  is normalized to unity. If that is not convenient for some reason, one has the more general relation

$$\langle A \rangle = \frac{\langle\psi|A|\psi\rangle}{\langle\psi|\psi\rangle}. \quad (3.3.3)$$



The expectation value of an observable is easily shown to be a real number.

Another quantity defined through measurements on a large pure ensemble of identical states  $|\psi\rangle$  is the **uncertainty** of an observable  $A$ , which we will denote by  $\Delta A$ . Here again the notation assumes that  $|\psi\rangle$  is understood by context. It is defined by

$$(\Delta A)^2 = \langle (A - \langle A \rangle)^2 \rangle = \langle \psi | (A - \langle \psi | A | \psi \rangle)^2 | \psi \rangle. \quad (3.3.4)$$

Intuitively, the uncertainty tells us how much the measurement of the observable is expected to fluctuate about its expectation value, over the course of many independent measurements, each performed starting in the same state  $|\psi\rangle$ . It is the same as the concept of standard deviation in statistics. By expanding eq. (3.3.4), one obtains the equivalent form that is most commonly used in practical calculations,

$$(\Delta A)^2 = \langle \psi | A^2 | \psi \rangle - (\langle \psi | A | \psi \rangle)^2, \quad (3.3.5)$$

which again assumes  $\langle \psi | \psi \rangle = 1$ .

There is a fundamental obstacle to the existence of states with arbitrarily small uncertainties for incompatible observables, imposed by the following result due to Howard P. Robertson and Schrödinger:

**Theorem 3.3.1. (Uncertainty relation)** *In any state  $|\psi\rangle$ , the uncertainties of two observables  $A$  and  $B$  obey*

$$(\Delta A)(\Delta B) \geq \frac{1}{2} |\langle [A, B] \rangle|. \quad (3.3.6)$$

Proof: Define observables  $\tilde{A} = A - \langle A \rangle$  and  $\tilde{B} = B - \langle B \rangle$ . These are Hermitian, because  $\langle A \rangle$  and  $\langle B \rangle$  are real numbers. It follows from the definition of uncertainty that

$$(\Delta A)^2 (\Delta B)^2 = \langle \psi | \tilde{A}^2 | \psi \rangle \langle \psi | \tilde{B}^2 | \psi \rangle = \langle \tilde{A} \psi | \tilde{A} \psi \rangle \langle \tilde{B} \psi | \tilde{B} \psi \rangle \geq |\langle \tilde{A} \psi | \tilde{B} \psi \rangle|^2, \quad (3.3.7)$$

where the Cauchy–Schwarz inequality eq. (2.2.4) was used at the end. Therefore, we have

$$(\Delta A)^2 (\Delta B)^2 \geq |\langle \psi | \tilde{A} \tilde{B} | \psi \rangle|^2 = \left| \frac{1}{2} \langle \psi | [\tilde{A}, \tilde{B}] | \psi \rangle + \frac{1}{2} \langle \psi | \{\tilde{A}, \tilde{B}\} | \psi \rangle \right|^2. \quad (3.3.8)$$

Using the Hermiticity of  $\tilde{A}$  and  $\tilde{B}$  yet again,  $\langle \psi | [\tilde{A}, \tilde{B}] | \psi \rangle$  is a pure imaginary number and  $\langle \psi | \{\tilde{A}, \tilde{B}\} | \psi \rangle$  is a pure real number. Therefore, the squared magnitude of the sum is equal to the sum of the square magnitudes, so

$$(\Delta A)^2 (\Delta B)^2 \geq \frac{1}{4} |\langle \psi | [\tilde{A}, \tilde{B}] | \psi \rangle|^2 + \frac{1}{4} |\langle \psi | \{\tilde{A}, \tilde{B}\} | \psi \rangle|^2. \quad (3.3.9)$$

Now, because the last term is certainly non-negative, we can drop it without affecting the validity of the inequality. Furthermore,  $[\tilde{A}, \tilde{B}]$  is just equal to  $[A, B]$ . So, eq. (3.3.6) follows from taking the square root of both sides of eq. (3.3.9).  $\square$

A famous special case is obtained by taking  $A = X$  and  $B = P$  for a particle moving in one dimension. Because  $[X, P] = i\hbar$ , the uncertainties must satisfy the **Heisenberg position-momentum uncertainty relation**, named after Werner Heisenberg (who originally proposed a weaker version of it) and derived rigorously first by Earle H. Kennard and shortly after by Hermann Weyl,

$$(\Delta X)(\Delta P) \geq \hbar/2, \quad (3.3.10)$$

in any state. It might seem that an even stronger version might be possible, since we simply discarded the non-negative last term in eq. (3.3.9). However, for the case of position and momentum, we will show later, in section 6.1, that there do exist states (those with Gaussian wavefunctions), in which eq. (3.3.10) is saturated, in other words equality holds. The same wavefunctions will reappear in section 7.4. So, eq. (3.3.10) is the strongest possible general version of the position-momentum uncertainty relation.

For a particle moving in three dimensions, one finds in the same way that each of  $(\Delta X)(\Delta P_x)$  and  $(\Delta Y)(\Delta P_y)$  and  $(\Delta Z)(\Delta P_z)$  cannot be less than  $\hbar/2$ . However, since  $X$  and  $P_y$  commute, there is no uncertainty relation for the product  $(\Delta X)(\Delta P_y)$ . This means that, in principle, one could simultaneously specify the exact values of a particle's coordinate along some direction and the momentum component in an orthogonal direction.

### 3.4 How states change

According to Postulate 6, the time evolution of a quantum state obeys a linear first-order differential equation, the Schrödinger equation. Let the initial condition for the state at time  $t = t_0$  be  $|\psi(t_0)\rangle$ . We then define the **time-evolution operator**  $U(t, t_0)$  such that the state ket at time  $t$  is

$$|\psi(t)\rangle = U(t, t_0) |\psi(t_0)\rangle. \quad (3.4.1)$$

It follows from this definition, and Schrödinger's eq. (3.1.10), that the operator  $U(t, t_0)$  satisfies

$$i\hbar \frac{d}{dt} U(t, t_0) = H U(t, t_0). \quad (3.4.2)$$

Our goal is to solve this differential equation for  $U(t, t_0)$  as a function of  $t$ , subject to the boundary condition  $U(t_0, t_0) = I$ . As we will see, it is a unitary operator if  $H(t)$  is Hermitian.

First, consider the case that the Hamiltonian does not have any dependence on  $t$ . In that case, the solution is simply

$$U(t - t_0) \equiv U(t, t_0) = e^{-i(t-t_0)H/\hbar}, \quad (3.4.3)$$

which only depends on the time difference  $t - t_0$ , not the individual times. To check this claim, note that it clearly satisfies the initial condition at  $t = t_0$ , and that

$$\begin{aligned} i\hbar \frac{d}{dt} |\psi(t)\rangle &= i\hbar \frac{d}{dt} [e^{-i(t-t_0)H/\hbar} |\psi(t_0)\rangle] = i\hbar [(-iH/\hbar)e^{-i(t-t_0)H/\hbar}] |\psi(t_0)\rangle \\ &= H |\psi(t)\rangle \end{aligned} \quad (3.4.4)$$

recovers the Schrödinger equation, as required. It is crucial in the preceding derivation that the operator  $H$  does not depend on time, so that it can be treated just like a number in the exponential, as it obviously commutes with itself.

The unitarity of  $U(t - t_0)$  in eq. (3.4.3) is simple to prove, given that  $H$  is Hermitian. From the rules for taking adjoints,

$$U(t - t_0)^\dagger = [e^{-i(t-t_0)H/\hbar}]^\dagger = e^{i(t-t_0)H^\dagger/\hbar} = e^{i(t-t_0)H/\hbar} = U(t - t_0)^{-1} = U(t_0 - t). \quad (3.4.5)$$

The last equality is a bonus, which shows that evolving a state backward in time is the inverse operation of evolving it forward in time by the same amount, as one might expect. Since  $U(t - t_0)$  is a unitary operator, time evolution can be regarded as equivalent to a change of orthobasis.

While eq. (3.4.3) is a neat formal solution of the Schrödinger equation, in practice it leaves more to do, because the exponential of an operator as an infinite series can be non-trivial to evaluate in matrix elements. To make further progress, we can apply the spectral decomposition trick of eq. (2.7.5) to the operator  $U(t - t_0)$  as given by eq. (3.4.3). To accomplish this, first consider the eigenvalue problem for the Hamiltonian, which we are still assuming is Hermitian and does not depend on time. This eigenvalue equation is called the **time-independent Schrödinger equation**, and is written as

$$H |\psi_E\rangle = E |\psi_E\rangle, \quad (3.4.6)$$

where  $|\psi_E\rangle$  does not depend on  $t$ . Suppose that this equation has been solved completely for all energy eigenvalues  $E$  and all corresponding orthobasis eigenstates  $|\psi_E\rangle = |E, u_E\rangle$ , where  $u_E$  is a degeneracy label. Then, using completeness of the energy orthobasis,

$$U(t - t_0) = \sum_E \sum_{u_E} e^{-i(t-t_0)E/\hbar} |E, u_E\rangle \langle E, u_E|, \quad (3.4.7)$$

where we have turned the operator  $H$  into the number  $E$  when acting on each of its eigenstates. Applying this to eq. (3.4.1) gives

$$|\psi(t)\rangle = \sum_E \sum_{u_E} e^{-i(t-t_0)E/\hbar} |E, u_E\rangle \langle E, u_E | \psi(t_0)\rangle, \quad (3.4.8)$$

the spectral decomposition of the state with respect to energy.

Finding solutions of the time-independent Schrödinger equation is one of the main problems of quantum mechanics. This often amounts to solving a matrix eigenvalue equation and/or a differential equation for the wavefunction. As an example of the latter, suppose that the Hamiltonian is that of a particle moving in a potential  $V$  in one dimension, with

$$H = \frac{1}{2m} P^2 + V(X), \quad (3.4.9)$$

where  $P$  and  $X$  are the momentum and position operators. Taking the inner product of eq. (3.4.6) with the position eigenstate  $|x\rangle$ , and using the toolbox of Table 2.8.1 so that  $X \rightarrow x$  and  $P \rightarrow -i\hbar d/dx$  and  $|\psi_E\rangle \rightarrow \psi_E(x)$ , we obtain

$$\left( -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + V(x) - E \right) \psi_E(x) = 0 \quad (3.4.10)$$

in the position representation. Similarly, for a spinless particle moving in a potential  $V(\vec{r})$  in three dimensions, the time-independent Schrödinger differential equation is

$$\left( -\frac{\hbar^2}{2m} \nabla^2 + V(\vec{r}) - E \right) \psi_E(\vec{r}) = 0, \quad (3.4.11)$$

to be solved simultaneously for  $E$  and  $\psi_E(\vec{r})$ . The cases of multiple particles, and particles coupled to an electromagnetic field, will be discussed in sections 4.2 and 4.3.

Note that the unitary time-evolution operator in eq. (3.4.7) is nontrivial only because the phases are different for the eigenstates with different energies. As a special case, acting on an initial state ket  $|\psi_E(t_0)\rangle$  that happens to be an eigenstate of energy, time evolution just multiplies by a **global phase** (that is, a single phase that multiplies the whole state ket), namely  $e^{-i(t-t_0)E/\hbar}$ . Because such a global phase is not physically significant, the state has not really changed. For this reason, a Hamiltonian eigenstate is also known as a **stationary state**.

To illustrate this, consider the time evolution of a stationary state, starting from  $t = 0$ ,

$$|\psi_E(t)\rangle = e^{-itE/\hbar} |\psi_E(0)\rangle, \quad (3.4.12)$$

and suppose that at time  $t$  we measure some observable  $A$ , which is assumed to have no explicit time dependence built into it.<sup>†</sup> Recall from Postulate 4 that the probability of getting a particular

---

<sup>†</sup>By “explicit time dependence”, we just mean an explicit appearance of  $t$  in the definition of the operator. For example, the position operator  $X$  has no explicit time dependence, but the operator  $A = X \sin(\omega t)$  does depend explicitly on time, which we express as  $\partial A / \partial t = \omega X \cos(\omega t)$ .

measurement result  $\alpha$  is

$$\begin{aligned}\mathcal{P}(\alpha, t) &= \sum_{u_\alpha} |\langle \alpha, u_\alpha | \psi_E(t) \rangle|^2 = \sum_{u_\alpha} |e^{-itE/\hbar} \langle \alpha, u_\alpha | \psi_E(0) \rangle|^2 = \sum_{u_\alpha} |\langle \alpha, u_\alpha | \psi_E(0) \rangle|^2 \\ &= \mathcal{P}(\alpha, 0).\end{aligned}\tag{3.4.13}$$

In a stationary state, the probability of measuring each particular possible result  $\alpha$  stays the same for all time, as long as the time evolution is not interrupted by a measurement.

Returning to the case of a general state  $|\psi(t)\rangle$  and a general observable  $A$ , the time dependence of the expectation value obeys

$$\frac{d}{dt} \langle \psi | A | \psi \rangle = \langle \psi | A \left( \frac{d}{dt} | \psi \rangle \right) + \left( \frac{d}{dt} \langle \psi | \right) A | \psi \rangle + \langle \psi | \frac{\partial A}{\partial t} | \psi \rangle, \tag{3.4.14}$$

where  $\partial A / \partial t$  is the derivative of the explicit time dependence of  $A$ . Applying Schrödinger's equation and its Hermitian conjugate to the first two terms on the right, respectively, gives

$$\frac{d}{dt} \langle \psi | A | \psi \rangle = -\frac{i}{\hbar} \langle \psi | [A, H] | \psi \rangle + \langle \psi | \frac{\partial A}{\partial t} | \psi \rangle, \tag{3.4.15}$$

or, in the more compact notation of expectation values, just

$$\frac{d}{dt} \langle A \rangle = -\frac{i}{\hbar} \langle [A, H] \rangle + \left\langle \frac{\partial A}{\partial t} \right\rangle. \tag{3.4.16}$$

This general result is known as **Ehrenfest's Theorem**, after Paul Ehrenfest.

In the special case that  $|\psi\rangle$  is a stationary state,  $H|\psi\rangle = E|\psi\rangle$  and  $\langle \psi | H = E \langle \psi |$ , so  $\langle [A, H] \rangle$  evaluates to  $(E - E) \langle A \rangle = 0$ . In that case,  $\frac{d}{dt} \langle A \rangle = \langle \frac{\partial A}{\partial t} \rangle$ . The change in an expectation value in a stationary state is due only to the explicit time dependence of the observable operator.

Suppose that the Hamiltonian is that of a particle moving in a constant potential in one dimension, eq. (3.4.9), and consider  $A = P$ , the momentum operator. The operator  $P$  has no explicit time dependence, so  $\partial P / \partial t = 0$ . Also,  $[P, H] = [P, V(X)] = -i\hbar dV/dX$ . Therefore, the expectation values in a general state obey

$$\frac{d}{dt} \langle P \rangle = -\left\langle \frac{dV}{dX} \right\rangle, \tag{3.4.17}$$

which is the quantum mechanical version of Newton's second law. Similarly, for  $A = X$ , using  $\partial X / \partial t = 0$  and  $[X, H] = [X, P^2]/2m = i\hbar P/m$ , we get

$$\frac{d}{dt} \langle X \rangle = \langle P \rangle / m. \tag{3.4.18}$$

This generalizes in a straightforward way to a particle moving in three dimensions. Ehrenfest's Theorem says that the momentum and position expectation values in quantum mechanics obey

the same equations of motion as the corresponding quantities in classical mechanics. Equations (3.4.17) and (3.4.18) therefore confirm the choices of sign and magnitude in our definition of the momentum operator  $P$  in eq. (2.8.15).

As another application of Ehrenfest's Theorem, apply eq. (3.4.16) to the time-independent observable  $A = \frac{1}{2}(XP + PX)$ . Evaluating the commutator with the same Hamiltonian, one finds that for expectation values in a general time-dependent state, the **Virial Theorem** holds:

$$\frac{d}{dt} \langle A \rangle = \frac{1}{m} \langle P^2 \rangle - \langle XV'(X) \rangle. \quad (3.4.19)$$

For the special case of a stationary state, since we proved that all expectation values are independent of time, the left side vanishes and the Virial Theorem becomes

$$\frac{1}{m} \langle P^2 \rangle = \langle XV'(X) \rangle \quad (\text{in a stationary state}). \quad (3.4.20)$$

Specializing further to the case of a power-law potential  $V(X) = kX^n$ , this becomes

$$2 \langle T \rangle = n \langle V \rangle, \quad (\text{in a stationary state, if } V \propto X^n), \quad (3.4.21)$$

where  $T = P^2/2m$  is the kinetic energy operator. The generalization of the Virial Theorem to a particle moving in three dimensions is straightforward, and left to Exercise 3.6.

We now turn to the more difficult case that the Hamiltonian operator depends on time. First, consider time evolution over an infinitesimal interval from time  $t_0$  to time  $t_0 + \Delta t$ . Then, from the Schrödinger equation,

$$|\psi(t_0 + \Delta t)\rangle = \left[ 1 - \frac{i}{\hbar} \Delta t H(t_0) \right] |\psi(t_0)\rangle. \quad (3.4.22)$$

Note that because  $\Delta t$  is infinitesimal, it does not matter here whether we use  $H(t_0)$ , or  $H(t)$ , or  $H$  evaluated at some intermediate time, because the difference will be higher order in  $\Delta t$ . Up to terms of order  $(\Delta t)^2$ , we can rewrite this as an exponential,

$$|\psi(t_0 + \Delta t)\rangle = \exp \left[ -\frac{i}{\hbar} \Delta t H(t_0) \right] |\psi(t_0)\rangle. \quad (3.4.23)$$

An advantage of writing it this way is that the exponential is a unitary operator if  $H(t_0)$  is Hermitian, so that  $|\psi(t_0 + \Delta t)\rangle$  has the same norm as  $|\psi(t_0)\rangle$ . Now, if we evolve the state further from time  $t_0 + \Delta$  to  $t_0 + 2\Delta t$  in the same way, we have

$$|\psi(t_0 + 2\Delta t)\rangle = \exp \left[ -\frac{i}{\hbar} \Delta t H(t_0 + \Delta t) \right] \exp \left[ -\frac{i}{\hbar} \Delta t H(t_0) \right] |\psi(t_0)\rangle. \quad (3.4.24)$$

Here, the exponentials cannot easily be combined into a single exponential, because  $H(t_0 + \Delta t)$  and  $H(t_0)$  are different operators, and need not commute. Continuing in this way,

$$|\psi(t_0 + N\Delta t)\rangle = \left( \prod_{n=0}^{N-1} \exp \left[ -\frac{i}{\hbar} \Delta t H(t_0 + n\Delta t) \right] \right) |\psi(t_0)\rangle, \quad (3.4.25)$$

where it is important that the terms in the product, are understood to be arranged from higher to lower  $n$ , reading from left to right. Since each of the terms in the product is a unitary operator, the whole product is a unitary operator as well. Now we can take

$$N = (t - t_0)/\Delta t \rightarrow \infty, \quad (3.4.26)$$

to obtain

$$|\psi(t)\rangle = U(t, t_0) |\psi(t_0)\rangle, \quad (3.4.27)$$

where the unitary time-evolution operator is

$$U(t, t_0) = \lim_{N \rightarrow \infty} \prod_{n=0}^{N-1} \exp \left[ -\frac{i}{\hbar} \Delta t H(t_0 + n\Delta t) \right]. \quad (3.4.28)$$

From its construction, it satisfies

$$[U(t_2, t_1)]^\dagger = [U(t_2, t_1)]^{-1} = U(t_1, t_2), \quad (3.4.29)$$

$$U(t_3, t_2)U(t_2, t_1) = U(t_3, t_1). \quad (3.4.30)$$

This time-evolution operator can depend on both arguments separately; in general  $U(t, t_0) \neq U(t - t_0)$  if the Hamiltonian depends on time.

Formally, one can also write for eq. (3.4.28)

$$U(t, t_0) = T \exp \left[ -\frac{i}{\hbar} \int_{t_0}^t dt' H(t') \right], \quad (3.4.31)$$

where the  $T$  is a time-ordering symbol, an instruction to rearrange the Hamiltonians in the expansion of the exponential so that each  $H(t_j)$  appears to the right of  $H(t_k)$  whenever  $t_j < t_k$ . More explicitly, it takes the form of a **Dyson series**, named after Freeman J. Dyson,

$$U(t, t_0) = I + \sum_{N=1}^{\infty} \left( -\frac{i}{\hbar} \right)^N \int_{t_0}^t dt_N \int_{t_0}^{t_N} dt_{N-1} \cdots \int_{t_0}^{t_2} dt_1 H(t_N) H(t_{N-1}) \cdots H(t_1), \quad (3.4.32)$$

where each  $t_k$  integration has lower limit  $t_0$  and upper limit  $t_{k+1}$ , for  $k = 1, \dots, N$ , with  $t_{N+1}$  interpreted as  $t$ . Note that the nested upper limits of integration have neatly removed the need for the  $1/N!$  usually present in the series expansion of the exponential function. You can also check directly that eq. (3.4.32) satisfies the differential equation (3.4.2), by plugging it in.

In eq. (3.4.32) the ordering of the Hamiltonians is important because in general they do not commute at different times. In the special case that they do all commute, one can write

$$U(t, t_0) = \exp \left[ -\frac{i}{\hbar} \int_{t_0}^t dt' H(t') \right], \quad (3.4.33)$$

which agrees with the result eq. (3.4.3) when  $H$  does not depend on time at all. Another Dyson series, based on the interaction picture of quantum mechanics and useful for time-dependent perturbation theory, will be discussed in section 20.2.

We have seen that there are two very different ways that a state can change in quantum mechanics: Schrödinger time evolution governed by the Hamiltonian, and collapse of the state ket due to measurement. It is natural to ask whether the latter might be a special case of the former. At least within the standard formulation of quantum mechanics, as given by the postulates listed in section 3.1, the answer is “No!”. Hamiltonian time evolution is accomplished by multiplying the state by a *unitary* operator, while collapse of the state due to measurement is associated with multiplying by a projection operator, which is instead *Hermitian*. Thus, there is a fundamental difference between Hamiltonian time evolution and measurement collapse.

The time evolution due to the Hamiltonian is perfectly causal and deterministic; the state at a given time is uniquely determined by the state at earlier times, just as in classical physics, provided that a measurement of the type described in Postulates 4 and 5 has not taken place in the interim. In contrast, the collapse of the wavefunction associated with measurement is inherently probabilistic, rather than deterministic. This means that, unlike in classical physics, it is not possible, even in principle, to predict the future. But the situation is actually worse than that: we cannot even predict the past. Given complete knowledge of the present state of a quantum system, the state before the most recent measurement cannot be known, because Postulate 5 says that the act of measurement changes the state in a way that destroys information, in an irreversible way. If you measured an observable  $A$  and got a result  $\alpha$ , leaving the system in a state  $|\alpha\rangle$ , then this tells you that the state of the system  $|\psi\rangle$  before the measurement must have had a non-zero matrix element  $\langle\alpha|\psi\rangle$ , but that is all. This is clearly very incomplete information.

The insistence on a fundamental distinction between the measuring agent and the quantum system being measured, as required by Postulate 5, was developed and promoted by Niels Bohr and collaborators, and is often referred to as the **Copenhagen interpretation** of quantum mechanics. According to the Copenhagen interpretation, the measuring agent apparently can be thought of in classical terms, or at least we do not ask questions about its quantum behavior. This seems troubling, as one can always imagine treating any particular measuring apparatus as itself a quantum system undergoing unitary time evolution. For this reason, many people have advocated modifying the postulates of quantum mechanics to modify or completely eliminate Postulate 5 dealing with the collapse of the state due to measurement.

In particular, Hugh Everett, in his 1957 PhD thesis, proposed that the state ket always undergoes unitary time evolution, so that collapse of the state due to measurement is an illusion



of limited human powers of discernment. In this **Everett interpretation** of quantum mechanics, all possible alternatives for every measurement outcome are actually realized, through superposition, in some parts of the great state ket of the universe. This can be imagined as a continuous branching of possibilities that has inspired the alternative name **many-worlds interpretation**, and many science fiction story ideas. It is an observed fact that parts of the Everett superposition corresponding to non-classical-like outcomes for macroscopic systems seem to have negligible amplitudes. This can be explained by a phenomenon called decoherence, discussed in Chapter 25, which involves entanglement correlations between subsystems.

It is difficult not to be sympathetic to the no-collapse view. Indeed, we could draw a big sphere of radius several hundred light years around the Earth, and think of the contents (including us, all other observers who might be able to communicate with us, and all of our measuring devices) as one big quantum system evolving strictly according to unitary time evolution. In any case, there seems to be no reasonable scientific principle that could tell us exactly where we should put the boundary separating the quantum system from the measuring apparatus that supposedly undergoes collapse.

However, from a practical point of view, Postulate 5 is indispensable, because it provides a straightforward, consistent, and reliable way of making predictions for the actual experiments that we do in the real world. No matter how philosophically compelling it might be to discard the measurement collapse of the wavefunction, it is not scientifically necessary, with the exception of some interesting and ambitious proposals to treat the quantum dynamics of the universe as a whole. So far, the postulates of quantum mechanics as given in section 3.1 have stood the test of time, successfully providing accurate predictions of every experimental phenomenon with which they have been confronted.

### 3.5 Mixed ensembles and the density matrix operator

The expectation value and uncertainty for an operator were defined for a single quantum state in section 3.3, using the concept of a pure ensemble. However, it is often more realistic to suppose that in a large ensemble of quantum systems of the same type, some fraction of them  $p_1$  will be in a state  $|\psi_1\rangle$ , a fraction  $p_2$  will be in a different state  $|\psi_2\rangle$ , etc. Such a large collection of systems of the same type, but in different states, is called a **mixed ensemble**. If we choose one of the systems at random from a mixed ensemble, there is a probability  $p_I$  that it will be in the state  $|\psi_I\rangle$ , with  $p_I \geq 0$  for each  $I$ , and

$$\sum_I p_I = 1. \quad (3.5.1)$$

The kets  $|\psi_I\rangle$  are assumed to have unit norm in the following, but there is no reason why they must be linearly independent or orthogonal to each other. There is not even any constraint on the number of distinct states  $|\psi_I\rangle$  that are found in the ensemble, and it could be larger than the dimension of the state space. The mixed ensemble generalizes the concept of a pure ensemble, for which one of the  $p_I$  is equal to 1 and all others are 0.

Suppose we choose one system of the mixed ensemble at random, and measure an observable  $A$  with eigenvalues  $\alpha$  and orthonormal eigenstates  $|\alpha, u_\alpha\rangle$ . Then, the probability of getting a particular result  $\alpha$  is equal to the sum over  $|\psi_I\rangle$  of the product of the probability of choosing a system in that state and the probability that a measurement in that state will give  $\alpha$ . Applying Postulate 4 to evaluate the latter probability,

$$\mathcal{P}(\alpha)_{\text{mixed}} = \sum_I p_I \sum_{u_\alpha} |\langle \alpha, u_\alpha | \psi_I \rangle|^2. \quad (3.5.2)$$

We can similarly compute the average result obtained by measuring the observable  $A$  many times on systems chosen at random from the mixed ensemble,

$$\overline{A} = \sum_I p_I \langle \psi_I | A | \psi_I \rangle. \quad (3.5.3)$$

We use an overline notation to denote this **mixed ensemble average**, to distinguish it from the expectation value associated with measuring  $A$  in a single state in a pure ensemble.

There are two very different types of probabilities at work in eqs. (3.5.2) and (3.5.3). First, there are the **ensemble probabilities**  $p_I$ , which simply reflect the fact that the mixed ensemble is populated by different states. These ensemble probabilities would exist even if our systems were classical. Second, we have the probabilities associated with the inherently non-deterministic nature of measurement in quantum systems, which are manifested in  $\sum_{u_\alpha} |\langle \alpha, u_\alpha | \psi_I \rangle|^2$  and in the expectation value  $\langle \psi_I | A | \psi_I \rangle$ . The results for  $\mathcal{P}(\alpha)_{\text{mixed}}$  and  $\overline{A}$  incorporate both types of probabilities.

One should not confuse the concepts of a pure ensemble based on a *superposition* of quantum states and a mixed ensemble containing a *population* of the same quantum states. A simple example will illustrate the distinction. Consider a state space with two orthobasis kets  $|1\rangle$  and  $|2\rangle$ . Suppose that initially we have a pure ensemble, with all systems in the superposition state

$$|\psi\rangle = c_1|1\rangle + c_2|2\rangle, \quad (3.5.4)$$

where  $c_1$  and  $c_2$  are complex numbers subject to the normalization condition  $|c_1|^2 + |c_2|^2 = 1$ . Now we can conduct a measurement to ask if a system is in the state  $|1\rangle$ . The probability of finding the result 1 (yes) is  $|c_1|^2$ , and the probability of finding the result 0 (no) is  $|c_2|^2$ . If we do this measurement on each and every system in the pure ensemble, we will afterwards have a

mixed ensemble, with  $p_1 = |c_1|^2$  for the state  $|1\rangle$ , and  $p_2 = |c_2|^2$  for the state  $|2\rangle$ . This illustrates that one way to prepare a mixed ensemble is to conduct measurements on the members of a pure ensemble.

Continuing with this example, if we now make the same measurement again on the mixed ensemble, we will get the same results as for the pure ensemble; the probability is still  $|c_1|^2$  to find the state  $|1\rangle$ . But now consider instead the probability that measurement of some other observable  $A$  will yield the result  $\alpha$ , with corresponding eigenket  $|\alpha\rangle$ . For the pure ensemble with state  $|\psi\rangle$ , this is

$$\mathcal{P}(\alpha)_{\text{pure}} = |\langle\alpha|\psi\rangle|^2 = |c_1\langle\alpha|1\rangle + c_2\langle\alpha|2\rangle|^2, \quad (3.5.5)$$

but for the mixed ensemble, we find instead, from eq. (3.5.2),

$$\mathcal{P}(\alpha)_{\text{mixed}} = |c_1|^2|\langle\alpha|1\rangle|^2 + |c_2|^2|\langle\alpha|2\rangle|^2. \quad (3.5.6)$$

The key difference is that in the pure ensemble result there are interference terms that do not appear in the mixed ensemble result. There is relative phase information present in the pure ensemble that is absent in the mixed ensemble.

A standard realization of this same example is found in the double-slit diffraction experiment. Suppose that we have a source of particles that can impact on a plane detection screen, after having passed through one of two very narrow slits in a diffraction screen, as shown in Figure 1.3.2. The source of particles is said to be **coherent** if we can describe the state of a given particle as a superposition like eq. (3.5.4), where  $|1\rangle$  represents the state in which the particle travels through slit 1, and  $|2\rangle$  represents the state in which it travels through slit 2. If we let the operator  $A$  in the above discussion be the position operator  $X$  on the detection screen perpendicular to the slit directions, then the probability to find the particle between positions  $x$  and  $x + dx$  is given by the continuous version of eq. (3.5.5),

$$d\mathcal{P}(x) = |c_1\psi_1(x) + c_2\psi_2(x)|^2 dx. \quad (3.5.7)$$

The individual wavefunctions for the different slit contributions interfere destructively or constructively, depending on  $x$ , to produce an interference pattern as shown in Figure 1.3.2. But, now suppose that there are detectors that measure whether the particle goes through slit 1 or slit 2. This measurement results in a mixed ensemble of particles at the detection screen, and we will have instead the continuous version of eq. (3.5.6),

$$d\mathcal{P}(x) = \left(|c_1|^2|\psi_1(x)|^2 + |c_2|^2|\psi_2(x)|^2\right)dx, \quad (3.5.8)$$

in which the interference terms (also known as coherences) are absent. The act of measuring which slit each particle goes through destroys the interference pattern. This “which-slit” measurement might effectively be done by the environment in which the experiment takes place,

rather than an official detector monitored by a certified experimentalist. This reduction of a pure ensemble of coherent superpositions to a mixed ensemble of populations is an example of decoherence, which will be discussed further in Chapter 25.

All of the physical information about a mixed ensemble is encoded in an elegant way in the **density matrix operator**, or just the **density operator**, due to John von Neumann. It is defined in terms of the ensemble states and probabilities by

$$\rho = \sum_I p_I |\psi_I\rangle \langle \psi_I|. \quad (3.5.9)$$

In the special case of a pure ensemble,  $\rho$  is simply the projection operator for the state  $|\psi\rangle$ . More generally, it is a sum of projection operators weighted by the frequencies of occurrence of states within the ensemble. In terms of the density operator, the result of eq. (3.5.2) for the probability of getting the result  $\alpha$  for a single measurement of  $A$  can be written as

$$\mathcal{P}(\alpha)_{\text{ensemble}} = \sum_{u_\alpha} \langle \alpha, u_\alpha | \rho | \alpha, u_\alpha \rangle. \quad (3.5.10)$$

We can rewrite this by choosing an arbitrary orthobasis of kets  $\{|\phi_k\rangle\}$ , and then using completeness followed by a rearrangement,

$$\mathcal{P}(\alpha)_{\text{ensemble}} = \sum_{u_\alpha} \langle \alpha, u_\alpha | \left( \sum_{k=1}^d |\phi_k\rangle \langle \phi_k| \right) \rho | \alpha, u_\alpha \rangle = \sum_{k=1}^d \langle \phi_k | \rho P_\alpha | \phi_k \rangle = \text{Tr}[\rho P_\alpha], \quad (3.5.11)$$

where  $P_\alpha$  is the projection operator for the result  $\alpha$ , defined in eq. (3.1.4), and at the end we used the definition of the trace of an operator in eq. (2.6.57). Recall that the trace is independent of the choice of orthobasis.

The average of the results of many measurements of  $A$  in a mixed ensemble, as computed in eq. (3.5.3), can also be re-expressed in terms of the density operator. Again using completeness followed by a rearrangement,

$$\overline{A} = \sum_I p_I \langle \psi_I | A \left( \sum_{k=1}^d |\phi_k\rangle \langle \phi_k| \right) | \psi_I \rangle = \sum_{k=1}^d \langle \phi_k | \rho A | \phi_k \rangle = \text{Tr}[\rho A]. \quad (3.5.12)$$

As a special case,

$$\text{Tr}[\rho] = 1, \quad (3.5.13)$$

which simply re-expresses the conservation of probability from eq. (3.5.1). It is also straightforward to show  $\text{Tr}[\rho^2] \leq 1$ , with equality only in the special case that the ensemble is a pure one, in which case one also has  $\rho^2 = \rho$ .

From its definition in eq. (3.5.9),  $\rho$  is clearly a Hermitian operator. So, according to Theorem 2.6.6, one can find a special orthobasis consisting of its eigenkets, call them  $|\varphi_k\rangle$ , with eigenvalues  $p_k$  that are real because of Theorem 2.6.3. Therefore, no matter what states  $|\psi_I\rangle$  were involved in the original preparation of the mixed ensemble, or how many such states there were, we can always rewrite the density operator as

$$\rho = \sum_{k=1}^d p_k |\varphi_k\rangle\langle\varphi_k|, \quad (3.5.14)$$

which is very similar in appearance to eq. (3.5.9), but with the important difference that the index  $k$  now takes on a limited number of values up to the dimension  $d$  of the state space. The  $p_k$  can be interpreted as the ensemble probabilities for the orthobasis. This illustrates the more general fact that the density operator is not tied to any specific set of states  $|\psi_I\rangle$ , even if we used one when preparing the mixed ensemble. The density operator can also be used to summarize our (incomplete) information about a single system chosen at random from the mixed ensemble, called a **mixed state**.

A pure ensemble is one extreme special case of a mixed ensemble, in which all systems are in the same state. The opposite extreme is the **completely random ensemble**, which we can define by choosing *any* orthobasis  $|\varphi_k\rangle$ , and writing

$$\rho = \frac{1}{d} \sum_{k=1}^d |\varphi_k\rangle\langle\varphi_k|. \quad (3.5.15)$$

Here  $1/d$  is the ensemble probability for each of the orthobasis states. This density operator is proportional to the identity operator, so it is actually independent of the choice of orthobasis, and is the unique one associated with maximum randomness of states in the ensemble.

The extent to which an ensemble of quantum systems is randomized can more generally be quantified by the **von Neumann entropy**,<sup>†</sup>

$$\sigma = -\text{Tr}[\rho \ln \rho] = -\sum_{k=1}^d p_k \ln p_k, \quad (3.5.16)$$

where the final result is in terms of the orthobasis ensemble probabilities in eq. (3.5.14), which are defined to be the eigenvalues of  $\rho$ . When  $p_k = 0$ , one should interpret  $p_k \ln p_k$  as 0. The von Neumann entropy is the quantum mechanical analog of the **Shannon entropy** introduced (two decades later!) by Claude E. Shannon in the study of classical information and communication theory, but the Shannon entropy has different properties when combining classical subsystems.

---

<sup>†</sup>Many sources replace the natural logarithm in eq. (3.5.16) by the logarithm base 2, which is especially convenient for quantum information applications where the systems of interest have 2-state orthobases. This simply changes the normalization of the entropy by a multiplicative factor  $1/\ln(2)$ , since  $\log_2(x) = \ln(x)/\ln(2)$ .

It also has the same form, up to a multiplicative factor, as the **Gibbs entropy** defined by Josiah Willard Gibbs in classical statistical mechanics and thermodynamics,

$$S = -k_B \sum_{k=1}^d p_k \ln p_k, \quad (3.5.17)$$

where  $k_B$  is Boltzmann's constant, and in this case the  $p_k$  are interpreted as the probabilities for microstates  $|\varphi_k\rangle$  to occur in the fluctuations of a system.

In our two extreme cases,

$$\sigma = 0 \quad (\text{pure ensemble}), \quad (3.5.18)$$

$$\sigma = -n \left[ \frac{1}{n} \ln(1/n) \right] = \ln(n) \quad (\text{completely random ensemble}), \quad (3.5.19)$$

where  $n$  is the number of orthobasis states available to the systems in the ensemble, usually the same as the dimension  $d$  of the state space. In the case of a mixed state, the entropy is a measure of our ignorance, and it is always between 0 and  $\ln(n)$ . The result  $S = k_B \ln(n)$  obtained for the special case of the completely random ensemble is the **Boltzmann entropy**.

In general, the density operator for a mixed state or a mixed ensemble depends on time. From the general form in eq. (3.5.9), we have

$$\frac{d\rho}{dt} = \sum_I p_I \left[ \left( \frac{d}{dt} |\psi_I\rangle \right) \langle \psi_I| + |\psi_I\rangle \left( \frac{d}{dt} \langle \psi_I| \right) \right], \quad (3.5.20)$$

and evaluating the time derivatives using the Schrödinger equation, we obtain

$$\frac{d\rho}{dt} = -\frac{i}{\hbar} [H, \rho]. \quad (3.5.21)$$

Note that this vanishes in the special case of a completely random ensemble; random ensembles stay random. It also vanishes in the case of a pure ensemble if the state is an energy eigenstate, but not if it is a superposition of states with different energies. The historical reason for the name “density operator” is that this equation is analogous to Liouville's Theorem in classical mechanics, which says that for an ensemble of classical systems, the phase-space density  $\rho_{\text{classical}}$  (the number of classical ensemble members per unit position and momentum) obeys

$$\frac{d\rho_{\text{classical}}}{dt} = \{H, \rho_{\text{classical}}\}_{\text{PB}}, \quad (3.5.22)$$

which has the classical Poisson bracket (and removal of the factor of  $-i/\hbar$ ) on the right-hand side replacing the commutator. This is an example of the classical–quantum correspondence principle to be discussed further in section 4.1.

Equation (3.5.21) gives the instantaneous change in the density operator as the ensemble of systems undergoes time evolution. More generally, in terms of the unitary time evolution operator that we defined in eq. (3.4.1),

$$\rho(t) = U(t, t_0) \rho(t_0) U(t, t_0)^\dagger, \quad (3.5.23)$$

with a short proof left as an exercise. Since  $\rho(t)$  and  $\rho(t_0)$  are related by a unitary transformation, they have the same eigenvalues, according to Theorem 2.6.9. It follows that under time evolution, a pure state stays pure, and a mixed state remains mixed. Furthermore, from the expression for the entropy in eq. (3.5.16) in terms of the eigenvalues  $p_k$ ,

$$\sigma(t) = \sigma(t_0) \quad (\text{unitary time evolution}), \quad (3.5.24)$$

in the absence of external measurements or other influences. So, we have found that the entropy of a closed system does not change with time.

Let us now work out what the density operator will be after a measurement on a mixed state. Suppose we have a density operator  $\rho$  as given in eq. (3.5.14) in terms of orthobasis states  $|\varphi_k\rangle$  with ensemble probabilities  $p_k$ , and we make a measurement on a randomly chosen ensemble state of an observable  $A$  and obtain the result  $\alpha$ . For each of the ensemble states  $|\varphi_k\rangle$ , Postulate 5 tells us that the state after the measurement will be

$$\frac{P_\alpha |\varphi_k\rangle}{\sqrt{\langle \varphi_k | P_\alpha | \varphi_k \rangle}}, \quad (3.5.25)$$

where  $P_\alpha$  is the projection operator for the result  $\alpha$ . Therefore, we can write the post-measurement density operator as

$$\rho_\alpha = \sum_{k=1}^d \mathcal{P}(k|\alpha) \left( \frac{P_\alpha |\varphi_k\rangle}{\sqrt{\langle \varphi_k | P_\alpha | \varphi_k \rangle}} \right) \left( \frac{\langle \varphi_k | P_\alpha}{\sqrt{\langle \varphi_k | P_\alpha | \varphi_k \rangle}} \right), \quad (3.5.26)$$

where  $\mathcal{P}(k|\alpha)$  is the conditional probability that the state selected from the ensemble was  $|\varphi_k\rangle$ , given that the result  $\alpha$  was obtained for  $A$ . To evaluate this, we use **Bayes' Theorem**, the fundamental result in the theory of conditional probabilities, which says

$$\mathcal{P}(k|\alpha) = \frac{\mathcal{P}(\alpha|k) p_k}{\mathcal{P}(\alpha)}. \quad (3.5.27)$$

Here, as given by Postulate 4,

$$\mathcal{P}(\alpha|k) = \langle \varphi_k | P_\alpha | \varphi_k \rangle \quad (3.5.28)$$

is the conditional probability that the result of a measurement of  $A$  is  $\alpha$ , assuming that the state was  $|\varphi_k\rangle$ , while  $\mathcal{P}(\alpha) = \text{Tr}[\rho P_\alpha]$  is the probability, obtained in eq. (3.5.11), that a measurement

of  $A$  in the ensemble resulted in  $\alpha$ . Putting these results together, eq. (3.5.26) for the density operator after the measurement becomes

$$\rho_\alpha = \frac{P_\alpha \rho P_\alpha}{\mathcal{P}(\alpha)} = \frac{P_\alpha \rho P_\alpha}{\text{Tr}[\rho P_\alpha]}. \quad (3.5.29)$$

This is the density operator version of Postulate 5. Note that it obeys  $\rho_\alpha^\dagger = \rho_\alpha$  and  $\text{Tr}[\rho_\alpha] = 1$ , so that the post-measurement density operator is still Hermitian and has trace 1.

We have now succeeded in finding the rules for measurement probabilities and averages, post-measurement collapse, and time evolution for a mixed state in terms of the density matrix operator  $\rho$ , in eqs. (3.5.11), (3.5.12), (3.5.21), (3.5.23), and (3.5.29). Since each of these results only depends on  $\rho$ , and not on the individual ensemble states or probabilities, we have justified the assertion that all of the physical properties of the mixed ensemble or mixed state are contained in the density operator.

A common situation is that a measurement of  $A$  has been made, but we do not know the result. Perhaps we were not looking, or we lost the data, or the measurement was made by something or someone else who is unwilling or unable to communicate with us. In that case, the density operator describing the mixed state after the measurement will be

$$\rho' = \sum_\alpha \mathcal{P}(\alpha) \rho_\alpha = \sum_\alpha P_\alpha \rho P_\alpha. \quad (3.5.30)$$

We will now show that the entropy always increases when this occurs, except in the trivial special case  $\rho' = \rho$ , which happens only if all states in the ensemble were eigenstates of  $A$  with the same eigenvalue. To do so, we will use the following fact from linear algebra about traces of functions of matrices, due to Oskar Klein, the proof of which is omitted.

**Theorem 3.5.1. (Klein's inequality, general)** *Suppose that the function  $f(x)$  is differentiable and strictly convex ( $f''(x) > 0$ ) for all  $0 < x < \infty$ , and that  $A$  and  $B$  are Hermitian matrices with non-negative eigenvalues. Then*

$$\text{Tr}[f(A) - f(B) + (B - A)f'(B)] \geq 0, \quad (3.5.31)$$

*with equality if and only if  $A = B$ .*

We now apply this to the case  $f(x) = x \ln x$ , and let  $A = \rho$  and  $B = \rho'$  be any two density operators; we are not yet assuming the special form of eq. (3.5.30). Then, using  $\text{Tr}[\rho] = \text{Tr}[\rho'] = 1$ , we obtain:

**Theorem 3.5.2. (Klein's inequality for density operators)** *Suppose that  $\rho$  and  $\rho'$  are any two density operators on a common state space. Then*

$$\text{Tr}[\rho(\ln \rho - \ln \rho')] \geq 0, \quad (3.5.32)$$

*with equality if and only if  $\rho = \rho'$ .*



Now consider the case of  $\rho'$  given by eq. (3.5.30), which arose from having made a measurement of  $A$  on a mixed state described by  $\rho$ , with unknown result. The entropy after the measurement is

$$\sigma' = -\text{Tr}[\rho' \ln \rho'] = -\sum_{\alpha} \text{Tr}[P_{\alpha} \rho P_{\alpha} \ln \rho'] = -\sum_{\alpha} \text{Tr}[\rho P_{\alpha} \ln \rho' P_{\alpha}], \quad (3.5.33)$$

where the cyclic property of the trace was used at the end. Since  $P_{\alpha}^2 = 1$ , eq. (3.5.30) shows that  $P_{\alpha}$  commutes with  $\rho'$ , which implies that  $P_{\alpha}$  also commutes with  $\ln \rho'$ . Therefore,  $P_{\alpha} \ln \rho' P_{\alpha} = P_{\alpha}^2 \ln \rho' = P_{\alpha} \ln \rho'$ . Then, using completeness in the form  $\sum_{\alpha} P_{\alpha} = I$ , we obtain

$$\sigma' = -\text{Tr}[\rho \ln \rho']. \quad (3.5.34)$$

Using  $\sigma = -\text{Tr}[\rho \ln \rho]$  and Klein's inequality (3.5.32), we finally obtain the claimed result,

$$\sigma' \geq \sigma. \quad (3.5.35)$$

The entropy increases whenever a non-trivial measurement is made but the result is unknown.

Note that this increase of the von Neumann entropy does not apply to a situation in which we made a measurement on a single mixed state and the result is known. In fact, if the known result of the measurement  $\alpha$  is a non-degenerate eigenvalue, then the resulting density operator describing the system will be that of a pure state, with vanishing entropy.

As an important example, consider an ensemble consisting of a bottle of, say,  $\sim 10^{24}$  gas molecules, each of which can be in states characterized by energy eigenvalues  $E$  and degeneracy labels  $u_E$ . The molecules interact with each other and with the bottle walls, but weakly enough that they can be considered an ensemble of independent quantum states of the same type. Intuitively, each interaction can be thought of as a sort of external measurement on the gas molecule, but the results of these measurements remain unknown, so that eq. (3.5.35) applies so as to make the entropy as large as it can be, subject to the constraint of energy conservation. Thus, when the molecules reach thermal equilibrium, the ensemble probability to find one of them in a particular orthobasis state  $|E, u_E\rangle$  can be determined by the statistical principle that the entropy should be maximized, subject to the constraint, due to energy conservation, that the ensemble average energy has a fixed value  $\overline{E}$ .

To see the implications of this, we write the density operator in the form

$$\rho = \sum_E \sum_{u_E} p_E |E, u_E\rangle \langle E, u_E|. \quad (3.5.36)$$

Here, we have already implemented the idea that maximizing the entropy will require the density operator to correspond to complete randomization within each subspace of fixed energy eigenvalue  $E$ , but the relative probabilities  $p_E$  for each energy level remain to be found. Equation

(3.5.36) gives

$$\sigma = - \sum_E g_E p_E \ln p_E, \quad (3.5.37)$$

where  $g_E$  is the degeneracy of the energy eigenvalue  $E$ . Now, to maximize  $\sigma$  subject to the constraints  $1 = \text{Tr}[\rho] = \sum_E g_E p_E$  and fixed  $\bar{E} = \sum_E g_E p_E E$ , it is simplest to use the method of Lagrange multipliers. The function to be extremized is

$$f(p_E, \alpha, \beta) = - \sum_E g_E p_E \ln p_E + \alpha \left(1 - \sum_E g_E p_E\right) + \beta \left(\bar{E} - \sum_E g_E p_E E\right), \quad (3.5.38)$$

where  $\alpha$  and  $\beta$  are the Lagrange multipliers for the trace constraint and the energy constraint, respectively, and  $\bar{E}$  is fixed. We then obtain, for each  $E$ ,

$$0 = \frac{\partial f}{\partial p_E} = -g_E (\ln p_E + 1) - \alpha g_E - \beta g_E E, \quad (3.5.39)$$

which has the solution

$$p_E = e^{-(\beta E + \alpha + 1)}. \quad (3.5.40)$$

The  $e^{-(\alpha+1)}$  factor is independent of  $E$ , and so can be absorbed into a common normalization factor; the important point is that we have derived that the canonical ensemble probabilities that maximize the entropy must be proportional to the **Boltzmann factor**,

$$p_E \propto e^{-\beta E}. \quad (3.5.41)$$

The Lagrange multiplier  $\beta$  is related to temperature by the definition

$$\beta = \frac{1}{k_B T}. \quad (3.5.42)$$

Since  $\beta$  has units of 1/energy, this definition shows that Boltzmann's constant is really just a conversion factor between energy and temperature. If we agreed to measure temperature in units of energy, then Boltzmann's constant would be 1.

The result of eq. (3.5.36) with  $p_E \propto e^{-\beta E}$  is called the **canonical ensemble**, with density operator

$$\rho = \frac{1}{Z} \sum_E \sum_{u_E} e^{-\beta E} |E, u_E\rangle \langle E, u_E|, \quad (3.5.43)$$

where the normalization factor  $Z$  is called the **partition function**. Thus, the canonical ensemble is completely randomized at each fixed energy level  $E$ , but with relative probabilities

between energy levels determined by the Boltzmann factor. The partition function can be computed using the requirement  $\text{Tr}[\rho] = 1$ , which gives

$$Z = \sum_E \sum_{u_E} e^{-\beta E} = \sum_E g_E e^{-\beta E}. \quad (3.5.44)$$

Equation (3.5.43) can be recognized as the spectral decomposition form [see eq. (2.7.5)] of

$$\rho = \frac{1}{Z} e^{-\beta H}. \quad (3.5.45)$$

This obviously commutes with  $H$ , so according to eq. (3.5.21),  $\rho$  is constant in time. For any observable  $A$  defined for each molecule, the canonical ensemble average is, from eq. (3.5.12),

$$\bar{A} = \frac{1}{Z} \text{Tr} [e^{-\beta H} A] = \frac{1}{Z} \sum_E \sum_{u_E} e^{-\beta E} \langle E, u_E | A | E, u_E \rangle. \quad (3.5.46)$$

In particular, the average energy for states in the canonical ensemble is

$$\bar{E} = \frac{1}{Z} \sum_E g_E E e^{-\beta E} = -\frac{\partial}{\partial \beta} \ln Z. \quad (3.5.47)$$

It is left as an exercise to check that, with the entropy definition  $S = k_B \sigma$ ,

$$\bar{E} - TS = -\beta \ln Z = F, \quad (3.5.48)$$

where  $F$  is called the **Helmholtz free energy** after Hermann von Helmholtz. Let us stop our discussion of the canonical ensemble here, before this book accidentally turns into one on statistical mechanics.

In all of the preceding, we have assumed for notational simplicity that the states  $|\psi_I\rangle$  appearing in the ensemble are discrete and countable. As usual, one can also consider a continuum of states, which entails turning summations into integrals. If the ensemble states  $|\psi_u\rangle$  are labeled by some continuous parameter  $u$  (instead of the discrete label  $I$ ), then the density operator is

$$\rho = \int du p(u) |\psi_u\rangle \langle \psi_u|, \quad (3.5.49)$$

where the probability density  $p(u)$  must satisfy the constraint  $\int du p(u) = 1$ .

## 3.6 Exercises

**Exercise 3.1.** Consider compatible observables  $A$  and  $B$ , whose (possibly degenerate) eigenvalues include  $\alpha$  and  $\beta$ , respectively. Suppose that they are both measured, with a negligible time delay so that Schrödinger time evolution does not come into play. Show that the probability of obtaining  $(\alpha, \beta)$  for  $(A, B)$  does not depend on the order in which they are measured.

**Exercise 3.2.** Show that if the position-space wave function  $\psi(\vec{r}) = \langle \vec{r} | \psi \rangle$  is real, then the expectation value of the momentum operator  $\vec{P}$  must vanish:  $\langle \vec{P} \rangle = 0$ . This helps to explain why the states for quantum mechanics must form a *complex* vector space.

**Exercise 3.3.** Consider a state  $|\psi\rangle$  with position wave function  $\langle \vec{r} | \psi \rangle = \psi(\vec{r})$  and momentum expectation value  $\langle \psi | \vec{P} | \psi \rangle = \vec{p}$ . Show that the state with position-space wave function  $e^{i\vec{k} \cdot \vec{r}} \psi(\vec{r})$  must have momentum expectation value  $\vec{p} + \hbar \vec{k}$ .

**Exercise 3.4.** Consider observables  $A$  and  $B$  with the following matrix representations on a state space with dimension 3,

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad B = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & -i & 0 \\ i & 0 & -i \\ 0 & i & 0 \end{pmatrix}. \quad (3.6.1)$$

These matrices are representations of the operators in an orthonormal basis  $|1\rangle, |0\rangle, |-1\rangle$ , which are labeled by the non-degenerate eigenvalues of  $A$ , which we can therefore call “the  $A$  basis”.

- Find the eigenvalues and the corresponding normalized eigenkets of  $B$  (in the  $A$  basis).
- In the state  $|-1\rangle$ , calculate the expectation value  $\langle B \rangle$  and the uncertainty  $\Delta B$ .
- If the particle is in the state with  $A = 1$ , and  $B$  is measured, what are the possible outcomes and their probabilities?
- If the particle is in the state with  $B = 0$ , and  $A$  is measured, what are the possible outcomes and their probabilities?
- Consider the state  $|\psi\rangle = \frac{1}{\sqrt{6}} |1\rangle + \frac{1}{\sqrt{6}} |0\rangle + \frac{2}{\sqrt{6}} |-1\rangle$ . If the operator  $A^2$  is measured and a result  $+1$  is obtained, what is the normalized state ket immediately after the measurement? What was the probability of this result? If  $A$  is then immediately measured, what are the possible outcomes and their respective probabilities?

**Exercise 3.5.** Consider a particle free to move throughout all space in 3-d. Suppose that its wavefunction in spherical coordinates is  $\psi(\vec{r}) = Ce^{-r/a}$ , where  $C$  and  $a$  are constants.

- If the wavefunction is normalized to unity, what is the magnitude of the constant  $C$ ?
- What is the probability that the particle will be found to be farther from the origin than  $a$ ?
- What are the expectation value and uncertainty of the radial coordinate operator  $R$ ?
- What are the expectation value and uncertainty of the momentum squared operator  $P^2$ ?

**Exercise 3.6.** Consider a particle moving in three dimensions, governed by the Hamiltonian

$$H = \frac{1}{2m} P^2 + V(\vec{R}), \quad (3.6.2)$$

where  $P^2 = \vec{P} \cdot \vec{P} = P_x^2 + P_y^2 + P_z^2$ . Apply Ehrenfest’s Theorem eq. (3.4.16) to the observable  $A = \frac{1}{2}(\vec{R} \cdot \vec{P} + \vec{P} \cdot \vec{R})$  to obtain the **3-d version of the Virial Theorem**. You should find that

for a general state,

$$\frac{d}{dt} \langle A \rangle = \frac{1}{m} \langle P^2 \rangle - \langle \vec{R} \cdot \vec{\nabla} V \rangle, \quad (3.6.3)$$

so that for a stationary state,

$$\frac{1}{m} \langle P^2 \rangle = \langle \vec{R} \cdot \vec{\nabla} V \rangle, \quad (\text{in a stationary state}), \quad (3.6.4)$$

and for the special case of a spherically symmetric power-law potential

$$2 \langle T \rangle = n \langle V \rangle, \quad (\text{in a stationary state, if } V \propto R^n), \quad (3.6.5)$$

where  $T = P^2/2m$  is the kinetic energy operator.

**Exercise 3.7.** Consider a particle of mass  $m$  moving in a 1-d potential, with Hamiltonian  $H = P^2/2m + V(X)$ . Suppose that the eigenstates and energy eigenvalues of  $H$  are denoted by  $|\psi_n\rangle$  and  $E_n$ , where  $n$  is a discrete label, so that  $H|\psi_n\rangle = E_n|\psi_n\rangle$ .

(a) Show that  $\langle \psi_n | P | \psi_k \rangle = \alpha \langle \psi_n | X | \psi_k \rangle$ , where  $\alpha$  is a quantity that you will determine, which depends on the difference between  $E_n$  and  $E_k$ . (Hint: consider the commutator  $[X, H]$ .)

(b) From the result of the previous part, show that

$$\langle \psi_n | P^2 | \psi_n \rangle = \beta \sum_k (E_n - E_k)^2 |\langle \psi_n | X | \psi_k \rangle|^2, \quad (3.6.6)$$

where  $\beta$  is a constant quantity that you will find. (Hint: completeness is your friend.)

(c) How does this rule generalize to cases with some continuous energy eigenvalues?

(d) Derive the corresponding results for a particle moving in three dimensions.

**Exercise 3.8.** Prove eq.(3.5.23) for the unitary time evolution of the density operator.

**Exercise 3.9.** Consider a quantum system with a state space of dimension  $d$ .

(a) Show that to specify a general pure state of the system requires  $2d - 2$  real parameters.

(b) Show that to specify the density matrix for a general mixed state of the system requires  $d^2 - 1$  real parameters.

**Exercise 3.10.** A spin-1/2 system with Hamiltonian  $H = \omega S_z$  has energy eigenstates  $|\uparrow\rangle$  and  $|\downarrow\rangle$  with  $S_z$  eigenvalues  $\hbar/2$  and  $-\hbar/2$  respectively. For the canonical ensemble of a large number of such spins at temperature  $T$ , what are the density operator  $\rho$  and the partition function  $Z$ ? What is the Gibbs entropy? What is the average result for a measurement of  $S_z$ ?

## 4 Canonical variables and the Hamiltonian

### 4.1 Quantum observables from classical dynamics

The postulates of quantum mechanics refer to observable operators with time evolution governed by a Hamiltonian, but do not specifically identify these objects. This is intentional, in order to be general. In many physical situations, one can determine the appropriate Hamiltonian and observables by considering a classical version, but this is not always true. Indeed, one should think of classical physics as an approximate limit obtained from quantum mechanics, and not the reverse. Still, in favorable circumstances the classical-limit properties of a system can be used to infer the basic observable operators, including the Hamiltonian, and their algebraic commutator properties in the quantum theory.

In the Lagrangian formulation of classical mechanics, one starts with some dynamical variables  $q_n$ , often called generalized coordinates, which we will label by an index  $n$ . The Lagrangian is a function of the  $q_n$  and their first time derivatives  $\dot{q}_n = dq_n/dt$ ,

$$L(q_n, \dot{q}_n, t). \quad (4.1.1)$$

The classical equations of motion for the system are then

$$\frac{\partial L}{\partial q_n} = \frac{d}{dt} \frac{\partial L}{\partial \dot{q}_n}, \quad (4.1.2)$$

for each  $n$ . A short calculation, found in any good classical physics textbook, shows that this follows from a variational principle involving the action obtained by integrating the Lagrangian with respect to time. However, we will postpone our own discussion of that until section 28.6. The reason for doing so is that rather than accept the variational principle as a postulate of classical mechanics, we will be able to *derive* it as a consequence of the path integral formulation of quantum mechanics.

The Hamiltonian formulation of classical mechanics recasts the Lagrangian formulation by defining a canonical momentum conjugate to each generalized coordinate,

$$p_n = \frac{\partial L}{\partial \dot{q}_n}. \quad (4.1.3)$$

Now one defines the Hamiltonian as

$$H(q_n, p_n, t) = \sum_n p_n \dot{q}_n - L(q_n, \dot{q}_n, t), \quad (4.1.4)$$

where it is important that the  $\dot{q}_n$  are to be completely eliminated in favor of the  $p_n$  using eq. (4.1.3). Thus  $H$  is a function only of the generalized coordinates and their canonical conjugate momenta, and not their time derivatives. The  $q_n$  and  $p_n$  are collectively called the **phase-space coordinates**. As shown in the same good classical mechanics textbook, the Lagrangian

equations of motion eq. (4.1.2) are equivalent to the Hamiltonian equations of motion,

$$\dot{q}_n = \frac{\partial H}{\partial p_n}, \quad \dot{p}_n = -\frac{\partial H}{\partial q_n}. \quad (4.1.5)$$

The generalized coordinates  $q_n$  need not be the rectangular coordinates of a particle, but rather could be any quantities that fully specify the classical state of the system at a given time. Likewise, the canonical momenta need not coincide with mechanical momenta (equal to the product of mass and velocity for particles).

Consider any quantity  $a(q_n, p_n, t)$ , built out of the generalized coordinates and their conjugate momenta. According to the same good classical physics textbook, the time evolution of  $a$  is

$$\frac{da}{dt} = \{a, H\}_{\text{PB}} + \frac{\partial a}{\partial t}, \quad (4.1.6)$$

where the **Poisson bracket** for any two functions on phase space  $a$  and  $b$  is defined as

$$\{a, b\}_{\text{PB}} \equiv \sum_n \left( \frac{\partial a}{\partial q_n} \frac{\partial b}{\partial p_n} - \frac{\partial b}{\partial q_n} \frac{\partial a}{\partial p_n} \right). \quad (4.1.7)$$

Dirac noted that the Poisson brackets  $\{a, b\}_{\text{PB}}$  of classical mechanics are closely analogous to the commutators  $[A, B]$  for the corresponding observables in the quantum theory. Both are antisymmetric under interchange of the observables, and at least for the position and momentum observables, one has the exact correspondence

$$\overbrace{\{q_n, p_k\}_{\text{PB}} = \delta_{nk}}^{\text{classical}} \longleftrightarrow \overbrace{[Q_n, P_k] = i\hbar\delta_{nk}}^{\text{quantum}}. \quad (4.1.8)$$

Commutators obtained in this way are called **canonical commutation relations**. Furthermore, eq. (4.1.6) has a striking resemblance to Ehrenfest's Theorem in quantum mechanics, which we found in eq. (3.4.16). Indeed, one finds from the latter equation that

$$\frac{d}{dt} \langle Q_n \rangle = \left\langle \frac{\partial H}{\partial P_n} \right\rangle, \quad \frac{d}{dt} \langle P_n \rangle = -\left\langle \frac{\partial H}{\partial Q_n} \right\rangle, \quad (4.1.9)$$

directly analogous to the Hamiltonian equations of motion (4.1.5).

For a single particle of mass  $m$  moving in three dimensions in a potential  $V$ , it is natural to choose the  $Q_n$  to be the usual rectangular coordinate operators  $X = R_x$ ,  $Y = R_y$ , and  $Z = R_z$ . Their conjugate **canonical momenta**  $P_x$ ,  $P_y$ , and  $P_z$ , satisfy the commutation relations already given in eq. (2.8.54),

$$[R_a, P_b] = i\hbar\delta_{ab}, \quad [R_a, R_b] = 0, \quad [P_a, P_b] = 0, \quad (4.1.10)$$

for  $a, b = x, y, z$ . Writing  $P^2 \equiv \vec{P} \cdot \vec{P}$ , the Hamiltonian operator is then

$$H = \frac{P^2}{2m} + V(\vec{R}). \quad (4.1.11)$$

The connection between classical and quantum physics just described is often called the **correspondence principle**, but it has some weaknesses. First, there are some quantum observables that do not have a classical counterpart at all, for example spin, also known as intrinsic angular momentum. Second, there is no guarantee that every generalized coordinate and its momentum will obey a canonical commutation relation. Although it is true for the rectangular coordinates of a particle, or a collection of particles, in other cases one might encounter corrections higher order in  $\hbar$ , or ambiguities in connecting the classical observables to the quantum ones. This is why we preferred to derive the position-momentum commutation relations by the method given in section 2.8. More generally, the most logical (but perhaps not the simplest) way to make the connection is to derive the classical theory as an approximation to the quantum theory, as we will do using the Feynman sum-over-paths approach in section 28.6.

## 4.2 The two-body problem

An important special case is that of two particles that are free except for a potential energy of interaction that depends only on their separation. This occurs, for example, in the hydrogen atom to be treated in Chapter 11, where the two particles are the electron and the (much heavier) proton. Another example is neutron-proton scattering, to be studied in section 23.8, where the masses are almost the same.

In general, the two-body Hamiltonian has the form

$$H = \frac{P_1^2}{2m_1} + \frac{P_2^2}{2m_2} + V(\vec{R}_1 - \vec{R}_2), \quad (4.2.1)$$

where we allow for the possibility that the potential energy depends on the vector displacement (not just its magnitude), and the two particles have masses  $m_1$  and  $m_2$  and canonical position and momentum operators  $(\vec{R}_1, \vec{P}_1)$  and  $(\vec{R}_2, \vec{P}_2)$ . The components of these observables satisfy commutation relations  $[R_{1a}, P_{1b}] = i\hbar\delta_{ab}$  and  $[R_{2a}, P_{2b}] = i\hbar\delta_{ab}$ , for  $a, b = x, y, z$ , with other combinations vanishing. In particular, each of the observables for particle 1 commutes with those of particle 2. As an orthobasis, one can choose the tensor product of the eigenkets of  $\vec{R}_1$  and  $\vec{R}_2$ ,

$$|\vec{r}_1, \vec{r}_2\rangle = |\vec{r}_1\rangle \otimes |\vec{r}_2\rangle, \quad (4.2.2)$$

defined to satisfy the eigenvalue equations

$$\vec{R}_1 |\vec{r}_1, \vec{r}_2\rangle = \vec{r}_1 |\vec{r}_1, \vec{r}_2\rangle, \quad \vec{R}_2 |\vec{r}_1, \vec{r}_2\rangle = \vec{r}_2 |\vec{r}_1, \vec{r}_2\rangle. \quad (4.2.3)$$

However, the solution of the Hamiltonian eigenvalue problem is complicated by the fact that the potential couples the two particle degrees of freedom together.



Fortunately, as in classical mechanics, such cases can be reduced to a simpler problem that is effectively the same as for two decoupled particles, by making a change of variables to separate the relative motion from that of the center of mass. The relative motion is described by

$$\vec{R} = \vec{R}_1 - \vec{R}_2, \quad \vec{P} = \frac{m_2 \vec{P}_1 - m_1 \vec{P}_2}{m_1 + m_2}, \quad (4.2.4)$$

and the motion of the center of mass by

$$\vec{R}_{\text{cm}} = \frac{m_1 \vec{R}_1 + m_2 \vec{R}_2}{m_1 + m_2}, \quad \vec{P}_{\text{tot}} = \vec{P}_1 + \vec{P}_2. \quad (4.2.5)$$

It is a short exercise to show that the pairs  $(\vec{R}, \vec{P})$  and  $(\vec{R}_{\text{cm}}, \vec{P}_{\text{tot}})$  each satisfy canonical commutation relations, and do not interfere with each other. Furthermore, if one defines the total mass  $M$  and the **reduced mass**  $\mu$  according to

$$M = m_1 + m_2, \quad \mu = \frac{m_1 m_2}{m_1 + m_2}, \quad (4.2.6)$$

then the Hamiltonian eq. (4.2.1) can be rewritten in the decoupled form

$$H = H_{\text{cm}} + H_{\text{rel}}, \quad H_{\text{cm}} = \frac{P_{\text{tot}}^2}{2M}, \quad H_{\text{rel}} = \frac{P^2}{2\mu} + V(\vec{R}). \quad (4.2.7)$$

The center-of-mass degrees of freedom have the same Hamiltonian as that of a completely free particle with mass  $M$ , whose eigenvalue problem is easy to solve (plane waves). The dynamics of  $H_{\text{rel}}$  are the same as for a single particle with mass equal to  $\mu$ , moving in the potential  $V(\vec{R})$ . One can now choose a new orthobasis as the tensor product of eigenkets of  $\vec{R}$  and  $\vec{R}_{\text{cm}}$ ,

$$|\vec{r}, \vec{r}_{\text{cm}}\rangle = |\vec{r}\rangle \otimes |\vec{r}_{\text{cm}}\rangle, \quad (4.2.8)$$

and look for stationary-state wavefunction solutions of the form

$$\Psi(\vec{r}, \vec{r}_{\text{cm}}) = \langle \vec{r}, \vec{r}_{\text{cm}} | \Psi \rangle = \frac{1}{(2\pi\hbar)^{3/2}} e^{i\vec{k}_{\text{tot}} \cdot \vec{r}_{\text{cm}}} \psi(\vec{r}), \quad (4.2.9)$$

where  $\vec{p}_{\text{tot}} = \hbar\vec{k}_{\text{tot}}$  is the eigenvalue of  $\vec{P}_{\text{tot}}$ , and the relative coordinate wavefunction satisfies

$$\left( -\frac{\hbar^2 \nabla^2}{2\mu} + V(\vec{r}) - E \right) \psi(\vec{r}) = 0, \quad (4.2.10)$$

and the total energy eigenvalue is  $E + \hbar^2 k_{\text{tot}}^2 / 2M$ . We can then solve the eigenvalue problem in eq. (4.2.10) for  $E$  and  $\psi(\vec{r})$  as if it were a single particle. If one of the particles is much heavier than the other, as in the case of electrons compared to atomic nuclei, then  $\mu$  is equal to the mass of the lighter particle, to a good approximation.

If the number of particles  $N$  is three or more, and fixed, then one can construct appropriate **Jacobi coordinates**, and their canonical momenta, by iteration. First, choose two of the particles and define their relative and center-of-mass coordinates and momenta, just as in the preceding. The two-particle center-of-mass coordinates and momenta are then combined in the same way with those of a third particle, to give another relative coordinate/momentum pair and a three-particle center-of-mass coordinate and momentum. The three-particle center-of-mass coordinate and momentum are combined with those of a fourth particle, etc. In the end one will have a single center-of-mass coordinate and momentum for the whole system with total mass  $M = \sum_{i=1}^N m_i$ ,

$$\vec{R}_{\text{cm}} = \frac{1}{M} \sum_{i=1}^N m_i \vec{R}_i, \quad \vec{P}_{\text{tot}} = \sum_{i=1}^N \vec{P}_i, \quad (4.2.11)$$

and  $N - 1$  translation-invariant coordinate/momentum pairs, each satisfying canonical commutation relations. The kinetic energy terms for these momenta are all decoupled from each other, but with the complication that they have different effective masses even if the particle masses  $m_i$  are all the same. If there are no external forces, so that the potential energy is invariant under translations, then  $\vec{R}_{\text{cm}}$  will not appear in the Hamiltonian at all. The energy eigenstates will therefore have the form of a tensor product, of plane-wave free particle eigenstates of  $\vec{P}_{\text{tot}}$  and  $P_{\text{tot}}^2/2M$ , and eigenstates of the remaining, more complicated, part of the Hamiltonian.

In the case of multi-electron atoms, where one of the particles is a nucleus that is much more massive than the electrons, it is much more common to make the simple and good approximation that the nucleus is infinitely massive and fixed at the origin in the center-of-mass frame, and the remaining coordinates and momenta are (very close to) those of the individual electrons.

### 4.3 Charged particle in external electromagnetic fields

We now turn our attention to the dynamics of charged particles in external electromagnetic fields. We will follow the example of most quantum mechanics books by using Gaussian cgs unit normalizations for electrodynamics quantities, rather than the SI units that you may be more familiar with. This means that Maxwell's equations for the electric and magnetic fields are (with the SI versions indicated parenthetically, for comparison)

$$\vec{\nabla} \cdot \vec{E} = 4\pi\rho \quad (\text{SI: } \rho/\epsilon_0), \quad (4.3.1)$$

$$\vec{\nabla} \cdot \vec{B} = 0 \quad (\text{SI: } 0), \quad (4.3.2)$$

$$\vec{\nabla} \times \vec{E} = -\frac{1}{c} \frac{\partial \vec{B}}{\partial t} \quad (\text{SI: } -\frac{\partial \vec{B}}{\partial t}), \quad (4.3.3)$$

$$\vec{\nabla} \times \vec{B} = \frac{1}{c} \frac{\partial \vec{E}}{\partial t} + \frac{4\pi}{c} \vec{j} \quad (\text{SI: } \frac{1}{c^2} \frac{\partial \vec{E}}{\partial t} + \mu_0 \vec{j}). \quad (4.3.4)$$

It follows that the charge density  $\rho$  and the current density  $\vec{j}$  obey local charge conservation,

$$\vec{\nabla} \cdot \vec{j} + \frac{\partial \rho}{\partial t} = 0. \quad (4.3.5)$$

The electromagnetic energy density and Poynting vector (power per unit area) are

$$u_{\text{EM}} = \frac{1}{8\pi}(E^2 + B^2) \quad (\text{SI: } \frac{\epsilon_0}{2}E^2 + \frac{1}{2\mu_0}B^2), \quad (4.3.6)$$

$$\vec{S} = \frac{c}{4\pi}\vec{E} \times \vec{B} \quad (\text{SI: } \frac{1}{\mu_0}\vec{E} \times \vec{B}). \quad (4.3.7)$$

The electromagnetic fields are obtained as derivatives of the scalar and vector potentials,

$$\vec{E} = -\vec{\nabla}\Phi - \frac{1}{c}\frac{\partial \vec{A}}{\partial t}, \quad (4.3.8)$$

$$\vec{B} = \vec{\nabla} \times \vec{A}. \quad (4.3.9)$$

The fields  $\vec{E}$  and  $\vec{B}$  remain unchanged if one makes a simultaneous change in  $\Phi$  and  $\vec{A}$ , called a **gauge transformation**,

$$\Phi \rightarrow \Phi - \frac{1}{c}\frac{\partial \Lambda}{\partial t}, \quad \vec{A} \rightarrow \vec{A} + \vec{\nabla}\Lambda, \quad (4.3.10)$$

where  $\Lambda(\vec{r}, t)$  is an arbitrary function of position and time.

In classical electrodynamics, the Lagrangian for a nonrelativistic particle with mass  $m$  and charge<sup>†</sup>  $q$  and position  $\vec{r}(t)$ , moving in the potentials  $\Phi$  and  $\vec{A}$ , is

$$L = \frac{1}{2}m\left(\frac{d\vec{r}}{dt}\right)^2 + \frac{q}{c}\frac{d\vec{r}}{dt} \cdot \vec{A}(\vec{r}, t) - q\Phi(\vec{r}, t). \quad (4.3.11)$$

The equation of motion resulting from applying eq. (4.1.2) is the Lorentz force law,

$$m\frac{d^2\vec{r}}{dt^2} = q\left(\vec{E} + \frac{1}{c}\frac{d\vec{r}}{dt} \times \vec{B}\right). \quad (4.3.12)$$

The motion of a classical<sup>‡</sup> charged particle is thus determined only by the local values of  $\vec{E}$  and  $\vec{B}$ . Although the Lagrangian is written in terms of the potentials  $\Phi$  and  $\vec{A}$ , they are not

---

<sup>†</sup>The convention in this book is that the electric charge for a particle is given by  $q = Qe$ , where  $Q$  is a dimensionless number, while  $e$  is the proton's charge, numerically given by eq. (1.1.2), and positive. (Some other sources define  $e$  to be negative, referring to the electron.) Thus, for the electron,  $Q = -1$  and  $q = -e$ , and for the proton,  $Q = 1$  and  $q = e$ . All known particles have  $Q$  equal to integer multiples of  $1/3$ , and the ones unconfined by the strong nuclear force have integer  $Q$ . For example,  $Q = 2/3$  for up, charm, and top quarks, and  $Q = -1/3$  for down, strange, and bottom quarks, and  $Q = -1$  for the electron, muon, and tau lepton. Within the assumed structure of the Standard Model of particle physics, this remarkable charge quantization can be understood as a requirement of anomaly cancellation, a consistency constraint on quantum field theories with gauge interactions. Grand Unified Theories (based on non-Abelian gauge groups like  $SU(5)$ ,  $SO(10)$ , or  $E_6$ ) go further, elegantly explaining why all particles, known or unknown, must have integer values of  $3Q$ , but it is not presently known if these theories are correct.

<sup>‡</sup>In quantum mechanics, the potentials  $\Phi$  and  $\vec{A}$  affect charged particles in ways that are *not* encoded locally in the fields  $\vec{E}$  and  $\vec{B}$ . This is demonstrated by the Aharonov–Bohm effect, discussed in section 28.4.

physical observables, because the physics is equally well described if they are modified by any gauge transformation of the form eq. (4.3.10).

Applying the procedure in section 4.1 to eq. (4.3.11), the classical canonical momentum conjugate to  $\vec{r}$  is

$$\vec{p} = m \frac{d\vec{r}}{dt} + \frac{q}{c} \vec{A}, \quad (4.3.13)$$

and the classical Hamiltonian is

$$H = \frac{1}{2m} \left( \vec{p} - \frac{q}{c} \vec{A}(\vec{r}, t) \right)^2 + q\Phi(\vec{r}, t). \quad (4.3.14)$$

Like the Lagrangian, the Hamiltonian and canonical momentum are not written directly in terms of the fields  $\vec{E}$  and  $\vec{B}$ , but rather in terms of the potentials  $\Phi$  and  $\vec{A}$ , even though the latter are gauge-dependent. One must be careful to distinguish the canonical momentum  $\vec{p}$  of the particle from the **kinetic momentum** (also known as **mechanical momentum**), defined as the product of mass and velocity,

$$\vec{\pi} \equiv m \frac{d\vec{r}}{dt} = \vec{p} - \frac{q}{c} \vec{A}. \quad (4.3.15)$$

The kinetic momentum  $\vec{\pi}$  is a gauge-invariant observable, since  $d\vec{r}/dt$  can be calculated from the observed trajectory of the particle, and so cannot depend on the choice of gauge. In contrast, the canonical momentum  $\vec{p}$  is not a gauge-invariant observable, due to the appearance of  $\vec{A}$  in eq. (4.3.13).

In quantum mechanics, we promote the classical position  $\vec{r}$  and canonical momentum  $\vec{p}$  to operators, and thus the electromagnetic potentials become operators  $\Phi(\vec{R}, t)$  and  $\vec{A}(\vec{R}, t)$  that are functions of  $\vec{R}$ . So, naively, the quantum Hamiltonian should be

$$H = \frac{1}{2m} \left( \vec{P} - \frac{q}{c} \vec{A}(\vec{R}, t) \right)^2 + q\Phi(\vec{R}, t), \quad (4.3.16)$$

where  $\vec{R}$  and  $\vec{P}$  satisfy the usual canonical commutation relations of eq. (2.8.54). This implies that in the position representation, these canonical operators are represented by

$$\vec{R} \leftrightarrow \vec{r}, \quad \vec{P} \leftrightarrow -i\hbar \vec{\nabla}, \quad (4.3.17)$$

as in Table 2.8.2. One must be careful with the ordering of  $\vec{P}$  and  $\vec{A}$ , so that  $[\vec{P} - \frac{q}{c} \vec{A}(\vec{R}, t)]^2$  is interpreted as the symmetrized form  $P^2 - \frac{q}{c}(\vec{P} \cdot \vec{A} + \vec{A} \cdot \vec{P}) + \frac{q^2}{c^2} A^2$ , in order that  $H$  is Hermitian. However, eq. (4.3.16) is still not complete, because it does not include the (purely quantum) effect of intrinsic angular momentum, or spin.

We will discuss spin more thoroughly in 8.2, but for the present discussion we only need to know that the spin for a particle is an observable vector operator  $\vec{S}$ . The intrinsic magnetic

moment of a particle is always proportional to its spin (because, in the rest frame of the particle, there is no other special direction in which it could point):

$$\vec{\mu} = \gamma \vec{S}. \quad (4.3.18)$$

The constant of proportionality  $\gamma$  is a property of the particle type, called the **gyromagnetic ratio**. For the electron, the gyromagnetic ratio is often written as

$$\gamma_e = -\frac{g_e e}{2m_e c}, \quad (4.3.19)$$

where the dimensionless quantity  $g_e$  is called the  $g$ -factor of the electron. The Dirac equation of relativistic quantum mechanics predicts  $g_e = 2$ , as shown in section 27.4, but there are small corrections to this coming from the quantum field theory of relativistic quantum electrodynamics (QED). It has been predicted very precisely by calculations in perturbation theory in QED, and measured experimentally with comparable accuracy, with the results

$$g_e = 2.00231930436321(46) \quad (\text{QED prediction, 5th-order perturbation theory}), \quad (4.3.20)$$

$$g_e = 2.00231930436182(52) \quad (\text{experiment}), \quad (4.3.21)$$

a famous agreement of better than 12 significant digits between theory and experiment. The quantity  $(g_e - 2)/2$  is called the **anomalous magnetic moment** of the electron. In this book, we will usually simply use the approximation  $g_e = 2$ .

For the proton and the neutron, the gyromagnetic ratios are often written as

$$\gamma_p = \frac{g_p e}{2m_p c}, \quad (4.3.22)$$

$$\gamma_n = \frac{g_n e}{2m_p c}, \quad (4.3.23)$$

which again define dimensionless  $g$ -factors. Note that the neutron has a magnetic moment, even though it has no net charge, because it is a composite particle with charged constituents (quarks). The conventional definition of  $g_n$  for the neutron in eq. (4.3.23) uses the proton's charge and mass. The nucleon  $g$ -factors are also very accurately known experimentally,

$$g_p = 5.5856946893(16), \quad (4.3.24)$$

$$g_n = -3.82608545(90), \quad (4.3.25)$$

but the theoretical predictions of these quantities are not nearly as accurate. The reason is that, unlike the electron, the proton and neutron are complicated composite particles made up of quarks and gluons (and virtual quark/antiquark pairs) held together by the strong nuclear

force, (quantum chromodynamics, or QCD) for which perturbation theory does not converge, and non-perturbative methods are highly advanced but limited by finite computing power.

For electrons, protons, and neutrons, the magnitude of  $\vec{S}$  is always the same,  $\hbar/2$ . The magnitude of the electron's intrinsic magnetic moment is within about 0.1% of the **Bohr magneton**,

$$\mu_B = \frac{e\hbar}{2m_e c} = 5.78838 \times 10^{-5} \frac{\text{eV}}{\text{Tesla}} = 9.27401 \times 10^{-21} \frac{\text{ergs}}{\text{gauss}}, \quad (4.3.26)$$

and the proton and neutron have magnetic moment magnitudes equal to about 2.79 and 1.91 times the **nuclear magneton**,

$$\mu_N = \frac{e\hbar}{2m_p c} = 3.15245 \times 10^{-8} \frac{\text{eV}}{\text{Tesla}} = 5.05078 \times 10^{-24} \frac{\text{ergs}}{\text{gauss}}. \quad (4.3.27)$$

Because  $\mu_B/\mu_N = m_p/m_e \approx 1836$  is a large number, the magnetic moments of atoms with unpaired electrons are typically 3 orders of magnitude larger than nuclear magnetic moments.

The classical energy of interaction of a magnetic moment  $\vec{\mu}$  with an external magnetic field is  $-\vec{\mu} \cdot \vec{B}$ . So, we add this to the quantum Hamiltonian for a nonrelativistic particle, to get

$$H = \frac{1}{2m} \left( \vec{P} - \frac{q}{c} \vec{A}(\vec{R}, t) \right)^2 + q\Phi(\vec{R}, t) - \gamma \vec{S} \cdot \vec{B}, \quad (4.3.28)$$

where  $q$  is the electric charge and  $\gamma$  is the appropriate gyromagnetic ratio for the particle. Even this Hamiltonian is not complete, for it does not include relativistic effects suppressed by further powers of  $1/c$ . These will be discussed when needed for the fine and hyperfine contributions to the hydrogen atom, in sections 17.1 and 17.2, and derived in a more fundamental way from the Dirac equation in sections 27.4 and 27.5.

If the Hamiltonian eq. (4.3.28) is written in terms of the kinetic momentum operator

$$\vec{\Pi} = \vec{P} - \frac{q}{c} \vec{A}, \quad (4.3.29)$$

it will appear simpler, since it then does not depend on the vector potential  $\vec{A}$ ,

$$H = \frac{1}{2m} \Pi^2 + q\Phi(\vec{R}, t) - \gamma \vec{S} \cdot \vec{B}. \quad (4.3.30)$$

However, it is important that the kinetic momentum does not obey canonical commutation relations. In the position representation,  $\vec{P} \leftrightarrow -i\hbar \vec{\nabla}$ , so that  $\vec{\Pi} \leftrightarrow -i\hbar \vec{\nabla} - \frac{q}{c} \vec{A}(\vec{r})$ , and

$$[R_a, \Pi_b] = i\hbar \delta_{ab}, \quad (a, b = x, y, z), \quad (4.3.31)$$

just as for the canonical momentum, but

$$[\Pi_a, \Pi_b] = i \frac{q\hbar}{c} (\nabla_a A_b - \nabla_b A_a). \quad (4.3.32)$$

Rewriting this directly in terms of the magnetic field,

$$[\Pi_x, \Pi_y] = i\frac{q\hbar}{c}B_z, \quad [\Pi_y, \Pi_z] = i\frac{q\hbar}{c}B_x, \quad [\Pi_z, \Pi_x] = i\frac{q\hbar}{c}B_y. \quad (4.3.33)$$

This should be contrasted with the canonical commutator  $[P_a, P_b] = 0$ . If one chooses to write the Hamiltonian in terms of the kinetic momentum, the simplicity comes with a cost; the information about the vector potential and the magnetic field is hidden in the commutation relations for the components of  $\vec{\Pi}$  with each other. Since  $\vec{\Pi}$  is the product of mass and velocity for the particle, we see that in the presence of a magnetic field one cannot simultaneously specify the three components of the velocity of a charged particle, because they are not compatible observables.

The freedom to make gauge transformations as in eq. (4.3.10) always allows us to select **Coulomb gauge** (also known as **transverse gauge**), defined by

$$\vec{\nabla} \cdot \vec{A} = 0. \quad (4.3.34)$$

Since  $\vec{P}$  is given in the position representation by  $-i\hbar\vec{\nabla}$  acting on everything to its right, the Coulomb gauge condition implies  $\vec{P} \cdot \vec{A} = \vec{A} \cdot \vec{P}$ , with the consequent advantage that (unlike other gauge choices) there is no operator ordering issue with the cross-terms in the Hamiltonian, of the type mentioned after eq. (4.3.17). Thus, in Coulomb gauge, one is free to write

$$H = \frac{P^2}{2m} - \frac{q}{mc}\vec{A} \cdot \vec{P} + \frac{q^2}{2mc^2}A^2 + q\Phi - \gamma\vec{S} \cdot \vec{B}. \quad (4.3.35)$$

This form will be useful to us when we discuss absorption and emission of light, in Chapter 22.

## 4.4 Exercises

**Exercise 4.1.** The purpose of this problem is to illustrate the possible ambiguities in connecting classical observables to their quantum counterparts due to operator ordering. Consider a particle moving in one dimension with position  $x$  and momentum  $p$ .

(a) Consider the classical quantity  $x^2p^2$ . The quantum operator  $X^2P^2$  is not Hermitian, but two versions of it that are symmetrized in different ways to obtain Hermitian operators are  $A = \frac{1}{2}(X^2P^2 + P^2X^2)$  and  $B = \frac{1}{4}(XP + PX)^2$ . Show that  $A$  and  $B$  differ by a certain rational multiple of  $\hbar^2I$  that you will find.

(b) Similarly, consider the classical quantity  $x^3p^3$  and two candidate quantum Hermitian operator versions of it,  $C = \frac{1}{2}(X^3P^3 + P^3X^3)$  and  $D = \frac{1}{8}(XP + PX)^3$ . Show that these differ by an observable that you will discover. Simplify it and show that it is not proportional to the identity.

**Exercise 4.2.** Show that the pairs  $(\vec{R}, \vec{P})$  and  $(\vec{R}_{\text{cm}}, \vec{P}_{\text{tot}})$  defined in eqs. (4.2.4) and (4.2.5) satisfy canonical commutation relations. Then check that the Hamiltonian of eq. (4.2.1) decomposes into  $H_{\text{cm}}$  and  $H_{\text{rel}}$  as claimed in eq. (4.2.7).

## 5 Transformations, symmetries, and conservation laws

### 5.1 Continuous unitary transformations, symmetries, and Noether's principle

As we saw in section 2.5, a unitary operator can be viewed as implementing a change of orthobasis for the space of states. Certain unitary operators also have the interpretation of physical changes, or **transformations**, on a system. Notable examples include translations, rotations, inversion of the coordinate system, and displacements in time.

If a transformation leaves the Hamiltonian unchanged, then we say, as a definition, that the transformation is a **symmetry** of the quantum system. In this section we will show, in a general way, that there is always a conserved quantity corresponding to each continuous symmetry. As special cases, we will see that the conserved quantities associated with time translation, spatial translation, and rotation symmetries are, respectively, energy, momentum, and angular momentum.

Consider the unitary transformation operator

$$U(\alpha) = \exp(-i\alpha_a G_a), \quad (5.1.1)$$

where the  $G_a$  are a set of  $N$  Hermitian operators, typically observables without explicit time dependence, called the **generators** of the transformations, and the  $\alpha_a$  are  $N$  real numbers that parameterize the transformations. Here, and in the following, repeated indices  $a = 1, \dots, N$  are implicitly summed over. Theorem 2.4.1 confirms that since  $\alpha_a G_a$  is Hermitian,  $U(\alpha)$  is a unitary operator. Following the discussion surrounding eqs. (2.5.26)-(2.5.31), the transformations for the state ket and all observables  $A$  are defined by

$$|\psi\rangle \rightarrow |\psi'\rangle = U|\psi\rangle, \quad (5.1.2)$$

$$A \rightarrow A' = UAU^\dagger, \quad (5.1.3)$$

so that matrix elements are invariant under the transformation, because  $U^\dagger U = I$ . The inverse of the transformation parameterized by  $\alpha_a$  is parameterized by  $-\alpha_a$ ,

$$U(\alpha)^\dagger = U(\alpha)^{-1} = U(-\alpha). \quad (5.1.4)$$

The closure property says that the combination of two transformations  $\alpha_a$  and  $\beta_a$  should always be another transformation, parameterized by some set of real numbers  $\gamma_a$ ,

$$U(\beta)U(\alpha) = U(\gamma). \quad (5.1.5)$$

A continuous set of transformations obeying these properties has the structure of a **Lie group**, named after the mathematician Marius Sophus Lie.



Consider the following sequence of unitary transformations that are infinitesimally close to the identity,

$$\begin{aligned} U(\epsilon)U(\delta)U(\epsilon)^\dagger U(\delta)^\dagger &= (I - i\epsilon_a G_a + \cdots)(I - i\delta_b G_b + \cdots)(I + i\epsilon_c G_c + \cdots)(I + i\delta_d G_d + \cdots) \\ &= I - \epsilon_a \delta_b [G_a, G_b] + \cdots, \end{aligned} \quad (5.1.6)$$

where terms of higher order in either  $\epsilon_a$  or  $\delta_b$  have been dropped. The closure property implies that this must also be a unitary transformation infinitesimally close to the identity, so  $[G_a, G_b]$  must be a linear combination of generators. Therefore, we can write

$$[G_a, G_b] = if_{abc} G_c \quad (5.1.7)$$

for some set of numbers  $f_{abc}$ , called the **structure constants** of the Lie group of transformations. Equation (5.1.7) is called the **Lie algebra** of the group. Since the commutator of any two Hermitian operators  $[G_a, G_b]$  is anti-Hermitian, and the  $G_c$  are Hermitian, the quantities  $f_{abc}$  must all be real. This is the reason for the conventional factor of  $i$  in eq. (5.1.7). If the numbers  $f_{abc}$  are all 0, so that the generators all commute, then the group is said to be Abelian, otherwise it is non-Abelian. The mathematical study and classification of Lie groups is a rich and beautiful subject that we will not delve into further here.

For the remainder of this section, let us consider unitary transformations that are symmetries of a quantum system. According to our definition, this means that  $H$  does not change, so eq. (5.1.3) gives

$$U(\alpha)HU(\alpha)^\dagger = H. \quad (5.1.8)$$

Taking the special case that the parameters  $\alpha_a = \epsilon_a$  are infinitesimal, we have

$$(I - i\epsilon_a G_a + \cdots) H (I + i\epsilon_b G_b + \cdots) = H, \quad (5.1.9)$$

which implies  $\epsilon_a [H, G_a] = 0$ . Since this is supposed to hold for any  $\epsilon_a$ , we learn that

$$[H, G_a] = 0. \quad (5.1.10)$$

It follows from Theorem 2.7.1 that an orthobasis of energy eigenstates can also be chosen to be eigenstates of any subset of the symmetry generators  $G_a$  that are mutually commuting. (If this subset is maximal, it is called a Cartan subalgebra, after mathematician Élie Cartan).

Consider an eigenstate  $|g_a\rangle$  of one of the symmetry generators  $G_a$ , labeled by its eigenvalue  $g_a$ . Since the symmetry generators  $G_a$  commute with the Hamiltonian,  $|g_a\rangle$  will remain an eigenstate of  $G_a$  at later times, because

$$G_a (e^{-itH/\hbar} |g_a\rangle) = e^{-itH/\hbar} G_a |g_a\rangle = g_a (e^{-itH/\hbar} |g_a\rangle). \quad (5.1.11)$$

This holds even if the Hamiltonian depends on time, provided that  $[H(t), G_a] = 0$  for all  $t$ , as one can see by replacing the factors  $e^{-itH/\hbar}$  by the time-evolution operator  $U(t, t_0)$  from eq. (3.4.32).

Furthermore, the symmetry generators  $G_a$  are **conserved quantities**, also known as **constants of the motion**. The precise meaning of this is given by the following result.

**Theorem 5.1.1. (Noether's principle)** *Suppose that an observable  $G_a$  has no explicit time dependence, and that it commutes with the Hamiltonian, which may be time-dependent. Then, for an arbitrary state, the probability of each possible outcome  $g_a$  for a measurement of  $G_a$  is independent of time.*

Proof: The key point is that  $[H(t), G_a] = 0$  implies that, for any particular time  $t = t_0$ , one can find an orthobasis of common eigenkets  $|g_a, E, u\rangle$ , defined such that  $G_a |g_a, E, u\rangle = g_a |g_a, E, u\rangle$  and  $H(t_0) |g_a, E, u\rangle = E |g_a, E, u\rangle$ . Here  $u = u_{g_a, E}$  are possible degeneracy labels for  $g_a$  and  $E$ , with the subscripts dropped for typographical simplicity. These orthobasis kets are fixed, determined by the Hamiltonian at the time  $t_0$ . Now, Postulate 4 tells us that if the system is in a state  $|\psi(t)\rangle$ , then the probability of measuring  $G_a$  at time  $t$  and getting the result  $g_a$  is

$$\mathcal{P}(g_a, t) = \sum_{E, u} \langle g_a, E, u | \psi(t) \rangle \langle \psi(t) | g_a, E, u \rangle. \quad (5.1.12)$$

Here  $E$  and  $u$  are summed over, as the degeneracy labels for  $g_a$ . Taking the time derivative, and then using the time-dependent Schrödinger equation  $\frac{d}{dt} |\psi\rangle = -\frac{i}{\hbar} H(t) |\psi\rangle$  and its adjoint  $\frac{d}{dt} \langle \psi| = \frac{i}{\hbar} \langle \psi| H(t)$ , we get

$$\begin{aligned} \frac{d}{dt} \mathcal{P}(g_a, t) = & -\frac{i}{\hbar} \sum_{E, u} \left( \langle g_a, E, u | H(t) | \psi(t) \rangle \langle \psi(t) | g_a, E, u \rangle \right. \\ & \left. - \langle g_a, E, u | \psi(t) \rangle \langle \psi(t) | H(t) | g_a, E, u \rangle \right). \end{aligned} \quad (5.1.13)$$

Since  $H(t_0) |g_a, E, u\rangle = E |g_a, E, u\rangle$  and  $\langle g_a, E, u | H(t_0) = E \langle g_a, E, u |$ , the two terms on the right side of eq. (5.1.13) simply cancel if we evaluate them at  $t = t_0$ . Thus, we obtain

$$\left. \frac{d}{dt} \mathcal{P}(g_a, t) \right|_{t=t_0} = 0. \quad (5.1.14)$$

This shows that, for an arbitrary state,  $\mathcal{P}(g_a, t)$  has a vanishing time derivative at any given  $t = t_0$ , so it must be constant in time, as claimed.  $\square$

The expectation value of an operator in a state is the sum of its measurement outcomes weighted by the probabilities; see eq. (3.3.1). Therefore, Theorem 5.1.1 immediately implies a weaker but still interesting and important result, that the expectation value of a symmetry generator  $G_a$  in an arbitrary state does not depend on time:

$$\frac{d}{dt} \langle G_a \rangle = 0. \quad (5.1.15)$$

This also follows directly from Ehrenfest's Theorem eq. (3.4.16) with  $G_a$  playing the role of  $A$ . Theorem 5.1.1 and eq. (5.1.15) are quantum mechanical versions of Emmy Noether's celebrated 1918 theorem in classical mechanics, that every continuous symmetry implies a conservation law. The observables  $G_a$  that generate the symmetries of the Hamiltonian are conserved quantities.

The most straightforward special case is that of **time translation** symmetry, for which the generator is the Hamiltonian itself, as expressed by the Schrödinger equation (3.1.10) in Postulate 6. If the Hamiltonian does not depend explicitly on time, then the energy is a conserved quantity, since  $[H, H] = 0$  is trivially satisfied. In that case, for any state the probability of measuring the energy to be any particular value is independent of time, and the expectation value of the energy does not change in time.

## 5.2 Translations

For a particle moving in one dimension with position operator  $X$  and momentum operator  $P$ , consider the operator

$$T(a) = e^{-iaP/\hbar}, \quad (5.2.1)$$

where  $a$  is a constant length. Because  $P$  is Hermitian, Theorem 2.4.1 tells us that  $T(a)$  is a unitary operator,

$$T(a)^\dagger = T(a)^{-1} = T(-a). \quad (5.2.2)$$

Using  $[X, P] = i\hbar$ , one can apply Theorem 2.4.4 to obtain the commutator

$$[X, T(a)] = aT(a). \quad (5.2.3)$$

From this we get, acting on a position eigenstate  $|x\rangle$ ,

$$X(T(a)|x\rangle) = T(a)(X + a)|x\rangle = (x + a)(T(a)|x\rangle), \quad (5.2.4)$$

This shows that  $T(a)|x\rangle$  is an eigenstate of  $X$  with eigenvalue  $x + a$ , which means that it must be equal to  $|x + a\rangle$  up to a multiplicative constant. If  $|x\rangle$  is normalized according to the Dirac condition, then so will be  $T(a)|x\rangle$ , since  $T(a)$  is unitary. Therefore, the constant is just a phase, and  $T(a)|x\rangle = e^{i\theta}|x + a\rangle$  for some  $\theta$ . The only way to resolve the ambiguity represented by  $\theta$  is to arbitrarily choose a value for it, and  $\theta = 0$  is as good as any, so we define

$$|x + a\rangle = T(a)|x\rangle. \quad (5.2.5)$$

In words,  $T(a)$  operates by changing a state in which the particle is known to be at  $x$  into a state where it is known to be at  $x + a$ . We therefore call  $T(a)$  a **translation operator**.

Together, eqs. (5.2.5) and (5.2.2) imply

$$\langle x|T(a) = \langle x-a|, \quad (5.2.6)$$

so that for any state  $|\psi\rangle$ , the wavefunction obeys

$$\psi(x-a) = \langle x|T(a)|\psi\rangle. \quad (5.2.7)$$

This can be interpreted as shifting the particle's position to the right by  $a$  (the “active view”), or equivalently as leaving the particle's position unchanged but shifting the coordinate system to the left by  $a$  (the “passive view”).

If we do a translation on any normalized state  $|\psi\rangle$ ,

$$|\psi\rangle \rightarrow T(a)|\psi\rangle, \quad (5.2.8)$$

then the expectation value of the position is changed according to

$$\langle\psi|X|\psi\rangle \rightarrow \langle\psi|T(a)^\dagger XT(a)|\psi\rangle = \langle\psi|(X+a)|\psi\rangle = \langle\psi|X|\psi\rangle + a. \quad (5.2.9)$$

The effect of  $T(a)$  on momentum eigenstates is just to multiply by a phase,

$$T(a)|p\rangle = e^{-iap/\hbar}|p\rangle, \quad (5.2.10)$$

and the expectation value of momentum is unaffected by the transformation,

$$\langle\psi|P|\psi\rangle \rightarrow \langle\psi|T(a)^\dagger PT(a)|\psi\rangle = \langle\psi|P|\psi\rangle. \quad (5.2.11)$$

Following the general example of a unitary transformation of an operator in eq. (5.1.3), we can also define for an arbitrary observable  $A$  the translated version  $A' = T(a)AT(a)^\dagger$ . Now if we do a simultaneous transformation of both states and operators,

$$|\psi\rangle \rightarrow |\psi'\rangle = T(a)|\psi\rangle, \quad (5.2.12)$$

$$A \rightarrow A' = T(a)AT(a)^\dagger, \quad (5.2.13)$$

then matrix elements will be unchanged. Using  $[X, T(a)] = aT(a)$ , the translated position operator is

$$X' = T(a)XT(a)^\dagger = X - a, \quad (5.2.14)$$

while  $[P, T(a)] = 0$  implies that the momentum operator does not change,

$$P' = T(a)PT(a)^\dagger = P. \quad (5.2.15)$$

From the rule  $T(a)|x\rangle = |x+a\rangle$  found in eq. (5.2.5), it follows that translations simply add,

$$T(a)T(b) = T(a+b). \quad (5.2.16)$$

One consequence of this is that any finite translation can be viewed as the combination of many infinitesimal translations. For an infinitesimal translation, the exponential can be truncated to linear order,

$$T(\epsilon) = 1 - i\frac{\epsilon}{\hbar}P. \quad (5.2.17)$$

This is described in words by saying that the momentum operator is the generator of translations. Since the momentum operator commutes with itself, in the general language of section 5.1 where  $T(a)$  plays the role of  $U(\alpha)$  and  $P$  plays the role of  $G_a$ , the Lie algebra for translations is simply

$$[P, P] = 0. \quad (5.2.18)$$

This is an Abelian algebra; the antisymmetric structure constants defined in general by eq. (5.1.7) trivially vanish in this case, because there is only one generator,  $P$ .

Now suppose that a quantum mechanical system has translations as a symmetry, by which we mean that the Hamiltonian is invariant. In particular, for infinitesimal translations, we require  $T(\epsilon)HT(\epsilon)^\dagger = H$ , from which it follows that

$$\left(1 - i\frac{\epsilon}{\hbar}P\right)H\left(1 + i\frac{\epsilon}{\hbar}P\right) - H = i\frac{\epsilon}{\hbar}[H, P] + \mathcal{O}(\epsilon^2) \quad (5.2.19)$$

must vanish, so

$$[H, P] = 0. \quad (5.2.20)$$

In general, this requires the potential  $V$  to have no dependence on  $X$ . The Hamiltonian could be that of a free particle with  $H = P^2/2m$ , but it could also have some extra terms that may involve other functions of  $P$  or other degrees of freedom (for example, spin), but not  $X$ . In any case, eq. (5.2.20) implies that there must be an orthobasis of common eigenstates of  $P$  and  $H$ . An eigenstate of  $P$  with momentum  $p$  remains so at later times, since

$$P\left(e^{-itH/\hbar}|p\rangle\right) = p\left(e^{-itH/\hbar}|p\rangle\right). \quad (5.2.21)$$

Also, Ehrenfest's Theorem eq. (3.4.16) says

$$\frac{d}{dt}\langle P \rangle = 0, \quad (5.2.22)$$

and Theorem 5.1.1 says even more, that the probability to measure the momentum within any given range will be constant in time. Note that these statements are true for any state,

including an arbitrary superposition of free particle eigenstates with different momenta, provided that  $[H, P] = 0$ . A non-trivial illustration will be provided at the end of section 6.2.

For a particle moving in three dimensions, the preceding discussion can be adapted in a straightforward way. The translation operator for a vector distance  $\vec{a}$  is

$$T(\vec{a}) = e^{-i\vec{a} \cdot \vec{P}/\hbar}, \quad (5.2.23)$$

and it satisfies

$$T(\vec{a})T(\vec{b}) = T(\vec{a} + \vec{b}), \quad (5.2.24)$$

$$T(\vec{a})^{-1} = T(\vec{a})^\dagger = T(-\vec{a}). \quad (5.2.25)$$

The operation on position and momentum eigenkets is

$$T(\vec{a})|\vec{r}\rangle = |\vec{r} + \vec{a}\rangle, \quad \langle\vec{r}|T(\vec{a}) = \langle\vec{r} - \vec{a}|, \quad (5.2.26)$$

$$T(\vec{a})|\vec{p}\rangle = e^{-i\vec{a} \cdot \vec{p}/\hbar}|\vec{p}\rangle, \quad \langle\vec{p}|T(\vec{a}) = e^{i\vec{a} \cdot \vec{p}/\hbar}\langle\vec{p}|, \quad (5.2.27)$$

and the remaining discussion for the one-dimensional case likewise follows through for the three-dimensional case with  $P$  replaced by  $\vec{P}$ . Although there are now three generators  $P_x$ ,  $P_y$ , and  $P_z$ , they all commute with each other, so the structure constants are all 0, and the Lie group is Abelian. Note that in the case of a charged particle moving in an electromagnetic field as discussed in section 4.3, it is the canonical momentum  $\vec{P}$  that generates translations, not the kinetic momentum  $\vec{\Pi}$ .

Consider a quantum system describing two distinguishable particles labeled 1 and 2. Then one can have translation invariance even with a non-zero potential, provided that the Hamiltonian has the form considered in our discussion of the two-body problem in section 4.2,

$$H = \frac{P_1^2}{2m_1} + \frac{P_2^2}{2m_2} + V(\vec{R}_1 - \vec{R}_2). \quad (5.2.28)$$

The individual translation operators for particles 1 and 2 are

$$T_1(\vec{a}) = e^{-i\vec{a} \cdot \vec{P}_1/\hbar}, \quad T_2(\vec{a}) = e^{-i\vec{a} \cdot \vec{P}_2/\hbar}. \quad (5.2.29)$$

These are not symmetries of the Hamiltonian unless the potential  $V(\vec{R}_1 - \vec{R}_2)$  is neglected. However, defining the total momentum operator  $\vec{P}_{\text{tot}} = \vec{P}_1 + \vec{P}_2$  as in section 4.2, one can check that each component of  $\vec{P}_{\text{tot}}$  commutes with each component of  $\vec{R}_1 - \vec{R}_2$ , so

$$[H, \vec{P}_{\text{tot}}] = 0. \quad (5.2.30)$$

Therefore, the total translation operator

$$T(\vec{a}) = T_1(\vec{a})T_2(\vec{a}) = \exp\left[-i\vec{a} \cdot \vec{P}_{\text{tot}}/\hbar\right] \quad (5.2.31)$$

leaves the full Hamiltonian eq. (5.2.28) invariant,

$$T(\vec{a})HT(\vec{a})^\dagger = H, \quad (5.2.32)$$

and is therefore a symmetry of the system. [Note that this would not be true if one added individual potentials  $V_1(\vec{R}_1)$  or  $V_2(\vec{R}_2)$  to the Hamiltonian.] Equation (5.2.32) simply reflects the feature that the total translation moves both particles 1 and 2 by the same amount, and so does not change the separation between them, nor either of their momenta. It follows that one can find an orthonormal basis of simultaneous eigenstates of  $H$  and  $\vec{P}_{\text{tot}} = \vec{P}_1 + \vec{P}_2$ , and that an eigenstate of  $\vec{P}_{\text{tot}}$  remains so at later times, and that in an arbitrary state the probability of measuring a given result for  $\vec{P}_{\text{tot}}$  does not change in time, and its expectation value is conserved.

### 5.3 Rotations

In this section, we will explore the connection between rotations and the angular momentum operators that generate them. Classically, the angular momentum of a particle about the point chosen as the coordinate system origin is defined by

$$\vec{l} = \vec{r} \times \vec{p}. \quad (5.3.1)$$

In quantum mechanics, we promote this to a vector operator, and define the **orbital angular momentum operator** for a particle as

$$\vec{L} = \hat{x}L_x + \hat{y}L_y + \hat{z}L_z = \vec{R} \times \vec{P}, \quad (5.3.2)$$

where the components

$$L_x = YP_z - ZP_y, \quad L_y = ZP_x - XP_z, \quad L_z = XP_y - YP_x. \quad (5.3.3)$$

are each observables. There is no problem with operator ordering to worry about here, because  $[Y, P_z] = [Z, P_y] = [Z, P_x] = [X, P_z] = [X, P_y] = [Y, P_x] = 0$ .

The commutator algebra for the angular momentum components  $L_x$ ,  $L_y$ , and  $L_z$  can be computed using the commutators of the position and momentum operators that they are built out of. For example,

$$[L_x, L_y] = [YP_z, ZP_x] + [ZP_y, XP_z] = -i\hbar YP_x + i\hbar XP_y = i\hbar L_z. \quad (5.3.4)$$

Similarly,

$$[L_y, L_z] = i\hbar L_x, \quad [L_z, L_x] = i\hbar L_y. \quad (5.3.5)$$

The last three equations can be summarized as<sup>†</sup>

$$[L_a, L_b] = i\hbar\epsilon_{abc}L_c, \quad (a, b, c = x, y, z). \quad (5.3.6)$$

Here,  $\epsilon_{abc}$  is the totally antisymmetric **Levi-Civita symbol**, with

$$\epsilon_{xyz} = \epsilon_{yzx} = \epsilon_{zxy} = 1, \quad \epsilon_{xzy} = \epsilon_{yxz} = \epsilon_{zyx} = -1, \quad (5.3.7)$$

and all other components  $\epsilon_{abc} = 0$ . It obeys the identities

$$\epsilon_{abc}\epsilon_{dec} = \delta_{ad}\delta_{be} - \delta_{ae}\delta_{bd}, \quad (5.3.8)$$

$$\epsilon_{abc}\epsilon_{dbc} = 2\delta_{ad}, \quad (5.3.9)$$

$$\epsilon_{abc}\epsilon_{abc} = 6. \quad (5.3.10)$$

In the general language of section 5.1, the role of the generators  $G_a$  is taken by  $L_a/\hbar$  for rotations, as we are about to show. Comparing eq. (5.3.6) to eq. (5.1.7), we see that the Lie algebra of angular momentum operators is non-Abelian, with structure constants  $f_{abc} = \epsilon_{abc}$ . Note that there is always an arbitrary normalization in the definition of the generators; it was convenient to include the factor of  $1/\hbar$  here so that the structure constants are dimensionless.

Rotations are defined by the property that, as changes in coordinates, they leave invariant the distances of points from the origin. The composition of two rotations is another rotation; in mathematical language, rotations correspond to the Lie group called  $SO(3)$ . In quantum mechanics, we can think of rotations as a unitary change of basis corresponding to the change in coordinates; a rotation by an angle  $\alpha$  about the axis defined by a unit vector  $\vec{n}$  is implemented by a unitary operator  $U(\vec{\alpha})$ , where  $\vec{\alpha} = \hat{n}\alpha$ . The fact that rotations form a group means that for any  $\vec{\alpha}$  and  $\vec{\beta}$ ,

$$U(\vec{\beta})U(\vec{\alpha}) = U(\vec{\gamma}), \quad (5.3.11)$$

for some  $\vec{\gamma}$ . In particular, arbitrary finite rotations can be constructed from the limit of a large number of infinitesimal rotations.

Let us now show that (for the case of a single spinless particle)  $L_z$  is the generator of rotations about the  $z$ -axis (analogously to how  $P_z$  is the generator of translations along the  $z$  direction). For such a rotation by an angle  $\alpha$ , the coordinates transform as

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} \rightarrow \begin{pmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}. \quad (5.3.12)$$

---

<sup>†</sup>Here, and from now on, we use the **repeated index summation convention**, which says that repeated indices are implicitly summed over, except when they appear on both sides of an equation. Thus, in eq. (5.3.6),  $c$  is summed over, but  $a$  and  $b$  are not.



If  $\alpha = \epsilon$  is infinitesimal, then this becomes  $x \rightarrow x - \epsilon y$  and  $y \rightarrow y + \epsilon x$ , and of course  $z \rightarrow z$ . We therefore define an infinitesimal unitary rotation operator by its action on the position eigenkets,

$$U(\epsilon \hat{z}) |x, y, z\rangle = |x - \epsilon y, y + \epsilon x, z\rangle, \quad (5.3.13)$$

which implies

$$\langle x, y, z | U(\epsilon \hat{z})^\dagger = \langle x - \epsilon y, y + \epsilon x, z |, \quad (5.3.14)$$

with  $U(\epsilon \hat{z})^\dagger = U(\epsilon \hat{z})^{-1} = U(-\epsilon \hat{z})$ . It follows that, for any state  $|\psi\rangle$ ,

$$\begin{aligned} \langle x, y, z | U(\epsilon \hat{z}) |\psi\rangle &= \langle x + \epsilon y, y - \epsilon x, z | \psi\rangle = \psi(x + \epsilon y, y - \epsilon x, z) \\ &= \psi(x, y, z) + \epsilon y \frac{\partial}{\partial x} \psi(x, y, z) - \epsilon x \frac{\partial}{\partial y} \psi(x, y, z), \end{aligned} \quad (5.3.15)$$

where the last equality is the result of the Taylor series expansion to linear order in  $\epsilon$ . Now, since  $\partial/\partial x \leftrightarrow iP_x/\hbar$  and  $\partial/\partial y \leftrightarrow iP_y/\hbar$  in the position representation, we have

$$\langle x, y, z | U(\epsilon \hat{z}) |\psi\rangle = \langle x, y, z | \left[ I - \frac{i}{\hbar} \epsilon (XP_y - YP_x) \right] |\psi\rangle, \quad (5.3.16)$$

for every state  $|\psi\rangle$ , so comparing with the definition of  $L_z$  in eq. (5.3.3), we find

$$U(\epsilon \hat{z}) = I - \frac{i}{\hbar} \epsilon L_z. \quad (5.3.17)$$

This establishes that  $L_z$  generates rotations about the  $z$  axis, and that

$$L_z \leftrightarrow -i\hbar \left( x \frac{\partial}{\partial y} - y \frac{\partial}{\partial x} \right) \quad (5.3.18)$$

in the position wavefunction representation.

There is nothing special about the  $z$ -axis in the preceding discussion, so repeating the preceding process for infinitesimal rotations about the  $x$  and  $y$  axes, one obtains

$$U(\epsilon \hat{x}) = I - \frac{i}{\hbar} \epsilon L_x, \quad U(\epsilon \hat{y}) = I - \frac{i}{\hbar} \epsilon L_y, \quad (5.3.19)$$

with  $L_x$  and  $L_y$  given by eqs. (5.3.3), and position representations

$$L_x \leftrightarrow -i\hbar \left( y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y} \right), \quad L_y \leftrightarrow -i\hbar \left( z \frac{\partial}{\partial x} - x \frac{\partial}{\partial z} \right). \quad (5.3.20)$$

Thus  $L_x$ ,  $L_y$ , and  $L_z$  are generators for rotations about the  $x$ ,  $y$ , and  $z$  axes, respectively (assuming that there is no intrinsic angular momentum).

The unitary rotation operator for a non-infinitesimal angle  $\alpha$  can be built from the limit of a large number  $N$  of infinitesimal operators with  $\epsilon = \alpha/N$ , acting sequentially, so

$$U(\alpha \hat{z}) = \lim_{N \rightarrow \infty} \left( I - \frac{i}{\hbar} \frac{\alpha}{N} L_z \right)^N = \exp \left( -\frac{i}{\hbar} \alpha L_z \right), \quad (5.3.21)$$

where we have used the definition of the exponential of an operator in eq. (2.4.27). For a rotation by an angle  $\alpha$  about an arbitrary axis  $\hat{n}$ , this generalizes to

$$U(\alpha\hat{n}) = \exp\left(-\frac{i}{\hbar}\vec{\alpha} \cdot \vec{L}\right), \quad (5.3.22)$$

where  $\vec{\alpha} = \hat{n}\alpha$ .

For the position representation, we could also use cylindrical or spherical coordinates instead of rectangular coordinates, so that the position eigenkets are  $|r, \phi, z\rangle$  or  $|r, \theta, \phi\rangle$ , respectively. For spherical coordinates in particular, this has the advantage that one of the coordinates,  $r$ , is not involved in the differential operators that represent the angular momenta. In spherical coordinates, eq. (5.3.15) becomes

$$\langle r, \theta, \phi | U(\epsilon\hat{z}) | \psi \rangle = \psi(r, \theta, \phi) + \epsilon \frac{\partial}{\partial \phi} \psi(r, \theta, \phi), \quad (5.3.23)$$

so that as an equivalent to eq. (5.3.18),

$$L_z \leftrightarrow -i\hbar \frac{\partial}{\partial \phi}. \quad (5.3.24)$$

For the special case of rotations about the  $z$  axis, this also holds in cylindrical coordinates.

Let us now solve the eigenvalue problem for the operator  $L_z$ , using the position representation. Because eq. (5.3.24) only involves the coordinate  $\phi$ , the following derivation works equally well in spherical or cylindrical coordinates. We start with

$$L_z |l_z\rangle = l_z |l_z\rangle, \quad (5.3.25)$$

where  $l_z$  is the eigenvalue, also used as a label for the eigenstate, and we have suppressed any degeneracy labels. In the position representation, this becomes

$$-i\hbar \frac{\partial}{\partial \phi} \psi(\phi) = l_z \psi(\phi), \quad (5.3.26)$$

also suppressing the dependence on other coordinates. The solutions are

$$\psi(\phi) = ce^{il_z\phi/\hbar}, \quad (5.3.27)$$

where  $c$  is a non-zero normalization constant. The range of  $\phi$  is the continuous interval  $0 \leq \phi \leq 2\pi$ . Therefore, for  $L_z$  to be a Hermitian operator, by following exactly the same derivation that led to eq. (2.8.23) with  $P$  replaced by  $L_z$ , we find that for any wavefunctions  $\psi_1$  and  $\psi_2$

$$[\psi_1(2\pi)]^* \psi_2(2\pi) = [\psi_1(0)]^* \psi_2(0) \quad (5.3.28)$$

must hold. A sufficient condition for this to be satisfied is that all wavefunctions obey  $\psi(2\pi) = \psi(0)$ , and more generally  $\psi(\phi + 2\pi) = \psi(\phi)$ . It is necessary to impose this periodic boundary

condition anyway, in order for the wavefunction to have a unique expansion in terms of position eigenstate kets. It follows that  $e^{il_z 2\pi/\hbar} = 1$ , which requires that

$$l_z = \hbar m, \quad (5.3.29)$$

where  $m$  is an integer. The use of the letter  $m$  is traditional in this role, and it is sometimes called the **magnetic quantum number**. We therefore use it as a label for the corresponding eigenstates of  $L_z$ , and write, from now on,

$$L_z|m\rangle = \hbar m|m\rangle. \quad (5.3.30)$$

All of the preceding was derived with the assumption that the angular momentum was associated with the motion of a single particle. More generally, a Hilbert space can describe more than one particle. Furthermore, each of the particles may also have an intrinsic angular momentum, or spin, which has no classical counterpart, and is not associated at all with the quantum mechanical position wavefunction of the particle. These distinct types of angular momenta can also be combined to form new angular momenta.

In general, we define an angular momentum operator

$$\vec{J} = \hat{x}J_x + \hat{y}J_y + \hat{z}J_z \quad (5.3.31)$$

to be one that satisfies a commutator algebra that has the same form as for orbital angular momentum. Specifically,

$$[J_x, J_y] = i\hbar J_z \quad [J_y, J_z] = i\hbar J_x \quad [J_z, J_x] = i\hbar J_y, \quad (5.3.32)$$

or equivalently

$$[J_a, J_b] = i\hbar \epsilon_{abc} J_c. \quad (5.3.33)$$

Because the components of an angular momentum operator  $\vec{J}$  do not commute with each other, they are not compatible, and one cannot find a complete orthobasis of  $\vec{J}$  eigenstates. The only solution to the eigenvalue equation  $\vec{J}|\vec{j}\rangle = \vec{j}|\vec{j}\rangle$  has  $\vec{j} = 0$ . Thus, an angular momentum vector  $\vec{J}$  is not an observable, although each of its components is. If we choose eigenstates of  $J_z$ , they cannot also be eigenstates of  $J_x$  or  $J_y$ , except in the very special case that all of the eigenvalues are 0. However, you can check that the **angular momentum squared operator**

$$J^2 = J_x^2 + J_y^2 + J_z^2 \quad (5.3.34)$$

does commute with  $J_z$ . This means that  $J^2$  and  $J_z$  have common eigenkets and can be part of a CSCO. The simultaneous eigenvalue problem for  $J^2$  and  $J_z$  will be worked out in Chapter 8.

For each particle, the intrinsic angular momentum, or spin  $\vec{S}$ , is a special case of  $\vec{J}$  defined by the property that it has fixed magnitude, in the sense that all particles of a given type have the same eigenvalue of  $S^2$ , which can never change. For any single particle, the total angular momentum operator is simply the sum of the orbital and spin parts,  $\vec{J} = \vec{L} + \vec{S}$ . (Note that we use the same symbol  $\vec{J}$  as for a generic angular momentum operator defined in the previous paragraph.)

For the two-body problem of section 4.2, you can check that the total orbital angular momentum operator (not including the spins) can be written in two equivalent ways as

$$\vec{L}_{\text{tot}} = \vec{L}_1 + \vec{L}_2 = \vec{L} + \vec{L}_{\text{cm}}, \quad (5.3.35)$$

where  $\vec{L}_1 = \vec{R}_1 \times \vec{P}_1$  and  $\vec{L}_2 = \vec{R}_2 \times \vec{P}_2$  for the individual particles, and

$$\vec{L} = \vec{R} \times \vec{P}, \quad \vec{L}_{\text{cm}} = \vec{R}_{\text{cm}} \times \vec{P}_{\text{tot}}, \quad (5.3.36)$$

define the relative and center-of-mass contributions to the angular momenta. It is often sensible to restrict to the subspace of states with vanishing total momentum in the center-of-mass frame, consisting of eigenstates of  $\vec{P}_{\text{tot}}$  with eigenvalue 0. If we do so, then  $\vec{L}_{\text{cm}}$  vanishes identically, and the relative angular momentum  $\vec{L}$  is equal to the total angular momentum  $\vec{L}_{\text{tot}}$ .

More generally, for a given choice of origin, every three-dimensional physical system has a **total angular momentum** operator  $\vec{J}$ , which adds both orbital and spin angular momentum contributions for all of the particles that are present. By definition,  $\vec{J}$  is the operator that acts on the full Hilbert space of states to generate rotations about any axis  $\hat{n}$  by any angle  $\alpha$ , with

$$U(\vec{\alpha}) = \exp\left(-i\alpha\hat{n} \cdot \vec{J}/\hbar\right), \quad (5.3.37)$$

where  $\vec{\alpha} = \hat{n}\alpha$ . To rotate a state  $|\psi\rangle$ , the unitary transformation is

$$|\psi\rangle \rightarrow |\psi'\rangle = U(\vec{\alpha})|\psi\rangle. \quad (5.3.38)$$

We also define rotated operators

$$A \rightarrow A' = U(\vec{\alpha})AU(\vec{\alpha})^\dagger, \quad (5.3.39)$$

so that, due to the unitarity of  $U(\vec{\alpha})$ , matrix elements of the rotated operators between rotated states are the same as the original matrix elements,  $\langle\chi'|A'|\psi'\rangle = \langle\chi|A|\psi\rangle$ .

Suppose that the Hamiltonian for a system has the symmetry of invariance under rotations generated by an angular momentum component  $\hat{n} \cdot \vec{J}$ , so that rotations about the unit vector axis  $\hat{n}$  are a symmetry of the system, and

$$[H, \hat{n} \cdot \vec{J}] = 0. \quad (5.3.40)$$

(It is traditional to choose the coordinate system so that  $\hat{n} = \hat{z}$ , unless there is a good reason not to, but let us be more general.) One can then choose an orthobasis of simultaneous eigenstates of  $H$  and  $\hat{n} \cdot \vec{J}$ , and as shown on general grounds in section 5.1,  $\hat{n} \cdot \vec{J}$  is a conserved quantity. This means that eigenstates of  $\hat{n} \cdot \vec{J}$  will remain so under time evolution, and for an arbitrary state the probability to obtain a given outcome for a measurement of  $\hat{n} \cdot \vec{J}$  is constant, and

$$\frac{d}{dt} \langle \hat{n} \cdot \vec{J} \rangle = 0. \quad (5.3.41)$$

If the Hamiltonian is invariant under all rotations, it is convenient to choose a CSCO to include  $H, J^2, J_z$ , since these observables commute with each other.

## 5.4 Parity

Another kind of coordinate transformation is **parity** (also known as **space inversion**), defined as the replacement of each rectangular position coordinate by minus itself. Unlike translations and rotations, parity is a discrete group known as  $Z_2$ ; acting twice with parity gives back the identity operation, and there are no infinitesimal parity transformations.

Let us start by defining the **parity operator**  $\Pi$  for a particle moving in one dimension by giving its action on the position-eigenstate orthobasis kets,

$$\Pi |x\rangle = |-x\rangle. \quad (5.4.1)$$

(Note that  $|-x\rangle$  is the ket that describes a particle known to be at the point  $-x$ , and is not at all the same thing as  $-|x\rangle$ , which still describes a particle known to be at the point  $x$ .) Since  $\Pi(\Pi|x\rangle) = |x\rangle$ , we have  $\Pi\Pi = I$ , so that parity is its own inverse,

$$\Pi^{-1} = \Pi. \quad (5.4.2)$$

Also, taking the Hermitian adjoint of eq. (5.4.1) gives  $\langle x|\Pi^\dagger = \langle -x|$ , so

$$\langle x'|\Pi^\dagger|x\rangle = \langle -x'|x\rangle = \delta(x+x') = \langle x'|-x\rangle = \langle x'|\Pi|x\rangle. \quad (5.4.3)$$

Since this is true for every  $|x\rangle$  and  $|x'\rangle$  in the position orthobasis, it must be that

$$\Pi^\dagger = \Pi. \quad (5.4.4)$$

Comparing eqs. (5.4.2) and (5.4.4), we see that the parity operator is both unitary (an invertible map from an orthobasis to another orthobasis) and Hermitian (an observable). Since all Hermitian operators have real eigenvalues, and all unitary operators have eigenvalues with magnitude 1, the only possible eigenvalues of  $\Pi$  are 1 and  $-1$ .

Given a wavefunction  $\langle x|\psi\rangle = \psi(x)$  for an arbitrary state  $|\psi\rangle$ , we have

$$\langle x|\Pi|\psi\rangle = \langle -x|\psi\rangle = \psi(-x). \quad (5.4.5)$$

Applying this to the case of a momentum eigenstate,

$$\langle x|\Pi|p\rangle = \langle -x|p\rangle = \frac{1}{\sqrt{2\pi\hbar}}e^{-ipx/\hbar} = \langle x|-p\rangle, \quad (5.4.6)$$

from which we learn that

$$\Pi|p\rangle = |-p\rangle. \quad (5.4.7)$$

Similarly, it is easy to show that

$$\Pi X \Pi = -X, \quad \Pi P \Pi = -P, \quad (5.4.8)$$

so that the position and momentum operators are both said to be odd under parity. More generally, one defines the **parity of an operator**  $A$  as

$$\pi_A = \pm 1, \quad \text{if} \quad \Pi A \Pi = \pm A. \quad (5.4.9)$$

However, not all operators have definite parity in this sense.

If  $A$  is even under parity,  $\pi_A = +1$ , it follows that

$$\Pi A - A \Pi = 0, \quad (5.4.10)$$

so that  $\Pi$  and  $A$  are compatible operators, and can have common eigenstates. For example, the Hamiltonian  $H = P^2/2m + V(X)$  is parity-even if, and only if,  $V(X)$  is an even function of  $X$ . In the special case of a free particle with  $V = 0$ , the simultaneous eigenstates of  $H$  and  $\Pi$  are

$$|E, +1\rangle = \frac{1}{\sqrt{2}}(|p\rangle + |-p\rangle), \quad |E, -1\rangle = \frac{1}{\sqrt{2}}(|p\rangle - |-p\rangle), \quad (5.4.11)$$

where  $p = \sqrt{2mE}/\hbar$ . However, since  $P$  does not commute with  $\Pi$ , one cannot find simultaneous eigenstates of them, and indeed the parity eigenstates  $|E, \pm 1\rangle$  are not eigenstates of momentum.

Just as for translations, parity generalizes straightforwardly to three dimensions. Define

$$\Pi|\vec{r}\rangle = |-\vec{r}\rangle, \quad (5.4.12)$$

from which it follows that

$$\Pi|\vec{p}\rangle = |-\vec{p}\rangle, \quad (5.4.13)$$

and position and momentum operators carry odd parity,

$$\Pi \vec{R} \Pi = -\vec{R}, \quad \Pi \vec{P} \Pi = -\vec{P}. \quad (5.4.14)$$

Angular momentum operators, as defined in the previous section, always have even parity,

$$\Pi \vec{J} \Pi = \vec{J}. \quad (5.4.15)$$

In the case of orbital angular momentum,  $\Pi \vec{L} \Pi = \vec{L}$  follows directly from the definition of the components of  $\vec{L}$  in eq. (5.3.3). For more general angular momentum operators (including spin), the even parity can be inferred from the general commutator algebra structure in eq. (5.3.33). If the potential energy depends only on the radial coordinate, so that the Hamiltonian has the form  $H = P^2/2m + V(R)$ , then this  $H$  commutes not only with  $L^2$  and  $L_z$ , but also with  $\Pi$ . Such a system therefore admits simultaneous eigenstates of all four operators. In that case, parity is a symmetry of the system, and eigenstates of parity will remain so under time evolution.

A powerful application of parity is to the identification of selection rules for matrix elements. Suppose that an operator  $A$  has definite parity  $\pi_A$  as defined by eq. (5.4.9), and that the states  $|\psi\rangle$  and  $|\phi\rangle$  are parity eigenstates with eigenvalues  $\pi_\psi$  and  $\pi_\phi$ , respectively. Now we note that

$$\langle \phi | A | \psi \rangle = \pi_A \langle \phi | \Pi A \Pi | \psi \rangle = \pi_A \pi_\phi \pi_\psi \langle \phi | A | \psi \rangle. \quad (5.4.16)$$

The product  $\pi_A \pi_\phi \pi_\psi$  is either  $+1$  or  $-1$ . In the former case, eq. (5.4.16) tells us nothing, but in the latter case, we obtain:

**Theorem 5.4.1. (Parity selection rule)** *If states  $|\psi\rangle$  and  $|\phi\rangle$  and an operator  $A$  all have definite parities, and  $\pi_A \pi_\phi \pi_\psi = -1$ , then the matrix element  $\langle \phi | A | \psi \rangle$  must vanish.*

This result has many practical applications, including neatly explaining the absence of certain atomic transitions. Before investing your valuable time in calculating a quantity, it is always a good idea to consider first whether it must vanish due to a matrix element selection rule.

## 5.5 Gauge transformations

Physical systems that involve electromagnetic interactions with charged particles can be formulated in terms of potentials, subject to gauge transformations

$$\Phi \rightarrow \Phi - \frac{1}{c} \frac{\partial \Lambda}{\partial t}, \quad \vec{A} \rightarrow \vec{A} + \vec{\nabla} \Lambda, \quad (5.5.1)$$

where  $\Lambda(\vec{r}, t)$  is an arbitrary function. As discussed in section 4.3, the physical situation described by a pair of potentials  $(\Phi, \vec{A})$  is equally well described by any pair  $(\Phi', \vec{A}')$  related to them by a gauge transformation.

Before proceeding, we pause to emphasize that gauge transformations have a completely different character than the other transformations discussed in the preceding sections. If we do a translation or rotation on a system, we have transformed the system into a physically distinct one. A particle that has been translated by 1 centimeter is in a different place, clearly distinguishable by experiment. To say that a system has translation symmetry means that we can physically move the whole system in an experimentally measurable way without changing the Hamiltonian that describes its time evolution. In contrast, the whole point of gauge transformations is that although they *do* change the Hamiltonian, they do so *without* changing the actual physical situation at all. There is no way an experiment can tell whether or not we have chosen to write the Hamiltonian in Coulomb gauge! Although it is common to refer to “gauge symmetries”, gauge transformations really are not symmetries in the same way that translations or rotations can be. Instead, they are a manifestation of the fact that the potentials have an arbitrariness, in the form of redundancies in our description that can be eliminated by fixing the gauge.

Consider a classical charged particle in an electromagnetic field. Clearly, if we do a gauge transformation the position and the velocity of the charged particle are unaffected, since they can be measured experimentally. However, the classical canonical momentum  $\vec{p}$  defined in eq. (4.3.13) does change with a gauge transformation, because it involves not just the velocity but also the gauge-dependent potential  $\vec{A}$ .

In the quantum description, the state ket used to describe the system is similarly gauge-dependent. Naively, this might seem like a problem, but it is not, because the state ket by itself is not a physically measurable observable. For example, we already have pointed out that multiplying a ket by a *constant* complex phase does not change probabilities. In the following, we will show that the gauge transformation of the ket describing a single particle of charge  $q$  is realized as a unitary transformation

$$U_{\Lambda} = \exp \left[ i \frac{q}{\hbar c} \Lambda(\vec{R}, t) \right], \quad (5.5.2)$$

which imparts a *position-dependent* complex phase to the state ket,

$$|\psi\rangle \rightarrow |\psi'\rangle = U_{\Lambda} |\psi\rangle, \quad (5.5.3)$$

and which must be accompanied by changes in the electromagnetic potentials,

$$\Phi(\vec{R}, t) \rightarrow \Phi(\vec{R}, t)' = \Phi(\vec{R}, t) - \frac{1}{c} \frac{\partial}{\partial t} \Lambda(\vec{R}, t), \quad (5.5.4)$$

$$\vec{A}(\vec{R}, t) \rightarrow \vec{A}(\vec{R}, t)' = \vec{A}(\vec{R}, t) + \vec{\nabla} \Lambda(\vec{R}, t). \quad (5.5.5)$$

Thus, a gauge transformation in quantum mechanics is defined as the simultaneous changes in eqs. (5.5.3)–(5.5.5) with the same  $\Lambda$ . Primes are used to indicate the state ket and operators



after the gauge transformation. For any operator  $\Omega$ , the definition of the gauge transformation

$$\Omega \rightarrow \Omega' \quad (5.5.6)$$

is obtained by applying eq. (5.5.4) and (5.5.5) to the potentials appearing explicitly in the definition of  $\Omega$ . We now show that physical predictions are unaffected by such a transformation.

We start by noting that a prerequisite for an operator  $\Omega$  to be a **gauge-invariant observable** is that its expectation values should be the same before and after the gauge transformation,

$$\langle \psi | \Omega | \psi \rangle = \langle \psi' | \Omega' | \psi' \rangle. \quad (5.5.7)$$

Requiring this to be true for all state kets  $|\psi\rangle$ , and using eq. (5.5.3), we obtain the defining property of a gauge-invariant observable,

$$\Omega' = U_\Lambda \Omega U_\Lambda^\dagger. \quad (5.5.8)$$

Not all Hermitian operators that would otherwise satisfy the requirements of being an observable will have this property.

For example, consider the canonical operators  $\vec{R}$  and  $\vec{P}$ . Since they do not depend explicitly on the potentials, we have

$$\vec{R}' = \vec{R}, \quad \vec{P}' = \vec{P}. \quad (5.5.9)$$

Meanwhile, using the definition of  $U_\Lambda$  in eq. (5.5.2), we find

$$U_\Lambda \vec{R} U_\Lambda^\dagger = \vec{R}, \quad (5.5.10)$$

$$U_\Lambda \vec{P} U_\Lambda^\dagger = \vec{P} - \frac{q}{c} \vec{\nabla} \Lambda. \quad (5.5.11)$$

The first equation is an immediate consequence of the fact that  $U_\Lambda$  involves only  $\vec{R}$  and not  $\vec{P}$ , and the second follows from applying Theorem 2.4.4. Comparison of eqs. (5.5.9)–(5.5.11) with eq. (5.5.8) shows that  $\vec{R}$  is a gauge-invariant observable, but the canonical momentum operator  $\vec{P}$  is not. In contrast, it follows from eq. (5.5.5) that the gauge transformation of the kinetic momentum

$$\vec{\Pi} = \vec{P} - \frac{q}{c} \vec{A} \quad (5.5.12)$$

is  $\vec{\Pi}' = \vec{P} - \frac{q}{c} (\vec{A} + \vec{\nabla} \Lambda) = \vec{\Pi} - \frac{q}{c} \vec{\nabla} \Lambda$ . Using eq. (5.5.11), this can be rewritten as

$$\vec{\Pi}' = U_\Lambda \vec{\Pi} U_\Lambda^\dagger. \quad (5.5.13)$$

This establishes that the kinetic momentum  $\vec{\Pi}$  is a gauge-invariant observable, according to our defining requirement of eq. (5.5.8).

Since this was perhaps a bit tricky and unexpected, we reiterate. Even though the canonical momentum operator  $\vec{P}$  does not change when we do a gauge transformation, its expectation values do change, so it is not a gauge-invariant observable. And, although the kinetic momentum operator  $\vec{\Pi}$  changes when we do a gauge transformation, its expectation values do not, so it is a gauge-invariant observable. These are the quantum versions of the statements made about the gauge dependences of the classical quantities  $\vec{p}$  and  $\vec{\pi}$  following eq. (4.3.15).

The Hamiltonian operator in eq. (4.3.28) certainly changes when we do a gauge transformation, since it depends explicitly on the potentials  $\Phi$  and  $\vec{A}$ . Using eqs. (5.5.4) and (5.5.5) gives the gauge transformation of  $H$ ,

$$H \rightarrow H' = \frac{1}{2m} \left( \vec{P} - \frac{q}{c} \vec{A} - \frac{q}{c} \vec{\nabla} \Lambda \right)^2 + q\Phi - \frac{q}{c} \frac{\partial \Lambda}{\partial t} - \gamma \vec{S} \cdot \vec{B}. \quad (5.5.14)$$

Equations (5.5.10) and (5.5.11) allow us to rewrite this in the convenient form

$$H' = U_{\Lambda} H U_{\Lambda}^{\dagger} + i\hbar \left( \frac{\partial U_{\Lambda}}{\partial t} \right) U_{\Lambda}^{\dagger}. \quad (5.5.15)$$

Now, given the Schrödinger equation

$$i\hbar \frac{d}{dt} |\psi(t)\rangle = H |\psi(t)\rangle, \quad (5.5.16)$$

it is straightforward to use  $|\psi(t)'\rangle = U_{\Lambda} |\psi(t)\rangle$  and eq. (5.5.15) to obtain

$$i\hbar \frac{d}{dt} |\psi(t)'\rangle = H' |\psi(t)'\rangle. \quad (5.5.17)$$

This shows the key result that the Schrödinger equation for time evolution is also satisfied if  $H$  and  $|\psi(t)\rangle$  are replaced by their gauge-transformed counterparts.

Another consequence of eq. (5.5.15) is that the Hamiltonian in eq. (4.3.28) is not, in general, a gauge-invariant observable, due to the presence of the last term. However, the requirement  $H' = U_{\Lambda} H U_{\Lambda}^{\dagger}$  is satisfied for the subset of gauge transformations such that  $\partial \Lambda / \partial t = 0$ . Thus, if we limit the gauge transformations to those that do not depend explicitly on time, then  $H$  is a gauge-invariant observable in that restricted sense. Also, if the potentials  $\Phi$  and  $\vec{A}$  do not depend on time, then according to the results at the end of section 5.1,  $H$  is a conserved quantity, the total energy of the particle.

If the gauge non-invariance of  $H$  for time-dependent  $\Lambda$  bothers you, note that a  $\Lambda$  that is linear in time and independent of position just adds a constant term to  $\Phi$  and thus to  $H$ . This corresponds to the classical freedom to add a constant to the energy without affecting the equations of motion at all.

The defining requirement for gauge-invariant observables, eq. (5.5.8), was obtained by requiring that expectation values do not depend on the choice of gauge. Now we will use eq. (5.5.8) to

prove the stronger result that all probabilities for measurements of such observables are gauge independent. Consider a gauge-invariant observable  $\Omega$ , and choose an orthobasis of its eigenkets with degeneracy labels  $u_\omega$ , so that

$$\Omega|\omega, u_\omega\rangle = \omega|\omega, u_\omega\rangle. \quad (5.5.18)$$

According to Postulates 3 and 4, the probabilities for allowed results of the measurement of  $\Omega$  in a state  $|\psi\rangle$  are

$$\mathcal{P}(\omega) = \sum_{u_\omega} |\langle\omega, u_\omega|\psi\rangle|^2. \quad (5.5.19)$$

Now, we observe that

$$\Omega' U_\Lambda |\omega, u_\omega\rangle = U_\Lambda \Omega U_\Lambda^\dagger U_\Lambda |\omega, u_\omega\rangle = U_\Lambda \Omega |\omega, u_\omega\rangle = \omega U_\Lambda |\omega, u_\omega\rangle, \quad (5.5.20)$$

which shows that the states  $U_\Lambda |\omega, u_\omega\rangle$  are eigenkets of  $\Omega'$ , with the same eigenvalues  $\omega$  and the same degeneracies. Therefore, we can compute the probability to obtain the result  $\omega$  from a measurement of  $\Omega'$  in the gauge-transformed description as

$$\mathcal{P}(\omega)' = \sum_{u_\omega} |\langle\omega, u_\omega|U_\Lambda^\dagger|\psi'\rangle|^2 = \sum_{u_\omega} |\langle\omega, u_\omega|U_\Lambda^\dagger U_\Lambda|\psi\rangle|^2 = \sum_{u_\omega} |\langle\omega, u_\omega|\psi\rangle|^2 = \mathcal{P}(\omega). \quad (5.5.21)$$

Thus, we have succeeded in our goal of showing that the predictions for measurements of gauge-invariant observables are not changed by the gauge transformation. This is in accord with the general principle that gauge transformations affect our equations in intermediate steps of calculations, but do not change the final results that reflect physical reality.

For simplicity, in the preceding we have treated the case of a single particle with charge  $q$ . In the case of more than one particle, with charges  $q_n$  and masses  $m_n$ , the same discussion goes through, with Hamiltonian

$$H = \sum_n \left( \frac{1}{2m_n} \left[ \vec{P}_n - \frac{q_n}{c} \vec{A}(\vec{R}_n, t) \right]^2 + q_n \Phi(\vec{R}_n, t) - \gamma_n \vec{S}_n \cdot \vec{B}(\vec{R}_n, t) \right), \quad (5.5.22)$$

subject to gauge transformations as given by eqs. (5.5.3)–(5.5.5), but with

$$U_\Lambda = \exp \left[ \frac{i}{\hbar c} \sum_n q_n \Lambda(\vec{R}_n, t) \right]. \quad (5.5.23)$$

The gauge transformation of the wavefunction in the position representation,  $\psi(\vec{r}_1, \dots, \vec{r}_n) = \langle \vec{r}_1, \dots, \vec{r}_n | \psi \rangle$ , is therefore given by

$$\psi(\vec{r}_1, \dots, \vec{r}_n) \rightarrow \exp \left[ \frac{i}{\hbar c} \sum_n q_n \Lambda(\vec{r}_n, t) \right] \psi(\vec{r}_1, \dots, \vec{r}_n). \quad (5.5.24)$$

This consists of multiplication by a phase that may depend on time and the particle positions. Thus, gauge invariance can be viewed as the statement that multiplying the wavefunction for a system of charged particles by a phase of this special form can be compensated by a redefinition of the electromagnetic potentials. In practice, one may avoid this redundancy in the description of the physical system by fixing the gauge. This just means that one chooses a specific form of the potentials, or at least imposes some condition on them that will not be satisfied if one does an arbitrary gauge transformation.

It is also possible to turn things around, and view the gauge transformations as the starting point that determines the interactions. In quantum mechanics, the physics is always independent of a change in the conventional choice of global phase of the wavefunction. If we generalize this to a local change in phase, as in eq. (5.5.24), then we are led to introduce the electromagnetic potentials to compensate, and thus to the necessity of interactions of charged particles with electromagnetic fields. Although it is beyond our scope here, this approach allows the interactions of the strong and weak nuclear forces to be determined by their non-Abelian gauge transformation groups,  $SU(3)$  and  $SU(2) \times U(1)$ , respectively.

## 5.6 Currents and local conservation of probability

From Postulate 4, and the completeness of position eigenstates, we know that the probability to find a particle within an infinitesimal volume  $d^3\vec{r}$  is given by the Born rule,

$$d\mathcal{P} = \rho(\vec{r}, t) d^3\vec{r}, \quad (5.6.1)$$

where the probability density per unit volume is

$$\rho(\vec{r}, t) = |\langle \vec{r} | \psi(t) \rangle|^2 = |\psi(\vec{r}, t)|^2. \quad (5.6.2)$$

Assuming the total probability of finding the particle somewhere is fixed and equal to 1, there must be a law of conservation of probability. The local form of this law is a differential equation

$$\frac{\partial \rho}{\partial t} = -\vec{\nabla} \cdot \vec{J}, \quad (5.6.3)$$

where  $\vec{J}$  is a probability current density. The left side of this equation is the rate at which probability density is accumulating at a point, which the right side tells us is the negative of a source for the vector field  $\vec{J}$ .

To prove eq. (5.6.3), and identify the current density, start with the Schrödinger equation in the position representation with wavefunction  $\psi$ , and multiply by  $-i\psi^*/\hbar$ ,

$$\psi^* \frac{\partial}{\partial t} \psi = \frac{i\hbar}{2m} \psi^* \nabla^2 \psi - \frac{i}{\hbar} V \psi^* \psi. \quad (5.6.4)$$

Adding this to its complex conjugate, we get

$$\frac{\partial \rho}{\partial t} = \frac{i\hbar}{2m} (\psi^* \nabla^2 \psi - \psi \nabla^2 \psi^*), \quad (5.6.5)$$

in which the contribution proportional to  $V(\vec{r})$  has canceled due to the fact that the potential is real. Now if we define the probability current density by

$$\vec{J} \equiv \frac{i\hbar}{2m} (\psi \vec{\nabla} \psi^* - \psi^* \vec{\nabla} \psi), \quad (5.6.6)$$

then we have

$$\vec{\nabla} \cdot \vec{J} = \frac{i\hbar}{2m} (\psi \nabla^2 \psi^* - \psi^* \nabla^2 \psi), \quad (5.6.7)$$

where two terms proportional to  $\vec{\nabla} \psi^* \cdot \vec{\nabla} \psi$  have canceled. Comparison of eqs. (5.6.5) and (5.6.7) establishes the local conservation of probability, eq. (5.6.3).

In the case of a charged particle, an important modification is needed if the vector potential  $\vec{A}$  is non-zero. Starting with the Hamiltonian in eq. (4.3.28), one can show by steps similar to those followed in eqs. (5.6.4)–(5.6.7) that the current density satisfying  $\vec{\nabla} \cdot \vec{J} = -\partial \rho / \partial t$  is

$$\vec{J} = \frac{1}{m} \text{Re} \left[ \psi^* \left( -i\hbar \vec{\nabla} - \frac{q}{c} \vec{A} \right) \psi \right], \quad (5.6.8)$$

generalizing eq. (5.6.6). Here,  $-i\hbar \vec{\nabla} - \frac{q}{c} \vec{A}$  is the position representation of the kinetic momentum operator,  $\vec{\Pi}$ , which was introduced in eq. (4.3.29) of section 4.3. This expression for  $\vec{J}$  is invariant under gauge transformations, and is related to the electric current density  $\vec{j}$  by  $\vec{j} = q\vec{J}$ , where  $q$  is the electric charge of the particle.

As an example, consider a free particle in a plane-wave simultaneous eigenstate of momentum and energy with eigenvalues  $\vec{p} = \hbar \vec{k}$  and  $E = \hbar^2 k^2 / 2m$ ,

$$\psi(\vec{r}, t) = C e^{i\vec{k} \cdot \vec{r} - iEt/\hbar}. \quad (5.6.9)$$

In terms of the complex normalization constant  $C$ , the probability and current densities are just constants in both time and position,

$$\rho = |C|^2, \quad \vec{J} = \frac{i\hbar}{2m} (-i\vec{k} - i\vec{k}) |C|^2 = \frac{\vec{p}}{m} \rho. \quad (5.6.10)$$

If the domain of the particle is all space, then the wavefunction is not normalizable to unity for any finite  $C$ , but at least the ratio of the current density to the probability density is well-defined and equal to the velocity eigenvalue of the particle. Although the probability density is constant, it is constantly flowing in the direction of  $\vec{v} = \vec{p}/m$ . So, in the case of a free particle,

$$\frac{\partial \rho}{\partial t} = 0, \quad -\vec{\nabla} \cdot \vec{J} = 0, \quad (5.6.11)$$

satisfying the local conservation of probability in the most trivial possible way.

In the case of one-dimensional problems, the probability density and current are

$$\rho = |\psi(x, t)|^2, \quad J = \frac{i\hbar}{2m} \left( \psi \frac{\partial \psi^*}{\partial x} - \psi^* \frac{\partial \psi}{\partial x} \right), \quad (5.6.12)$$

and the statement of local conservation of probability is

$$\frac{\partial \rho}{\partial t} = -\frac{\partial J}{\partial x}. \quad (5.6.13)$$

The current density vector is just a number in the one-dimensional case, and is positive for probability density flowing to the right, and negative for flow to the left.

A key ingredient in establishing conservation of probability was the reality of the potential  $V$  in the position representation, or more generally the hermiticity of the Hamiltonian operator. This is also directly related to the unitarity of the time evolution operator  $U(t, t_0)$  that we studied in section 3.4. However, in some situations, one can think of probability as *not* being conserved. For example, suppose that we are using quantum mechanics to model the behavior of an unstable particle. If our Hilbert space only describes the unstable particle, and not the other particles that it decays into, then we should expect that the total integrated probability to find it somewhere should decrease with time. To describe situations like this, we can break the law for a good cause, by taking the Hamiltonian to be non-Hermitian.

To see how this works, suppose that we defy the authority of Postulate 6 by taking  $H$  to have complex eigenvalues, with corresponding normalized eigenkets  $|\phi_n\rangle$ , so that

$$H |\phi_n\rangle = (E_n - i\Gamma_n/2) |\phi_n\rangle, \quad (5.6.14)$$

where  $E_n$  and  $\Gamma_n$  are real numbers. Of course, this is only possible if  $H$  is not Hermitian, due to Theorem 2.6.3. If the state of the system at time  $t = 0$  is one of them,  $|\psi(0)\rangle = |\phi_n\rangle$ , with unit norm, then the time evolution predicted by the Schrödinger equation will be non-unitary,

$$|\psi(t)\rangle = e^{-i(E_n - i\Gamma_n/2)t/\hbar} |\phi_n\rangle. \quad (5.6.15)$$

The norm of this ket as a function of time is therefore

$$\langle \psi(t) | \psi(t) \rangle = e^{-\Gamma_n t/\hbar}. \quad (5.6.16)$$

We can then interpret the squared norm of the ket as the probability that the particle exists at time  $t > 0$ , given that it existed at time  $t = 0$ . [Compare to eq. (3.1.3) with  $P_\alpha = I$ .] The mean lifetime of our unstable state  $|\phi_n\rangle$  is thus  $\tau = \hbar/\Gamma_n$ , where  $\Gamma_n/2$  is the negative of the imaginary part of the Hamiltonian eigenvalue.

The unstable states could correspond to particles that undergo spontaneous decay, such as the neutron, the muon, or many atomic nuclei. The non-Hermitian-Hamiltonian formalism could also be applied to a set of atomic states, if our quantum treatment does not include the complete set of states to which they could decay, including photons released in the process. It could even apply in a description of a single absolutely stable particle like the electron, if it can be removed from the system under discussion by a process like electron-capture inverse beta decay, which in some nuclei occurs as  $e^-p \rightarrow \nu_e n$ . This would correspond to an effective potential  $V(\vec{r})$  with negative imaginary parts localized at the nuclei. By repeating the steps of eqs. (5.6.4)-(5.6.7), one can see that, in the case of a non-Hermitian potential, the equation that governs the local probability density and current is

$$\frac{\partial \rho}{\partial t} = -\vec{\nabla} \cdot \vec{J} + \frac{2}{\hbar} \text{Im}[V(\vec{r})]\rho. \quad (5.6.17)$$

In all such cases of particles decaying to other particles, or interacting in such a way as to change their numbers, the use of a non-Hermitian Hamiltonian is really a sign of an incomplete description. There is always a more fundamental description in which the *complete* Hamiltonian will be Hermitian. Quantum field theories provide the natural way to incorporate processes that change the numbers of particles, in such a way that the postulates of quantum mechanics hold, including unitary time evolution as predicted by Postulate 6.

## 5.7 Exercises

**Exercise 5.1.** Show that the components of the position, momentum, and angular momentum operators for a particle satisfy

$$[L_a, R_b] = i\hbar\epsilon_{abc}R_c, \quad [L_a, P_b] = i\hbar\epsilon_{abc}P_c, \quad (5.7.1)$$

for all  $a, b, c = 1, 2, 3$ , where  $R_1 = X$ ,  $R_2 = Y$ , and  $R_3 = Z$ , and  $P_1 = P_x$ ,  $P_2 = P_y$ , and  $P_3 = P_z$ . Use these results, and eq. (5.1.3), to obtain the transformed position and momentum operators  $\vec{R}'$  and  $\vec{P}'$  resulting from a rotation  $U = e^{-i\alpha L_z/\hbar}$ , to first order in  $\alpha$ .

**Exercise 5.2.** For a system of two particles labeled 1 and 2 as discussed in section 4.2, show that the total angular momentum can be written in the way claimed in eq. (5.3.35) in terms of  $\vec{L}$  and  $\vec{L}_{\text{cm}}$  defined in eq. (5.3.36). Show that  $\vec{L}$  and  $\vec{L}_{\text{cm}}$  each obey the defining requirement of an angular momentum operator in eq. (5.3.33).

**Exercise 5.3.** Derive eq. (5.6.8), the probability current density  $\vec{J}$  in the presence of electromagnetic potentials  $\Phi$  and  $\vec{A}$ . Show that it is invariant under gauge transformations.

## 6 Particle moving in one dimension

### 6.1 Gaussian wavefunctions

Consider a particle moving in one dimension in the domain  $-\infty < x < \infty$ . Suppose that at some moment in time, the position wavefunction has a Gaussian form,

$$\langle x|\psi\rangle = \psi(x) = N \exp \left[ -(x-a)^2/4\sigma^2 \right]. \quad (6.1.1)$$

Here  $a$  and  $\sigma$  are constants with units of [length], corresponding respectively to the center and the width of the probability density  $|\psi(x)|^2$ . More precisely, the full width at half maximum (FWHM) of the Gaussian peak of the probability density is  $2\sqrt{2 \ln 2} \sigma \approx 2.35\sigma$ . The factor  $N$  is a normalization. If we require the ket  $|\psi\rangle$  to have unit norm, we need

$$1 = |N|^2 \int_{-\infty}^{\infty} dx \exp \left[ -(x-a)^2/2\sigma^2 \right] = |N|^2 \sqrt{2\pi}\sigma, \quad (6.1.2)$$

so we can choose  $N = e^{i\theta}/(2\pi\sigma^2)^{1/4}$ , where  $\theta$  is any phase. If  $\theta$  is a constant, then it is a global phase and has no physical significance at all, so let us choose the next simplest possibility, that  $\theta$  depends linearly on  $x$ . The Gaussian wavefunction that we choose to study is thus

$$\psi(x) = \frac{1}{(2\pi\sigma^2)^{1/4}} e^{ikx} \exp \left[ -(x-a)^2/4\sigma^2 \right] \quad (6.1.3)$$

for some constant  $k$ , whose interpretation will become clear soon. We will not need to assume anything in particular about the Hamiltonian of the system in this section.

According to Postulate 4, the probability of finding the particle between  $x$  and  $x + dx$  is

$$d\mathcal{P}(x) = |\langle x|\psi\rangle|^2 dx = \frac{1}{\sqrt{2\pi}\sigma} \exp \left[ -(x-a)^2/2\sigma^2 \right] dx. \quad (6.1.4)$$

Therefore, the probability of finding the particle in a range  $b < x < c$  is

$$\mathcal{P}(b < x < c) = \frac{1}{\sqrt{2\pi}\sigma} \int_b^c dx \exp \left[ -(x-a)^2/2\sigma^2 \right]. \quad (6.1.5)$$

We can also find the expectation value of  $X$  in the state  $|\psi\rangle$ ,

$$\begin{aligned} \langle X \rangle &= \langle \psi|X|\psi \rangle = \int_{-\infty}^{\infty} dx \langle \psi|X|x \rangle \langle x|\psi \rangle = \int_{-\infty}^{\infty} dx x |\psi(x)|^2 \\ &= \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^{\infty} dx x \exp \left[ -(x-a)^2/2\sigma^2 \right] = \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^{\infty} du (u+a) \exp \left[ -u^2/2\sigma^2 \right] \\ &= a. \end{aligned} \quad (6.1.6)$$

The second equality uses the completeness relation, the third uses  $X|x\rangle = x|x\rangle$ , and the fourth uses the common trick of “completing the square”, which means that we define a new integration



variable  $u$  so that the exponential in the integrand depends only on  $u^2$  (with no linear term in  $u$ ). Here, the change of variable was  $x = u + a$ . Similarly, we have

$$\langle X^2 \rangle = \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^{\infty} du (u + a)^2 \exp[-u^2/2\sigma^2] = a^2 + \sigma^2. \quad (6.1.7)$$

Therefore, the uncertainty of  $X$  for this state is

$$\Delta X = \sqrt{\langle X^2 \rangle - \langle X \rangle^2} = \sigma. \quad (6.1.8)$$

So far, the constant  $k$  has not made any difference at all.

The momentum wavefunction for the same state is

$$\tilde{\psi}(p) = \langle p|\psi \rangle = \int_{-\infty}^{\infty} dx \langle p|x \rangle \langle x|\psi \rangle = \frac{1}{\sqrt{2\pi\hbar}(2\pi\sigma^2)^{1/4}} \int_{-\infty}^{\infty} dx e^{-ipx/\hbar} e^{ikx} e^{-(x-a)^2/4\sigma^2}, \quad (6.1.9)$$

where we have used the complex conjugate of eq. (2.8.39). We now use the completing-the-square trick again; this time the appropriate change of variables is  $x = u + b$ , where we choose  $b = a + i2\sigma^2(k - p/\hbar)$ , so

$$\int_{-\infty}^{\infty} dx e^{-ipx/\hbar} e^{ikx} e^{-(x-a)^2/4\sigma^2} = e^{(b^2-a^2)/4\sigma^2} \int_{-\infty}^{\infty} du e^{-u^2/4\sigma^2} = 2\sqrt{\pi}\sigma e^{(b^2-a^2)/4\sigma^2}. \quad (6.1.10)$$

It follows that

$$\tilde{\psi}(p) = \left( \frac{2\sigma^2}{\pi\hbar^2} \right)^{1/4} e^{ia(k-p/\hbar)} e^{-\sigma^2(p/\hbar-k)^2}. \quad (6.1.11)$$

The last factor shows that a state with a Gaussian position wavefunction also has a Gaussian momentum wavefunction, centered at  $p = \hbar k$ . The momentum wavefunction also contains a phase that depends on  $p$ ; this factor encodes the information about the center of the position wavefunction Gaussian peak,  $a$ .

It is now clear that our state depends on three physically significant parameters: the average momentum  $\hbar k$ , the position center  $a$ , and the position width  $\sigma$ , which is also the uncertainty in  $X$ . There is a duality between the position and momentum wavefunctions, for if one of them has a Gaussian magnitude with a linear phase, then so does the other, with parameters that are related by comparing eqs. (6.1.3) and (6.1.11). In particular, the widths of the position and momentum Gaussian wavefunctions are inversely proportional.

Using eq. (6.1.11), one can now obtain

$$\langle P \rangle = \langle \psi|P|\psi \rangle = \int_{-\infty}^{\infty} dp \langle \psi|P|p \rangle \langle p|\psi \rangle = \int_{-\infty}^{\infty} dp p |\tilde{\psi}(p)|^2 = \hbar k. \quad (6.1.12)$$

Similarly,

$$\langle P^2 \rangle = \int_{-\infty}^{\infty} dp p^2 |\tilde{\psi}(p)|^2 = \hbar^2(k^2 + 1/4\sigma^2). \quad (6.1.13)$$

It follows that the momentum uncertainty in the state with a Gaussian position wavefunction is

$$\Delta P = \sqrt{\langle P^2 \rangle - \langle P \rangle^2} = \hbar/2\sigma, \quad (6.1.14)$$

and so the product of the position and momentum uncertainties is

$$(\Delta X)(\Delta P) = \hbar/2. \quad (6.1.15)$$

This result does not depend on the center of the Gaussian  $a$ , or its width  $\sigma$ , or its average momentum  $\hbar k$ . Thus, Gaussian wavefunctions always realize the minimum possible product of uncertainties consistent with the general position-momentum uncertainty relation, eq. (3.3.10).

## 6.2 Time evolution of free-particle state in one dimension

For a free particle with mass  $m$  moving in one dimension, the Schrödinger equation is

$$i\hbar \frac{d}{dt} |\psi\rangle = H |\psi\rangle = \frac{P^2}{2m} |\psi\rangle, \quad (6.2.1)$$

where  $P$  is the momentum operator. Since  $[H, P] = 0$ , we know that there is an orthobasis of stationary states that are also eigenstates of  $P$  with eigenvalue  $p$ . Acting on such states,  $E = P^2/2m = p^2/2m$ . Therefore, for a given  $E$  there are exactly two solutions,  $p = \pm\sqrt{2mE}$ , and the plane-wave stationary states can be labeled

$$|E, R\rangle = |p = \sqrt{2mE}\rangle, \quad |E, L\rangle = |p = -\sqrt{2mE}\rangle, \quad (6.2.2)$$

where  $R, L$  is a degeneracy label that tells us whether the particle is moving right or left.

The time-dependent wavefunction for a stationary state with momentum  $p$  is, combining eq. (2.8.39) with the time-evolution phase factor  $e^{-iEt}$  gives

$$\psi_p(x, t) = \frac{1}{\sqrt{2\pi\hbar}} e^{i(kx - \omega t)}, \quad (6.2.3)$$

where  $k = p/\hbar$  and  $\omega = E/\hbar$ . The position of constant phase is  $x = \omega t/k = Et/p = pt/2m$ , so the **phase velocity** of one of these waves is

$$v_{\text{phase}} = \omega/k = p/2m, \quad (6.2.4)$$

which is half of the classical velocity  $p/m$ .

To understand the classical speed of propagation, one must consider the **group velocity** for wavepacket superpositions of states with a continuous distribution of  $k$ . It is a general feature of wave kinematics that, in the presence of dispersion (that is,  $\omega$  depending nonlinearly on  $k$ ), the velocity for a wavepacket is not  $\omega/k$ , but instead

$$v_{\text{group}} = \partial\omega/\partial k. \quad (6.2.5)$$

Let us pause from our discussion of the free particle to see how this arises in a general context. Consider a packet of waves described by the wavefunction

$$\psi(x, t) = \int dk a(k) e^{i(\omega t - kx)}, \quad (6.2.6)$$

where, in general, the angular frequency  $\omega(k)$  and the amplitude function  $a(k)$  both depend on the wavenumber in some arbitrary way. Now suppose that  $a(k)$  is such that only a narrow range of  $k$  near  $k_0$  contribute appreciably, so that we can write  $k = k_0 + \Delta k$  and

$$\omega(k) = \omega_0 + \Delta k \left. \frac{\partial \omega}{\partial k} \right|_{k=k_0} + \mathcal{O}(\Delta k)^2, \quad (6.2.7)$$

where  $\omega_0 = \omega(k_0)$ . Then eq. (6.2.6) can be rewritten in the form

$$\psi(x, t) = e^{i(\omega_0 t - k_0 x)} A(x, t), \quad (6.2.8)$$

which is the product of a global phase and a factor describing the shape of the magnitude of the wavepacket,

$$A(x, t) = \int d(\Delta k) a(k_0 + \Delta k) \exp \left[ i \left( \frac{\partial \omega}{\partial k} t - x \right) \Delta k + \dots \right]. \quad (6.2.9)$$

Here the partial derivative is understood to be evaluated at  $k = k_0$ . In this approximation, the magnitude  $|\psi(x, t)| = |A(x, t)|$  depends on position and time only through the combination

$$\frac{\partial \omega}{\partial k} t - x, \quad (6.2.10)$$

which immediately implies that the velocity of the wavepacket is indeed  $v_{\text{group}} = \partial \omega / \partial k$ .

Applying this to the special case of a free particle, with  $\omega = E/\hbar$ , we have

$$v_{\text{group}} = \frac{\partial}{\partial k} \left( \frac{\hbar k^2}{2m} \right) = \hbar k / m = p / m, \quad (6.2.11)$$

which is the classical value. For wavepackets in quantum mechanics with momentum sharply peaked near  $p$ , this group velocity corresponds to the motion of the expectation value of the position, as can be seen in general from eq. (3.4.18). Before the end of this section, we will verify this for the particular case of a Gaussian superposition of plane waves.

Using eq. (3.4.7), the unitary time evolution operator is

$$U(t) = \int_{-\infty}^{\infty} dp |p\rangle \langle p| e^{-itp^2/2m\hbar}. \quad (6.2.12)$$

The matrix element of this operator between different position eigenstates is therefore

$$\langle x | U(t) | x' \rangle = \int_{-\infty}^{\infty} dp \langle x | p \rangle \langle p | x' \rangle e^{-itp^2/2m\hbar} = \frac{1}{2\pi\hbar} \int_{-\infty}^{\infty} dp e^{ip(x-x')/\hbar} e^{-itp^2/2m\hbar}. \quad (6.2.13)$$

This integral is again done by the completing-the-square trick, this time by changing the integration variable to  $p' = p - m(x - x')/t$ , with the result

$$\langle x|U(t)|x'\rangle = \left(\frac{m}{2\pi\hbar it}\right)^{1/2} e^{im(x-x')^2/2\hbar t}. \quad (6.2.14)$$

By now applying the completeness relation, the time dependence of the wavefunction for a free particle reduces to an integral involving the wavefunction at time  $t = 0$ ,

$$\begin{aligned} \psi(x, t) &= \langle x|\psi(t)\rangle = \langle x|U(t)|\psi(0)\rangle = \int_{-\infty}^{\infty} dx' \langle x|U(t)|x'\rangle \langle x'|\psi(0)\rangle \\ &= \left(\frac{m}{2\pi\hbar it}\right)^{1/2} \int_{-\infty}^{\infty} dx' e^{im(x-x')^2/2\hbar t} \psi(x', 0). \end{aligned} \quad (6.2.15)$$

In principle, the time evolution of an arbitrary free-particle state has therefore been solved.

As a good example for which the integral can be done analytically, consider the motion and spreading with time of the wavefunction of a Gaussian wavepacket state. We start at time  $t = 0$  with a state  $|\psi(0)\rangle$  that has

$$\psi(x', 0) = \frac{1}{(2\pi\sigma^2)^{1/4}} e^{ip_0 x'/\hbar} e^{-x'^2/4\sigma^2}, \quad (6.2.16)$$

which as we saw in the previous section can be interpreted as having center at  $x' = 0$ , width  $\sigma$ , and average momentum  $p_0$ . At time  $t$ , eq. (6.2.15) gives

$$\psi(x, t) = \left(\frac{m}{2\pi\hbar it}\right)^{1/2} \frac{1}{(2\pi\sigma^2)^{1/4}} \int_{-\infty}^{\infty} dx' \exp \left[ -\frac{x'^2}{4\sigma^2} + i\frac{m(x-x')^2}{2\hbar t} + i\frac{p_0 x'}{\hbar} \right]. \quad (6.2.17)$$

By the usual completing-the-square integration variable change trick, this becomes, after some algebraic manipulation,

$$\psi(x, t) = \frac{1}{\sqrt{\sqrt{2\pi}(\sigma + i\hbar t/2m\sigma)}} \exp \left[ -\frac{(x - p_0 t/m)^2}{4\sigma^2 + 2i\hbar t/m} \right] \exp \left[ i\frac{p_0}{\hbar} \left( x - \frac{p_0 t}{2m} \right) \right], \quad (6.2.18)$$

or, after further rearrangement,

$$\psi(x, t) = \frac{e^{i\theta(x, t)}}{\sqrt{\sqrt{2\pi}(\sigma + i\hbar t/2m\sigma)}} \exp \left[ -\frac{(x - p_0 t/m)^2}{4\sigma^2 + \hbar^2 t^2/m^2 \sigma^2} \right], \quad (6.2.19)$$

where

$$\theta(x, t) = \frac{p_0}{\hbar} \left( x - \frac{p_0 t}{2m} \right) + \frac{\hbar t (x - p_0 t/m)^2}{2m(4\sigma^2 + \hbar^2 t^2/m^2 \sigma^2)}. \quad (6.2.20)$$

The probability density at time  $t$  is therefore

$$|\psi(x, t)|^2 = \frac{1}{\sqrt{2\pi(\sigma^2 + \hbar^2 t^2/4m^2 \sigma^2)}} \exp \left[ -\frac{(x - p_0 t/m)^2}{2(\sigma^2 + \hbar^2 t^2/4m^2 \sigma^2)} \right]. \quad (6.2.21)$$

This is a Gaussian function of  $x$  whose peak moves to the right at exactly the constant speed  $p_0/m$  of naive classical expectation, but whose width grows with time  $t$ . Using the results of the previous section, we obtain

$$\langle X \rangle = p_0 t / m, \quad (6.2.22)$$

$$\Delta X = \sqrt{\sigma^2 + \hbar^2 t^2 / 4m^2 \sigma^2}. \quad (6.2.23)$$

The speed at which the expectation value  $\langle X \rangle$  moves is also the group velocity  $v_{\text{group}} = \partial\omega/\partial k$ . At large  $t$ ,  $\Delta X \approx \hbar t / 2m\sigma$  grows linearly with time, and ironically is larger for smaller  $\sigma$ ; the more we try to confine the wavefunction initially, the more spread out the particle's probability density support will be at late times. The explanation is that, due to the uncertainty relation eq. (3.3.10), a highly constrained particle position has larger amplitudes for momenta that deviate from the central value.

One can also compute the momentum wavefunction as a function of time, by applying eq. (2.8.41) to eq. (6.2.18). After another integration and more algebraic juggling, one finds

$$\tilde{\psi}(p, t) = \left( \frac{2\sigma^2}{\pi\hbar^2} \right)^{1/4} \exp \left[ -\sigma^2 (p - p_0)^2 / \hbar^2 \right] e^{ip^2 t / 2\hbar m}. \quad (6.2.24)$$

This is remarkably simple; the time dependence is entirely in the complex phase, so

$$|\tilde{\psi}(p, t)|^2 = \sqrt{\frac{2}{\pi}} \frac{\sigma}{\hbar} \exp \left[ -2\sigma^2 (p - p_0)^2 / \hbar^2 \right]. \quad (6.2.25)$$

This does not depend on time at all, despite the fact that the width of the support of the position wavefunction grows with time. This is an illustration of Theorem 5.1.1; because the momentum is a conserved quantity for a free particle Hamiltonian, the probability density to measure the momentum between  $p$  and  $p + dp$  is a constant in time.

### 6.3 Properties of stationary states in one-dimensional potentials

Consider a particle moving in one dimension in a potential that is given classically by  $V(x)$ , so that the Hamiltonian operator is

$$H = \frac{P^2}{2m} + V(X). \quad (6.3.1)$$

In general, we want to find stationary states,

$$H |\psi_E\rangle = E |\psi_E\rangle. \quad (6.3.2)$$

To accomplish this, we use the position representation, in which  $X \rightarrow x$  and  $P \rightarrow -i\hbar d/dx$ , by multiplying on the left by  $\langle x|$ . Then the wavefunction  $\psi_E(x) = \langle x|\psi_E\rangle$  obeys the eigenvalue

differential equation

$$\frac{d^2}{dx^2}\psi_E(x) = \frac{2m}{\hbar^2}[V(x) - E]\psi_E(x). \quad (6.3.3)$$

This is the time-independent Schrödinger equation for a spinless particle in one dimension.

Before exploring some special cases, it is helpful to make some general statements about this problem. First, consider a range of positions  $x$  over which the potential  $V(x)$  is less than the energy eigenvalue  $E$ . This is called a classically allowed range, because in classical mechanics the kinetic energy contribution is always positive, so that  $E > V(x)$ . In a classically allowed range, the wavefunction is oscillatory provided that  $V(x)$  varies sufficiently smoothly with  $x$ . To illustrate this, consider the case of a nearly constant  $V$ , and define the wavenumber  $k = \sqrt{2m(E - V)}/\hbar$ . Then, within that range, the Schrödinger equation becomes (dropping the subscript  $E$  on the wavefunction from here on)  $\psi'' \approx -k^2\psi$ , which has the general solution

$$\psi \approx ae^{ikx} + be^{-ikx}, \quad (\text{for nearly constant } k^2 = 2m(E - V)/\hbar^2 > 0), \quad (6.3.4)$$

where  $a$  and  $b$  are constants.

Conversely, classically forbidden ranges of  $x$  are those in which  $E < V(x)$ . If  $V$  is nearly constant, we can define a real quantity  $\kappa = \sqrt{2m(V - E)}/\hbar$ , and the Schrödinger equation becomes  $\psi'' \approx \kappa^2\psi$ . The corresponding general solution is a sum of real exponentials,

$$\psi \approx ae^{-\kappa x} + be^{\kappa x}, \quad (\text{for nearly constant } \kappa^2 = 2m(V - E)/\hbar^2 > 0). \quad (6.3.5)$$

If the classically forbidden range includes  $x = \infty$ , then one must have  $b = 0$  in order to have a sensible wavefunction without exponential growth at large distances. If the classically forbidden range instead includes  $x = -\infty$ , then one must have  $a = 0$  for the same reason.

It is often the case that the potential approaches a constant at large distances. Unbound states are those that have  $E > V(x)$ , and therefore oscillatory behavior, at either  $x = \infty$ , or  $x = -\infty$ , or both. Bound states are those that have  $E < V(x)$  at both  $x = \pm\infty$ , implying exponentially falling wavefunctions at large distances.

A stationary state with energy  $E$  may have both classically allowed and classically forbidden ranges of  $x$ , depending on the potential. The points  $x$  with  $V(x) = E$  that separate them are called classical turning points, because the corresponding classical trajectory for a particle must turn back at those points.

It is also possible that  $V(x)$  has some special points where it may not be smooth, or may even diverge. We would like to know what can be said about the behavior of the wavefunction  $\psi(x)$  at such special points. Consider a particular special point  $x_0$ . Integrating eq. (6.3.3) with respect to  $x$  over a small neighborhood of that point gives

$$\int_{x_0-\epsilon}^{x_0+\epsilon} dx \frac{d}{dx} \left( \frac{d\psi}{dx} \right) = \psi'(x_0 + \epsilon) - \psi'(x_0 - \epsilon) = \frac{2m}{\hbar^2} \int_{x_0-\epsilon}^{x_0+\epsilon} dx [V(x) - E] \psi(x), \quad (6.3.6)$$

where the first equality follows from the fundamental theorem of calculus.

Suppose that  $V(x)$  is bounded near  $x = x_0$ . Then the right side of eq. (6.3.6) tends to 0 as  $\epsilon \rightarrow 0$ , so we can conclude that the first derivative of  $\psi(x)$  must be continuous at  $x_0$ . Note that this holds even if  $V(x)$  is discontinuous, as long as it stays finite in the neighborhood of  $x = x_0$ . It follows that  $\psi(x)$  itself is also continuous at  $x = x_0$ .

Another possibility is that  $V(x) = C\delta(x - x_0) + \dots$ , where  $C$  is a constant, and the ellipses represent a contribution that is possibly discontinuous but bounded near  $x = x_0$ . In that case, eq. (6.3.6) reveals that

$$\lim_{\epsilon \rightarrow 0} [\psi'(x_0 + \epsilon) - \psi'(x_0 - \epsilon)] = \frac{2m}{\hbar^2} C \psi(x_0), \quad (6.3.7)$$

so that the derivative of the wavefunction at  $x_0$  is discontinuous by an amount that we now know. Equation (6.3.7) is consistent with  $\psi'(x)$  being bounded in a neighborhood of  $x = x_0$ , even if it is not continuous. Then, integrating  $\int_{x_0 - \epsilon}^{x_0 + \epsilon} dx \psi'(x)$ , we obtain the continuity of the wavefunction,

$$\lim_{\epsilon \rightarrow 0} [\psi(x_0 + \epsilon) - \psi(x_0 - \epsilon)] = 0. \quad (6.3.8)$$

More generally, **continuity of the wavefunction** is a requirement that we always impose on physically sensible states. The idea is that because  $|\psi(x)|^2$  represents the probability density, its value at  $x = x_0$  has a unique physical meaning and so must not depend on whether  $x$  approaches  $x_0$  from above or below.

Yet another possibility is that  $V(x) = \infty$  for a whole range  $x < x_0$  but it is finite for  $x > x_0$ . In that case, the particle is forbidden to enter the region of infinite potential, and so  $\psi(x) = 0$  for  $x \leq x_0$ . Now, continuity of the probability density tells us that  $\psi(x_0) = 0$ , which acts as a boundary condition for  $\psi(x)$  in the range  $x \geq x_0$ . Of course, the same holds if the inequalities are reversed: if  $V(x) = \infty$  for  $x > x_0$  but it is finite for  $x < x_0$ , then  $\psi(x) = 0$  for  $x \geq x_0$ .

Bound states in one-dimensional potentials in quantum mechanics never have degenerate energy eigenvalues. To prove it, suppose there are two states  $|\psi_1\rangle$  and  $|\psi_2\rangle$  with the same energy eigenvalue  $E$ . In the position representation, the wavefunctions satisfy

$$-\frac{\hbar^2}{2m}\psi_1'' + V\psi_1 = E\psi_1, \quad -\frac{\hbar^2}{2m}\psi_2'' + V\psi_2 = E\psi_2. \quad (6.3.9)$$

Multiplying the first equation by  $\psi_2$  and the second by  $\psi_1$  and taking the difference gives  $\psi_2\psi_1'' - \psi_1\psi_2'' = 0$ , or

$$\frac{d}{dx}(\psi_2\psi_1' - \psi_1\psi_2') = 0, \quad (6.3.10)$$

so that integrating with respect to  $x$  gives

$$\psi_2\psi_1' - \psi_1\psi_2' = c \quad (6.3.11)$$

where  $c$  is a constant of integration. Now, by definition the bound states have  $\psi_1 = \psi_2 = 0$  when  $x = \pm\infty$ . So, evaluating eq. (6.3.11) at  $x = \infty$  in particular, we learn that  $c$  must be 0. Therefore  $\psi'_1/\psi_1 = \psi'_2/\psi_2$ , or

$$\frac{d}{dx}(\ln \psi_1) = \frac{d}{dx}(\ln \psi_2), \quad (6.3.12)$$

which integrates to

$$\ln(\psi_1) = \ln(\psi_2) + C \quad (6.3.13)$$

where  $C$  is another constant of integration. So,  $\psi_1 = e^C \psi_2$ , and since  $e^C$  is a non-zero constant,  $\psi_1$  and  $\psi_2$  are proportional and therefore represent the same state. Note that in this proof it is crucial that we are working in one dimension. In two or three dimensions, bound states certainly can have degenerate energies, as we will see by finding examples in Chapter 10.

## 6.4 Particle in a one-dimensional box

Consider a particle of mass  $m$  confined to a box of length  $L$ , so that its potential energy as a function of  $x$  is

$$V(x) = \begin{cases} 0 & \text{for } |x| < L/2, \\ \infty & \text{for } |x| > L/2. \end{cases} \quad (6.4.1)$$

Since this is an even function of  $x$ , we know from the discussion in section 5.4 that there must be simultaneous eigenstates of energy and parity. Let us find them.

In the region  $|x| < L/2$ , eq. (6.3.3) becomes

$$\psi'' = -\frac{2mE}{\hbar^2}\psi. \quad (6.4.2)$$

The general solution to this differential equation is

$$\psi(x) = Ae^{ikx} + Be^{-ikx}, \quad (6.4.3)$$

where  $A$  and  $B$  are constants and

$$k = \sqrt{2mE}/\hbar. \quad (6.4.4)$$

Because the particle is completely confined to the box region, one must have  $\psi(x) = 0$  for  $|x| \geq L/2$ , and so by continuity of the wavefunction,  $\psi(L/2) = 0$  and  $\psi(-L/2) = 0$ , or

$$Ae^{ikL/2} + Be^{-ikL/2} = 0, \quad (6.4.5)$$

$$Ae^{-ikL/2} + Be^{ikL/2} = 0. \quad (6.4.6)$$



This system of equations will have a non-trivial solution for  $A$  and  $B$  if and only if the matrix

$$M = \begin{pmatrix} e^{ikL/2} & e^{-ikL/2} \\ e^{-ikL/2} & e^{ikL/2} \end{pmatrix} \quad (6.4.7)$$

is not invertible, which means  $\text{Det}(M) = e^{ikL} - e^{-ikL} = 2i \sin(kL) = 0$ . The solutions to the particle-in-a-box eigenvalue problem therefore must have  $k = n\pi/L$ , where  $n$  is an integer. It follows that  $B = -e^{ikL}A = (-1)^{n+1}A$ .

For odd  $n$ , we have  $B = A$ , and so for  $-L/2 \leq x \leq L/2$ ,

$$\psi_n(x) = 2A \cos(n\pi x/L). \quad (6.4.8)$$

To normalize the states to unity, we require

$$1 = \int_{-\infty}^{\infty} dx |\psi_n(x)|^2 = 4|A|^2 \int_{-L/2}^{L/2} dx \cos^2(n\pi x/L) = 2|A|^2 L, \quad (6.4.9)$$

and so we can choose  $A = 1/\sqrt{2L}$ , yielding the even-parity solutions

$$\psi_n(x) = \sqrt{\frac{2}{L}} \cos(n\pi x/L) \quad (n = 1, 3, 5, \dots). \quad (6.4.10)$$

Note that we do not need to include negative  $n$ , because those just have the same wavefunction up to a physically irrelevant sign, and are therefore the same states. Similarly, for even  $n$ , one finds the odd-parity solutions

$$\psi_n(x) = \sqrt{\frac{2}{L}} \sin(n\pi x/L) \quad (n = 2, 4, 6, \dots). \quad (6.4.11)$$

Here, not only do we not need negative  $n$ , but also the case  $n = 0$  is excluded, because it would result in  $\psi_0(x) = 0$ , which would correspond to the null ket, which is not a physical state. For both odd and even  $n$ , eq. (6.4.4) gives the allowed energy levels:

$$E_n = \frac{\hbar^2 \pi^2 n^2}{2mL^2} \quad (n = 1, 2, 3, \dots). \quad (6.4.12)$$

These are non-degenerate and discrete, as is true for any bound-state solutions in one dimension.

The ground state energy for the particle in a box,  $E_1 = \hbar^2 \pi^2 / 2mL^2$ , becomes very large as the confining box is taken smaller ( $L \rightarrow 0$ ). This can be seen to be in accord with the uncertainty relation eq. (3.3.10), as follows. First, note that  $\langle P \rangle = 0$  for each of the stationary states; this can be obtained either by direct computation in terms of the wavefunction, or seen as a consequence of the parity selection rule discussed at the end of section 5.4. It follows that

$$\langle H \rangle = \frac{1}{2m} \langle P^2 \rangle = \frac{1}{2m} (\Delta P)^2. \quad (6.4.13)$$

Now the uncertainty relation eq. (3.3.10) says that  $(\Delta P)^2 \geq (\hbar/2\Delta X)^2$ , and it is also clear that  $\langle X \rangle = 0$ , again either by direct computation or as a consequence of the parity selection rule. Therefore,  $(\Delta X)^2 = \langle X^2 \rangle$ , and we have a bound

$$\langle H \rangle \geq \frac{\hbar^2}{8m^2 \langle X^2 \rangle}. \quad (6.4.14)$$

Since the particle is confined to a box of length  $L$ , a crude, conservative estimate valid for any state is that  $\langle X^2 \rangle \leq (L/2)^2$ . Using this to compare our estimate of the lower bound on  $\langle H \rangle$  from the uncertainty relation to eq. (6.4.12) with  $n = 1$ , we see that the latter is a factor of  $\pi^2$  larger. For the ground state, a much better estimate of  $\langle X^2 \rangle$  comes from actually computing it, with the result

$$\langle X^2 \rangle = L^2 \left( \frac{1}{12} - \frac{1}{2\pi^2} \right), \quad (6.4.15)$$

so

$$\langle H \rangle \geq \frac{\hbar^2 \pi^2}{2mL^2} \left( \frac{3}{\pi^2 - 6} \right). \quad (6.4.16)$$

Comparing again with eq. (6.4.12) with  $n = 1$ , we see that the ground state energy is a factor of  $\pi^2/3 - 2$ , or about 1.29, larger than the estimate eq. (6.4.16) of its lower bound following from the uncertainty relation.

## 6.5 Bound states for the one-dimensional square well

As a generalization of the previous section, consider a particle of mass  $m$  in a finite square-well potential with linear width  $L$  and energy depth  $V_0$ :

$$V(x) = \begin{cases} V_0 & \text{for } x \leq -L/2 & \text{(region I),} \\ 0 & \text{for } -L/2 < x < L/2 & \text{(region II),} \\ V_0 & \text{for } x \geq L/2 & \text{(region III).} \end{cases} \quad (6.5.1)$$

Before proceeding, we note that the special case  $V_0 \rightarrow \infty$  should give the results of the previous section. Again, we expect to find energy eigenstate solutions with definite parity, because the potential is invariant under  $x \rightarrow -x$ .

The strategy for finding the stationary states is to first solve the differential equation (6.3.3) separately in each of the three regions I, II, and III, and then stitch these solutions together using eqs. (6.3.7) and (6.3.8) as boundary conditions at the points  $x = \pm L/2$ .

In region II, the differential equation is exactly the same as for the particle in a box,

$$\psi'' = -\frac{2mE}{\hbar^2} \psi, \quad (6.5.2)$$

and so the general solution is an arbitrary linear combination of  $e^{ikx}$  and  $e^{-ikx}$ , where  $k = \sqrt{2mE}/\hbar$ , as before. So, we can write

$$\psi_{\text{II}}(x) = A \cos(kx) + B \sin(kx). \quad (6.5.3)$$

In regions I and III, the differential equation is

$$\psi'' = \frac{2m(V_0 - E)}{\hbar^2} \psi. \quad (6.5.4)$$

Assume that  $E < V_0$ , so that we will have a bound state. Then, defining

$$\kappa = \sqrt{2m(V_0 - E)}/\hbar, \quad (6.5.5)$$

the general solutions in regions I and III are

$$\psi_{\text{I}}(x) = Ce^{\kappa x} + De^{-\kappa x}, \quad (6.5.6)$$

$$\psi_{\text{III}}(x) = Fe^{\kappa x} + Ge^{-\kappa x}. \quad (6.5.7)$$

The coefficient  $D$  must be 0, because otherwise  $\psi_{\text{I}}(x)$  would blow up for  $x \rightarrow -\infty$ , giving a non-normalizable unphysical solution. Similarly, from requiring  $\psi_{\text{III}}(x)$  to be well-behaved at  $x \rightarrow \infty$ , we get  $F = 0$ . It remains to solve for  $A$ ,  $B$ ,  $C$ , and  $G$ , using the boundary conditions at the points  $x = -L/2$  and  $L/2$ .

Let us first look for wavefunction solutions that are even under the parity transformation  $x \rightarrow -x$ . It follows that  $B = 0$  and  $G = C$ , and they must have the form

$$\psi(x) = \begin{cases} Ce^{\kappa x} & \text{for } x \leq -L/2, \\ A \cos(kx) & \text{for } -L/2 \leq x \leq L/2, \\ Ce^{-\kappa x} & \text{for } x \geq L/2. \end{cases} \quad (6.5.8)$$

Now we can apply the requirements that the wavefunction and its first derivatives are both continuous at  $x = L/2$ , as proved on general grounds in eqs. (6.3.7) and (6.3.8). In the present case, these conditions amount to

$$A \cos(kL/2) = Ce^{-\kappa L/2}, \quad (6.5.9)$$

$$-Ak \sin(kL/2) = -C\kappa e^{-\kappa L/2}. \quad (6.5.10)$$

By taking the ratio of these equations, one obtains  $k \tan(kL/2) = \kappa$ . It is convenient to define dimensionless quantities  $X = kL/2$  and  $Y = \kappa L/2$ , so that

$$X \tan X = Y, \quad (\text{even parity}), \quad (6.5.11)$$

$$X^2 + Y^2 = mV_0 L^2 / 2\hbar^2, \quad (6.5.12)$$




Figure 6.5.1: Graphical solutions for the bound energy eigenstates of a particle of mass  $m$  in a 1-d square well potential with depth  $V_0$  and width  $L$ . The dashed circles are  $X^2 + Y^2 = R^2 = mV_0L^2/2\hbar^2$ , for sample values  $R = 1.4, 4$ , and  $8$ . Even-parity solutions are the intersections with  $Y = X \tan X$  (darker, green curves) and odd-parity solutions are the intersections with  $Y = -X \cot X$  (lighter, red, curves). The energy eigenvalues for each solution are  $E = 2\hbar^2 X^2/mL^2$ . The cases  $R = 1.4, 4$ , and  $8$  are seen to have 1, 3, and 6 bound state solutions, respectively.

where eq. (6.5.12) follows from eq. (6.5.5) and  $E = \hbar^2 k^2/2m$ .

It is not possible to solve the simultaneous transcendental equations (6.5.11) and (6.5.12) analytically, but one can use graphical methods to understand the solutions and then obtain numerical results. In Figure 6.5.1 we graph in the  $X, Y$  plane the curves  $Y = X \tan X$  (darker solid curves) and the circle eq. (6.5.12) (dashed), for some sample values (1.4, 4, and 8) of the dimensionless radius  $R = \sqrt{mV_0L^2/2\hbar^2}$ . For a given value of  $R$ , the intersections provide the numerical eigenvalue solutions for  $X$  and  $Y$ , and thus for  $k$  and  $\kappa$ . We only need to consider positive  $X$  and  $Y$ , because  $\kappa$  is positive, and  $k$  can be taken positive without loss of generality.

The graphical approach makes clear that there is always at least one bound-state solution, no matter how small  $V_0$  (and thus  $R$ ) is. The ground state, which we will call  $|\psi_1\rangle$  with energy eigenvalue  $E_1$ , is the solution with smallest  $X$ , which always has even parity and occurs for  $0 < X < \pi/2$ , or  $0 < k < \pi/L$ , and so

$$0 < E_1 < \hbar^2 \pi^2 / 2mL^2. \quad (6.5.13)$$

The lower bound corresponds to small  $R$ , which means the limit of small  $V_0 \ll \hbar^2/mL^2$ . Conversely, the upper bound is relevant for large  $R$  and thus  $V_0 \gg \hbar^2/mL^2$ , the limit of the particle completely confined to a box as treated in the previous section. As  $V_0$  is increased (for fixed  $m, L$ ), the radius of the dashed circle grows, so the number of solutions increases, but for any finite  $V_0$  the number of bound state solutions is always finite.

Consider the lone bound state solution in the limit of small  $V_0$ , which means  $X$  and  $Y$  are also very small. By expanding  $X \tan X$  to quadratic order in  $X^2$ , and solving eqs. (6.5.11) and

(6.5.12) in that limit, we obtain

$$E_1 = \frac{\hbar^2 k^2}{2m} \approx V_0 \left( 1 - \frac{mV_0 L^2}{2\hbar^2} + \dots \right). \quad (6.5.14)$$

Because this energy is only slightly lower than  $V_0$ , the state is very weakly bound, but it always exists, no matter how small  $V_0$  is.

Now consider the stationary states that are odd under parity ( $x \rightarrow -x$ ), with wavefunctions that are therefore of the form

$$\psi(x) = \begin{cases} -Ce^{\kappa x} & \text{for } x \leq -L/2, \\ B \sin(kx) & \text{for } -L/2 \leq x \leq L/2, \\ Ce^{-\kappa x} & \text{for } x \geq L/2. \end{cases} \quad (6.5.15)$$

The continuity of the wavefunction and its derivative give

$$B \sin(kL/2) = Ce^{-\kappa L/2}, \quad (6.5.16)$$

$$Bk \cos(kL/2) = -C\kappa e^{-\kappa L/2}. \quad (6.5.17)$$

Defining  $X$  and  $Y$  in exactly the same way as before, we now have

$$-X \cot X = Y, \quad (\text{odd parity}) \quad (6.5.18)$$

in place of eq. (6.5.18), with the same equation for  $X^2 + Y^2 = R^2$ . The graph of  $Y = -X \cot X$  is shown in Figure 6.5.1 as the lighter solid lines. This time, we see that if  $V_0$  is small enough, there will not be any odd-parity bound-state solution. In order for there to exist at least one bound-state solution with an odd wavefunction, the dashed circle must have a large enough radius to intersect with the curve  $Y = -X \cot X$  for positive  $X$  and  $Y$ , specifically,  $\sqrt{\frac{mV_0 L^2}{2\hbar^2}} > \pi/2$ , so that  $V_0 > \pi^2 \hbar^2 / 2mL^2$ . In this case, the first excited state will have  $\pi/2 < X < \pi$ , so  $\hbar^2 \pi^2 / 2mL^2 < E_2 < 2\hbar^2 \pi^2 / 2mL^2$ .

Combining the information for even and odd parity states (which have odd and even  $n$ , respectively) from Figure 6.5.1 we can see that the energy levels alternate between even and odd parity, and the bound state  $|\psi_n\rangle$  exists if and only if the potential well is deep enough, which requires that  $R^2$  is sufficiently large,

$$\frac{mV_0 L^2}{2\hbar^2} > \frac{\pi^2}{4}(n-1)^2. \quad (6.5.19)$$

If  $|\psi_n\rangle$  does exist as a bound state, then  $\pi(n-1)/2 < X_n < \pi n/2$ , which implies

$$\frac{\hbar^2 \pi^2 (n-1)^2}{2mL^2} < E_n < \frac{\hbar^2 \pi^2 n^2}{2mL^2} \quad (n = 1, 2, 3, \dots). \quad (6.5.20)$$

Here  $E_n$  will approach its upper bound in the limit of large  $V_0$ , in agreement with the result found in section 6.4.

The square well potential also has a continuum of unbound states with energies  $E \geq V_0$ , with wavefunctions whose magnitudes approach a constant for large distances  $|x|$ . These are also known as scattering states. We will discuss the problem of scattering from the one-dimensional square well in section 6.6.

## 6.6 Scattering problems in one dimension

**Scattering theory** deals with unbound quantum states with continuous energy eigenvalues. The particles described by these states can be thought of as originating far away, in a measured or controlled configuration, typically a superposition of free plane waves moving in a common direction. In a localized region of space, the particles then interact with a potential, or with another group of particles, after which they escape to large distances again where they can be measured. In practice, one can learn about the potential, or interactions between particles, by studying the asymptotic forms of the unbound states. In this section we will consider potential scattering problems in the simplified realm of one dimension, and turn to the problem of scattering in three dimensions in Chapter 23.

Consider two asymptotic regions I and II with constant potentials, and an intermediate region III where the potential can be arbitrary, so

$$V(x) = \begin{cases} 0 & \text{region I } (x < a), \\ U(x) & \text{region III } (a < x < b), \\ V_0 & \text{region II } (x > b), \end{cases} \quad (6.6.1)$$

as illustrated in Figure 6.6.1. The constant potential in region I, which will contain the incident particles, is taken to be 0, by subtraction from  $V(x)$  if necessary. This entails no loss of generality, because as usual the effect of a constant contribution to the energy can be absorbed into a global phase that is the same for all states. The potential  $V_0$  in region II can be either positive or negative. We will be interested only in unbound states with energy  $E > 0$ , although there may also be bound states if the potential  $U(x)$  in region III goes negative for some  $x$ .

The stationary states for this potential have wavefunctions that might be complicated in region III, but they are definitely simple in regions I and II. Assuming that  $E > V_0$ , one class




Figure 6.6.1: A generic one-dimensional scattering potential of the type in eq. (6.6.1). The potential  $V_0$  in region II on the far right can be either positive or negative.

of such stationary states is

$$\psi(x, t) = (Ae^{ikx} + Be^{-ikx}) e^{-iEt/\hbar}, \quad (\text{region I}), \quad (6.6.2)$$

$$\psi(x, t) = Ce^{ik'x} e^{-iEt/\hbar}, \quad (\text{region II}). \quad (6.6.3)$$

Here the component  $A$  corresponds to an initial beam of particles moving to the right in region I from  $x = -\infty$ , and  $B$  corresponds to a reflected component going back to  $x = -\infty$ . In region II, there is only a transmitted component moving to the right; as a boundary condition we are imposing that there is no left-moving component that would correspond to incident particles arriving from  $x = +\infty$ . Note that in 1-d scattering problems, we are not interested in eigenstates of parity; the symmetry is broken because the incident particles are coming from the left.

Now, for a given energy  $E$  and normalization  $A$  treated as known inputs characteristic of the incident beam of particles, we can solve for  $k$ ,  $k'$  and then, in principle, for  $B$  and  $C$ . The easy part is that from the Schrödinger equation,

$$\frac{\hbar^2 k^2}{2m} = E, \quad \frac{\hbar^2 k'^2}{2m} = E - V_0. \quad (6.6.4)$$

To solve for  $B$  and  $C$  requires the scattering potential  $U(x)$  to be specified. Once  $B$  and  $C$  are known, eq. (5.6.12) can be used to find the probability and current densities in each region. In region I, the probability density is

$$\rho = |A|^2 + |B|^2 + 2\text{Re}[A^*B] \cos(2kx) + 2\text{Im}[A^*B] \sin(2kx). \quad (6.6.5)$$

The last two terms will each give 0 after averaging over a range of  $x$  that is large compared to  $1/k$ . The current density in region I is

$$J = \frac{\hbar k}{m}(|A|^2 - |B|^2), \quad (6.6.6)$$

where the cross-terms between  $A$  and  $B$  canceled completely, even without doing any averaging. Meanwhile, in region II, the probability density and current are

$$\rho = |C|^2, \quad (6.6.7)$$

$$J = \frac{\hbar k'}{m}|C|^2. \quad (6.6.8)$$

The interpretation of the probability densities of eqs. (6.6.5) and (6.6.7) is

$$\text{density of particles in } \begin{pmatrix} \text{incident} \\ \text{reflected} \\ \text{transmitted} \end{pmatrix} \text{ beam} = \begin{pmatrix} |A|^2 \\ |B|^2 \\ |C|^2 \end{pmatrix}, \quad (6.6.9)$$

while the interpretation of the current densities of eqs. (6.6.6) and (6.6.8) is

$$\text{flux} = \text{particles/time in } \begin{pmatrix} \text{incident} \\ \text{reflected} \\ \text{transmitted} \end{pmatrix} \text{ beam} = \begin{pmatrix} \hbar k |A|^2/m \\ -\hbar k |B|^2/m \\ \hbar k' |C|^2/m \end{pmatrix} = \begin{pmatrix} J_A \\ J_B \\ J_C \end{pmatrix}. \quad (6.6.10)$$

The effect of the scattering potential on the incident mono-energetic beam can therefore be given in terms of reflection and transmission ratios, reminiscent of quantities in classical optics,

$$R = \frac{|J_B|}{|J_A|} = \frac{|B|^2}{|A|^2}, \quad T = \frac{|J_C|}{|J_A|} = \frac{k' |C|^2}{k |A|^2}. \quad (6.6.11)$$

In specific problems, we can always set  $A = 1$  at the beginning, since only the ratios  $B/A$  and  $C/A$  are needed, and  $B$  and  $C$  will always be proportional to  $A$  due to the linearity of the wavefunction in the Schrödinger equation.

To find  $R$  or  $T$  we must specify the scattering potential  $V(x)$ . However, in general, probability conservation implies that they are related by

$$R + T = 1. \quad (6.6.12)$$

To prove this, we note that for a stationary state,

$$\frac{\partial J}{\partial x} = -\frac{\partial \rho}{\partial t} = -\frac{\partial}{\partial t} |\psi(x, t)|^2 = -\frac{\partial}{\partial t} |\psi(x, t_0) e^{-i(t-t_0)E/\hbar}|^2 = 0, \quad (6.6.13)$$

so, by the fundamental theorem of calculus,

$$0 = \int_{-\infty}^{\infty} dx \frac{\partial J}{\partial x} = J(\infty) - J(-\infty). \quad (6.6.14)$$

This can be rewritten as

$$J_C = |J_A| - |J_B|, \quad (6.6.15)$$

from which eq. (6.6.12) follows immediately. Since  $R$  and  $T$  are manifestly positive, they must also be between 0 and 1, inclusive, with the extremes reached only in very special cases. This probabilistic situation is in contrast to classical mechanics, where  $T$  is always exactly either 1 or 0, depending on whether or not the particle has enough energy to overcome the potential barrier set by the maximum value of the potential.

To find  $R$  and  $T$  in specific examples, it is necessary to solve the Schrödinger equation in region III and to apply boundary conditions in the form of continuity of the wavefunction, and its derivative if the potential is finite, at the boundaries between adjacent regions. If the potential at a boundary has a delta function, then eq. (6.3.7) can be used to obtain the boundary condition for the first derivative of the wavefunction.



As a first example, consider a step-function potential

$$V(x) = \begin{cases} 0 & \text{region I } (x < 0), \\ V & \text{region II } (x > 0), \end{cases} \quad (6.6.16)$$

for which region III does not exist, and regions I and II meet at  $x = 0$ . Continuity of the wavefunction eq. (6.6.2)-(6.6.3) and its first derivative at  $x = 0$  give

$$1 + B = C, \quad (6.6.17)$$

$$ik + (-ik)B = ik'C. \quad (6.6.18)$$

Here we have taken the opportunity to set  $A = 1$ , since we will be interested in the ratios  $R$  and  $T$ . The solutions for  $B$  and  $C$  are

$$B = \frac{k - k'}{k + k'}, \quad C = \frac{2k}{k + k'}. \quad (6.6.19)$$

The reflection and transmission coefficients are therefore

$$R = \left| \frac{1 - k'/k}{1 + k'/k} \right|^2, \quad T = \frac{4k'/k}{(1 + k'/k)^2}, \quad (6.6.20)$$

where

$$k'/k = \sqrt{1 - V/E}. \quad (6.6.21)$$

Note that we need  $E > V$  in order for  $k'$  and  $T$  to be real. Otherwise, the wavefunction for  $x > 0$  is instead an exponential of the form  $\psi_{\text{II}}(x) = Ce^{-\kappa x}$  with  $\hbar^2\kappa^2/2m = V - E$ , and one finds  $R = 1$ . Since the flux decreases exponentially in region II rather than maintaining constant magnitude,  $T = 0$  in that case; the particles are all reflected, although they still have a non-zero probability to be found at any point  $x > 0$ , proportional to  $e^{-2\kappa x}$ . Region II in this case is said to be a classically forbidden region.

As a second example, consider the symmetric rectangular barrier potential

$$V(x) = \begin{cases} 0 & \text{region I } (x < -a/2), \\ V & \text{region III } (|x| < a/2), \\ 0 & \text{region II } (x > a/2), \end{cases} \quad (6.6.22)$$

as illustrated in Figure 6.6.2.

Let us first consider the case that  $E > V > 0$ , so that transmission past the barrier is classically allowed. Then we can write the wavefunctions in the three regions as

$$\psi_{\text{I}}(x) = e^{ikx} + Be^{-ikx}, \quad (6.6.23)$$

$$\psi_{\text{III}}(x) = De^{ik'x} + Fe^{-ik'x}, \quad (6.6.24)$$

$$\psi_{\text{II}}(x) = Ce^{ikx}, \quad (6.6.25)$$




Figure 6.6.2: A one-dimensional scattering potential of the type in eq. (6.6.22). The potential can be either a barrier (if  $V > 0$ , as shown) or a well (if  $V < 0$ ).

where  $k' = \sqrt{2m(E - V)}/\hbar$  and  $k = \sqrt{2mE}/\hbar$  are real numbers, and again we choose  $A = 1$ . There are four remaining unknowns,  $B, C, D, F$ , and only two of these enter into the measurable quantities  $R = |B|^2$  and  $T = |C|^2$ . Continuity of the wavefunction and its first derivative at  $x = -a/2$ , using  $\psi_I$  and  $\psi_{III}$ , give us

$$e^{-ika/2} + Be^{ika/2} = De^{-ik'a/2} + Fe^{ik'a/2}, \quad (6.6.26)$$

$$ike^{-ika/2} - ikBe^{ika/2} = ik'De^{-ik'a/2} - ik'Fe^{ik'a/2}, \quad (6.6.27)$$

Also, at  $x = a/2$  we find from  $\psi_{II}$  and  $\psi_{III}$ ,

$$Ce^{ika/2} = De^{ik'a/2} + Fe^{-ik'a/2}, \quad (6.6.28)$$

$$ikCe^{ika/2} = ik'De^{ik'a/2} - ik'Fe^{-ik'a/2}. \quad (6.6.29)$$

It is convenient to first solve eqs. (6.6.28) and (6.6.29) for  $D$  and  $F$  in terms of  $C$ , and plug the results into eqs. (6.6.26) and (6.6.27) which then involve only  $B$  and  $C$  as unknowns. The resulting eq. (6.6.26) and eq. (6.6.27) then combine to give

$$B = i \frac{k'^2 - k^2}{2kk'} \sin(k'a) C. \quad (6.6.30)$$

Now, since we also know  $R + T = |B|^2 + |C|^2 = 1$ , we can solve to get

$$T = |C|^2 = \left[ 1 + \left( \frac{k'^2 - k^2}{2kk'} \right)^2 \sin^2(k'a) \right]^{-1} \quad (6.6.31)$$

for the transmission coefficient. Then,  $R$  is just  $1 - T$ .

To express the result directly in terms of the incident energy and the height of the potential barrier, we can now plug in  $k' = k\sqrt{1 - V/E}$ , to get

$$T = \left[ 1 + \frac{V^2}{4E(E - V)} \sin^2 \left( \frac{a}{\hbar} \sqrt{2m(E - V)} \right) \right]^{-1}, \quad (E \geq V). \quad (6.6.32)$$

As a check, if  $V = 0$ , then there is no barrier, and  $T = 1$  and  $R = 0$ . In the high-energy limit,  $E \gg V$  gives  $T \approx 1$ , and the barrier is almost transparent, as the potential is too weak to have

much effect on the incident particles. It is perhaps more surprising that even if  $V$  and  $E$  are comparable, we also get  $T = 1$  whenever  $\sin(k'a) = 0$ , which occurs if  $k' = n\pi/a$  for any integer  $n$ . This shows that the barrier is transparent to the incident mono-energetic particle beam if its width  $a$  is equal to an integer times half of the de Broglie wavelength  $2\pi/k'$  inside the barrier. The discrete energies at which this occurs, called **scattering resonances**, are

$$E = E_n + V, \quad E_n = \frac{\hbar^2 \pi^2 n^2}{2ma^2}, \quad (n = 1, 2, 3, \dots). \quad (6.6.33)$$

These  $E_n$  coincide with the energy eigenvalues for a particle in a box of width  $a$ , but with walls of infinite potential height, as found in section 6.4. Finally, note that if  $E \approx V$ , then we have  $\sin^2 \left( a\sqrt{2m(E-V)}/\hbar \right) \approx 2a^2m(E-V)/\hbar^2$ , so

$$T \approx \left( 1 + \frac{mVa^2}{2\hbar^2} \right)^{-1}, \quad (E \approx V). \quad (6.6.34)$$

This is the transmission coefficient for the case that classical transmission past the barrier is just barely possible; for  $\hbar \rightarrow \infty$  it approaches 1.

Now we turn to the case that  $E < V$ , so that classically the particle would not be expected to make it past the barrier. Then, within the barrier region III, we have instead of eq. (6.6.24),

$$\psi_{\text{III}}(x) = De^{-\kappa x} + Fe^{\kappa x}, \quad (6.6.35)$$

where

$$\kappa = \sqrt{2m(V-E)}/\hbar. \quad (6.6.36)$$

All of the subsequent algebra is the same, but with  $k' \rightarrow i\kappa$ , so that the sine function is replaced by a hyperbolic sine, and

$$T = \left[ 1 + \frac{V^2}{4E(V-E)} \sinh^2 \left( \frac{a}{\hbar} \sqrt{2m(V-E)} \right) \right]^{-1}, \quad (E \leq V). \quad (6.6.37)$$

In this case,  $T < 1$  always, but it is never 0. This is an example of **quantum tunneling**. Classically, there would be no transmission at all, but the Schrödinger equation gives a non-zero probability for the particle to enter the classically forbidden region and emerge on the right side of the barrier. For  $E \ll V$ , one finds that  $T \rightarrow 0$ ; the transmission due to tunneling becomes exponentially small in the limit of an incident energy much smaller than the barrier potential height. The results for the transmission coefficient  $T$  are shown as a function of the incident particle energy  $E$  in Figure 6.6.3, for two different values of the barrier height  $V$ .

So far we have assumed that the potential barrier height in Figure 6.6.2 is positive, with  $V > 0$ . Now let us consider the case of scattering from a potential well, so that  $V < 0$  in




Figure 6.6.3: Transmission ratio  $T$  for scattering from a one-dimensional rectangular barrier of width  $a$  and height  $V = 5\hbar^2/ma^2$  (left) or  $V = 50\hbar^2/ma^2$  (right), as a function of the energy  $E$  of the incident particles of mass  $m$ . For  $E/V < 1$ , the transmission is classically forbidden, and the non-zero  $T$  is an example of quantum tunneling. For  $E/V > 1$ , the barrier is perfectly transparent if its width is an integer multiple of half of the particle's de Broglie wavelength inside the barrier. Transparency also occurs in the limit  $E \gg V$ .

eq. (6.6.22). In that case, all of the same algebra goes through as before, with  $V = -|V|$ . So, we have

$$T = \left[ 1 + \frac{V^2}{4E(E + |V|)} \sin^2 \left( \frac{a}{\hbar} \sqrt{2m(E + |V|)} \right) \right]^{-1}, \quad (E > 0 > V). \quad (6.6.38)$$

Classically, there would never be reflection, as the particle incident from the left would have enough momentum and energy to follow a trajectory that takes it ineluctably to  $x = +\infty$ . The prediction of quantum mechanics in the low-energy limit is very different, as eq. (6.6.38) gives  $T \rightarrow 0$  for  $E \rightarrow 0$ . In the high-energy limit, one finds  $T \rightarrow 1$  for  $E \rightarrow \infty$ , in agreement with the classical expectation. Just as we found for  $V > 0$ , there are scattering resonances that occur when  $E = E_n - |V|$  where  $E_n = \hbar^2 \pi^2 n^2 / 2ma^2$  are the binding energies of bound states in a box (with walls at infinite potential) of width  $a$ . The results for the transmission coefficient  $T$  are shown as a function of the incident particle energy  $E$  in Figure 6.6.4, for two different values of the well depth  $|V|$ . For larger negative  $V$ , the resonance energies are more sharply defined.

More generally, including in three-dimensional problems, an attractive potential can become almost transparent to scattering for certain resonance energies, which are determined by the geometry of the potential and the de Broglie wavelengths of the incident particles. This phenomenon is known as the **Ramsauer–Townsend effect**, as it was first observed independently by Carl Ramsauer and John S. Townsend in 1921, before its subsequent explanation by quantum mechanics, in the scattering of electrons from noble gas atoms Ar, Kr, and Xe. The qualitative




Figure 6.6.4: Transmission ratio  $T$  for scattering from a one-dimensional well of width  $a$  and  $V = -50\hbar^2/ma^2$  (left) or  $V = -5 \times 10^4 \hbar^2/ma^2$  (right). In the very low energy limit, the particles are entirely reflected. Transparency occurs in the limit of large  $E$ , and also when  $E$  is equal to a resonance energy, given by the eigenvalues  $E_n = \frac{\hbar^2 \pi^2 n^2}{2ma^2}$  of a particle confined to a box of width  $a$ .

explanation for this is that close to the positively charged nucleus, the scattering electrons see a spherically symmetric attractive potential well that is cut off at larger distances by the screening of the nuclear charge due to the atomic electrons. This geometry leads to a strong suppression of the scattering cross-section, corresponding to near transparency for incident electron energies  $E \approx 0.7$  eV. The preceding results for the first peak in  $T$  in the one-dimensional well scattering problem with large negative  $V$  are a rough qualitative model for this phenomenon.

## 6.7 Particle acted on by a constant force

Consider the problem of a particle acted on by a constant force. (For example, this could be a charged particle in a uniform electric field.) Let us take the force to have magnitude  $f$ , and to point in the negative  $x$  direction, so that the classical potential energy is  $V = fx$ . We will treat this as a one-dimensional problem; restoring the effects of the  $y$  and  $z$  degrees of freedom just adds a constant to the energy. Our Hamiltonian operator is therefore

$$H = \frac{P^2}{2m} + fX. \quad (6.7.1)$$

Our goal is to find the stationary states of this one-dimensional problem.

This problem is much easier to solve by starting in the momentum representation rather than the position representation. To do it, act with  $\langle p|$  on the time-independent Schrödinger equation  $H|\psi_E\rangle = E|\psi_E\rangle$ . According to the momentum-representation rules  $|\psi_E\rangle \rightarrow \tilde{\psi}_E(p)$  and

$P \rightarrow p$  and  $X \rightarrow i\hbar d/dp$ , as in Table 2.8.1, one obtains the first-order differential equation

$$\left( \frac{p^2}{2m} + i\hbar f \frac{d}{dp} - E \right) \tilde{\psi}_E(p) = 0, \quad (6.7.2)$$

which can be rearranged into

$$\frac{d}{dp} \ln(\tilde{\psi}_E(p)) = \frac{i}{\hbar f} \left( \frac{p^2}{2m} - E \right). \quad (6.7.3)$$

Integrating both sides, and then exponentiating, gives the solution, for any real energy  $E$ ,

$$\tilde{\psi}_E(p) = \tilde{\psi}_E(0) \exp \left[ \frac{i}{\hbar f} \left( \frac{p^3}{6m} - Ep \right) \right], \quad (6.7.4)$$

where the arbitrary normalization factor  $\tilde{\psi}_E(0)$  arises as the constant of integration.

Let us choose to fix the normalization factor so as to satisfy the Dirac orthonormality condition with respect to energy. To that end, consider

$$\langle \psi_{E'} | \psi_E \rangle = \int_{-\infty}^{\infty} dx \psi_{E'}^*(x) \psi_E(x) = \int_{-\infty}^{\infty} dp \tilde{\psi}_{E'}^*(p) \tilde{\psi}_E(p). \quad (6.7.5)$$

Using eq. (6.7.4), the last expression can be rewritten as

$$\tilde{\psi}_{E'}^*(0) \tilde{\psi}_E(0) \int_{-\infty}^{\infty} dp \exp \left[ \frac{ip}{\hbar f} (E' - E) \right], \quad (6.7.6)$$

where the  $p^3$  terms in the exponential have conveniently canceled. The integral can be evaluated in terms of a Dirac delta function using eq. (2.2.20), giving

$$\langle \psi_{E'} | \psi_E \rangle = \tilde{\psi}_{E'}^*(0) \tilde{\psi}_E(0) 2\pi \hbar f \delta(E' - E). \quad (6.7.7)$$

We now see that to achieve Dirac orthonormality of the energy eigenstates,  $\langle \psi_{E'} | \psi_E \rangle = \delta(E' - E)$ , we should choose the normalization  $\tilde{\psi}_E(0) = 1/\sqrt{2\pi \hbar f}$ .

Having successfully found the momentum wavefunction, we can get the position wavefunction by taking the inverse Fourier transform. Applying eq. (2.8.42),

$$\psi_E(x) = \frac{1}{\sqrt{2\pi \hbar}} \int_{-\infty}^{\infty} dp e^{ipx/\hbar} \tilde{\psi}_E(p) = \frac{1}{\pi \hbar \sqrt{f}} \int_0^{\infty} dp \cos \left( \frac{1}{\hbar f} \left[ \frac{p^3}{6m} - Ep + fxp \right] \right). \quad (6.7.8)$$

The integral is not trivial, but can be put into a standard form by defining a dimensionless integration variable  $s = p/(2\hbar m f)^{1/3}$  and a shifted and rescaled dimensionless position coordinate

$$y = \left( \frac{2mf}{\hbar^2} \right)^{1/3} (x - E/f). \quad (6.7.9)$$

The resulting integral over  $s$  is proportional to the **Airy function**, defined by

$$\text{Ai}(y) = \frac{1}{\pi} \int_0^{\infty} ds \cos(sy + s^3/3). \quad (6.7.10)$$




Figure 6.7.1: The Airy function  $\text{Ai}(y)$ , proportional to the position wavefunction for a particle in a linear potential with classical allowed region  $y < 0$ , forbidden region  $y > 0$ , and turning point  $y = 0$ .

This improper integral can be shown to converge, in the sense that it can be defined more formally either as  $\text{Ai}(y) = \lim_{N \rightarrow \infty} \frac{1}{\pi} \int_0^N ds \cos(sy + s^3/3)$ , or by means of a convergence factor as  $\text{Ai}(y) = \lim_{\epsilon \rightarrow 0} \frac{1}{\pi} \int_0^\infty ds e^{-\epsilon s} \cos(sy + s^3/3)$ . The Airy function is the solution bounded at  $y = \infty$  of the Airy differential equation

$$\left( \frac{d^2}{dy^2} - y \right) \text{Ai}(y) = 0. \quad (6.7.11)$$

It is named for George B. Airy, who first used it to describe optical caustics (like rainbows). The final result for the position wavefunction of the stationary state with energy  $E$  is

$$\psi_E(x) = \left( \frac{2m}{\hbar^2 \sqrt{f}} \right)^{1/3} \text{Ai}(y). \quad (6.7.12)$$

The point  $y = 0$  (also known as  $x = E/f$ ) is the classical turning point. This means that a classical particle of energy  $E$  coming from the left will follow a trajectory confined to  $y < 0$ , because it turns around at  $y = 0$  where the momentum vanishes. The quantum wavefunction we have found can be thought of as a special kind of scattering problem in which an incident particle is always reflected, with  $T = 0$  and  $R = 1$ .

The Airy function is depicted in Figure 6.7.1. It decreases rapidly in the classically forbidden region  $y > 0$ , with an approximation for large positive  $y$ ,

$$\text{Ai}(y) \approx \frac{1}{2\sqrt{\pi}y^{1/4}} \exp\left(-\frac{2}{3}y^{3/2}\right), \quad (y \gg 0). \quad (6.7.13)$$

In the classically allowed region, it instead oscillates, but with a wavelength and amplitude that both decrease for larger  $-y$ . The asymptotic form for large negative  $y$  is

$$\text{Ai}(y) \approx \frac{1}{\sqrt{\pi}(-y)^{1/4}} \sin\left(\frac{2}{3}(-y)^{3/2} + \frac{\pi}{4}\right), \quad (y \ll 0). \quad (6.7.14)$$

The approximate values of the first four zeros of  $\text{Ai}(y)$  and its first derivative are

$$y \approx -2.33811, -4.08795, -5.52056, -6.78671, \dots \quad [\text{zeros of } \text{Ai}(y)], \quad (6.7.15)$$

$$y \approx -1.01879, -3.24820, -4.82010, -6.16331, \dots \quad [\text{zeros of } \text{Ai}'(y)]. \quad (6.7.16)$$

The solution that we have found is useful for understanding the tunneling penetration of a particle into a classically forbidden region in which the potential grows linearly, for example a charged particle confined by a constant electric field. The form of the exponential fall of the wavefunction is as given in eq. (6.7.13). Of course, this solution is only an idealization, because in reality the potential energy is always bounded from above. However, if the cutoff of the potential is far in the forbidden region, then this makes little difference.

A common practical application is to stitch the solutions of the Airy differential equation, including the orthogonal solution  $\text{Bi}(y)$  that is not bounded as  $y \rightarrow \infty$ , onto other solutions associated with other potentials in other regions, by matching the wavefunction (and its first derivative, if the potential is finite). For example, this is used in one approach to the WKB approximation method, named for Gregor Wentzel, Hendrik Kramers, and Léon Brillouin, which is discussed in other books.

## 6.8 Exercises

**Exercise 6.1.** A particle of mass  $m$  moves in 1 dimension in the presence of an attractive delta-function potential  $V(x) = -a\delta(x)$ , where  $a$  is a positive constant.

- (a) Show that there is always exactly one bound state solution, with energy  $E = -ma^2/2\hbar^2$ , and obtain its normalized position and momentum wavefunctions  $\psi(x)$  and  $\tilde{\psi}(p)$ .
- (b) For the bound state, sketch  $\psi(x)$ , and compute  $\langle X \rangle$ ,  $\langle X^2 \rangle$ , and  $\Delta X$ .
- (c) For the bound state, sketch  $\tilde{\psi}(p)$ , and compute  $\langle P \rangle$ ,  $\langle P^2 \rangle$ , and  $\Delta P$ .
- (d) What is  $(\Delta X)(\Delta P)$ ? How does it compare to the result for a Gaussian wavefunction?

**Exercise 6.2.** For the particle in a 1-d box of length  $L$  discussed in section 6.4, consider the wavefunction  $\psi_n(x)$  for each energy level  $n$ .

- (a) Find the probability that the particle will be found within a distance  $L/4$  of the center of the box. (Your answer should contain no trigonometric functions, and may have a different form for even and odd  $n$ .) For which  $n$  is this probability largest, and for which is it smallest?
- (b) Find the uncertainties  $\Delta X$  and  $\Delta P$  for each energy level  $n$ . Show that their product is always consistent with the uncertainty principle.

**Exercise 6.3.** Consider a particle of mass  $m$  moving in a potential that has a power-law behavior  $V(x) = (x/a)^n V_0$  at large positive  $x$ , where  $V_0$ ,  $a$ , and  $n$  are positive constants. Show that the



wavefunction for a stationary state falls off at large  $x$  like  $\psi \sim N \exp(-c(x/a)^\nu)$ , where  $c$  and  $\nu$  are constants that you will find in terms of the quantities  $\hbar$ ,  $m$ ,  $V_0$ ,  $a$ , and  $n$ .

**Exercise 6.4.** Consider scattering of particles with mass  $m$  and energy  $E$  from a potential  $V(x) = V_0\delta(x/a)$ , where  $a$  is a length scale and  $V_0$  is a constant potential energy.

- (a) Find the transmission and reflection coefficients  $T$  and  $R$  by matching stationary-state wavefunctions at  $x = 0$ .
- (b) Check your result for  $T$  by comparing to the behavior of eq. (6.6.37) for small  $a$ .

**Exercise 6.5.** Consider a particle of mass  $m$  moving in one dimension in a potential

$$V(x) = -aV_0[\delta(x-a) + \delta(x+a)], \quad (6.8.1)$$

where  $a > 0$  and  $V_0 > 0$ . (Note that  $V_0$  has units of energy.)

- (a) What is the form of the wave function for a bound stationary state with even parity? Normalize your answer so that  $\psi(x) = e^{-\kappa x}$  for large positive  $x$ .
- (b) Find an equation that determines the energies for even-parity bound states, and determine graphically how many such states there are. [Hint: the equation can be written in the form  $\kappa a = (\text{polynomial in } e^{-\kappa a})$ . Sketch the shape of the right-hand side as a function of  $\kappa a$ .]
- (c) Repeat parts (a) and (b) for odd-parity bound states. For what values of  $V_0$  are there no such states? [Hint: this time your equation should have the form  $\kappa a = (\text{polynomial in } e^{-\kappa a})$ . Sketch the right-hand side, and consider its slope at  $\kappa a = 0$ .]
- (d) Solve for the even-parity bound state energy analytically in the limit of  $V_0 \ll \hbar^2/ma^2$ .
- (e) Find the even- and odd-parity bound-state energies in the limit  $V_0 \gg \hbar^2/ma^2$ . (They are equal to each other in that limit.)

**Exercise 6.6.** For the potential in Exercise 6.5, let us look for stationary scattering states with  $E = \hbar^2 k^2/2m > 0$  and wavefunctions of the form

$$\psi(x) = \begin{cases} e^{ikx} + Be^{-ikx} & (x \leq -a) \\ De^{ikx} + Fe^{-ikx} & (-a \leq x \leq a) \\ Ce^{ikx} & (x \geq a). \end{cases} \quad (6.8.2)$$

- (a) Derive four equations that relate the coefficients  $B$ ,  $C$ ,  $D$ , and  $F$ . Simplify the notation for your work below by writing them in terms of the dimensionless quantity  $n = maV_0/\hbar^2 k$ .
- (b) Solve for  $C$ , and use it to find the transmission coefficient  $T$ . Some partial results:  $C = [1 - iN_1 n + n^2(e^{iN_2 ka} - N_3)]^{-1}$ , where  $N_1$ ,  $N_2$ , and  $N_3$  are certain positive integers that you will discover, and  $T = [P_1 + P_2 \cos(N_2 ka) + P_3 \sin(N_2 ka)]^{-1}$  where  $P_1$ ,  $P_2$ , and  $P_3$  are certain polynomials in  $n$  with integer coefficients.
- (c) Check that your transmission coefficient has the expected behavior when the energy  $E$  becomes very large. How does it behave when  $E$  is very small?

**Exercise 6.7.** Consider a quantum mechanical “bouncing ball”, consisting of a particle of mass  $m$  moving only vertically in the potential

$$V(x) = \begin{cases} \infty & (\text{for } x < 0), \\ mgx & (\text{for } x > 0), \end{cases} \quad (6.8.3)$$

where  $x$  is the height above an impenetrable surface at  $x = 0$ , and  $g$  is the acceleration due to gravity.

(a) Find the stationary-state wavefunctions  $\psi_n(x)$  and energies  $E_n$ , in terms of the Airy function  $\text{Ai}(y)$  and its zeros  $y_n$  for  $n = 1, 2, 3, \dots$

(b) For the ground state, estimate the numerical energy and characteristic height, defined as the point where  $\psi_1(x)$  is maximum, when  $m = 1$  gram and when  $m = 1.7 \times 10^{-24}$  grams (the mass of a neutron or proton or a hydrogen atom). Use eqs. (6.7.15) and (6.7.16) and  $g = 9.8 \text{ m/s}^2$ .

## 7 The harmonic oscillator

### 7.1 The unreasonable effectiveness of the harmonic oscillator

A harmonic oscillator is any system in which the restoring force is proportional to the displacement from equilibrium. Consider the case of a particle of mass  $m$  moving in one dimension. In terms of energy, the classical harmonic oscillator is defined by

$$\text{kinetic energy} = \frac{p^2}{2m}, \quad \text{potential energy} = V(x) = \frac{1}{2}m\omega^2 x^2, \quad (7.1.1)$$

where the angular frequency  $\omega$  has units of 1/time and parameterizes the strength of the restoring force, according to

$$\text{force} = -\frac{\partial V}{\partial x} = -m\omega^2 x. \quad (7.1.2)$$

The classical Hamiltonian is the sum of the kinetic and potential energies,

$$H = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 x^2, \quad (7.1.3)$$

which leads to the phase-space equations of motion

$$\dot{x} = \frac{\partial H}{\partial p} = p/m, \quad \dot{p} = -\frac{\partial H}{\partial x} = -m\omega^2 x, \quad (7.1.4)$$

with the general solution

$$x(t) = x_0 \cos(\omega t + \phi_0), \quad p(t) = -m\omega x_0 \sin(\omega t + \phi_0), \quad (7.1.5)$$

where  $x_0$  and  $\phi_0$  are constants determined by the initial conditions.

A wise theorist, Sidney Coleman, was fond of remarking that the job of a physicist consists of “treating the harmonic oscillator in ever-increasing levels of abstraction”. This may be a slight exaggeration, but there are at least three reasons why the quantum harmonic oscillator is particularly worthy of study.

First, the harmonic oscillator is a problem that can actually be solved exactly in closed form in quantum mechanics. Most problems in quantum mechanics do not have this property.

Second, it arises quite often as a good approximation to more complicated problems. To understand this, consider a more general potential  $V(x)$ , which we assume to have a local minimum at some point  $x = x_0$ , and to be smooth in a neighborhood of that point. Then we can expand it in a Taylor series for small  $x - x_0$ ,

$$V(x) = V(x_0) + (x - x_0) \left. \frac{dV}{dx} \right|_{x=x_0} + \frac{1}{2}(x - x_0)^2 \left. \frac{d^2V}{dx^2} \right|_{x=x_0} + \frac{1}{6}(x - x_0)^3 \left. \frac{d^3V}{dx^3} \right|_{x=x_0} + \cdots \quad (7.1.6)$$

The first term is a constant, and so can be absorbed into a redefinition of the zero point of energy; as a constant part of the Hamiltonian, it gives the same overall phase to all states, and therefore can be ignored. The second term is zero, since by definition  $dV/dx$  vanishes at a minimum of the potential. Thus, if the terms in the expansion are getting smaller, the leading behavior is captured by the quadratic term. After redefining coordinates according to  $x \rightarrow x + x_0$ , we can hope to approximate

$$V(x) \approx \frac{x^2}{2} \left( \frac{d^2V}{dx^2} \Big|_{x=0} \right) \quad (7.1.7)$$

where the constant quantity in parentheses is defined to be  $m\omega^2$ .

The same argument generalizes to a particle moving in a three dimensional potential. Let us suppose that the  $x, y, z$  coordinates have already been redefined so that the minimum of the potential is at the origin  $(x, y, z) = (0, 0, 0)$ . Then we have

$$V(x, y, z) = V(0, 0, 0) + \frac{1}{2}x_a x_b \mathcal{V}_{ab} + \cdots, \quad (7.1.8)$$

where by convention repeated indices  $a, b, \dots$  are implicitly summed over  $1, 2, 3$ , with  $x_1 = x$ ,  $x_2 = y$ ,  $x_3 = z$ , and

$$\mathcal{V}_{ab} = \frac{\partial^2 V}{\partial x_a \partial x_b} \Big|_{x_c=0} \quad (7.1.9)$$

is a real symmetric matrix. A linear term involving  $\frac{\partial V}{\partial x_a} \Big|_{x_b=0}$  vanishes because we are at the minimum of the potential. A theorem in linear algebra says that a real symmetric matrix can always be diagonalized by some orthogonal matrix  $O$ , according to

$$\mathcal{V} = O^T \tilde{\mathcal{V}} O, \quad (7.1.10)$$

where  $\tilde{\mathcal{V}} = \text{diag}(\tilde{\mathcal{V}}_1, \tilde{\mathcal{V}}_2, \tilde{\mathcal{V}}_3)$ , so that

$$\mathcal{V}_{ab} = O_{ca} \tilde{\mathcal{V}}_c O_{cb}, \quad (7.1.11)$$

and the orthogonality condition on  $O$  is written as

$$O_{ac} O_{bc} = O_{ca} O_{cb} = \delta_{ab}. \quad (7.1.12)$$

Dropping the constant  $V(0, 0, 0)$ , and assuming the higher-order terms in the expansion can be neglected, the quantum mechanical Hamiltonian can be approximated as

$$H = \frac{1}{2m}(P_x^2 + P_y^2 + P_z^2) + \frac{1}{2}O_{ca} \tilde{\mathcal{V}}_c O_{cb} X_a X_b. \quad (7.1.13)$$

Now we can do a change of variables, by defining

$$\tilde{X}_c = O_{cb}X_b, \quad \tilde{P}_c = O_{cb}P_b. \quad (7.1.14)$$

A key feature of this change of variables is that  $\tilde{X}_a$  and  $\tilde{P}_a$  have canonical commutators, since

$$[\tilde{X}_c, \tilde{P}_d] = O_{cb}O_{da}[X_b, P_a] = i\hbar O_{ca}O_{da} = i\hbar\delta_{cd}, \quad (7.1.15)$$

where the second equality used the canonical commutation relation for the original variables,  $[X_b, P_a] = i\hbar\delta_{ab}$ , and eq. (7.1.12) was used to get the last equality. Also,

$$\tilde{P}_a\tilde{P}_a = O_{ca}P_aO_{cb}P_b = \delta_{ab}P_aP_b = P_aP_a = P_x^2 + P_y^2 + P_z^2. \quad (7.1.16)$$

Therefore, the Hamiltonian in terms of the new canonical variables is

$$H = \sum_{a=1}^3 \left( \frac{\tilde{P}_a^2}{2m} + \frac{1}{2}m\omega_a^2\tilde{X}_a^2 \right), \quad (7.1.17)$$

where  $m\omega_a^2 = \tilde{\mathcal{V}}_a$ . This shows that  $H$  is the sum of three independent harmonic oscillator Hamiltonians, with possibly different natural frequencies. The general case is called the anisotropic three-dimensional harmonic oscillator. The special case  $\omega_x = \omega_y = \omega_z$  is called the isotropic three-dimensional harmonic oscillator; we will study it in more detail later in section 10.5, using its spherical coordinate wavefunction.

The preceding illustrates why many systems can be treated as if they were, effectively, systems of harmonic oscillators, with various higher-order effects that one may hope to either neglect or treat as perturbations. Of course, there are some important problems that are not approximated well by the harmonic oscillator. The free particle has no restoring force at all. The potential of a particle in a box or a square well is not close to its Taylor series expansion. The hydrogen atom has a potential with minimum  $-\infty$  at  $r = 0$ , and again the Taylor series expansion fails. Fortunately, these three cases are also exactly solvable, and one can say that together with the harmonic oscillator they are the four important examples of potential problems in nonrelativistic quantum mechanics that should be familiar to a well-educated physicist.

A third reason for the importance of the harmonic oscillator is that, as we will see, it has the unique property of equally spaced energy levels. This means that its operator algebra is suited for describing multiple energy excitations that can be added independently of each other. This turns out to have special relevance in quantum field theories, where free-particle modes with different momenta can be described using an infinite collection of harmonic oscillators, with couplings between them that describe particle interactions.

The energy eigenvalue problem of a harmonic oscillator in three dimensions can be solved as the tensor product of three one-dimensional harmonic oscillators. In the following two sections,

we will use two very different methods to derive the solutions of the one-dimensional harmonic oscillator with Hamiltonian

$$H = \frac{P^2}{2m} + \frac{1}{2}m\omega^2 X^2. \quad (7.1.18)$$

But first, we make a general observation: the eigenstates of this  $H$  must all have positive energy. This should be intuitively plausible, since both contributions to the classical energy are manifestly positive. To prove it formally in the quantum theory, note that in a stationary state  $|\psi\rangle$  with energy  $E$ ,

$$E = \langle\psi|H|\psi\rangle = \frac{1}{2m} \langle\psi|P^2|\psi\rangle + \frac{1}{2}m\omega^2 \langle\psi|X^2|\psi\rangle \quad (7.1.19)$$

$$= \frac{1}{2m} \langle P\psi|P\psi\rangle + \frac{1}{2}m\omega^2 \langle X\psi|X\psi\rangle > 0. \quad (7.1.20)$$

Here we have applied the fact that  $P$  and  $X$  are Hermitian, so that for example  $\langle\psi|P = \langle P^\dagger\psi| = \langle P\psi|$ , and then used the positivity of the inner product for non-null kets.

## 7.2 Position and momentum representations: the differential equations approach

In this section, we will solve for the energy eigenstates of the harmonic oscillator in the position representation. This involves solving a differential equation for the wavefunction. It must be admitted that this procedure is less elegant than the algebraic (energy representation) approach given in the following section. So why do we bother with it? Besides the goal of building character in the student, the differential equation method is important to learn because of its greater applicability; it can be used to solve the eigenvalue problem for many other Hamiltonians for which algebraic approaches are not available.

The position-representation version of the time-independent Schrödinger equation for the harmonic oscillator  $H|\psi\rangle = E|\psi\rangle$  is

$$\langle x| \left( \frac{P^2}{2m} + \frac{1}{2}m\omega^2 X^2 \right) |\psi\rangle = E \langle x|\psi\rangle, \quad (7.2.1)$$

or, in terms of the wavefunction  $\psi(x) = \langle x|\psi\rangle$ , after using  $X \rightarrow x$  and  $P \rightarrow -i\hbar d/dx$ ,

$$\frac{d^2\psi}{dx^2} + \frac{2m}{\hbar^2} \left( E - \frac{1}{2}m\omega^2 x^2 \right) \psi = 0. \quad (7.2.2)$$

It is convenient to introduce dimensionless variables corresponding to the position and energy,

$$y = x/b, \quad \mathcal{E} = E/\hbar\omega, \quad (7.2.3)$$

where we have defined a constant length scale

$$b = \sqrt{\hbar/m\omega}. \quad (7.2.4)$$

In terms of  $y$  and  $\mathcal{E}$ , eq. (7.2.2) becomes

$$\frac{d^2\psi}{dy^2} + (2\mathcal{E} - y^2)\psi = 0, \quad (7.2.5)$$

which we will now solve as an eigenvalue problem for  $\mathcal{E}$  and  $\psi(y)$ . Since  $H$  commutes with the parity operator  $\Pi$ , our solutions will be simultaneous eigenstates of energy and parity.

To help us further understand the character of the solutions, first consider the behavior at large  $|y|$ , where we can neglect the constant  $\mathcal{E}$  compared to  $y^2$ , so that  $d^2\psi/dy^2 \approx y^2\psi$ . Let us try a solution of the form

$$\psi = Ay^n e^{-\alpha y^2}, \quad (7.2.6)$$

where  $n$  and  $\alpha$  are constants. Then

$$\frac{d\psi}{dy} = A (ny^{n-1} - 2\alpha y^{n+1}) e^{-\alpha y^2} \approx -2\alpha y\psi, \quad (7.2.7)$$

where at large  $|y|$  the first term in parentheses is neglected compared to the second. Repeating this yields

$$\frac{d^2\psi}{dy^2} \approx (2\alpha y)^2\psi. \quad (7.2.8)$$

Therefore the guess eq. (7.2.6) indeed works for large  $y$ , with  $\alpha = \pm 1/2$ , regardless of the value of the constant  $n$ . We can reject  $\alpha = -1/2$  on physical grounds, since the wavefunction would blow up at large  $|y|$  and would not be normalizable. A purported wavefunction that grows exponentially with  $|y|$  must be unphysical because it would imply that no matter how far from the origin you look, the probability that the particle would be found farther away would be infinitely larger.

This motivates trying a solution of the form

$$\psi = u(y) e^{-y^2/2}, \quad \text{with} \quad u = \sum_{j=0}^{\infty} c_j y^j, \quad (7.2.9)$$

where the  $c_j$  are constants to be determined. No negative powers are included in this guess, because we expect that  $\psi$  will be well-behaved as  $y \rightarrow 0$ , where the potential smoothly vanishes.

Plugging eq. (7.2.9) into eq. (7.2.5) gives

$$\sum_{j=0}^{\infty} c_j [j(j-1)y^{j-2} + (2\mathcal{E} - 1 - 2j)y^j] = 0. \quad (7.2.10)$$

Now we use a trick on the first term. Since it vanishes for  $j = 0$  and  $j = 1$ , we rewrite it as  $\sum_{j=2}^{\infty} c_j j(j-1)y^{j-2}$ , and then let  $j = k+2$  so that it becomes  $\sum_{k=0}^{\infty} c_{k+2}(k+2)(k+1)y^k$ , and finally rename  $k \rightarrow j$ , so that it becomes  $\sum_{j=0}^{\infty} c_j(j+2)(j+1)y^j$ . Now it can be nicely reunited with the second term, combining like powers of  $y$  to give

$$\sum_{j=0}^{\infty} y^j [c_{j+2}(j+1)(j+2) + c_j(2\mathcal{E} - 1 - 2j)] = 0. \quad (7.2.11)$$

Because each power  $y^j$  in the sum is independent, their coefficients must vanish separately for each  $j$ , so we have a recurrence relation,

$$c_{j+2} = \frac{2j+1-2\mathcal{E}}{(j+1)(j+2)} c_j. \quad (7.2.12)$$

This shows that we only need to know two constants  $c_0$  and  $c_1$ , because if  $c_0$  is known, then so are  $c_2, c_4, c_6, \dots$ , and if  $c_1$  is known, then so are  $c_3, c_5, c_7$ , etc. So, the solution is

$$\begin{aligned} u(y) = & c_0 \left[ 1 + \left( \frac{1-2\mathcal{E}}{2} \right) y^2 + \left( \frac{1-2\mathcal{E}}{2} \right) \left( \frac{5-2\mathcal{E}}{12} \right) y^4 + \dots \right] \\ & + c_1 \left[ y + \left( \frac{3-2\mathcal{E}}{6} \right) y^3 + \left( \frac{3-2\mathcal{E}}{6} \right) \left( \frac{7-2\mathcal{E}}{20} \right) y^5 + \dots \right], \end{aligned} \quad (7.2.13)$$

which, naively, appears to be an infinite series.

However, there is something horribly wrong with the solution if it is really an infinite series. For sufficiently large  $|y|$ , the series will be dominated by terms with large powers  $j$ , where eq. (7.2.12) appears to give

$$\frac{c_{j+2}}{c_j} \approx \frac{2}{j + \mathcal{E} + 5/2}, \quad (7.2.14)$$

up to contributions to the denominator that vanish as  $j \rightarrow \infty$ . If arbitrarily large powers  $j$  are present, the function  $u(y)$  will necessarily grow too fast as  $|y| \rightarrow \infty$ . To see this, consider for comparison the function  $f(y) = y^p e^{y^2}$ , with series expansion

$$f(y) = \sum_{k=0}^{\infty} y^{2k+p}/k!. \quad (7.2.15)$$

Now, writing  $j = 2k + p$ , this series has terms  $C_j y^j$  where  $C_{j+2}/C_j = 1/(k+1) = 2/(j-p+2)$ . Comparing to eq. (7.2.14), we see that if  $u(y)$  is really a non-terminating series in powers of  $y$ , then  $u(y) \sim y^p e^{y^2}$  for  $p = -\mathcal{E} - 1/2$ , and the wavefunction behaves for large  $|y|$  like

$$\psi(y) \sim (y^p e^{y^2}) e^{-y^2/2} = y^p e^{y^2/2}. \quad (7.2.16)$$

These are recognized as the unphysical solutions with  $\alpha = -1/2$  that we had already rejected.



For a sensible physical solution, there is only one way out: the series for  $u(y)$  must terminate. To find a basis of such solutions, first consider  $c_0 \neq 0$  with  $c_1 = 0$ , so that only even-index coefficients are present. If one of them vanishes,  $c_{2k} = 0$  for some  $k$ , then all higher coefficients  $c_j$  with  $j > 2k$  will also vanish, according to the recurrence relation. Likewise, we can take  $c_1 \neq 0$  with  $c_0 = 0$ , so that only odd-index coefficients are present. If one of them vanishes,  $c_{2k+1} = 0$  for some  $k$ , then all  $c_j$  with  $j > 2k + 1$  will also vanish.

In either case, the condition for the series in  $u(y)$  to terminate, yielding a physical solution, is that the numerator in the recurrence relation eq. (7.2.12) must vanish for some non-negative integer  $j = n$ . Therefore, the allowed energy eigenvalues are  $\mathcal{E} = n + 1/2$ , or

$$E_n = (n + 1/2)\hbar\omega, \quad (n = 0, 1, 2, \dots). \quad (7.2.17)$$

It follows that  $u(y)$  is a polynomial of degree  $n$  in  $y$ , and contains only even (odd) powers of  $y$  if  $n$  is even (odd). For any given  $n$ , they can be constructed from the recurrence relation, eq. (7.2.12), up to an overall multiplicative constant given by either  $c_0$  or  $c_1$ . The resulting  $u(y) = H_n(y)$  for a given  $n$  are called **Hermite polynomials**. From eqs. (7.2.5) and (7.2.9) with  $\mathcal{E} = n + 1/2$ , they satisfy the differential equation

$$\left( \frac{d^2}{dy^2} - 2y \frac{d}{dy} + 2n \right) H_n = 0. \quad (7.2.18)$$

A general expression for the Hermite polynomials is

$$H_n(y) = e^{y^2/2} \left( y - \frac{d}{dy} \right)^n e^{-y^2/2}. \quad (7.2.19)$$

This can be verified by plugging it into the differential equation (7.2.18). Here we are relying on the fact that stationary bound states in 1-dimensional problems are non-degenerate, as we found at the end of section 6.3, and therefore unique (up to a normalization constant) for a given  $n$ . An even nicer derivation of eq. (7.2.19) will be found near the end of the section 7.3.

The first few Hermite polynomials are

$$H_0(y) = 1, \quad H_1(y) = 2y, \quad (7.2.20)$$

$$H_2(y) = -2 + 4y^2, \quad H_3(y) = -12y + 8y^3, \quad (7.2.21)$$

$$H_4(y) = 12 - 48y^2 + 16y^4, \quad H_5(y) = 120y - 160y^3 + 32y^5. \quad (7.2.22)$$

Using eq. (7.2.19), one can establish the identities

$$H_n(-y) = (-1)^n H_n(y), \quad (7.2.23)$$

$$\frac{d}{dy} H_n(y) = 2n H_{n-1}(y), \quad (7.2.24)$$

$$H_{n+1}(y) = 2y H_n(y) - 2n H_{n-1}(y). \quad (7.2.25)$$

It is useful to define the function  $F(y, t) = \sum_{n=0}^{\infty} \frac{t^n}{n!} H_n(y)$ , which can be evaluated as follows. First, using eq. (7.2.24), one finds the differential equation  $\partial F / \partial y = 2tF$ , which has the solution  $F(t, y) = F(t, 0)e^{2ty}$ . Then using eq. (7.2.25), one finds  $\partial F / \partial t = 2(y - t)F$ , which yields  $\partial F(t, 0) / \partial t = -2tF(t, 0)$ , which has the solution  $F(t, 0) = F(0, 0)e^{-t^2}$ . Since  $F(0, 0) = H_0(0) = 1$ , we arrive at the **generating function for Hermite polynomials**,

$$F(y, t) = \sum_{n=0}^{\infty} \frac{t^n}{n!} H_n(y) = \exp(2yt - t^2). \quad (7.2.26)$$

One use of the generating function is to find the orthonormality properties of the Hermite polynomials. Consider the product of two generating functions with  $e^{-y^2}$ ,

$$e^{-y^2} \sum_{n=0}^{\infty} \frac{t^n}{n!} H_n(y) \sum_{m=0}^{\infty} \frac{s^m}{m!} H_m(y) = e^{-y^2 + 2y(t+s) - s^2 - t^2}. \quad (7.2.27)$$

Now integrating both sides with respect to  $y$ , we get

$$\sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{t^n}{n!} \frac{s^m}{m!} \left( \int_{-\infty}^{\infty} dy e^{-y^2} H_n(y) H_m(y) \right) = \sqrt{\pi} e^{2ts} \quad (7.2.28)$$

The right side can be expanded as

$$\sqrt{\pi} e^{2ts} = \sqrt{\pi} \sum_{n=0}^{\infty} \frac{2^n t^n s^n}{n!} = \sqrt{\pi} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \delta_{nm} \frac{2^n t^n s^m}{n!} \quad (7.2.29)$$

Comparing the coefficients of  $t^n s^m$ , we arrive at the orthonormality condition

$$\int_{-\infty}^{\infty} dy e^{-y^2} H_n(y) H_m(y) = \delta_{nm} \sqrt{\pi} 2^n n!. \quad (7.2.30)$$

The orthonormal energy basis wavefunctions for the harmonic oscillator can now be written

$$\psi_n(x) = \left( \frac{m\omega}{\pi\hbar} \right)^{1/4} \sqrt{\frac{1}{2^n n!}} H_n(y) e^{-y^2/2}, \quad (7.2.31)$$

where  $y = x/b = x\sqrt{m\omega/\hbar}$ , and the multiplicative constant has been chosen so that

$$\int_{-\infty}^{\infty} dx (\psi_n(x))^* \psi_m(x) = \delta_{nm}. \quad (7.2.32)$$

(Actually, the wavefunctions are all real, so the complex conjugation does nothing in this case.) In particular, the normalized ground state wavefunction is a pure Gaussian,

$$\psi_0(x) = \left( \frac{m\omega}{\pi\hbar} \right)^{1/4} \exp(-m\omega x^2 / 2\hbar). \quad (7.2.33)$$

The wavefunctions  $\psi_n$  and the corresponding probability densities  $|\psi_n|^2$  are shown for  $n = 0, 1, 2, 3, 4$ , and 16 in Figure 7.2.1.




Figure 7.2.1: Harmonic oscillator stationary-state wavefunctions  $\psi_n$  in units of  $(m\omega/\hbar)^{1/4}$  (left column) and probability densities  $|\psi_n|^2$  in units of  $\sqrt{m\omega/\hbar}$  (right column) for  $n = 0, 1, 2, 3, 4$ , and 16, as functions of  $y = x\sqrt{m\omega/\hbar}$ .

The wavefunctions are related to the energy eigenstate kets  $|n\rangle$  of the harmonic oscillator by

$$\langle x|n\rangle = \psi_n(x), \quad (7.2.34)$$

with

$$H|n\rangle = \hbar\omega(n + 1/2)|n\rangle. \quad (7.2.35)$$

This shows that the energy levels of the one-dimensional harmonic oscillator are equally spaced and non-degenerate, with a quantum of energy given by  $\Delta E = \hbar\omega$ . There is also a **zero-point energy** of the ground state,  $E_0 = \hbar\omega/2$ , in agreement with our earlier proof that the energies had to be positive. From eq. (7.2.23), the states with even  $n$  have even parity, and those with odd  $n$  have odd parity.

Another interesting relation satisfied by the harmonic oscillator wavefunctions, and thus the Hermite polynomials, is

$$\sum_{n=0}^{\infty} \psi_n(x') \psi_n(x) = \delta(x - x'). \quad (7.2.36)$$

This can be derived immediately from  $\langle x|x'\rangle = \delta(x - x')$  by applying the completeness relation  $\sum_{n=0}^{\infty} |n\rangle \langle n| = 1$ .

Having found the position wavefunctions for the harmonic oscillator, we now turn to the momentum representation. One way to evaluate the momentum wavefunctions of the energy eigenstates  $|n\rangle$  is to use the completeness relation:

$$\tilde{\psi}_n(p) = \langle p|n\rangle = \int_{-\infty}^{\infty} dx \langle p|x\rangle \langle x|n\rangle = \int_{-\infty}^{\infty} dx \frac{1}{\sqrt{2\pi\hbar}} e^{-ipx/\hbar} \psi_n(x), \quad (7.2.37)$$

but it may not be immediately obvious how to evaluate the integral for general  $n$ . Fortunately, we can gain some insight by noting that the Hamiltonian in eq. (7.1.18) has the very special property of being invariant under the simultaneous substitutions

$$X \leftrightarrow P, \quad m\omega \leftrightarrow \frac{1}{m\omega}. \quad (7.2.38)$$

Since the energy eigenstates are non-degenerate, their momentum wavefunctions must be given, up to a phase  $e^{i\varphi_n}$  to be determined, by making these same substitutions in eq. (7.2.31). Therefore, defining a dimensionless variable proportional to momentum,

$$v = p/\sqrt{\hbar\omega m}, \quad (7.2.39)$$

it must be that

$$\tilde{\psi}_n(p) = e^{i\varphi_n} \frac{1}{(\pi\hbar\omega m)^{1/4}} \sqrt{\frac{1}{2^n n!}} H_n(v) e^{-v^2/2}. \quad (7.2.40)$$

The phase factor  $e^{i\varphi_n}$  must be chosen consistently with the phase convention of the ket  $|n\rangle$  that has already been fixed by eqs. (7.2.31) and (7.2.34). For the lowest few  $n = 0, 1, 2, 3, \dots$ , you can do the integral in eq. (7.2.37) to check that eq. (7.2.40) is indeed true with

$$e^{i\varphi_n} = (-i)^n, \quad (7.2.41)$$

and we will prove it for general  $n$  at the end of the next section.

### 7.3 Energy representation: the algebraic approach

In this section, we will use a different, and more elegant, method to solve for the stationary states of the harmonic oscillator. In this approach, due to Dirac, we use algebraic methods rather than differential equations, working directly in the energy basis.

We begin by defining dimensionless creation and annihilation operators (also known as raising and lowering or destruction operators, or together as ladder operators) by

$$a = \sqrt{\frac{m\omega}{2\hbar}}X + i\frac{1}{\sqrt{2\hbar\omega m}}P, \quad (7.3.1)$$

$$a^\dagger = \sqrt{\frac{m\omega}{2\hbar}}X - i\frac{1}{\sqrt{2\hbar\omega m}}P. \quad (7.3.2)$$

As the notation indicates, these are not Hermitian operators and so are not observables, but rather are Hermitian adjoints of each other. Equivalently, one can write the relationship as

$$X = \sqrt{\frac{\hbar}{2m\omega}}(a^\dagger + a), \quad (7.3.3)$$

$$P = i\sqrt{\frac{\hbar\omega m}{2}}(a^\dagger - a). \quad (7.3.4)$$

Using the canonical commutation relations  $[X, P] = i\hbar$ , we find that

$$[a, a^\dagger] = 1. \quad (7.3.5)$$

We also can compute

$$a^\dagger a = \frac{m\omega}{2\hbar}X^2 + \frac{1}{2\hbar\omega m}P^2 + \frac{i}{2\hbar}[X, P] = H/\hbar\omega - 1/2,$$

so that the Hamiltonian is simply<sup>‡</sup>

$$H = \hbar\omega(a^\dagger a + 1/2). \quad (7.3.6)$$

---

<sup>‡</sup>One could also write the equivalent form  $H = \hbar\omega(a^\dagger a + aa^\dagger)/2$ . In choosing to write eq. (7.3.6), we have followed the systematic protocol known as **normal ordering**. To “normal order” an operator means to rewrite it by moving all  $a$  operators to the right and all  $a^\dagger$  operators to the left, using  $aa^\dagger = a^\dagger a + 1$  (which is equivalent to the commutation relation) as many times as necessary.

Using eq. (7.3.5), one then obtains

$$[H, a] = -\hbar\omega a, \quad [H, a^\dagger] = \hbar\omega a^\dagger. \quad (7.3.7)$$

Now suppose that we have an energy eigenstate  $|E\rangle$ , with energy eigenvalue  $E$ . Consider the Hamiltonian acting on the state  $a^\dagger|E\rangle$ :

$$H(a^\dagger|E\rangle) = a^\dagger H|E\rangle + [H, a^\dagger]|E\rangle = (E + \hbar\omega)a^\dagger|E\rangle. \quad (7.3.8)$$

This shows that  $a^\dagger|E\rangle$  is an eigenstate of  $H$  with energy  $E + \hbar\omega$ . Repeating this  $n$  times, the state  $(a^\dagger)^n|E\rangle$  must be an energy eigenstate with energy  $E + n\hbar\omega$ . Similarly,

$$H(a|E\rangle) = (E - \hbar\omega)a|E\rangle, \quad (7.3.9)$$

so the state  $a^n|E\rangle$  is apparently an energy eigenstate with energy  $E - n\hbar\omega$ .

Thus, naively it might appear that, given  $|E\rangle$ , we can construct an infinite chain of energy eigenstates with both arbitrarily low and arbitrarily high energies,

$$\dots, |E - 2\hbar\omega\rangle, |E - \hbar\omega\rangle, |E\rangle, |E + \hbar\omega\rangle, |E + 2\hbar\omega\rangle, \dots \quad (7.3.10)$$

But this cannot be true, since we proved at the end of section 7.1 that all of the energy eigenvalues of the harmonic oscillator are positive. The only way out is that all of the kets in the chain with negative energy are actually the null ket. Let us rename the state with lowest non-negative energy as  $|0\rangle$ , where the label 0 is not the energy, but signifies that this is the ground state. It must satisfy

$$a|0\rangle = 0, \quad (7.3.11)$$

so that all of the kets  $a^n|0\rangle = 0$  for  $n \geq 1$  are null and do not actually exist as physical states. Since  $a^\dagger a|0\rangle = 0$ , we have

$$(H/\hbar\omega - 1/2)|0\rangle = 0, \quad (7.3.12)$$

so the ground state must have  $E_0 = \hbar\omega/2$ , a result that we had also found in the previous section using the differential equation approach.

In section 6.3, we proved that bound states in one-dimensional quantum mechanics never have degenerate energies. This implies that there is a unique state  $|0\rangle$  with  $E_0 = \hbar\omega/2$ , and unique states with  $E_n = \hbar\omega(n+1/2)$  for  $n = 0, 1, 2, \dots$ , just as we had found by the differential equations method in eq. (7.2.17). Up to normalization, the energy eigenstates  $|n\rangle$  are proportional to  $(a^\dagger)^n|0\rangle$ . Since they are non-degenerate eigenkets of a Hermitian operator ( $H$ ), they can be normalized to form an orthobasis,

$$\langle k|n\rangle = \delta_{nk}. \quad (7.3.13)$$

Let  $|0\rangle$  have norm 1. Then we can define the other orthobasis kets by recurrence, using

$$|n\rangle = c_n a^\dagger |n-1\rangle, \quad (7.3.14)$$

where  $c_n$  are normalization constants to be determined. We have

$$\langle n|n\rangle = |c_n|^2 \langle n-1|aa^\dagger|n-1\rangle = n|c_n|^2 \langle n-1|n-1\rangle, \quad (7.3.15)$$

where the second equality makes use of

$$aa^\dagger = a^\dagger a + 1 = H/\hbar\omega + 1/2 \quad (7.3.16)$$

and then  $H|n-1\rangle = \hbar\omega(n-1/2)|n-1\rangle$ . Equation (7.3.15) shows that  $\langle n|n\rangle = \langle n-1|n-1\rangle = 1$  requires  $c_n = 1/\sqrt{n}$ , where we have made an arbitrary choice of phase. From this, we use eqs. (7.3.14) and (7.3.16) to deduce that

$$a^\dagger|n\rangle = \sqrt{n+1}|n+1\rangle, \quad (7.3.17)$$

$$a|n\rangle = \sqrt{n}|n-1\rangle, \quad (7.3.18)$$

so that  $a^\dagger$  raises the energy of the state (or creates an energy quantum), and  $a$  lowers the energy (or destroys an energy quantum). Taking the Hermitian conjugate gives

$$\langle n|a = \sqrt{n+1} \langle n+1|, \quad (7.3.19)$$

$$\langle n|a^\dagger = \sqrt{n} \langle n-1|. \quad (7.3.20)$$

It follows that the matrix elements of  $a^\dagger$  and  $a$  in the energy eigenstate orthobasis are

$$\langle k|a^\dagger|n\rangle = \sqrt{n+1} \delta_{k,n+1}, \quad \langle k|a|n\rangle = \sqrt{n} \delta_{k,n-1}. \quad (7.3.21)$$

Another consequence is that the  $n$ th excited state can be written in terms of  $n$  creation operators acting on the ground state,

$$|n\rangle = \frac{(a^\dagger)^n}{\sqrt{n!}}|0\rangle. \quad (7.3.22)$$

The Hamiltonian is sometimes written as  $H = \hbar\omega(N+1/2)$ , where  $N = a^\dagger a$  is called the number operator. It is clearly Hermitian, and satisfies

$$N|n\rangle = n|n\rangle, \quad (7.3.23)$$

so it is the observable that just measures the number of energy quanta in the state.

The matrix elements of the position and momentum operators in the energy eigenstate basis follow immediately from eqs. (7.3.3) and (7.3.4) using (7.3.21), with the results

$$\langle k|X|n\rangle = \sqrt{\frac{\hbar}{2m\omega}} \left( \sqrt{n+1} \delta_{k,n+1} + \sqrt{n} \delta_{k,n-1} \right), \quad (7.3.24)$$

$$\langle k|P|n\rangle = i\sqrt{\frac{\hbar\omega m}{2}} \left( \sqrt{n+1} \delta_{k,n+1} - \sqrt{n} \delta_{k,n-1} \right). \quad (7.3.25)$$

In particular, their expectation values in any energy eigenstate  $|n\rangle$  vanish,

$$\langle n|X|n\rangle = 0, \quad \langle n|P|n\rangle = 0. \quad (7.3.26)$$

This is an example of the parity selection rule, Theorem 5.4.1, since  $X$  and  $P$  each have odd parity. One can also compute, using eqs. (7.3.17) and (7.3.18),

$$\begin{aligned} X^2|n\rangle &= \frac{\hbar}{2m\omega} (a^\dagger + a)(a^\dagger + a)|n\rangle \\ &= \frac{\hbar}{2m\omega} \left( \sqrt{(n+1)(n+2)}|n+2\rangle + (2n+1)|n\rangle + \sqrt{n(n-1)}|n-2\rangle \right), \end{aligned} \quad (7.3.27)$$

with the consequence that

$$\langle n|X^2|n\rangle = \frac{\hbar}{m\omega} (n+1/2). \quad (7.3.28)$$

In a very similar way,

$$\langle n|P^2|n\rangle = \hbar\omega m(n+1/2). \quad (7.3.29)$$

As a check,

$$\langle n|H|n\rangle = \frac{1}{2m} \langle n|P^2|n\rangle + \frac{m\omega^2}{2} \langle n|X^2|n\rangle = \hbar\omega(n+1/2), \quad (7.3.30)$$

in agreement with the result for  $E_n$  that we found in eq. (7.2.17). Furthermore, the uncertainties in position and momentum,  $\Delta X = \sqrt{\langle n|X^2|n\rangle}$  and  $\Delta P = \sqrt{\langle n|P^2|n\rangle}$  each grow with  $n$ , proportionally to  $\sqrt{E_n}$  or  $\sqrt{n+1/2}$ .

To further illustrate the power and convenience of the algebraic method, suppose that for some reason we needed to evaluate the matrix element  $\langle 3|X^3|2\rangle$ . We have, using eq. (7.3.3),

$$\langle 3|X^3|2\rangle = \left( \frac{\hbar}{2m\omega} \right)^{3/2} \langle 3|(a^\dagger + a)^3|2\rangle \quad (7.3.31)$$

$$= \left( \frac{\hbar}{2m\omega} \right)^{3/2} \langle 3| \left( \cancel{a^{\dagger 3}} + a^{\dagger 2}a + a^\dagger a a^\dagger + a a^{\dagger 2} + \cancel{a^\dagger a^2} + \cancel{a a^\dagger a} + \cancel{a^2 a^\dagger} + \cancel{a^3} \right) |2\rangle. \quad (7.3.32)$$

Here, we have crossed out terms that can be immediately seen to give no contribution by simply counting quanta created and destroyed. For the first term, we start in the ket on the far right



with 2 quanta, and create 3 more, so it is proportional to  $|5\rangle$ , which by orthonormality has vanishing inner product with the bra  $\langle 3|$  on the left. Likewise, the fifth, sixth, and seventh terms cannot contribute because we start with 2 quanta and then (in various orders) create 1 more but destroy 2, resulting in a state with 1, which again has vanishing inner product with the 3-quanta bra. And, the last term immediately vanishes because  $a^3|2\rangle = 0$ . The remaining three terms do not vanish, but can be easily evaluated with nothing but simple arithmetic, by applying eqs. (7.3.17) and (7.3.18) repeatedly. In contrast, evaluation of the same matrix element in the position representation approach of section 7.2 would read

$$\begin{aligned} \langle 3|X^3|2\rangle &= \int_{-\infty}^{\infty} dx \left( \frac{m\omega}{\pi\hbar 2^6 (3!)^2} \right)^{1/4} H_3 \left( x\sqrt{m\omega/\hbar} \right) e^{-m\omega x^2/2\hbar} x^3 \\ &\quad \left( \frac{m\omega}{\pi\hbar 2^4 (2!)^2} \right)^{1/4} H_2 \left( x\sqrt{m\omega/\hbar} \right) e^{-m\omega x^2/2\hbar}. \end{aligned} \quad (7.3.33)$$

This is certainly doable, but less pleasant.

Let us now see how to connect the energy and position representations, by writing  $a$  and  $a^\dagger$  as differential operators in the latter. In the position representation,  $X \rightarrow x$  and  $P \rightarrow -i\hbar d/dx$ , so from eqs. (7.3.1) and (7.3.2) we get

$$a = \frac{1}{\sqrt{2}} \left( y + \frac{d}{dy} \right), \quad a^\dagger = \frac{1}{\sqrt{2}} \left( y - \frac{d}{dy} \right), \quad (7.3.34)$$

where  $y = x/b = x\sqrt{m\omega/\hbar}$  as before. Therefore, the condition  $a|0\rangle = 0$  gives

$$\left( y + \frac{d}{dy} \right) \psi_0(y) = 0. \quad (7.3.35)$$

This implies  $d\psi_0/\psi_0 = -ydy$ , or  $d(\ln \psi_0) = -d(y^2/2)$ , so that upon integration,  $\ln \psi_0 = -y^2/2 + \ln A_0$ , where  $\ln A_0$  is a constant of integration. Thus,

$$\psi_0(y) = A_0 e^{-y^2/2} = A_0 e^{-m\omega x^2/2\hbar}, \quad (7.3.36)$$

in agreement with eq. (7.2.33) after fixing the normalization constant  $A_0 = (m\omega/\pi\hbar)^{1/4}$ . Now, combining eqs. (7.3.22) and (7.3.34), we have

$$\psi_n(x) = \langle x|n\rangle = \frac{1}{\sqrt{n!}} \left[ \frac{1}{\sqrt{2}} \left( y - \frac{d}{dy} \right) \right]^n \left( \frac{m\omega}{\pi\hbar} \right)^{1/4} e^{-y^2/2}. \quad (7.3.37)$$

Comparing this with eq. (7.2.31) yields the general form for the Hermite polynomials given in eq. (7.2.19), as promised.

The energy and momentum representations can be connected in a similar way. The momentum representations are  $P \rightarrow p$  and  $X \rightarrow i\hbar d/dp$ , from which one finds

$$a = \frac{i}{\sqrt{2}} \left( v + \frac{d}{dv} \right), \quad a^\dagger = -\frac{i}{\sqrt{2}} \left( v - \frac{d}{dv} \right), \quad (7.3.38)$$

where  $v = p/\sqrt{\hbar\omega m}$  is the dimensionless rescaled momentum, as in eqs. (7.2.39) and (7.2.40). Evaluating eq. (7.2.37) for the special case  $n = 0$ , we then find

$$\tilde{\psi}_0(p) = \frac{1}{(\pi\hbar\omega m)^{1/4}} e^{-v^2/2} \quad (7.3.39)$$

for the ground state. Using eq. (7.3.38) in eq. (7.3.22), we obtain

$$\tilde{\psi}_n(p) = \langle p|n\rangle = \frac{1}{\sqrt{n!}} \left[ -\frac{i}{\sqrt{2}} \left( v - \frac{d}{dv} \right) \right]^n \frac{1}{(\pi\hbar\omega m)^{1/4}} e^{-v^2/2}. \quad (7.3.40)$$

Now comparing to eq. (7.2.19), we obtain eq. (7.2.40) with  $e^{i\varphi_n} = (-i)^n$ , as promised.

## 7.4 Coherent states of the harmonic oscillator

Consider a macroscopic harmonic oscillator of the type one might encounter in a lab experiment in an introductory physics course, with mass  $m = 0.2$  kg,  $\omega = 10$  radians/second, and amplitude  $x_0 = 0.1$  meters. Classically, the energy can be obtained as the potential energy at the extremum of the displacement, when  $p = 0$  and  $x = x_0$ , so that  $E = m\omega^2 x_0^2/2 = 0.1$  J. The energy quantum associated with the oscillator is  $\hbar\omega = 1.055 \times 10^{-34}$  J. Therefore, we expect macroscopic oscillator states to have enormous numbers of energy quanta, roughly  $n = E/\hbar\omega \approx 10^{32}$  in this example.

Is the energy eigenstate  $|n = 10^{32}\rangle$  classical-like? The answer is clearly no, since in any energy eigenstate,  $\langle X\rangle(t) = 0$  and  $\langle P\rangle(t) = 0$  for all times  $t$  by the parity selection rule, while the classical trajectory, eq. (7.1.5), involves macroscopic oscillations. We would like to find quantum states that are approximately classical, by which we mean that  $\langle X\rangle(t) \approx x_{\text{cl}}(t)$  and  $\langle P\rangle(t) \approx p_{\text{cl}}(t)$  should be as close as possible to the solutions in eq. (7.1.5). The states that accomplish this are called **coherent** (or **quasi-classical**) states, and the argument just given ensures that they cannot be energy eigenstates, or parity eigenstates.

Instead, with the wisdom of foresight, let us try kets  $|\alpha\rangle$  that are eigenstates of the lowering operator  $a$  with eigenvalue  $\alpha$ ,

$$a|\alpha\rangle = \alpha|\alpha\rangle. \quad (7.4.1)$$

Since  $a$  is not Hermitian, we have no reason to expect that the allowed eigenvalues  $\alpha$  will be real, and indeed it will turn out to be very important that they are complex in general. For the same reason, we also cannot expect that the set of all coherent states  $|\alpha\rangle$  for different  $\alpha$  will form an orthobasis or even be orthogonal, and again they are not.

To construct the coherent states that satisfy eq. (7.4.1), let us try an arbitrary linear combination of energy eigenstates,

$$|\alpha\rangle = \sum_{n=0}^{\infty} c_n |n\rangle, \quad (7.4.2)$$

with coefficients  $c_n$  to be determined. Then,

$$a|\alpha\rangle = \sum_{n=0}^{\infty} c_n a|n\rangle = \sum_{n=1}^{\infty} c_n \sqrt{n} |n-1\rangle, \quad (7.4.3)$$

where in the second equality we have used eq. (7.3.18), and started the sum from  $n = 1$  by exploiting the fact that the  $n = 0$  term vanishes. If we now relabel  $n \rightarrow n + 1$ , and require that the result

$$a|\alpha\rangle = \sum_{n=0}^{\infty} c_{n+1} \sqrt{n+1} |n\rangle \quad (7.4.4)$$

is equal to  $\alpha \sum_{n=0}^{\infty} c_n |n\rangle$ , we obtain a recurrence relation

$$c_{n+1} = \frac{\alpha}{\sqrt{n+1}} c_n. \quad (7.4.5)$$

Starting with  $c_0$ , we get  $c_1 = \alpha c_0$ ,  $c_2 = \alpha^2 c_0 / \sqrt{2}$ , etc., or in general  $c_n = \alpha^n c_0 / \sqrt{n!}$ . Therefore, up to normalization,

$$|\alpha\rangle = c_0 \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} |n\rangle. \quad (7.4.6)$$

To fix  $c_0$ , we require  $\langle\alpha|\alpha\rangle = 1$ , or

$$1 = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} c_0^* \frac{\alpha^{*m}}{\sqrt{m!}} c_0 \frac{\alpha^n}{\sqrt{n!}} \langle m|n\rangle = |c_0|^2 \sum_{n=0}^{\infty} \frac{|\alpha|^{2n}}{n!} = |c_0|^2 e^{|\alpha|^2}, \quad (7.4.7)$$

where the second equality uses the orthonormality  $\langle m|n\rangle = \delta_{nm}$  to collapse the sum over  $m$ . Therefore,  $c_0 = e^{-|\alpha|^2/2}$  (up to the usual arbitrary global phase), and the normalized coherent state with complex eigenvalue  $\alpha$  is

$$|\alpha\rangle = e^{-|\alpha|^2/2} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} |n\rangle. \quad (7.4.8)$$

Using eq. (7.3.22), this can be rewritten as

$$|\alpha\rangle = e^{-|\alpha|^2/2} \sum_{n=0}^{\infty} \frac{(\alpha a^\dagger)^n}{n!} |0\rangle = e^{-|\alpha|^2/2} e^{\alpha a^\dagger} |0\rangle. \quad (7.4.9)$$

If a harmonic oscillator is in a coherent state  $|\alpha\rangle$ , and the energy is measured, all results  $E_n = \hbar\omega(n + 1/2)$  can occur, with probabilities

$$\mathcal{P}_n = |\langle n|\alpha\rangle|^2 = \frac{|\alpha|^{2n}}{n!} e^{-|\alpha|^2}. \quad (7.4.10)$$

These do not depend on the phase of  $\alpha$ , and form a Poisson distribution with mean  $|\alpha|^2$ . Since

$$\mathcal{P}_n = \frac{|\alpha|^2}{n} \mathcal{P}_{n-1}, \quad (7.4.11)$$

we see that the probability increases with  $n$  as long as  $n < |\alpha|^2$ , but decreases thereafter. Therefore, the most probable result of an energy measurement is of order  $\hbar\omega|\alpha|^2$ . This shows that for a macroscopic oscillator like the one posed at the beginning of this section,  $|\alpha| \sim \sqrt{n} \sim 10^{16}$  in a coherent state. The expectation value of the energy is

$$\langle \alpha | H | \alpha \rangle = \hbar\omega \left[ (\langle \alpha | a^\dagger ) (a | \alpha \rangle) + 1/2 \right] = \hbar\omega \left[ (\langle \alpha | \alpha^* ) (\alpha | \alpha \rangle) + 1/2 \right] = \left( |\alpha|^2 + \frac{1}{2} \right) \hbar\omega, \quad (7.4.12)$$

giving a similar measure of the average energy.

Next let us compute the uncertainty in the energy. First, we need

$$\langle \alpha | H^2 | \alpha \rangle = (\hbar\omega)^2 \left[ \langle \alpha | a^\dagger a a^\dagger a | \alpha \rangle + \langle \alpha | a^\dagger a | \alpha \rangle + \frac{1}{4} \langle \alpha | \alpha \rangle \right] \quad (7.4.13)$$

$$= (\hbar\omega)^2 (|\alpha|^4 + 2|\alpha|^2 + 1/4). \quad (7.4.14)$$

Therefore

$$\Delta H = \sqrt{\langle H^2 \rangle - \langle H \rangle^2} = \hbar\omega |\alpha|. \quad (7.4.15)$$

This is a very small energy uncertainty compared to the energy expectation value,

$$\Delta H / \langle H \rangle = 1/|\alpha| \ll 1, \quad (7.4.16)$$

since we found  $1/|\alpha|$  of order  $10^{-16}$  in our numerical example. Thus, the energy distribution is peaked extremely sharply about the classical value.

We can likewise compute the expectation value and uncertainty of the position. First,

$$\langle \alpha | X | \alpha \rangle = \sqrt{\frac{\hbar}{2m\omega}} \langle \alpha | (a^\dagger + a) | \alpha \rangle = \sqrt{\frac{\hbar}{2m\omega}} (\alpha^* + \alpha) = \sqrt{\frac{2\hbar}{m\omega}} \text{Re}[\alpha]. \quad (7.4.17)$$

Also, one has

$$\langle \alpha | X^2 | \alpha \rangle = \frac{\hbar}{2m\omega} \langle \alpha | (a^\dagger + a)^2 | \alpha \rangle = \frac{\hbar}{2m\omega} \langle \alpha | (a^{\dagger 2} + 2a^\dagger a + a^2 + 1) | \alpha \rangle \quad (7.4.18)$$

$$= \frac{\hbar}{2m\omega} [(\alpha^* + \alpha)^2 + 1]. \quad (7.4.19)$$

It follows that

$$\Delta X = \sqrt{\langle \alpha | X^2 | \alpha \rangle - \langle \alpha | X | \alpha \rangle^2} = \sqrt{\frac{\hbar}{2m\omega}}. \quad (7.4.20)$$

Remarkably, this does not depend on  $\alpha$  at all, and in fact is exactly the same as for the ground state, making it ridiculously tiny by macroscopic standards. The interpretation of  $\Delta X / \langle X \rangle \ll 1$

is that the position wavefunction is extremely sharply peaked about its expectation value. In a similar way, one can compute the expectation value and the uncertainty of the momentum:

$$\langle \alpha | P | \alpha \rangle = \sqrt{2m\hbar\omega} \operatorname{Im}[\alpha], \quad (7.4.21)$$

$$\Delta P = \sqrt{\frac{m\hbar\omega}{2}}. \quad (7.4.22)$$

The latter is again independent of  $\alpha$  and is the same as in the ground state. Putting these results together, the product of uncertainties in position and momentum is

$$(\Delta X)(\Delta P) = \hbar/2, \quad (7.4.23)$$

which is exactly as small as it could possibly be, consistent with the uncertainty relation.

The preceding results suggest that there is some close relation between the ground state and the coherent states, since they have exactly the same uncertainties in position and momentum. To see this, let us work out the wavefunction of the coherent state  $|\alpha\rangle$ . We start with

$$\psi_\alpha(x) = \langle x | \alpha \rangle = e^{-|\alpha|^2/2} \langle x | e^{\alpha a^\dagger} | 0 \rangle \quad (7.4.24)$$

from eq. (7.4.9). Next, we write

$$\alpha a^\dagger = A + B \quad (7.4.25)$$

where, from the definition of  $a^\dagger$  in eq. (7.3.2),

$$A = \alpha \sqrt{\frac{m\omega}{2\hbar}} X, \quad B = -i \frac{\alpha}{\sqrt{2\hbar\omega m}} P. \quad (7.4.26)$$

Since  $[A, B] = \alpha^2/2$  is a constant, we can use the Baker–Campbell–Hausdorff formula eq. (2.4.35), repeated here for convenience as  $e^{A+B} = e^A e^B e^{-[A,B]/2}$ , to obtain

$$\psi_\alpha(x) = e^{-|\alpha|^2/2} e^{-\alpha^2/4} \exp\left(\sqrt{\frac{m\omega}{2\hbar}} \alpha x\right) \langle x | \exp\left(-i \frac{\alpha}{\sqrt{2\hbar\omega m}} P\right) | 0 \rangle. \quad (7.4.27)$$

The next step is to notice that the exponential operator inside the matrix element has the same form as the translation operator  $T(a)$  defined in eq. (5.2.1), with  $a$  replaced by the (complex, in general) number  $\alpha\sqrt{\hbar/2\omega m}$ . Therefore, we can invoke eq. (5.2.7) to find

$$\langle x | \exp\left(-i \frac{\alpha}{\sqrt{2\hbar\omega m}} P\right) | 0 \rangle = \psi_0\left(x - \alpha \sqrt{\frac{\hbar}{2\omega m}}\right), \quad (7.4.28)$$

where  $\psi_0(x)$  is the ground state wavefunction, found in eq. (7.2.33). Using this in eq. (7.4.27), and rearranging the exponentials, gives

$$\psi_\alpha(x) = e^{-(|\alpha|^2 + \alpha^2)/2} \left(\frac{m\omega}{\pi\hbar}\right)^{1/4} \exp\left[-\frac{m\omega}{2\hbar} \left(x - \alpha \sqrt{\frac{2\hbar}{\omega m}}\right)^2\right]. \quad (7.4.29)$$

Now, writing  $\alpha$  in terms of its real and imaginary parts, and then expressing them in terms of the position and momentum expectation values using eqs. (7.4.17) and (7.4.21), the wavefunction takes the simple form

$$\psi_\alpha(x) = e^{i\theta} e^{i\langle P \rangle x / \hbar} \psi_0(x - \langle X \rangle), \quad (7.4.30)$$

where  $\theta = -\text{Re}[\alpha]\text{Im}[\alpha]$ . Up to the physically irrelevant global phase provided by  $\theta$ , the coherent state wavefunction is the same as the ground state, but displaced by the position expectation value and multiplied by the position-dependent phase factor that encodes the momentum expectation value.

So far, we have considered a coherent state  $|\alpha\rangle$  at a fixed time  $t$ . Now let us investigate the time evolution of the state and its properties. Since the Hamiltonian is independent of time, the solution of the Schrödinger equation tells us that at time  $t$  the state is

$$e^{-iHt/\hbar}|\alpha\rangle = e^{-iHt/\hbar} e^{-|\alpha|^2/2} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} |n\rangle = e^{-|\alpha|^2/2} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} e^{-i\omega(n+1/2)t} |n\rangle \quad (7.4.31)$$

$$= e^{-i\omega t/2} e^{-|\alpha|^2/2} \sum_{n=0}^{\infty} \frac{(\alpha e^{-i\omega t})^n}{\sqrt{n!}} |n\rangle \quad (7.4.32)$$

$$= e^{-i\omega t/2} |\alpha e^{-i\omega t}\rangle. \quad (7.4.33)$$

This shows that the coherent state just evolves to another coherent state in which  $\alpha$  has changed<sup>†</sup> to  $\alpha e^{-i\omega t}$ . Thus, in the time-dependent coherent state, the magnitude of  $\alpha$  does not change, and the phase of  $\alpha$  rotates linearly with time, at the natural angular frequency of the harmonic oscillator.

Suppose that at time  $t = 0$ , we start with  $\alpha = |\alpha|e^{-i\phi_0}$ , so that at a general time  $t$ ,

$$\alpha = |\alpha|e^{-i(\omega t + \phi_0)} \quad (7.4.34)$$

Then, using the results from eqs. (7.4.17) and (7.4.21), we get

$$\langle X \rangle(t) = \sqrt{\frac{2\hbar}{m\omega}} \text{Re} [|\alpha|e^{-i(\omega t + \phi_0)}] = x_0 \cos(\omega t + \phi_0), \quad (7.4.35)$$

$$\langle P \rangle(t) = \sqrt{2m\hbar\omega} \text{Im} [|\alpha|e^{-i(\omega t + \phi_0)}] = -m\omega x_0 \sin(\omega t + \phi_0), \quad (7.4.36)$$

where we have defined

$$x_0 = \sqrt{\frac{2\hbar}{m\omega}} |\alpha|. \quad (7.4.37)$$

---

<sup>†</sup>The state ket has also acquired an irrelevant global phase  $e^{-i\omega t/2}$ . In contrast, the complex phase  $e^{-i\omega t}$  multiplying  $\alpha$  (inside the ket symbol) is certainly physically relevant, as is clear from eqs. (7.4.17) and (7.4.21).

This establishes the connection between the classical trajectory of the harmonic oscillator and the parameter  $\alpha$  of the coherent state that most closely resembles it. Not only is the coherent state for a harmonic oscillator as close as possible to the classical limit, but we see, by comparing to eq. (7.1.5), that the expectation values of the position and momentum evolve in exactly the same way as their classical counterparts. The uncertainties do not change with time. To summarize the properties of a coherent state,

$$\langle H \rangle = \hbar\omega(|\alpha|^2 + 1/2) \quad \Delta H = \hbar\omega|\alpha|, \quad (7.4.38)$$

$$\langle X \rangle(t) = x_{\text{cl}}(t), \quad \Delta X = \sqrt{\hbar/2m\omega}, \quad (7.4.39)$$

$$\langle P \rangle(t) = p_{\text{cl}}(t), \quad \Delta P = \sqrt{m\hbar\omega/2}. \quad (7.4.40)$$

The harmonic oscillator potential evidently has a remarkable “focusing” property, such that the coherent state wavefunctions do not spread out at all as they evolve in time, unlike the case for the free particle Gaussian states as seen in section 6.2.

There is a simple (and even practical) way to prepare a coherent state for the harmonic oscillator. Suppose we temporarily apply a constant force  $f$ , acting in the positive  $x$  direction, so that the new Hamiltonian is

$$H_f = \frac{P^2}{2m} + \frac{1}{2}m\omega^2 X^2 - fX. \quad (7.4.41)$$

By defining a shifted position operator

$$X' = X - \frac{f}{m\omega^2}, \quad (7.4.42)$$

the Hamiltonian can be rewritten as

$$H_f = \frac{P^2}{2m} + \frac{1}{2}m\omega^2 X'^2 - \frac{f^2}{2m\omega^2}. \quad (7.4.43)$$

The constant shift in the position operator does not affect the commutation relation,

$$[X', P] = i\hbar, \quad (7.4.44)$$

so the solution of the eigenvalue problem for  $H_f$  proceeds exactly as for the Hamiltonian with  $f = 0$ . The energy eigenvalues will be the same, but lowered by the constant amount  $-f^2/2m\omega^2$ . More importantly for our present purposes, the ground state  $|0\rangle_f$  of  $H_f$  is defined by the property that it is annihilated by the operator

$$a' = \sqrt{\frac{m\omega}{2\hbar}}X' + i\frac{1}{\sqrt{2\hbar\omega m}}P = a - \frac{f}{\sqrt{2\hbar\omega^3 m}}, \quad (7.4.45)$$

where  $a$  is the annihilation operator for  $H$  with  $f = 0$ . This implies that the state  $|0\rangle_f$  is exactly a coherent state of the original Hamiltonian  $H$ :

$$a|0\rangle_f = \alpha|0\rangle_f, \quad (7.4.46)$$

where the eigenvalue

$$\alpha = \frac{f}{\sqrt{2\hbar\omega^3m}} \quad (7.4.47)$$

is a real number.

To summarize, a coherent state of the harmonic oscillator can be prepared as follows. First, apply a constant force  $f$ , shifting the equilibrium position of the mass to the point  $x_0 = f/m\omega^2$ . Next, allow the oscillator to settle into the ground state of the new Hamiltonian  $H_f$ , with the same uncertainties in position and momentum as the ground state of  $H$ . Finally, we release the mass by removing the force  $f$ . At the instant that the mass is released, it will find itself in a coherent state of  $H$ , with  $\alpha$  initially real and given by eq. (7.4.47). It will therefore evolve in time as we have already seen, remaining in a coherent state as  $\alpha$  acquires a non-trivial phase  $e^{-i\omega t}$ . The subsequent behavior is as close as possible to what would happen in classical mechanics if we displaced the oscillator and then released it from rest.

## 7.5 Three-dimensional harmonic oscillator

We now return to the case of a three-dimensional harmonic oscillator, with Hamiltonian given by eq. (7.1.17). Rewriting it without the tildes (which just denoted a particular choice of rectangular coordinates), this can be rewritten as

$$H = H_x + H_y + H_z, \quad (7.5.1)$$

where

$$H_x = \frac{P_x^2}{2m} + \frac{1}{2}m\omega_x^2 X^2 = \hbar\omega_x(a_x^\dagger a_x + 1/2), \quad (7.5.2)$$

with exactly analogous expressions for  $H_y$  and  $H_z$ . Here, we have made use of the experience of section 7.3 to write

$$a_x = \sqrt{\frac{m\omega_x}{2\hbar}}X + i\frac{1}{\sqrt{2\hbar\omega_x m}}P_x, \quad (7.5.3)$$

$$a_y = \sqrt{\frac{m\omega_y}{2\hbar}}Y + i\frac{1}{\sqrt{2\hbar\omega_y m}}P_y, \quad (7.5.4)$$

etc. Now  $H_x$  and  $H_y$  and  $H_z$  all commute with each other, and are Hermitian. One can therefore find an orthobasis consisting of their common eigenstates, denoted  $|n_x, n_y, n_z\rangle$ , where operators



with the  $x$  subscript just ignore the  $n_y$  and  $n_z$  labels, and similarly for operators with the  $y$  and  $z$  subscripts. Thus,

$$H_x |n_x, n_y, n_z\rangle = E_x |n_x, n_y, n_z\rangle, \quad H_y |n_x, n_y, n_z\rangle = E_y |n_x, n_y, n_z\rangle, \quad (7.5.5)$$

$$H_z |n_x, n_y, n_z\rangle = E_z |n_x, n_y, n_z\rangle. \quad (7.5.6)$$

(This can also be viewed as the tensor product of three one-dimensional harmonic oscillators.) From section 7.3, we have already solved the eigenvalue problem for each  $H_j$ , and so we know that the allowed energies for each of  $j = x, y, z$  are

$$E_j = \hbar\omega_j(n_j + 1/2), \quad n_j = 0, 1, 2, \dots, \quad (7.5.7)$$

and that the corresponding eigenkets satisfy, for example:

$$a_x |n_x, n_y, n_z\rangle = \sqrt{n_x} |n_x - 1, n_y, n_z\rangle, \quad (7.5.8)$$

$$a_x^\dagger |n_x, n_y, n_z\rangle = \sqrt{n_x + 1} |n_x + 1, n_y, n_z\rangle, \quad (7.5.9)$$

$$a_y |n_x, n_y, n_z\rangle = \sqrt{n_y} |n_x, n_y - 1, n_z\rangle, \quad (7.5.10)$$

etc. Since the labels  $n_x$ ,  $n_y$ , and  $n_z$  uniquely specify the states of an orthobasis,  $H_x$ ,  $H_y$ , and  $H_z$  form a CSCO for this problem.

The state  $|n_x, n_y, n_z\rangle$  has total energy eigenvalue

$$E = E_x + E_y + E_z = \hbar\omega_x(n_x + 1/2) + \hbar\omega_y(n_y + 1/2) + \hbar\omega_z(n_z + 1/2), \quad (7.5.11)$$

and its wavefunction is just a product of the wavefunctions for the one-dimensional harmonic oscillator,

$$\psi_{n_x, n_y, n_z}(\vec{r}) = \langle \vec{r} | n_x, n_y, n_z \rangle = \psi_{n_x, \omega_x}(x) \psi_{n_y, \omega_y}(y) \psi_{n_z, \omega_z}(z), \quad (7.5.12)$$

where each of the  $\psi_{n_j, \omega_j}$  functions can be obtained from eq. (7.2.31) in the obvious way.

In the special case of the isotropic 3-dimensional harmonic oscillator,  $\omega_x = \omega_y = \omega_z = \omega$ , the system is invariant under arbitrary rotations about the origin. This symmetry is reflected in a degeneracy of the energy spectrum, which is now

$$E_n = \hbar\omega(n + 3/2), \quad (7.5.13)$$

for states  $|n, u_n\rangle$ , where  $n = n_x + n_y + n_z$  and  $u_n = 1, \dots, g_n$  is a degeneracy label for the energy eigenvalue  $E_n$ . To find the degeneracy  $g_n$  of each energy level, we need to know how many ways there are to choose three non-negative integers that add up to  $n$ . The ground state with  $n = 0$  is unique, as it can only be achieved with the lone combination  $n_x = n_y = n_z = 0$ , so  $g_0 = 1$ .

The first excited energy level can be achieved in three different ways, by taking one of  $n_x, n_y, n_z$  to be 1, and the others 0, so  $g_1 = 3$ . Similar straightforward counting reveals that  $g_2 = 6$ , and  $g_3 = 10$ . More generally, this combinatorics problem can be solved using the dots and lines trick: one arranges  $n$  dots in a row, and splits them into three groups by placing two vertical lines as shown below.

$$\underbrace{\bullet \cdots \bullet}_{n_x} | \underbrace{\bullet \cdots \bullet}_{n_y} | \underbrace{\bullet \cdots \bullet}_{n_z} \quad (7.5.14)$$

The degeneracy of the energy eigenvalue  $E_n$  is therefore the number of ways of arranging a row of  $n + 2$  objects, of which  $n$  are identical and 2 are identical,

$$g_n = \frac{(n+2)!}{n! 2!} = \frac{1}{2}(n+1)(n+2). \quad (7.5.15)$$

There is another natural choice of CSCO that one can use for the isotropic three-dimensional harmonic oscillator, consisting of the total Hamiltonian  $H$  and two observables associated with the angular momentum. We will discuss this approach in section 10.5.

## 7.6 Exercises

**Exercise 7.1.** For a particle of mass  $m$  in a 1-d harmonic oscillator potential with angular frequency  $\omega$ , with minimum at the origin, use ladder operators to calculate the following matrix elements for energy eigenstates  $|n\rangle$  and  $|k\rangle$  where  $n$  and  $k$  are non-negative integers. (Your answers should make use of the Kronecker delta symbol.)

$$\langle k|a|n\rangle, \quad \langle k|a^\dagger|n\rangle, \quad \langle k|X|n\rangle, \quad \langle k|P|n\rangle, \quad \langle k|X^2|n\rangle, \quad \langle k|P^2|n\rangle, \quad \langle k|H|n\rangle.$$

Make note of how the parity selection rules apply to each of your results.

**Exercise 7.2.** Consider a particle of mass  $m$  in a 1-d harmonic oscillator potential with angular frequency  $\omega$ , with minimum at the origin.

- (a) Use ladder operators to calculate  $\langle n|X^k|n\rangle$  and  $\langle n|P^k|n\rangle$ , for  $k = 0, 1, 2, 3, 4, 5$ , with  $n$  arbitrary. Make note of how the parity selection rule applies to your results.
- (b) Find the expectation values of the kinetic and potential energies for the state  $|n\rangle$ , and show that they are equal. [This illustrates the Virial Theorem; see eq. (3.4.21).]
- (c) What is  $(\Delta X)(\Delta P)$  for the state  $|n\rangle$ ?

**Exercise 7.3.** Use the raising and lowering identities (7.3.17) and (7.3.18) and the representation of  $a$  and  $a^\dagger$  in eq. (7.3.34) to derive the Hermite polynomial identities (7.2.24) and (7.2.25).

**Exercise 7.4.** Consider a particle of mass  $m$  in a 1-d harmonic oscillator potential with angular frequency  $\omega$ , with minimum at the origin. For the observable  $A = PX + XP$ , find:

- (a) The expression in simplest form for  $A$  in terms of the usual ladder operators.
- (b) The expectation value for  $A$  for the  $n$ th excited energy eigenstate  $|n\rangle$ .
- (c) The uncertainty for  $A$  in the state  $|n\rangle$ .

**Exercise 7.5.** At time  $t = 0$ , a harmonic oscillator is in the state  $|\psi(0)\rangle = \frac{1}{\sqrt{2}}|0\rangle + \frac{1}{\sqrt{2}}|1\rangle$  in the usual energy basis.

- (a) Find the state at later times,  $|\psi(t)\rangle$ , in the energy basis.
- (b) By direct calculation using the results of part (a) (without appealing to Ehrenfest's Theorem), find the expectation values  $\langle X \rangle$  and  $\langle P \rangle$  as functions of time  $t$ .
- (c) Check that your results from part (b) obey Ehrenfest's Theorem.
- (d) At time  $t$ , compute the probability that a measurement of the position yields  $x > 0$ . Evaluate the minimum and the maximum probabilities numerically.

**Exercise 7.6.** At time  $t = 0$ , a harmonic oscillator has wavefunction  $\psi(x, 0) = cx^2 \exp(-m\omega x^2/2\hbar)$ , where  $c$  is a positive real constant.

- (a) By requiring the wavefunction to be normalized, find the constant  $c$ .
- (b) If the energy is measured, what are the possible outcomes and their probabilities?
- (c) Find the state ket  $|\psi(t)\rangle$  in the energy basis, as a function of time.
- (d) Compute the expectation values  $\langle X \rangle$ ,  $\langle P \rangle$ ,  $\langle X^2 \rangle$ , and  $\langle P^2 \rangle$ , and the uncertainties  $\Delta X$  and  $\Delta P$ , each as a function of time.
- (e) At time  $t$ , what is the probability that a measurement of the position yields  $x > 0$ ?

**Exercise 7.7.** Consider a particle of mass  $m$  moving in 1 dimension in the potential

$$V(x) = \begin{cases} \frac{1}{2}m\omega^2 x^2 & (\text{for } x > 0), \\ \infty & (\text{for } x < 0). \end{cases} \quad (7.6.1)$$

Find all of the energy eigenvalues and the corresponding unit-normalized wavefunctions. [Hint: try to make use of calculations already done, rather than new ones.]

**Exercise 7.8.** Consider the coherent states of the harmonic oscillator in section 7.4.

- (a) For two different coherent states  $|\alpha\rangle$  and  $|\beta\rangle$ , compute  $\langle \beta | \alpha \rangle$  and  $|\langle \beta | \alpha \rangle|^2$  in simplest form. (Your answers should be exponentials. Therefore, they cannot vanish, which shows that the coherent states do *not* satisfy orthogonality.)
- (b) Suppose that the oscillator is in the state  $|\alpha\rangle$  at time  $t = 0$ . Find, in simplest form, the probability  $P(t)$  to find it again in the state  $|\alpha\rangle$  at a later time  $t$ . At what times is  $P(t) = 1$ ?
- (c) Suppose that  $|\alpha| \gg 1$ , as for a macroscopic oscillator. For what small length of time after

$t = 0$  does the probability  $P(t)$  remain greater than 0.5? [Hint: expand  $\cos(\omega t)$  to order  $t^2$ .]

(d) What is the minimum value of  $P(t)$ , and at what time or times is it achieved?

(e) Estimate your answers to parts (c) and (d) numerically for a macroscopic oscillator with  $m = 0.2$  kg,  $\omega = 10 \text{ sec}^{-1}$ , and  $x_0 = 0.1$  meters.

**Exercise 7.9.** We found in section 7.4 that the eigenstates of the lowering operator  $a$  are of great interest, so it is natural to wonder about the possible eigenstates of the raising operator. Show that  $a^\dagger$  has no normalizable eigenstates.

**Exercise 7.10.** Consider a spin-less particle of mass  $m$ , moving in three dimensions, with

$$H = \frac{1}{2m}(P_x^2 + P_y^2 + P_z^2) + \frac{1}{2}m\omega^2 \left( \frac{5}{2}X^2 + \frac{5}{2}Y^2 + 3XY + Z^2 \right). \quad (7.6.2)$$

(a) Use the change of coordinates:  $x = cx' + sy'$ ,  $y = -sx' + cy'$ ,  $z = z'$ , where  $c$  and  $s$  are the cosine and sine of an arbitrary rotation angle, to rewrite the Hamiltonian in terms of the operators  $X'$ ,  $Y'$ ,  $Z'$ ,  $P_{x'}$ ,  $P_{y'}$ , and  $P_{z'}$ . (You may use the fact that this is an orthogonal rotation on the coordinate system, so  $P_x^2 + P_y^2 + P_z^2 = P_{x'}^2 + P_{y'}^2 + P_{z'}^2$ .)

(b) Choose  $c$  and  $s$  (remembering that  $c^2 + s^2 = 1$ ), so that  $H$  will not contain a term proportional to  $X'Y'$ . Rewrite the Hamiltonian with this choice. (There is more than one valid choice here.)

(c) What are the three smallest energy eigenvalues of  $H$ , and what are their degeneracies?

**Exercise 7.11.** Two particles labeled 1 and 2 (they are distinguishable, but happen to have the same mass) are governed by a coupled harmonic oscillator Hamiltonian in one dimension,

$$H = \frac{1}{2m}(P_1^2 + P_2^2) + \frac{1}{2}m\omega^2 X_1^2 + \frac{1}{2}m\omega^2 X_2^2 + \frac{1}{2}m\Omega^2(X_1 - X_2)^2, \quad (7.6.3)$$

where the constant  $\omega$  parametrizes the restoring force for the particles to the origin, and  $\Omega$  describes the attractive force between the particles. The operators  $X_1, P_1$  and  $X_2, P_2$  satisfy the canonical commutation relations  $[X_1, P_1] = i\hbar$  and  $[X_2, P_2] = i\hbar$  and  $[X_1, P_2] = [X_2, P_1] = 0$  and  $[X_j, X_k] = 0$  and  $[P_j, P_k] = 0$  for all  $j, k = 1, 2$ . Consider the new operators

$$U = \frac{1}{\sqrt{2}}(X_1 + X_2), \quad V = \frac{1}{\sqrt{2}}(X_1 - X_2), \quad (7.6.4)$$

$$P_u = \frac{1}{\sqrt{2}}(P_1 + P_2), \quad P_v = \frac{1}{\sqrt{2}}(P_1 - P_2). \quad (7.6.5)$$

(a) Derive all of the commutation relations of all pairs of operators from the set  $U, V, P_u, P_v$ .

(b) Write the Hamiltonian in terms of the operators  $U, V, P_u, P_v$ .

(c) Define appropriate creation and destruction operators for the  $U, V, P_u, P_v$  system, so that the Hamiltonian has a simple form in terms of them. From this, infer the eigenvalues of the

Hamiltonian, and write down a suitable notation for the energy eigenbasis kets.

(d) Obtain the wavefunction of the ground state in the  $u, v$  representation  $\psi(u, v)$ , and use it to obtain the normalized ground state wavefunction  $\psi(x_1, x_2) = \langle x_1, x_2 | \psi \rangle$ .

**Exercise 7.12.** Consider the isotropic harmonic oscillator for a single particle in 2-d,

$$H = \frac{1}{2m}(P_x^2 + P_y^2) + \frac{1}{2}m\omega^2(X^2 + Y^2). \quad (7.6.6)$$

This can be viewed as the sum of independent oscillators for the  $x$  and  $y$  directions.

(a) Express  $H$  and the angular momentum operator  $L_z = XP_y - YP_x$  in simplest form in terms of the raising and lowering operators  $a_x^\dagger, a_x, a_y^\dagger, a_y$  for the independent  $x$  and  $y$  oscillations. Compute the commutator of  $L_z$  and  $H$ . Are they compatible?

(b) Consider the basis kets

$$|n_x, n_y\rangle = \frac{(a_x^\dagger)^{n_x}}{\sqrt{n_x!}} \frac{(a_y^\dagger)^{n_y}}{\sqrt{n_y!}} |0, 0\rangle, \quad (7.6.7)$$

where  $|0, 0\rangle$  is the ground state. Show that the  $|n_x, n_y\rangle$  are eigenstates of energy and parity, and give the corresponding eigenvalues. What is the degeneracy of the  $n$ th excited energy level?

(c) Show that the basis kets  $|n_x, n_y\rangle$  are not eigenstates of  $L_z$ .

(d) Define new raising and lowering operators

$$a_+^\dagger = (a_x^\dagger + ia_y^\dagger)/\sqrt{2}, \quad a_+ = (a_x - ia_y)/\sqrt{2}, \quad (7.6.8)$$

$$a_-^\dagger = (a_x^\dagger - ia_y^\dagger)/\sqrt{2}, \quad a_- = (a_x + ia_y)/\sqrt{2}. \quad (7.6.9)$$

Compute all of the non-zero commutators involving these four operators. Express  $H$  and  $L_z$  in terms of them, in normal-ordered form.

(e) Consider the basis kets

$$|n_+, n_-\rangle' = \frac{(a_+^\dagger)^{n_+}}{\sqrt{n_+!}} \frac{(a_-^\dagger)^{n_-}}{\sqrt{n_-!}} |0, 0\rangle. \quad (7.6.10)$$

Show that these are eigenstates of  $H$  and  $L_z$ , and give the corresponding eigenvalues. For the lowest three energy levels, write the eigenvalues of  $H$  and  $L_z$ , and express the corresponding eigenstates as linear combinations of the kets  $|n_x, n_y\rangle$ . Do  $H$  and  $L_z$  form a CSCO?

## 8 Angular momentum and its representations

### 8.1 The eigenvalue problem for angular momentum

At the end of section 5.3, we introduced the commutation relation algebra for the components of a generic angular momentum operator  $\vec{J} = \hat{x}J_x + \hat{y}J_y + \hat{z}J_z$ , defined to satisfy

$$[J_a, J_b] = i\hbar\epsilon_{abc}J_c, \quad (a, b, c = x, y, z), \quad (8.1.1)$$

with  $c$  implicitly summed over. Let us now find the simultaneous eigenvectors and the corresponding eigenvalues of the compatible observables  $J^2 = J_x^2 + J_y^2 + J_z^2$  and  $J_z$ .

We start by observing that the eigenvalues of  $J^2$  must be non-negative. To prove it, note that for any non-null ket  $|\psi\rangle$ ,

$$\langle\psi|J^2|\psi\rangle = \|J_x|\psi\rangle\|^2 + \|J_y|\psi\rangle\|^2 + \|J_z|\psi\rangle\|^2 \geq 0, \quad (8.1.2)$$

where  $\| |v\rangle \|$  denotes the norm of  $|v\rangle$ , and so the possibility of equality exists only because each of the kets  $J_x|\psi\rangle$ ,  $J_y|\psi\rangle$ , and  $J_z|\psi\rangle$  could be null. Now, if

$$J^2|\psi\rangle = \lambda|\psi\rangle, \quad (8.1.3)$$

then it follows that  $\langle\psi|J^2|\psi\rangle = \lambda\langle\psi|\psi\rangle \geq 0$ , so  $\lambda \geq 0$ . For reasons to become clear shortly, it turns out to be convenient to give  $\lambda$  the name  $\hbar^2 j(j+1)$ , by defining  $j = \sqrt{\lambda/\hbar^2 + 1/4} - 1/2$ . Since  $\lambda \geq 0$ , it follows that  $j \geq 0$  also.

Because  $J^2$  and  $J_z$  are compatible observables, Theorem 2.7.1 says that there must be an orthobasis consisting of common eigenstates,  $|j, m, u_{jm}\rangle$ , which satisfy orthonormality and completeness relations

$$\langle j', m', u'_{j'm'} | j, m, u_{jm} \rangle = \delta_{jj'} \delta_{mm'} \delta_{u'_{j'm'}, u_{jm}}, \quad (8.1.4)$$

$$\sum_j \sum_m \sum_{u_{jm}} |j, m, u_{jm}\rangle \langle j, m, u_{jm}| = I. \quad (8.1.5)$$

Here  $u_{jm}$  is a possible degeneracy label<sup>†</sup> that will be important in particular cases, but plays no role in the following discussion and so will be suppressed for simplicity. We therefore seek to solve the eigenvalue problem

$$J^2 |j, m\rangle = \hbar^2 j(j+1) |j, m\rangle, \quad (8.1.6)$$

$$J_z |j, m\rangle = \hbar m |j, m\rangle. \quad (8.1.7)$$

---

<sup>†</sup>Soon [just before eq. (8.1.28)], we will learn that the orthobasis can actually always be chosen in such a way that the  $u_{jm}$  do not depend on  $m$ . But we do not know that yet.

for the allowed eigenvalues  $j$  and  $m$ , recognizing that there may be more than one sector of such states, distinguished by the suppressed degeneracy label. In the special case of orbital angular momentum, we already found in section 5.3, by requiring the position wavefunction to be single-valued, that the allowed eigenvalues of  $L_z$  are  $\hbar m$ , where  $m$  is an integer. However, spin (intrinsic) angular momentum is not associated with a position wavefunction, so that argument does not apply, and  $m$  need not be an integer, as we will see.

It is very useful (and not just for the present purpose of solving the eigenvalue problem) to define the **angular momentum raising and lowering operators**, also sometimes known as **ladder operators** for angular momentum,

$$J_+ = J_x + iJ_y, \quad J_- = J_x - iJ_y, \quad (8.1.8)$$

which will play a role similar to  $a^\dagger$  and  $a$  for the harmonic oscillator. They are not Hermitian, but instead are Hermitian conjugates of each other,

$$(J_+)^\dagger = J_-. \quad (8.1.9)$$

The inverse relations of eq. (8.1.8) are

$$J_x = \frac{1}{2}(J_- + J_+), \quad J_y = \frac{i}{2}(J_- - J_+). \quad (8.1.10)$$

Some other useful identities are

$$[J_z, J_\pm] = \pm \hbar J_\pm, \quad (8.1.11)$$

$$[J_+, J_-] = 2\hbar J_z, \quad (8.1.12)$$

$$J_+ J_- = J^2 - J_z^2 + \hbar J_z, \quad (8.1.13)$$

$$J_- J_+ = J^2 - J_z^2 - \hbar J_z. \quad (8.1.14)$$

From the last two equations, we get

$$J^2 = \frac{1}{2}J_+ J_- + \frac{1}{2}J_- J_+ + J_z^2. \quad (8.1.15)$$

Each of  $J_+$ ,  $J_-$ , and  $J_z$  commute with the total angular momentum squared,

$$[J^2, J_+] = [J^2, J_-] = [J^2, J_z] = 0. \quad (8.1.16)$$

We now derive some useful facts by studying the ket  $J_+|j, m\rangle$ . Since  $J^2$  commutes with  $J_+$ ,

$$J^2 (J_+|j, m\rangle) = \hbar^2 j(j+1) (J_+|j, m\rangle). \quad (8.1.17)$$

Also,

$$J_z (J_+ |j, m\rangle) = [J_z, J_+] |j, m\rangle + J_+ J_z |j, m\rangle = \hbar J_+ |j, m\rangle + \hbar m J_+ |j, m\rangle \quad (8.1.18)$$

$$= \hbar(m+1) (J_+ |j, m\rangle). \quad (8.1.19)$$

Thus,  $J_+ |j, m\rangle$  is an eigenstate of both  $J^2$  and  $J_z$ , with eigenvalues  $\hbar^2 j(j+1)$  and  $\hbar(m+1)$ , respectively. Due to the fact that the  $|j, m, u_{j,m}\rangle$  were chosen as an orthobasis, we can conclude that either  $J_+ |j, m\rangle$  is proportional to  $|j, m+1\rangle$  within each sector labeled by  $u_{j,m}$ , or else it is the null ket. This is why  $J_+$  is called a raising operator; it raises the  $J_z$  eigenvalue. To find out whether  $J_+ |j, m\rangle$  is the null ket, we compute its squared norm,

$$\begin{aligned} \|J_+ |j, m\rangle\|^2 &= \langle j, m | J_- J_+ |j, m\rangle = \langle j, m | (J^2 - J_z^2 - \hbar J_z) |j, m\rangle \\ &= \langle j, m | [\hbar^2 j(j+1) - (\hbar m)^2 - \hbar(\hbar m)] |j, m\rangle \\ &= \hbar^2 [j(j+1) - m(m+1)]. \end{aligned} \quad (8.1.20)$$

From this we learn two useful things. First, the general properties of the inner product require that the squared norm is non-negative, so allowed  $j, m$  must satisfy  $j(j+1) - m(m+1) = (j-m)(j+m+1) \geq 0$ , and since we learned at the beginning of this section that  $j \geq 0$ , we can conclude that for every non-null ket  $|j, m\rangle$ ,

$$m \leq j. \quad (8.1.21)$$

The second useful result from eq. (8.1.20) is the constant of proportionality between  $J_+ |j, m\rangle$  and the unit-normalized ket  $|j, m+1\rangle$ ,

$$J_+ |j, m\rangle = \hbar \sqrt{j(j+1) - m(m+1)} |j, m+1\rangle. \quad (8.1.22)$$

Here we had to make an arbitrary and unavoidable choice of phase; eq. (8.1.22) can be taken as the definition of the relative phase between  $|j, m\rangle$  and  $|j, m+1\rangle$ . It follows from eq. (8.1.22) that  $J_+ |j, j\rangle$  is actually the null ket.

Everything in the previous paragraph can be repeated for  $J_- |j, m\rangle$ . We learn that, for every non-null ket  $|j, m\rangle$ ,

$$m \geq -j \quad (8.1.23)$$

due to the requirement of positive squared norm, and that  $J_- |j, -j\rangle$  is the null ket, and that

$$J_- |j, m\rangle = \hbar \sqrt{j(j+1) - m(m-1)} |j, m-1\rangle. \quad (8.1.24)$$

This justifies calling  $J_-$  the lowering operator for (the  $z$  component of) angular momentum.



In eq. (8.1.24), we have made another choice of phase, and we should be careful to make sure that it is consistent with the phase choice made in eq. (8.1.22). This can be checked by using eqs. (8.1.22) and (8.1.24) to compute  $J_+J_-|j, m\rangle$  and  $J_-J_+|j, m\rangle$ , and noting that they are equivalent to the results obtained using eqs. (8.1.13) and (8.1.14), respectively. A nice feature of eqs. (8.1.22) and (8.1.24) is that the coefficients on the right-hand sides are real and non-negative.

Combining eqs. (8.1.21) and (8.1.23) tells us that, for any given  $j$ , the only possible values of  $m$  for physical states  $|j, m\rangle$  are in the range

$$-j \leq m \leq j. \quad (8.1.25)$$

Intuitively, this is the quantum version of the classical statement that the magnitude of the  $z$ -component is less than the magnitude of the angular momentum vector.

We are now ready to prove that  $j = n/2$  for some integer  $n$ . If we compute  $(J_+)^p|j, m\rangle$ , it must be proportional to  $|j, m+p\rangle$ , according to eq. (8.1.22). For some finite  $p$ , this must be the ket  $|j, j\rangle$ , because otherwise we could keep acting with  $J_+$  to find non-null kets with arbitrarily large  $J_z$  eigenvalue, which would contradict eq. (8.1.25). (Here, we are relying on the fact that eq. (8.1.22) tells us that  $|j, j\rangle$  is the unique non-null ket that is annihilated by  $J_+$ .) Therefore,  $j = m+p$  for some non-negative integer  $p$ . Similarly, if we compute  $(J_-)^q|j, m\rangle$ , it must be proportional to  $|j, m-q\rangle$ , according to eq. (8.1.24). For some finite  $q$ , this must be the ket  $|j, -j\rangle$ ; otherwise, we would again contradict eq. (8.1.25). Therefore,  $-j = m-q$  for some non-negative integer  $q$ . Putting the results together gives  $2j = p+q$ , a non-negative integer. The allowed values are

$$j = 0, 1/2, 1, 3/2, 2, \dots \quad (8.1.26)$$

The fact that  $j = m+p$  for some integer  $p$ , together with eq. (8.1.25), also tells us which values of  $m$  can give non-null kets  $|j\rangle$ . For each  $j$ , they are the  $2j+1$  values

$$m = -j, -j+1, \dots, j-1, j. \quad (8.1.27)$$

For the special case that  $\vec{J} = \vec{L}$ , we already found in section 5.3 that  $m$  must be an integer, so in the case of orbital angular momentum the allowed values of  $j = l$  are also restricted to the non-negative integers  $0, 1, 2, \dots$ . The case of half-integer<sup>‡</sup>  $j$  must correspond to something other than orbital angular momenta.

For a given state  $|j, m, u_{jm}\rangle$ , the operations of  $J^2$ ,  $J_z$ ,  $J_+$ , and  $J_-$ , given by eqs. (8.1.6), (8.1.7), (8.1.22), and (8.1.24) are independent of  $u_{jm}$ , and can change  $m$ , but not  $j$ . This has two important implications.

---

<sup>‡</sup>The standard term “half-integer” means an odd integer divided by 2. It might be more logical to call this “half-odd-integer” or “integer-plus-half”, but it is difficult to fight tradition.

First, if we have a single representative state with a certain  $j$  and  $u_{jm}$ , then all other states with the same  $j$  and  $u_{jm}$  but other values of  $m$  are obtained by acting repeatedly with  $J_+$  or  $J_-$ . Therefore, the degeneracy label  $u_{jm}$  does not actually depend on  $m$ , as foreshadowed in the footnote following eq. (8.1.5). Thus, for any quantum system, the orthobasis of common eigenstates of  $J^2$  and  $J_z$  can be labeled as  $|j, m, u_j\rangle$ , with a slight simplification of eqs. (8.1.4) and (8.1.5),

$$\langle j', m', u'_j | j, m, u_j \rangle = \delta_{jj'} \delta_{mm'} \delta_{u'_j u_j}, \quad (8.1.28)$$

$$\sum_j \sum_m \sum_{u_j} |j, m, u_j\rangle \langle j, m, u_j| = I. \quad (8.1.29)$$

Typically,  $u_j$  represents the eigenvalue(s) of some observable(s) that, together with  $J^2$  and  $J_z$ , form a CSCO for the quantum system. If the Hamiltonian is invariant under rotations, then  $H$  can be taken to be one of those observables. The group of  $2j+1$  orthobasis states with common  $j$  and  $u_j$  and varying  $m = -j, \dots, j$  is called an **angular momentum multiplet**.

The second observation is that, because the operations of the angular momentum operators  $J^2$ ,  $J_z$ ,  $J_+$ , and  $J_-$  (and  $J_x$ ,  $J_y$ ) in the orthobasis  $|j, m, u_j\rangle$  do not depend on  $u_j$  at all, we can work out their matrix representations and operations on kets for each relevant value of  $j$  just once, and the results will be applicable to any quantum system with that  $j$ .

For a  $j = 0$  subspace, we have  $J^2 = 0$  and  $J_z = J_+ = J_- = J_x = J_y = 0$ , so the state space has only one non-null ket  $|j = 0, m = 0\rangle$ . All angular momentum operators acting on  $|j = 0, m = 0\rangle$  give the null ket, so that they are all represented by the  $1 \times 1$  matrix 0.

## 8.2 The $j = 1/2$ representation: Pauli matrices and spin

For a  $j = 1/2$  subspace, the orthobasis consists of two states with  $m = \pm 1/2$ , with corresponding two-component vector representations

$$|j = 1/2, m = 1/2\rangle \leftrightarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad |j = 1/2, m = -1/2\rangle \leftrightarrow \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (8.2.1)$$

The matrix representations of the angular momentum operators can now be constructed from eqs. (8.1.6), (8.1.7), (8.1.10), (8.1.22), and (8.1.24), with the results

$$J^2 \leftrightarrow \frac{3\hbar^2}{4} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad J_z \leftrightarrow \frac{\hbar}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (8.2.2)$$

$$J_x \leftrightarrow \frac{\hbar}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad J_y \leftrightarrow \frac{\hbar}{2} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}. \quad (8.2.3)$$

$$J_+ \leftrightarrow \hbar \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad J_- \leftrightarrow \hbar \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad (8.2.4)$$

Some special operator identities that hold only in the  $j = 1/2$  case are

$$J_x^2 = J_y^2 = J_z^2 = \hbar^2/4, \quad J_+^2 = J_-^2 = 0, \quad (j = 1/2). \quad (8.2.5)$$

As noted after eq. (8.1.27), the fact that  $m = \pm 1/2$  is not an integer implies that  $\vec{J}$  cannot be a pure orbital angular momentum.

The case of purely intrinsic angular momentum for a single particle is called **spin**, and we write  $\vec{J} = \vec{S}$  and  $j = s$  in that case. For  $s = 1/2$ , we say that the particle is (or has) spin-1/2, and we often rewrite eq. (8.2.1) using a simplified notation for the  $S_z$  eigenstates,

$$|\uparrow\rangle \leftrightarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad |\downarrow\rangle \leftrightarrow \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad (8.2.6)$$

referred to as spin-up and spin-down, with eigenvalues  $m_s = 1/2$  and  $-1/2$ , respectively.

From eqs. (8.2.2) and (8.2.3), one can deduce that the components of the spin operator written in ket-bra form are

$$S_x = \frac{\hbar}{2} (|\downarrow\rangle \langle\uparrow| + |\uparrow\rangle \langle\downarrow|), \quad (8.2.7)$$

$$S_y = \frac{i\hbar}{2} (|\downarrow\rangle \langle\uparrow| - |\uparrow\rangle \langle\downarrow|), \quad (8.2.8)$$

$$S_z = \frac{\hbar}{2} (|\uparrow\rangle \langle\uparrow| - |\downarrow\rangle \langle\downarrow|). \quad (8.2.9)$$

The standard notation for their matrix representation is

$$\vec{S} \leftrightarrow \frac{\hbar}{2} \vec{\sigma}, \quad (8.2.10)$$

where the components of the vector  $\vec{\sigma}$  are known as the **Pauli matrices**, defined by

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (8.2.11)$$

The Pauli matrices obey the commutation and anticommutation relations

$$[\sigma_a, \sigma_b] = 2i\epsilon_{abc}\sigma_c, \quad (8.2.12)$$

$$\{\sigma_a, \sigma_b\} = 2\delta_{ab} \quad (8.2.13)$$

for  $a, b = x, y, z$ , and

$$\text{Tr}[\sigma_a] = 0, \quad (8.2.14)$$

$$\text{Det}[\sigma_a] = -1. \quad (8.2.15)$$

For any spatial vector  $\vec{v} = \hat{x}v_x + \hat{y}v_y + \hat{z}v_z$ , we have

$$\vec{v} \cdot \vec{\sigma} = \begin{pmatrix} v_z & v_x - iv_y \\ v_x + iv_y & -v_z \end{pmatrix}, \quad (8.2.16)$$

and a formula useful for simplifications,

$$(\vec{v} \cdot \vec{\sigma})(\vec{w} \cdot \vec{\sigma}) = \vec{v} \cdot \vec{w} + i(\vec{v} \times \vec{w}) \cdot \vec{\sigma}, \quad (8.2.17)$$

with the  $2 \times 2$  identity matrix understood in the first term on the right side. In particular,

$$(\vec{v} \cdot \vec{\sigma})(\vec{v} \cdot \vec{\sigma}) = \vec{v} \cdot \vec{v} = v_x^2 + v_y^2 + v_z^2 = v^2 \quad (8.2.18)$$

is proportional to the identity matrix.

For a spin-1/2 particle like an electron, the complete state can be written as a linear combination of kets with  $S_z$  eigenvalues  $m_s = \hbar/2$  and  $-\hbar/2$ , as

$$|\psi\rangle = |\psi_\uparrow, \uparrow\rangle + |\psi_\downarrow, \downarrow\rangle, \quad (8.2.19)$$

where  $\psi_\uparrow$  and  $\psi_\downarrow$  represent the non-spin degrees of freedom corresponding to the classical motion of the particle in three dimensions. This state can also be represented as a two-component column vector of kets, called a **spinor**,

$$|\psi\rangle = \begin{pmatrix} |\psi_\uparrow\rangle \\ |\psi_\downarrow\rangle \end{pmatrix}. \quad (8.2.20)$$

Acting on these two-component spinor kets, the spin operator is again represented by  $\vec{S} \leftrightarrow \frac{\hbar}{2}\vec{\sigma}$ . The bra corresponding to eq. (8.2.20) is a two-component row spinor,

$$\langle\psi| = (\langle\psi_\uparrow| \quad \langle\psi_\downarrow|). \quad (8.2.21)$$

The inner product of two states  $|\psi\rangle$  and  $|\chi\rangle$  is

$$\langle\chi|\psi\rangle = (\langle\chi_\uparrow| \quad \langle\chi_\downarrow|) \begin{pmatrix} |\psi_\uparrow\rangle \\ |\psi_\downarrow\rangle \end{pmatrix} = \langle\chi_\uparrow|\psi_\uparrow\rangle + \langle\chi_\downarrow|\psi_\downarrow\rangle, \quad (8.2.22)$$

and unit normalization means

$$1 = \langle\psi|\psi\rangle = \langle\psi_\uparrow|\psi_\uparrow\rangle + \langle\psi_\downarrow|\psi_\downarrow\rangle. \quad (8.2.23)$$

For example, the two-component spinor position wavefunction for an electron is

$$\langle\vec{r}|\psi\rangle = \begin{pmatrix} \langle\vec{r}|\psi_\uparrow\rangle \\ \langle\vec{r}|\psi_\downarrow\rangle \end{pmatrix} = \begin{pmatrix} \psi_\uparrow(\vec{r}) \\ \psi_\downarrow(\vec{r}) \end{pmatrix}. \quad (8.2.24)$$

These correspond to orthobasis eigenstates of the CSCO consisting of the observables  $(\vec{R}, S_z)$ . If we impose the usual unit normalization condition for the state, the two spin-component wavefunctions are required to satisfy

$$1 = \int d^3\vec{r} |\psi_\uparrow(\vec{r})|^2 + |\psi_\downarrow(\vec{r})|^2, \quad (8.2.25)$$

but they are otherwise independent, in general.

### 8.3 Matrix representation for $j = 1$

For a  $j = 1$  subspace, the orthobasis of  $J_z$  eigenstates consists of three states  $|j, m\rangle$  with  $j = 1$  and  $m = 1, 0, -1$ , with corresponding vector representations

$$|1, 1\rangle \leftrightarrow \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad |1, 0\rangle \leftrightarrow \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad |1, -1\rangle \leftrightarrow \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}. \quad (8.3.1)$$

The matrix representations of the angular momentum operators can now be constructed from eqs. (8.1.6), (8.1.7), (8.1.10), (8.1.22), and (8.1.24), with the results

$$J^2 \leftrightarrow 2\hbar^2 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad J_z \leftrightarrow \hbar \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad (8.3.2)$$

$$J_+ \leftrightarrow \sqrt{2}\hbar \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad J_- \leftrightarrow \sqrt{2}\hbar \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad (8.3.3)$$

$$J_x \leftrightarrow \frac{\hbar}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad J_y \leftrightarrow \frac{\hbar}{\sqrt{2}} \begin{pmatrix} 0 & -i & 0 \\ i & 0 & -i \\ 0 & i & 0 \end{pmatrix}. \quad (8.3.4)$$

As an example of the use of these matrices, suppose that a system is in an eigenstate of  $J_z$ , and we want to know the possible results and their probabilities if  $J_x$  is measured. To answer this question, we first find the eigenvalues and eigenstates of  $J_x$ , expressed in the orthobasis of  $J_z$  eigenvectors of eq. (8.3.1). Since there is nothing special about the  $x$  direction as opposed to the  $z$  direction, we know even without computing the characteristic equation for  $J_x$  that its eigenvalues must be the same as  $J_z$ , namely  $\hbar$ ,  $0$ , and  $-\hbar$ . Then, solving for the eigenvectors of the matrix representation for  $J_x$ , one finds the normalized kets

$$|J_x = \hbar\rangle = \frac{1}{2}|1, 1\rangle + \frac{1}{\sqrt{2}}|1, 0\rangle + \frac{1}{2}|1, -1\rangle, \quad (8.3.5)$$

$$|J_x = 0\rangle = \frac{1}{\sqrt{2}}|1, 1\rangle - \frac{1}{\sqrt{2}}|1, -1\rangle, \quad (8.3.6)$$

$$|J_x = -\hbar\rangle = \frac{1}{2}|1, 1\rangle - \frac{1}{\sqrt{2}}|1, 0\rangle + \frac{1}{2}|1, -1\rangle. \quad (8.3.7)$$

As a check, these kets are mutually orthogonal, as required (Theorem 2.6.5) by the fact that they are eigenkets of a Hermitian operator with different eigenvalues. Then, for example, the probabilities that a measurement in the state  $|1, 1\rangle$  will yield the results  $J_x = \hbar$ ,  $J_x = 0$ , and  $J_x = -\hbar$  are, by applying Postulate 4,

$$\mathcal{P}_{J_x=\hbar} = |\langle J_x = \hbar | 1, 1 \rangle|^2 = 1/4, \quad (8.3.8)$$

$$\mathcal{P}_{J_x=0} = |\langle J_x = 0 | 1, 1 \rangle|^2 = 1/2, \quad (8.3.9)$$

$$\mathcal{P}_{J_x=-\hbar} = |\langle J_x = -\hbar | 1, 1 \rangle|^2 = 1/4. \quad (8.3.10)$$

## 8.4 Matrix representation for arbitrary $j$

For a subspace with arbitrary  $j$ , the orthobasis of  $J_z$  eigenstates consists of  $2j + 1$  states  $|j, m\rangle$  with  $m = j, j-1, \dots, -j+1, -j$ . The corresponding column vector representations of these states, in that order, are

$$|j, j\rangle \leftrightarrow \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \quad |j, j-1\rangle \leftrightarrow \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{pmatrix}, \quad \dots, \quad |j, -j\rangle \leftrightarrow \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix}. \quad (8.4.1)$$

Again, eqs. (8.1.6), (8.1.7), (8.1.22), and (8.1.24) provide the matrix elements of the angular momentum operators. The  $(2j + 1) \times (2j + 1)$  matrix representations have the forms

$$J^2 \leftrightarrow \hbar^2 j(j+1) \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix}, \quad J_z \leftrightarrow \hbar \begin{pmatrix} j & 0 & \dots & 0 \\ 0 & j-1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & -j \end{pmatrix}, \quad (8.4.2)$$

$$J_+ \leftrightarrow \hbar \begin{pmatrix} 0 & \bullet & 0 & \dots & 0 & 0 \\ 0 & 0 & \bullet & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & \bullet \\ 0 & 0 & 0 & \dots & 0 & 0 \end{pmatrix}, \quad J_- \leftrightarrow \hbar \begin{pmatrix} 0 & 0 & 0 & \dots & 0 & 0 \\ \bullet & 0 & 0 & \dots & 0 & 0 \\ 0 & \bullet & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & \bullet & 0 \end{pmatrix}, \quad (8.4.3)$$

where the  $\bullet$  are the only non-zero elements, which lie just above and below the main diagonal for  $J_+$  and  $J_-$ , respectively. Using eq. (8.1.10),  $J_x$  and  $J_y$  have matrix elements

$$(J_x)_{m',m} = \frac{\hbar}{2} \left[ \sqrt{j(j+1) - m(m-1)} \delta_{m',m-1} + \sqrt{j(j+1) - m(m+1)} \delta_{m',m+1} \right], \quad (8.4.4)$$

$$(J_y)_{m',m} = i \frac{\hbar}{2} \left[ \sqrt{j(j+1) - m(m-1)} \delta_{m',m-1} - \sqrt{j(j+1) - m(m+1)} \delta_{m',m+1} \right], \quad (8.4.5)$$

and so have the forms

$$J_x \leftrightarrow \hbar \begin{pmatrix} 0 & \bullet & 0 & \dots & 0 & 0 \\ \bullet & 0 & \bullet & \dots & 0 & 0 \\ 0 & \bullet & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & \bullet \\ 0 & 0 & 0 & \dots & \bullet & 0 \end{pmatrix}, \quad J_y \leftrightarrow i\hbar \begin{pmatrix} 0 & -\bullet & 0 & \dots & 0 & 0 \\ \bullet & 0 & -\bullet & \dots & 0 & 0 \\ 0 & \bullet & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & -\bullet \\ 0 & 0 & 0 & \dots & \bullet & 0 \end{pmatrix}, \quad (8.4.6)$$

where each pair of  $\bullet$ 's on opposite sides of the main diagonal are equal, consistent with the Hermiticity of these operators. Note that the  $\bullet$ 's are also all real and positive in the phase convention we have chosen.

## 8.5 Matrix representations for unitary rotation operators

In section 5.3, we discussed the unitary operators that act on the Hilbert space to generate rotations parameterized by  $\vec{\alpha} = \hat{n}\alpha$ , where the unit vector  $\hat{n}$  is the axis of rotation, and  $\alpha$  is the magnitude of the rotation angle,

$$U(\vec{\alpha}) = \exp\left(-i\vec{\alpha} \cdot \vec{J}/\hbar\right). \quad (8.5.1)$$

The matrix representations for these unitary operators are called **Wigner functions**, after Eugene P. Wigner, and traditionally denoted as  $D^{(j)}(\vec{\alpha})$ . They are defined by

$$\langle j', m' | U(\vec{\alpha}) | j, m \rangle = \delta_{jj'} D_{m'm}^{(j)}(\vec{\alpha}) \quad (m, m' = -j, \dots, j). \quad (8.5.2)$$

By completeness of the angular momentum eigenstates,

$$U(\vec{\alpha}) | j, m \rangle = \sum_{m'=-j}^j | j, m' \rangle D_{m'm}^{(j)}(\vec{\alpha}). \quad (8.5.3)$$

Note that each  $D^{(j)}$  is a  $(2j+1) \times (2j+1)$  dimensional matrix, which can mix different  $m$  values but keeps  $j$  fixed.

The set of matrices  $D_{m'm}^{(j)}$  for fixed  $j$  form an **irreducible representation** of the rotation group, with the following consequence. If you do one rotation  $\vec{\alpha}$  followed by another  $\vec{\beta}$ , the result is always some other rotation  $\vec{\gamma}$ ,

$$U(\vec{\gamma}) = U(\vec{\beta})U(\vec{\alpha}). \quad (8.5.4)$$

Then, applying completeness gives

$$D_{m'm}^{(j)}(\vec{\gamma}) = \sum_{m''=-j}^j D_{m'm''}^{(j)}(\vec{\beta}) D_{m''m}^{(j)}(\vec{\alpha}). \quad (8.5.5)$$

The unitarity of the operator  $U(\vec{\alpha})$  and the fact that  $U(\vec{\alpha})^{-1} = U(-\vec{\alpha})$  implies

$$D_{m'm}^{(j)}(-\vec{\alpha}) = \left[ D_{mm'}^{(j)}(\vec{\alpha}) \right]^*. \quad (8.5.6)$$

For small  $j$ , the exponentiation in eq. (8.5.1) can be done explicitly in the matrix representation. For the trivial  $j = 0$  case, we have  $D^{(0)}(\vec{\alpha}) = e^0 = 1$ , the unit  $1 \times 1$  matrix, independent of  $\alpha$ . This corresponds to the fact that states with 0 angular momentum are invariant under rotations.

For  $j = 1/2$ ,

$$D^{(1/2)}(\vec{\alpha}) = \exp(-i\vec{\alpha} \cdot \vec{\sigma}/2) = \sum_{k=0}^{\infty} \frac{1}{k!} (-i\vec{\alpha} \cdot \vec{\sigma}/2)^k. \quad (8.5.7)$$

Using eq. (8.2.18), we have  $(\vec{\alpha} \cdot \vec{\sigma})^2 = \alpha^2$ , so that the terms in eq. (8.5.7) have a recurring matrix structure, which allows us to resum the even and odd terms of the infinite sum separately,

$$D^{(1/2)}(\vec{\alpha}) = \cos(\alpha/2)I - i\hat{n} \cdot \vec{\sigma} \sin(\alpha/2), \quad (8.5.8)$$

where  $I$  is the  $2 \times 2$  unit matrix.

As an example, suppose we have a spin-1/2 that has been measured to be along the  $\hat{z}$  direction, so that the state ket is  $|\psi\rangle = |\uparrow\rangle$ , or in the matrix representation,

$$|\psi\rangle \leftrightarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \quad (8.5.9)$$

Now let us rotate this state by an angle  $\theta$  about the  $\hat{y}$  axis. The matrix representation of the rotation is

$$D^{(1/2)}(\hat{y}\theta) = \cos(\theta/2)I - i\sigma_y \sin(\theta/2) = \begin{pmatrix} \cos(\theta/2) & -\sin(\theta/2) \\ \sin(\theta/2) & \cos(\theta/2) \end{pmatrix}, \quad (8.5.10)$$

and so the rotated state has

$$|\psi'\rangle \leftrightarrow \begin{pmatrix} \cos(\theta/2) \\ \sin(\theta/2) \end{pmatrix}, \quad (8.5.11)$$

or  $|\psi'\rangle = \cos(\theta/2)|\uparrow\rangle + \sin(\theta/2)|\downarrow\rangle$ . You can check that this is indeed an eigenstate of the rotated operator

$$S'_z = \cos\theta S_z + \sin\theta S_x \leftrightarrow \frac{\hbar}{2} \begin{pmatrix} \cos\theta & \sin\theta \\ \sin\theta & -\cos\theta \end{pmatrix}, \quad (8.5.12)$$

with eigenvalue  $+\hbar/2$ . If we prepare the system in a spin eigenstate, then the probability to find the spin oriented in a direction at an angle  $\theta$  with respect to the original direction is

$$\mathcal{P} = |\langle\psi'|\psi\rangle|^2 = \cos^2(\theta/2). \quad (8.5.13)$$

As a check, this is 1 for  $\theta = 0$  or  $2\pi$ , and 0 for  $\theta = \pi$ . The probability to find the spin oriented in some particular direction at a right angle to the original spin direction is  $\cos^2(\pi/4) = 1/2$ .

Abstracting this to a slightly more general case, let  $|\hat{n}\rangle$  be the eigenstate of an arbitrary spin component  $\hat{n} \cdot \vec{S}$  with eigenvalue  $\hbar/2$ . Then the probabilities for outcomes of the measurement of a different spin component  $\hat{n}' \cdot \vec{S}$  are

$$\mathcal{P}(\hat{n}' \cdot \vec{S} = +\hbar/2) = |\langle\hat{n}'|\hat{n}\rangle|^2 = \cos^2(\theta/2), \quad (8.5.14)$$

$$\mathcal{P}(\hat{n}' \cdot \vec{S} = -\hbar/2) = |\langle-\hat{n}'|\hat{n}\rangle|^2 = \sin^2(\theta/2), \quad (8.5.15)$$

where  $\theta$  is the angle between  $\hat{n}$  and  $\hat{n}'$ .



A surprising feature of eq. (8.5.8) is that a rotation about any axis by an angle  $\alpha = 2\pi$  gives, instead of the identity matrix:

$$D^{(1/2)}(\hat{n}2\pi) = -I \quad (8.5.16)$$

for a spin-1/2 system. Thus, the  $j = 1/2$  state always acquires a minus sign when continuously rotated by an angle  $\alpha = 2\pi$ , even though such a  $2\pi$  rotation corresponds to no change at all for rotations of classical objects. If the state was in an eigenstate of a particular component  $\hat{m} \cdot \vec{J}$ , then it will still be after the  $2\pi$  rotation, but with a minus sign phase change.

For general  $j$ , the unitary matrix for a rotation about the  $z$  axis is easy to obtain, because  $J_z$  has a diagonal matrix representation, making the exponentiation easy,

$$D^{(j)}(\hat{z}\alpha) = \exp \left[ -i\alpha \begin{pmatrix} j & 0 & \cdots & 0 \\ 0 & j-1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & -j \end{pmatrix} \right] = \begin{pmatrix} e^{-i\alpha j} & 0 & \cdots & 0 \\ 0 & e^{-i\alpha(j-1)} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & e^{i\alpha j} \end{pmatrix}. \quad (8.5.17)$$

For example, in the  $j = 1$  case,

$$D^{(1)}(\hat{z}\alpha) = \text{diag}(e^{-i\alpha}, 1, e^{i\alpha}). \quad (8.5.18)$$

The special case of a rotation through an angle  $2\pi$  gives

$$D^{(j)}(\hat{z}2\pi) = (-1)^{2j} I, \quad (8.5.19)$$

which is equal to the identity matrix for integer  $j$ , but is equal to  $-I$  for half-integer  $j$ , generalizing what was found for  $j = 1/2$ . For this reason, it is impossible to define a single-valued continuous position wavefunction (like the spherical harmonics introduced in the next section) for non-integer  $j$ .

For rotations about other axes  $\hat{n} \neq \hat{z}$ , the algebraic form of matrices  $D^{(j)}(\vec{\alpha})$  for general  $j$  can be considerably more complicated. The resummation of the infinite sum in the exponential may depend on matrix recurrence relations of higher order. For an example of a harder case that can still be done straightforwardly in closed form, one can evaluate  $D^{(1)}(\hat{x}\alpha)$  by using

$$\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}^3 = 2 \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \quad (8.5.20)$$

to resum the exponential series, with the result:

$$D^{(1)}(\hat{x}\alpha) = \begin{pmatrix} \cos^2(\alpha/2) & -\frac{i}{\sqrt{2}} \sin(\alpha) & -\sin^2(\alpha/2) \\ -\frac{i}{\sqrt{2}} \sin(\alpha) & \cos(\alpha) & -\frac{i}{\sqrt{2}} \sin(\alpha) \\ -\sin^2(\alpha/2) & -\frac{i}{\sqrt{2}} \sin(\alpha) & \cos^2(\alpha/2) \end{pmatrix}. \quad (8.5.21)$$

Similar expressions for rotations about the  $\hat{y}$  and  $\hat{z}$  axis for  $j = 1$  are left as an exercise.

For general  $j$ , the numerical value of any  $D^{(j)}(\vec{\alpha})$  can always be obtained by exponentiation of the  $(2j+1) \times (2j+1)$  matrix corresponding to eq. (8.5.1). There is also a remarkable formula, due to Wigner, for rotations about the  $y$ -axis,

$$D_{m'm}^{(j)}(\hat{y}\beta) = \sum_k (-1)^{k-m+m'} [\cos(\beta/2)]^{2j-2k+m-m'} [\sin(\beta/2)]^{2k-m+m'} \frac{[(j+m)!(j-m)!(j+m')!(j-m')!]^{1/2}}{k!(k-m+m')!(j+m-k)!(j-m'-k)!}, \quad (8.5.22)$$

where the sum is over integers  $k$  for which all of the factorials in the denominator have non-negative arguments, in other words  $k$  is summed from  $\max(0, m-m')$  to  $\min(j+m, j-m')$ . The proof is omitted here.

The Wigner formula for the matrix  $D^{(j)}(\hat{y}\beta)$  is especially useful because it can be used to construct a general rotation matrix. The reason is that a general rotation can always be built up as a product of three sequential rotations about two fixed orthogonal axes. Suppose that we first rotate by an angle  $\alpha$  about the  $\hat{z}$  axis, then about the  $\hat{y}$  axis by an angle  $\beta$ , and finally by an angle  $\gamma$  about the  $\hat{z}$  axis again.<sup>†</sup> The change in coordinates induced by this rotation is

$$\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} c_\gamma & -s_\gamma & 0 \\ s_\gamma & c_\gamma & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} c_\beta & 0 & s_\beta \\ 0 & 1 & 0 \\ -s_\beta & 0 & c_\beta \end{pmatrix} \begin{pmatrix} c_\alpha & -s_\alpha & 0 \\ s_\alpha & c_\alpha & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}, \quad (8.5.23)$$

where  $c_\alpha = \cos \alpha$ ,  $s_\alpha = \sin \alpha$ , etc. The components of vector operators including  $\vec{R}$ ,  $\vec{P}$ , and  $\vec{J}$  rotate in the same way as eq. (8.5.23). The corresponding unitary rotation operator is

$$U(\alpha, \beta, \gamma) = U(\hat{z}\gamma) U(\hat{y}\beta) U(\hat{z}\alpha), \quad (8.5.24)$$

so that the Wigner rotation matrix for a multiplet with angular momentum  $j$  is

$$D^{(j)}(\alpha, \beta, \gamma) = D^{(j)}(\hat{z}\gamma) D^{(j)}(\hat{y}\beta) D^{(j)}(\hat{z}\alpha). \quad (8.5.25)$$

The matrices for the first and last rotations about the fixed  $z$  axis are simple, being just given by eq. (8.5.17).

---

<sup>†</sup>There are different conventional ways of defining the three **Euler angles** needed for a general rotation. In classical mechanics, it is traditional to choose the middle rotation to be about the  $\hat{x}$  axis. In quantum mechanics it is preferable to use the  $\hat{y}$  axis, because  $D^{(j)}(\hat{y}\beta)$  has purely real entries, as exhibited in eq. (8.5.22). There are also differing conventions for whether the axes of rotations are absolutely fixed, or whether the second and third rotation axes are the “body” axes, obtained by the previous rotations of the original (fixed)  $y$  and  $z$  axes. Here, our rotation axes are the absolutely fixed ones.

## 8.6 Spherical harmonic representation of orbital angular momentum

The orbital angular momentum operators  $L_x$ ,  $L_y$ , and  $L_z$  were introduced in section 5.3. We now consider the representation of these operators in terms of the position wavefunctions in spherical coordinates. Recall that in the position orthobasis,

$$\psi(r, \theta, \phi) = \langle r, \theta, \phi | \psi \rangle. \quad (8.6.1)$$

Using the definitions of  $L_x$ ,  $L_y$ , and  $L_z$  in eqs. (5.3.3), we get the rectangular coordinate representations of these as differential operators:

$$L_x \leftrightarrow -i\hbar \left( y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y} \right), \quad (8.6.2)$$

$$L_y \leftrightarrow -i\hbar \left( z \frac{\partial}{\partial x} - x \frac{\partial}{\partial z} \right), \quad (8.6.3)$$

$$L_z \leftrightarrow -i\hbar \left( x \frac{\partial}{\partial y} - y \frac{\partial}{\partial x} \right), \quad (8.6.4)$$

which can be converted into spherical coordinate differential operators, with the results

$$L_x \leftrightarrow i\hbar \left( \sin \phi \frac{\partial}{\partial \theta} + \frac{\cos \phi \cos \theta}{\sin \theta} \frac{\partial}{\partial \phi} \right), \quad (8.6.5)$$

$$L_y \leftrightarrow i\hbar \left( -\cos \phi \frac{\partial}{\partial \theta} + \frac{\sin \phi \cos \theta}{\sin \theta} \frac{\partial}{\partial \phi} \right), \quad (8.6.6)$$

$$L_z \leftrightarrow -i\hbar \frac{\partial}{\partial \phi}. \quad (8.6.7)$$

From these, we also obtain for the raising and lowering operators  $L_{\pm} = L_x \pm iL_y$  as defined by eq. (8.1.8),

$$L_+ \leftrightarrow \hbar e^{i\phi} \left( \frac{\partial}{\partial \theta} + i \frac{\cos \theta}{\sin \theta} \frac{\partial}{\partial \phi} \right), \quad (8.6.8)$$

$$L_- \leftrightarrow \hbar e^{-i\phi} \left( -\frac{\partial}{\partial \theta} + i \frac{\cos \theta}{\sin \theta} \frac{\partial}{\partial \phi} \right), \quad (8.6.9)$$

and, using  $L^2 = L_x^2 + L_y^2 + L_z^2 = (L_+ L_- + L_- L_+)/2 + L_z^2$  from eq. (8.1.15), we get

$$L^2 \leftrightarrow -\hbar^2 \left( \frac{\partial^2}{\partial \theta^2} + \frac{\cos \theta}{\sin \theta} \frac{\partial}{\partial \theta} + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2} \right). \quad (8.6.10)$$

For future reference, we note that  $L^2$  is closely related to the part of the Laplacian involving angular derivatives; this is useful because the kinetic energy term in the Hamiltonian involves  $\nabla^2$ . More precisely,

$$\nabla^2 = \frac{\partial^2}{\partial r^2} + \frac{2}{r} \frac{\partial}{\partial r} - \frac{1}{\hbar^2 r^2} L^2. \quad (8.6.11)$$

When acting on a wavefunction corresponding to an eigenstate of  $L^2$  with eigenvalue  $\hbar^2 l(l+1)$ , this becomes

$$\nabla^2 = \frac{\partial^2}{\partial r^2} + \frac{2}{r} \frac{\partial}{\partial r} - \frac{l(l+1)}{r^2}. \quad (8.6.12)$$

We now solve the eigenvalue problem for orbital angular momentum. In doing so, we can make use of a notable feature of eqs. (8.6.5)–(8.6.10): there is no  $r$  dependence in any of the angular momentum operator representations as differential operators on the wavefunctions. Thus we can consider wavefunctions for eigenstates of  $L^2$  and  $L_z$  in which the  $r$  dependence is factored out, with an orthobasis of position eigenkets that are a tensor product of radial and angular parts,

$$|r, \theta, \phi\rangle = |r\rangle \otimes |\theta, \phi\rangle, \quad (8.6.13)$$

with Dirac orthonormality conditions

$$\langle r' | r \rangle = \frac{1}{r^2} \delta(r - r'), \quad (8.6.14)$$

$$\langle \theta', \phi' | \theta, \phi \rangle = \delta(\phi - \phi') \delta(\cos \theta - \cos \theta'), \quad (8.6.15)$$

and completeness relations

$$\int_0^\infty dr r^2 |r\rangle \langle r| = I_r, \quad (8.6.16)$$

$$\int d\Omega |\theta, \phi\rangle \langle \theta, \phi| = I_{\theta, \phi}, \quad (8.6.17)$$

where  $I_r$  and  $I_{\theta, \phi}$  are the identity operators on the respective Hilbert spaces, with  $I = I_r \otimes I_{\theta, \phi}$ . In eq. (8.6.17), and from here on, we define

$$d\Omega = d\phi d(\cos \theta) \quad (8.6.18)$$

as the differential of solid angle in spherical coordinates, so that

$$\int d\Omega \cdots = \int_0^{2\pi} d\phi \int_{-1}^1 d(\cos \theta) \cdots \quad (8.6.19)$$

The Hilbert space spanned by the orthobasis of angular eigenkets  $|\theta, \phi\rangle$  is also spanned by an orthobasis of  $(L^2, L_z)$  eigenkets  $|l, m\rangle$  satisfying, from eqs. (8.1.6), (8.1.7), (8.1.22), and (8.1.24),

$$L_z |l, m\rangle = \hbar m |l, m\rangle, \quad (8.6.20)$$

$$L_+ |l, m\rangle = \hbar \sqrt{l(l+1) - m(m+1)} |l, m+1\rangle, \quad (8.6.21)$$

$$L_- |l, m\rangle = \hbar \sqrt{l(l+1) - m(m-1)} |l, m-1\rangle, \quad (8.6.22)$$

$$L^2 |l, m\rangle = \hbar^2 l(l+1) |l, m\rangle, \quad (8.6.23)$$

for  $l = 0, 1, 2, \dots$  and  $m = -l, \dots, l$ . The  $|l, m\rangle$  also satisfy orthonormality and completeness relations

$$\langle l', m' | l, m \rangle = \delta_{ll'} \delta_{mm'}, \quad (8.6.24)$$

$$\sum_{l=0}^{\infty} \sum_{m=-l}^l |l, m\rangle \langle l, m| = I_{\theta, \phi}. \quad (8.6.25)$$

We now define the angular wavefunctions

$$\langle \theta, \phi | l, m \rangle = Y_l^m(\theta, \phi), \quad (8.6.26)$$

called **spherical harmonics**.

The actions of the differential operators  $L_z$ ,  $L_+$ ,  $L_-$ , and  $L^2$  defined in eqs. (8.6.7)–(8.6.10) on the spherical harmonic wavefunctions are expressed as

$$L_z Y_l^m(\theta, \phi) = \hbar m Y_l^m(\theta, \phi), \quad (8.6.27)$$

$$L_+ Y_l^m(\theta, \phi) = \hbar \sqrt{l(l+1) - m(m+1)} Y_l^{m+1}(\theta, \phi), \quad (8.6.28)$$

$$L_- Y_l^m(\theta, \phi) = \hbar \sqrt{l(l+1) - m(m-1)} Y_l^{m-1}(\theta, \phi), \quad (8.6.29)$$

$$L^2 Y_l^m(\theta, \phi) = \hbar^2 l(l+1) Y_l^m(\theta, \phi). \quad (8.6.30)$$

These are the position representation differential operator versions of the Hilbert space operator equations (8.6.20)–(8.6.23). For simplicity, we are using the same symbols for the differential operators (acting on wavefunctions) and the corresponding Hilbert space operators (acting on kets and bras). From eqs. (8.6.17) and (8.6.24), the spherical harmonics satisfy the orthonormality condition

$$\int d\Omega Y_l^{m'}(\theta, \phi)^* Y_l^m(\theta, \phi) = \delta_{ll'} \delta_{mm'}, \quad (8.6.31)$$

and from eqs. (8.6.15) and (8.6.25), the completeness relation

$$\sum_{l=0}^{\infty} \sum_{m=-l}^l Y_l^m(\theta', \phi')^* Y_l^m(\theta, \phi) = \delta(\phi - \phi') \delta(\cos \theta - \cos \theta'). \quad (8.6.32)$$

The full Hilbert space is now spanned by the orthobasis of tensor product kets

$$|r\rangle \otimes |l, m\rangle, \quad (8.6.33)$$

which describe states in which the particle is known to be at a distance  $r$  from the origin, and in which  $L^2$  and  $L_z$  are also known to be  $\hbar^2 l(l+1)$  and  $\hbar m$  respectively. These orthobasis elements are an alternative to eq. (8.6.13).

Equation (8.6.27) together with eq. (8.6.7) gives the simple differential equation

$$-i\hbar \frac{\partial}{\partial \phi} Y_l^m = \hbar m Y_l^m, \quad (8.6.34)$$

which has the general solution

$$Y_l^m(\theta, \phi) = e^{im\phi} f_l^m(\theta), \quad (8.6.35)$$

where the functions  $f_l^m(\theta)$  are now to be determined. We know already that the largest possible value of  $m$  is  $l$ , so let us start with that case. Equation (8.6.28) implies  $L_+ Y_l^l = 0$ , or

$$\left( \frac{\partial}{\partial \theta} + i \frac{\cos \theta}{\sin \theta} \frac{\partial}{\partial \phi} \right) e^{il\phi} f_l^l(\theta) = 0 \quad (8.6.36)$$

after using eq. (8.6.8). This reduces to

$$\frac{df_l^l}{d(\sin \theta)} = l \frac{f_l^l}{\sin \theta}, \quad (8.6.37)$$

which has the general solution

$$f_l^l(\theta) = e^C (\sin \theta)^l, \quad (8.6.38)$$

where  $C$  is a constant of integration. Thus, we can write

$$Y_l^l(\theta, \phi) = (-1)^l \sqrt{\frac{(2l+1)!}{4\pi}} \frac{1}{2^l l!} e^{il\phi} (\sin \theta)^l, \quad (8.6.39)$$

where a multiplicative normalization factor has been included, with magnitude chosen in such a way that eq. (8.6.31) holds with  $m' = m = l' = l$ . The  $(-1)^l$  factor is a choice of convention.

The spherical harmonics for the remaining values  $m < l$  can now be obtained by acting repeatedly with  $L_-$ , using eq. (8.6.29), and comparing to eq. (8.6.9). First,

$$Y_l^{l-1}(\theta, \phi) = -\frac{e^{-i\phi}}{\sqrt{2l}} \left( \frac{\partial}{\partial \theta} + l \frac{\cos \theta}{\sin \theta} \right) Y_l^l(\theta, \phi). \quad (8.6.40)$$

Continuing in the same way, it can be shown by recursion that, for general  $-l \leq m \leq l$ ,

$$Y_l^m(\theta, \phi) = \frac{(-1)^l}{2^l l!} \sqrt{\frac{(2l+1)(l+m)!}{4\pi(l-m)!}} e^{im\phi} (\sin \theta)^{-m} \frac{d^{l-m}}{d(\cos \theta)^{l-m}} (\sin \theta)^{2l}. \quad (8.6.41)$$

The normalization factor in eq. (8.6.41) ensures that eqs. (8.6.31) and (8.6.32) are satisfied. The phase convention here is determined by the choices that we have already made in eqs. (8.1.22), (8.1.24), and (8.6.39), and is called the **Condon–Shortley phase convention**. (Other phase and normalization conventions for the spherical harmonics exist, so one must be careful when comparing results from different sources.)

An equivalent way of writing the spherical harmonics is

$$Y_l^m(\theta, \phi) = \sqrt{\frac{(2l+1)(l-m)!}{4\pi(l+m)!}} (-1)^m e^{im\phi} P_l^m(\cos\theta), \quad (8.6.42)$$

where the  $P_l^m(u)$  are called associated Legendre functions. They are solutions to the differential equations

$$\left[ (1-x^2) \frac{d^2}{dx^2} - 2x \frac{d}{dx} + l(l+1) - \frac{m^2}{1-x^2} \right] P_l^m(x) = 0. \quad (8.6.43)$$

For the special case  $m = 0$ , the solutions are the ordinary Legendre polynomials,

$$P_l(x) = P_l^0(x) = \frac{1}{2^l l!} \frac{d^l}{dx^l} (x^2 - 1)^l, \quad (8.6.44)$$

which have a generating function

$$\sum_{l=0}^{\infty} t^l P_l(x) = (1 - 2tx + t^2)^{-1/2}, \quad (8.6.45)$$

and satisfy the orthonormality relations

$$\int_{-1}^1 dx P_{l'}(x) P_l(x) = \frac{2}{2l+1} \delta_{ll'}. \quad (8.6.46)$$

The first few Legendre polynomials are

$$P_0(x) = 1, \quad P_1(x) = x, \quad P_2(x) = (3x^2 - 1)/2, \quad (8.6.47)$$

$$P_3(x) = (5x^3 - 3x)/2, \quad P_4(x) = (35x^4 - 30x^2 + 3)/8. \quad (8.6.48)$$

For non-negative  $m$ , the associated Legendre functions are then given by

$$P_l^m(x) = (1-x^2)^{m/2} \frac{d^m}{dx^m} P_l(x), \quad (m \geq 0), \quad (8.6.49)$$

and those with negative  $m$  are

$$P_l^{-m}(x) = \frac{(l-m)!}{(l+m)!} P_l^m(x), \quad (m < 0). \quad (8.6.50)$$

With the Condon–Shortley phase convention,

$$Y_l^{-m}(\theta, \phi) = (-1)^m Y_l^m(\theta, \phi)^*. \quad (8.6.51)$$

For large  $l$  and maximal  $m$ ,  $Y_l^{\pm l}(\theta, \phi) \propto (\sin\theta)^l$ , which is largest in magnitude for  $\theta \approx \pi$ . Thus, states with large angular momentum about the  $z$  axis have probability densities that are peaked near the  $xy$  plane and are suppressed near the  $z$  axis.

Since  $Y_l^m(\theta, \phi)$  is proportional to  $e^{im\phi}$ , the only spherical harmonics that are independent of  $\phi$  are those with  $m = 0$ ,

$$Y_l^0(\theta, \phi) = \sqrt{\frac{2l+1}{4\pi}} P_l(\cos \theta). \quad (8.6.52)$$

In the special case  $\theta = 0$ , the coordinate  $\phi$  is arbitrary. This implies that  $Y_l^m(0, \phi)$  cannot possibly depend on  $\phi$ , so it can only be non-zero if  $m = 0$ . From eq. (8.6.52) and  $P_l(1) = 1$ , we therefore obtain the special value

$$Y_l^m(0, \phi) = \delta_{m0} \sqrt{\frac{2l+1}{4\pi}}. \quad (8.6.53)$$

The spherical harmonics that are most commonly encountered in practical problems are the ones for  $l \leq 2$ , which are

$$Y_0^0 = \frac{1}{\sqrt{4\pi}}, \quad (8.6.54)$$

$$Y_1^0 = \sqrt{\frac{3}{4\pi}} \cos \theta, \quad Y_1^{\pm 1} = \mp \sqrt{\frac{3}{8\pi}} e^{\pm i\phi} \sin \theta, \quad (8.6.55)$$

$$Y_2^0 = \sqrt{\frac{5}{16\pi}} (3 \cos^2 \theta - 1), \quad (8.6.56)$$

$$Y_2^{\pm 1} = \mp \sqrt{\frac{15}{8\pi}} e^{\pm i\phi} \sin \theta \cos \theta, \quad Y_2^{\pm 2} = \sqrt{\frac{15}{32\pi}} e^{\pm 2i\phi} \sin^2 \theta. \quad (8.6.57)$$

Note that  $Y_l^m$  is always a polynomial of degree  $l$  in  $\cos \theta$  and  $\sin \theta$ .

Using completeness of the orthobasis  $|l, m\rangle$  over the Hilbert space component corresponding to the angular coordinates, as expressed in eq. (8.6.25), any wavefunction can be expanded as

$$\psi(r, \theta, \phi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l (\langle r | \otimes \langle \theta, \phi |) |l, m\rangle \langle l, m | \psi \rangle \quad (8.6.58)$$

$$= \sum_{l=0}^{\infty} \sum_{m=-l}^l Y_l^m(\theta, \phi) (\langle r | \otimes \langle l, m |) | \psi \rangle. \quad (8.6.59)$$

Defining functions  $F_{l,m}(r) = (\langle r | \otimes \langle l, m |) | \psi \rangle$ , this can be rewritten as

$$\psi(r, \theta, \phi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l F_{l,m}(r) Y_l^m(\theta, \phi). \quad (8.6.60)$$

To find the coefficient functions  $F_{l,m}(r)$  for a given  $\psi(r, \theta, \phi)$ , multiply both sides of eq. (8.6.60) by  $Y_{l'}^{m'}(\theta, \phi)^*$ , then integrate  $d\Omega$ , then use the orthonormality condition eq. (8.6.31) to reduce the double sum to a single term with  $l' = l$  and  $m' = m$ , and finally rename  $(l', m') \rightarrow (l, m)$ . The result is

$$F_{l,m}(r) = \int d\Omega Y_l^m(\theta, \phi)^* \psi(r, \theta, \phi). \quad (8.6.61)$$



Of course, eqs. (8.6.60) and (8.6.61) also apply if  $\psi$  and  $F$  have no dependence on  $r$ .

If the wavefunction  $\psi(r, \theta, \phi)$  is normalized, so that  $\int_0^\infty dr r^2 \int d\Omega |\psi(r, \theta, \phi)|^2 = 1$ , then one can use the orthonormality condition eq. (8.6.31) again to obtain

$$\sum_{l=0}^{\infty} \sum_{m=-l}^l \int_0^\infty dr r^2 |F_{l,m}(r)|^2 = 1. \quad (8.6.62)$$

It follows that the probability of simultaneously measuring  $L^2$  and  $L_z$  and getting the results  $\hbar^2 l(l+1)$  and  $\hbar m$  is, by using Postulate 4 with  $r$  playing the role of the degeneracy label,

$$\mathcal{P}_{l,m} = \int_0^\infty dr r^2 \left| (\langle r | \otimes \langle l, m |) |\psi\rangle \right|^2 = \int_0^\infty dr r^2 |F_{l,m}(r)|^2. \quad (8.6.63)$$

As a corollary, the probability of measuring just  $L^2$  and getting  $\hbar^2 l(l+1)$  is obtained by summing this over  $m$ , so  $\mathcal{P}_l = \sum_{m=-l}^l \mathcal{P}_{l,m}$ . And, the probability of measuring  $L_z$  and getting the result  $\hbar m$  is  $\mathcal{P}_m = \sum_{l=|m|}^\infty \mathcal{P}_{l,m}$ . Here, the sum over  $l$  starts at  $|m|$ , because for smaller  $l$  there are no states that have  $L_z$  eigenvalue  $\hbar m$ .

In many cases, it is not necessary to actually do the radial integral in eq. (8.6.63) in order to evaluate the probabilities; instead one can make use of ratios between the various possibilities. As a simple but essential example, any wavefunction that is a function of  $r$  only (with no  $\theta$  or  $\phi$  dependence) is proportional to  $Y_0^0$ , so one has  $l = m = 0$ , and the measurements of the compatible orbital angular momentum observables  $L^2$  and  $L_z$  are both certain to give 0.

For a slightly less trivial example, consider a wavefunction

$$\psi(\vec{r}) = C \sin^2 \theta \cos^2 \phi f(r), \quad (8.6.64)$$

where  $C$  and  $\alpha$  are constants. (Note that we are not providing any context about whether this state is related to any particular Hamiltonian.) If  $L^2$  and/or  $L_z$  is measured, what are the possible outcomes and their probabilities? To answer this, we seek to write the wavefunction as a linear combination of spherical harmonics multiplied by functions of  $r$  only. A useful clue is that the wavefunction is quadratic in sines and cosines of  $\theta$  and  $\phi$ , so one should expect that it will involve  $l = 2$  and  $l = 0$ . Indeed, one finds that

$$\sin^2 \theta \cos^2 \phi = \sqrt{\pi} \left[ \frac{2}{3} Y_0^0 - \frac{2}{3\sqrt{5}} Y_2^0 + \sqrt{\frac{2}{15}} Y_2^2 + \sqrt{\frac{2}{15}} Y_2^{-2} \right]. \quad (8.6.65)$$

This implies that the measurements of  $(L^2, L_z)$  can yield only the four possible pairs  $(0, 0)$  and  $(6\hbar^2, 0)$  and  $(6\hbar^2, 2\hbar)$  and  $(6\hbar^2, -2\hbar)$ , with probability ratios, respectively,

$$\mathcal{P}_{l=0,m=0} : \mathcal{P}_{l=2,m=0} : \mathcal{P}_{l=2,m=2} : \mathcal{P}_{l=2,m=-2} = \left| \frac{2}{3} \right|^2 : \left| -\frac{2}{3\sqrt{5}} \right|^2 : \left| \sqrt{\frac{2}{15}} \right|^2 : \left| \sqrt{\frac{2}{15}} \right|^2. \quad (8.6.66)$$

By requiring that the sum of the probabilities is 1, it follows that

$$\mathcal{P}_{l=0,m=0} = 5/9, \quad \mathcal{P}_{l=2,m=0} = 1/9, \quad \mathcal{P}_{l=2,m=2} = \mathcal{P}_{l=2,m=-2} = 1/6. \quad (8.6.67)$$

Note that the radial wavefunction here was irrelevant for the angular momentum measurement probabilities, since it is common factor.

One often needs to consider wavefunctions for states that are pure eigenstates of  $L^2$  and  $L_z$ , with fixed eigenvalues  $\hbar^2 l(l+1)$  and  $\hbar m$ , respectively. These will have the form

$$\psi(r, \theta, \phi) = F(r) Y_l^m(\theta, \phi). \quad (8.6.68)$$

According to eq. (8.6.12), the Laplacian acting on such a wavefunction is

$$\nabla^2 [F(r) Y_l^m(\theta, \phi)] = \left[ \frac{d^2 F}{dr^2} + \frac{2}{r} \frac{dF}{dr} - \frac{l(l+1)}{r^2} F \right] Y_l^m(\theta, \phi). \quad (8.6.69)$$

As a consequence of the rotational invariance of the Laplacian operator, this maintains the form of an eigenfunction of  $L^2$  and  $L_z$  with the same eigenvalues.

We now state and prove an extraordinarily useful formula:

**Theorem 8.6.1. (Spherical harmonics addition formula)** *Consider any two unit vectors  $\hat{n}$  and  $\hat{n}'$ , characterized by their spherical coordinate angles  $(\theta, \phi)$  and  $(\theta', \phi')$ , respectively. Let us call the angle between these vectors  $\gamma$ , so that*

$$\hat{n} \cdot \hat{n}' = \cos \gamma = \cos \theta \cos \theta' + \sin \theta \sin \theta' \cos(\phi - \phi'). \quad (8.6.70)$$

*Then, for each  $l$ ,*

$$P_l(\cos \gamma) = \frac{4\pi}{2l+1} \sum_{m=-l}^l Y_l^m(\theta', \phi')^* Y_l^m(\theta, \phi). \quad (8.6.71)$$

Proof: Since  $P_l(\cos \gamma)$  is a function of  $\theta$  and  $\phi$ , it can be written as an expansion in spherical harmonics with arguments  $\theta$  and  $\phi$ . Since the differential operator  $L^2$  is invariant under rotations, the spherical harmonics involved in the linear combination all must have the same  $l$ . This is because we can always rotate to a coordinate system in which  $\hat{n}'$  is the new  $z$  direction, so that the polar angular coordinate of  $\hat{n}$  is  $\gamma$ . In that coordinate system,  $P_l(\cos \gamma)$  is proportional to a spherical harmonic with  $m = 0$ , an eigenfunction of the differential operator  $L^2$  with eigenvalue  $\hbar^2 l(l+1)$ . Likewise,  $P_l(\cos \gamma)$  is also a function of  $\theta'$  and  $\phi'$ , so it can also be written as a linear combination of spherical harmonics of those angles, with the same  $l$ . Therefore, it must be that  $P_l(\cos \gamma)$  is a sum of terms of the form  $Y_l^{m'}(\theta', \phi')^* Y_l^m(\theta, \phi)$ . Each of these terms is proportional

to  $e^{i(m\phi-m'\phi')}$ , but if one chooses  $\phi' = \phi$ , then  $\cos \gamma$  is independent of  $\phi$ , so only terms with  $m' = m$  can occur. Therefore, we must have

$$P_l(\cos \gamma) = \sum_{m=-l}^l a_{lm} Y_l^m(\theta', \phi')^* Y_l^m(\theta, \phi), \quad (8.6.72)$$

and the remaining task is to evaluate the coefficients  $a_{lm}$ .

Because  $P_l(\cos \gamma)$  is real, and is unchanged if we exchange  $(\theta, \phi) \leftrightarrow (\theta', \phi')$ , the coefficients must satisfy  $a_{lm} = a_{lm}^*$ , so they are real. To learn more, consider the special case  $(\theta, \phi) = (\theta', \phi')$ , so that  $\cos \gamma = 1$ . Then, since the Legendre polynomials satisfy  $P_l(1) = 1$ , eq. (8.6.72) gives

$$1 = \sum_{m=-l}^l a_{lm} |Y_l^m(\theta, \phi)|^2. \quad (8.6.73)$$

Integrating with respect to  $d\Omega$ , and using the orthonormality condition eq. (8.6.31), we get

$$4\pi = \sum_{m=-l}^l a_{lm}. \quad (8.6.74)$$

Next, consider the square of eq. (8.6.72), which can be written

$$\left(P_l(\cos \gamma)\right)^2 = \sum_{m=-l}^l a_{lm} Y_l^m(\theta', \phi')^* Y_l^m(\theta, \phi) \sum_{m'=-l}^l a_{lm'} Y_l^{m'}(\theta', \phi') Y_l^{m'}(\theta, \phi)^*. \quad (8.6.75)$$

Let us integrate over all angles  $(\theta', \phi')$ . To evaluate the integral of the left side, it is convenient to again use coordinates such that  $\gamma$  is the polar angle, and making use of eq. (8.6.46), we have

$$\int d\Omega' \left(P_l(\cos \gamma)\right)^2 = 2\pi \int_{-1}^1 d(\cos \gamma) \left(P_l(\cos \gamma)\right)^2 = \frac{4\pi}{2l+1}. \quad (8.6.76)$$

Meanwhile, the  $d\Omega'$  integral of the right side of eq. (8.6.75) is evaluated by using the orthonormality condition eq. (8.6.31) again, after which only the terms with  $m' = m$  contribute in the double sum. Comparing the two sides, we get

$$\frac{4\pi}{2l+1} = \sum_{m=-l}^l a_{lm}^2 Y_l^m(\theta, \phi)^* Y_l^m(\theta, \phi). \quad (8.6.77)$$

Integrating this with respect to  $d\Omega$ , and using orthonormality once again, we find

$$\frac{(4\pi)^2}{2l+1} = \sum_{m=-l}^l a_{lm}^2. \quad (8.6.78)$$

We now have enough information to solve for the coefficients. From eqs. (8.6.74) and (8.6.78), we discover that

$$\sum_{m=-l}^l \left(a_{lm} - \frac{4\pi}{2l+1}\right)^2 = 0. \quad (8.6.79)$$

Since the left side is a sum of squares, the unique solution is obtained when each term vanishes, so that  $a_{lm} = 4\pi/(2l+1)$  for all  $l, m$ . This concludes the proof of the spherical harmonics addition formula.  $\square$

As an application, consider the identity

$$\frac{1}{|\vec{r} - \vec{r}'|} = \sum_{l=0}^{\infty} \frac{r_{\min}^l}{r_{\max}^{l+1}} P_l(\cos \gamma), \quad (8.6.80)$$

with

$$r_{\max} = \max(r, r'), \quad r_{\min} = \min(r, r'). \quad (8.6.81)$$

Equation (8.6.80) follows from the generating function eq. (8.6.45), and may be familiar from the multipole expansion for problems with azimuthal symmetry in electrostatics. The spherical harmonics addition formula (8.6.71) allows us to rewrite it as

$$\frac{1}{|\vec{r} - \vec{r}'|} = \sum_{l=0}^{\infty} \frac{4\pi}{2l+1} \frac{r_{\min}^l}{r_{\max}^{l+1}} \sum_{m=-l}^l Y_l^m(\theta', \phi')^* Y_l^m(\theta, \phi), \quad (8.6.82)$$

a form that will be useful to us later.

## 8.7 Parity of angular momentum eigenstates

As we noted in section 5.3, angular momentum operators have even parity, which is another way of saying that they commute with the parity operator  $\Pi$ . According to Theorem 2.7.1, this means that the eigenstates of angular momentum operators  $J^2, J_z$  can also be chosen to be parity eigenstates.

First, let us work out the parity eigenvalues of the orbital angular momentum eigenstates  $|l, m\rangle$ . If  $(x, y, z) \rightarrow (-x, -y, -z)$ , the spherical coordinate transformations are

$$r \rightarrow r, \quad \theta \rightarrow \pi - \theta, \quad \phi \rightarrow \phi + \pi, \quad (8.7.1)$$

so that

$$\cos \theta \rightarrow -\cos \theta, \quad \sin \theta \rightarrow \sin \theta, \quad e^{im\phi} \rightarrow (-1)^m e^{im\phi}. \quad (8.7.2)$$

Using these, it follows immediately from eq. (8.6.41) that

$$Y_l^m(\pi - \theta, \phi + \pi) = (-1)^l Y_l^m(\theta, \phi). \quad (8.7.3)$$

Therefore,

$$\langle \theta, \phi | \Pi | l, m \rangle = \langle \pi - \theta, \phi + \pi | l, m \rangle = (-1)^l Y_l^m(\theta, \phi) = (-1)^l \langle \theta, \phi | l, m \rangle, \quad (8.7.4)$$

so

$$\Pi |l, m\rangle = (-1)^l |l, m\rangle. \quad (8.7.5)$$

This shows that eigenstates of orbital angular momentum are always even (odd) under parity if the quantum number  $l$  is even (odd). This is true regardless of the quantum number  $m$ , which can be understood from the fact that the states  $|l, m\rangle$  are all obtained from  $|l, l\rangle$  by acting with  $L_-$ , which commutes with  $\Pi$ . [Therefore, the parity eigenvalue  $(-1)^l$  could actually have been obtained by considering the spherical harmonics for only one sample value of  $m$  for each  $l$ , for example  $m = l$  using eq. (8.6.39), rather than the more general formula of eq. (8.6.41).] Since parity in spherical coordinates does not change the radial coordinate, the parity eigenvalue of an angular momentum eigenstate also does not depend on the radial wavefunction or radial quantum numbers.

Let us next consider the parity of eigenkets  $|s, m_s\rangle$  of intrinsic angular momentum operators  $S^2$  and  $S_z$ . From the fact that the lowering operator  $S_-$  commutes with parity and relates kets with different  $m_s$ , we know that the parity eigenvalue of  $|s, m_s\rangle$  cannot depend on  $m_s$ . We can therefore write

$$\Pi |s, m_s\rangle = \eta |s, m_s\rangle, \quad (8.7.6)$$

where  $\eta$  is known as the **intrinsic parity** of the particle in question. For any given particle type, the intrinsic parity can be chosen to be either  $+1$  or  $-1$ , as an arbitrary convention. This conventional choice cannot be of any practical significance at all if the number of particles of each type does not change. This is because a change in the conventional choice of intrinsic parity just amounts to flipping the parity for every bra and every ket, therefore not affecting matrix elements. So, when the number of particles does not change, one might as well just always choose  $\eta = +1$ , and the parity of a single particle  $L^2$  eigenstate is  $(-1)^l$ , regardless of the spin. For a system of  $N$  particles with individual orbital angular momentum quantum numbers  $l_i$ , the parity eigenvalue is, from eq. (8.7.5),

$$\pi = (-1)^{\sum_{i=1}^N l_i}. \quad (8.7.7)$$

This is the case for electrons in an atom, where the parity and the angular momenta are all defined with respect to the origin chosen to be the location of the fixed nucleus.

However, if the Hamiltonian is invariant under parity and can cause changes in the numbers of particles, it is natural and useful to adopt a convention in which the intrinsic parities of particles and antiparticles are chosen in a consistent way so that parity is conserved. Such Hamiltonians arise in quantum field theory, where the electromagnetic and strong nuclear interactions conserve

parity, while the weak nuclear interactions do not but can often be treated as a perturbation. A generally accepted convention is to choose  $\eta = +1$  for spin-1/2 particles (the electron, proton, neutron, muon, and quarks, for example). Then the structure of kinetic terms in relativistic quantum field theory can be used to show that their antiparticles (the positron, antiproton, antineutron, antimuon, and antiquarks) must be assigned  $\eta = -1$  in the same convention.

For a composite particle  $c$ , made out of two particles  $a$  and  $b$  with intrinsic parities  $\eta_a$  and  $\eta_b$  in a state with relative angular momentum quantum number  $L$ , the consistent intrinsic parity assignment is<sup>†</sup>

$$\eta_c = (-1)^L \eta_a \eta_b. \quad (8.7.8)$$

Thus, mesons composed of a quark and an antiquark in an  $L = 0$  bound state (for example, pions and kaons) have intrinsic parity  $\eta = -1$ . For a bound state of three or more particles, the situation is more complicated, but the intrinsic parities of bound states can always be defined if the Hamiltonian is invariant under space inversion. In processes governed by the electromagnetic and strong nuclear forces, one can experimentally verify parity conservation and check the consistency of the intrinsic parity assignments. In doing so, the photon and the gluon have intrinsic parity  $-1$ , while the Higgs particle has intrinsic parity  $+1$ . The  $W$  and  $Z$  particles are not assigned a well-defined intrinsic parity, as they are the mediators of the parity-violating weak interactions.

## 8.8 Exercises

**Exercise 8.1.** Consider an eigenstate  $|j, m\rangle$  of  $J^2$  and  $J_z$ , as in eq. (8.1.6)-(8.1.7). Find the expectation value and the uncertainties of  $J_x$  and  $J_y$  in this state. For which state or states  $|j, m\rangle$  are these uncertainties maximized? For which are they minimized? Show that the uncertainty relation eq. (3.3.6) is satisfied.

**Exercise 8.2.** Consider a particle with spin 1/2, and let  $\hat{n} = \hat{x} \sin \beta + \hat{z} \cos \beta$  be a fixed unit vector, where  $\beta$  is a fixed angle.

(a) Consider the basis of eigenstates of  $S_z$ , denoted  $|\uparrow\rangle$  and  $|\downarrow\rangle$  for eigenvalues  $+\hbar/2$  and  $-\hbar/2$  respectively. In that basis, construct the matrix representation of  $\hat{n} \cdot \vec{S}$ . Find its eigenvalues and eigenvectors.

(b) Suppose that the spin is in the state  $|\uparrow\rangle$ . What is the probability that the measurement of  $\hat{n} \cdot \vec{S}$  yields the result  $+\hbar/2$ ?

---

<sup>†</sup>Note that this differs from eq. (8.7.7), since  $(-1)^L$  is not always equal to  $(-1)^{l_a+l_b}$ . This is because the angular momentum  $L$  and the intrinsic parity  $\eta_c$  in eq. (8.7.8) are defined with respect to the origin as the center-of-mass position of the two particles, rather than a fixed origin position as in eq. (8.7.7).

(c) Suppose that the measurement in part (b) has been carried out and the result for  $\hat{n} \cdot \vec{S}$  was indeed  $+\hbar/2$ . Immediately afterwards,  $S_z$  is measured. What is the probability that the measurement yields  $-\hbar/2$  ?

(d) Check that your results for parts (b) and (c) make sense when  $\beta = 0$  and  $\pi$  and  $\pi/2$ .

**Exercise 8.3.** Consider a spinless particle in a state with  $L^2$  eigenvalue  $2\hbar^2$  and  $L_z$  eigenvalue  $\hbar$ . As in the previous problem, let  $\hat{n} = \hat{x} \sin \beta + \hat{z} \cos \beta$ .

(a) Suppose that the angular momentum along the direction  $\hat{n}$  is measured. What are the possible results, and their probabilities?

(b) For each of the possible results in part (a), suppose that  $L_z$  is then measured. What are the possible results, and their probabilities?

(c) Check that your results make sense when  $\beta = 0$  and  $\pi$  and  $\pi/2$ .

**Exercise 8.4.** Consider a particle in a state with spherical coordinate wavefunction of the form  $\psi(r, \theta, \phi) = f(r) \cos^2 \theta$ .

(a) Write the wavefunction in terms of spherical harmonics  $Y_l^m(\theta, \phi)$ .

(b) If  $L^2$  and  $L_z$  are measured simultaneously, find the possible results and their probabilities.

**Exercise 8.5.** Professor Bumble measured the spin component of a spin-1/2 system at time  $t = 0$  and recorded the result as definitely  $+\hbar/2$ , but forgot whether it was  $S_x$ ,  $S_y$ , or  $S_z$ . Assigning equal probabilities to these possibilities, what is the density matrix operator  $\rho$  for the resulting mixed state at  $t = 0$ , in the basis of  $S_z$  eigenstates? What is the von Neumann entropy  $\sigma$ ? If the Hamiltonian is  $H = \omega S_z$ , use the unitary time evolution operator to find the probability of getting the result  $+\hbar/2$  for a measurement of  $S_x$  at time  $t$ .

## 9 Charged particle in a magnetic field

### 9.1 Spin precession in a constant magnetic field

One of the simplest, and yet most useful, quantum systems is a spin carrying a magnetic moment in a constant magnetic field. Let us assume that the particle carrying the spin has a fixed position, so that we do not need to worry about its kinetic energy and any position-dependent potentials. Specializing eq. (4.3.28), the Hamiltonian is therefore simply

$$H = -\gamma \vec{B} \cdot \vec{S}, \quad (9.1.1)$$

where  $\gamma$  is the gyromagnetic ratio of the particle. Since we only care about the magnetic field at the point where the particle is located, we can orient our coordinate system so that  $\vec{B} = B\hat{z}$ , where  $B$  is a constant number. An orthobasis of states for this system consists of the eigenstates of  $S_z$  with eigenvalues  $\hbar/2$  and  $-\hbar/2$ , denoted  $|\uparrow\rangle$  for spin up and  $|\downarrow\rangle$  for spin down. Then the Hamiltonian can be expressed as the spectral decomposition

$$H = \frac{1}{2}\hbar\omega_B \left( |\uparrow\rangle \langle\uparrow| - |\downarrow\rangle \langle\downarrow| \right), \quad (9.1.2)$$

where we have defined the **Larmor precession frequency**

$$\omega_B = -\gamma B. \quad (9.1.3)$$

(With this sign choice,  $\omega_B$  is positive for the electron when  $B$  is positive, since the electron's gyromagnetic ratio  $\gamma = -g_e e/2m_e c$  is a negative number. The same is true for the muon and the neutron. For the proton,  $\omega_B$  is negative when  $B$  is positive.) Thus the matrix representation for this orthobasis is

$$H = \frac{1}{2}\hbar\omega_B \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (9.1.4)$$

with energy eigenstates  $|\uparrow\rangle$  represented by  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$  with  $E_\uparrow = \hbar\omega_B/2$ , and  $|\downarrow\rangle$  represented by  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$  with  $E_\downarrow = -\hbar\omega_B/2$ . To decide which of these is the ground state requires knowing the product of the signs of  $B$  and  $\gamma$ . In any case, a transition between the two stationary states typically involves emitting or absorbing a photon with energy  $|E_\uparrow - E_\downarrow| = \hbar|\omega_B|$ . The Planck-Einstein relation, discussed in section 1.3, says that the wavelength of the photon will be  $\lambda = 2\pi c/|\omega_B| = 2\pi c/|\gamma B|$ , providing a way of determining the gyromagnetic ratio of the particle.

Another way of measuring the gyromagnetic ratio (or equivalently, the magnetic moment) is provided by the phenomenon of precession of the spin when the state is a linear combination of energy eigenstates. Suppose that at time  $t = 0$  the normalized initial state is

$$|\psi(0)\rangle = a |\uparrow\rangle + b |\downarrow\rangle, \quad (9.1.5)$$



with complex coefficients satisfying  $|a|^2 + |b|^2 = 1$ . The subsequent time evolution can be obtained either by solving the time-dependent Schrödinger equation, or by applying the unitary operator  $U(t) = e^{-itH/\hbar}$ , with the result

$$|\psi(t)\rangle = ae^{-i\omega_B t/2} |\uparrow\rangle + be^{i\omega_B t/2} |\downarrow\rangle. \quad (9.1.6)$$

If  $S_z$  is then measured, the possible results are the eigenvalues  $\pm\hbar/2$ , with probabilities

$$\mathcal{P}(S_z = \hbar/2) = |\langle\uparrow|\psi(t)\rangle|^2 = |a|^2, \quad (9.1.7)$$

$$\mathcal{P}(S_z = -\hbar/2) = |\langle\downarrow|\psi(t)\rangle|^2 = |b|^2, \quad (9.1.8)$$

which do not depend on time. (As a check, they sum to 1 because of the normalization condition.) It follows that the expectation value of  $S_z$  is

$$\langle S_z \rangle = \frac{\hbar}{2}(|a|^2 - |b|^2), \quad (9.1.9)$$

which is also independent of time.

If, instead, we measure  $S_x$ , then the possible results are again  $\pm\hbar/2$ , and the evaluation of the probabilities makes use of the corresponding eigenstates

$$|\uparrow_x\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle + |\downarrow\rangle), \quad (\text{for } S_x = \hbar/2), \quad (9.1.10)$$

$$|\downarrow_x\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle - |\downarrow\rangle), \quad (\text{for } S_x = -\hbar/2). \quad (9.1.11)$$

We therefore find

$$\mathcal{P}(S_x = \hbar/2) = |\langle\uparrow_x|\psi(t)\rangle|^2 = \left| \frac{1}{\sqrt{2}} (ae^{-i\omega_B t/2} + be^{i\omega_B t/2}) \right|^2 \quad (9.1.12)$$

$$= \frac{1}{2} (|a|^2 + |b|^2 + a^*be^{i\omega_B t} + ab^*e^{-i\omega_B t}) = \frac{1}{2} + \text{Re}(a^*be^{i\omega_B t}). \quad (9.1.13)$$

Similarly (or, just by requiring that the probabilities add to 1), one finds

$$\mathcal{P}(S_x = -\hbar/2) = |\langle\downarrow_x|\psi(t)\rangle|^2 = \frac{1}{2} - \text{Re}(a^*be^{i\omega_B t}). \quad (9.1.14)$$

The expectation value of  $S_x$  can be obtained either from the probability-weighted sum of eigenvalues as

$$\langle S_x \rangle = \frac{\hbar}{2}\mathcal{P}(S_x = \hbar/2) - \frac{\hbar}{2}\mathcal{P}(S_x = -\hbar/2), \quad (9.1.15)$$

or from matrix evaluation, using  $S_x = (\hbar/2)\sigma_x$ , as

$$\langle S_x \rangle = \begin{pmatrix} a^*e^{i\omega_B t/2} & b^*e^{-i\omega_B t/2} \end{pmatrix} \frac{\hbar}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} ae^{-i\omega_B t/2} \\ be^{i\omega_B t/2} \end{pmatrix} = \hbar \text{Re}(a^*be^{i\omega_B t}). \quad (9.1.16)$$




Figure 9.1.1: The expectation value of a spin precesses about a magnetic field  $\vec{B} = B\hat{z}$ , with a period  $2\pi/|\omega_B| = 2\pi/|\gamma B|$ , where  $\gamma$  is the gyromagnetic ratio of the particle. The polar angle  $\theta$  remains constant as the spin precesses, and is equal to  $\arctan(2ab/(a^2 - b^2))$  for an initial state  $|\psi\rangle = a|\uparrow\rangle + b|\downarrow\rangle$  with real  $a$  and  $b$ .

For a measurement of  $S_y$ , the probabilities are evaluated using the corresponding eigenstates

$$|\uparrow_y\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle + i|\downarrow\rangle), \quad (\text{for } S_y = \hbar/2), \quad (9.1.17)$$

$$|\downarrow_y\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle - i|\downarrow\rangle), \quad (\text{for } S_y = -\hbar/2), \quad (9.1.18)$$

with the results

$$\mathcal{P}(S_y = \hbar/2) = |\langle \uparrow_y | \psi(t) \rangle|^2 = \frac{1}{2} + \text{Im}(a^* b e^{i\omega_B t}), \quad (9.1.19)$$

$$\mathcal{P}(S_y = -\hbar/2) = |\langle \downarrow_y | \psi(t) \rangle|^2 = \frac{1}{2} - \text{Im}(a^* b e^{i\omega_B t}), \quad (9.1.20)$$

and the resulting expectation value is

$$\langle S_y \rangle = \hbar \text{Im}(a^* b e^{i\omega_B t}). \quad (9.1.21)$$

The oscillation of these quantities describes precession of  $\vec{S}$  about the  $\vec{B}$  direction.

For example, take  $a$  and  $b$  to be real for simplicity. Then our results for the expectation values of the spin components are summarized as

$$\langle \vec{S} \rangle = \frac{\hbar}{2} \left( 2ab [\cos(\omega_B t) \hat{x} + \sin(\omega_B t) \hat{y}] + (a^2 - b^2) \hat{z} \right). \quad (9.1.22)$$

At time  $t = 0$ , the spin expectation value lies in the  $xz$  plane. As time evolves, the angle  $\theta = \arctan(2ab/(a^2 - b^2))$  between  $\langle \vec{S} \rangle$  and  $\hat{z}$  remains constant. (See Figure 9.1.1.) The rotation of  $\langle \vec{S} \rangle$  is counterclockwise (as seen looking down from positive  $z$ ) if  $\gamma B$  is negative, since the angle  $\phi = \omega_B t$  of the expectation value of the spin vector projected onto the  $xy$  plane increases at the constant rate  $\omega_B = -\gamma B$ . This precession frequency is independent of the initial orientation of the spin, provided only that  $ab \neq 0$ . Thus, the rate of precession of the spin in a known magnetic field provides another way of measuring the gyromagnetic ratio. This effect has many practical applications, including (in more sophisticated setups) the measurement with exquisite accuracy of the magnetic moments of the electron and the muon and various nuclei.

It is useful to describe the spin precession more directly in terms of the state ket, rather than the expectation value, which after all is merely a statistical quantity. Inspired by the result of eq. (9.1.22), consider the time-dependent unit vector

$$\hat{n} = 2ab[\cos(\omega_B t)\hat{x} + \sin(\omega_B t)\hat{y}] + (a^2 - b^2)\hat{z}, \quad (9.1.23)$$

still taking  $a$  and  $b$  to be real. Note that  $\hat{n} \cdot \hat{n} = 1$  follows from the normalization condition  $a^2 + b^2 = 1$ . Now define the spin component along  $\hat{n}$ , as the Hermitian operator

$$S_{\hat{n}} = \hat{n} \cdot \vec{S} = \frac{\hbar}{2} \begin{pmatrix} a^2 - b^2 & 2abe^{-i\omega_B t} \\ 2abe^{i\omega_B t} & b^2 - a^2 \end{pmatrix}. \quad (9.1.24)$$

The last expression is really the matrix representation in the  $S_z$ -eigenstate basis, obtained using the Pauli matrices according to  $\vec{S} = (\hbar/2)\vec{\sigma}$ . It is now straightforward to check that the state ket  $|\psi(t)\rangle$  of eq. (9.1.6), represented by the column vector

$$\begin{pmatrix} ae^{-i\omega_B t/2} \\ be^{i\omega_B t/2} \end{pmatrix}, \quad (9.1.25)$$

satisfies the eigenvalue equation

$$S_{\hat{n}} |\psi(t)\rangle = \frac{\hbar}{2} |\psi(t)\rangle. \quad (9.1.26)$$

In other words, the time-dependent state is simply the one in which measurement of  $S_{\hat{n}}$  is always certain to give the result  $\hbar/2$ .

## 9.2 Magnetic spin resonance and Rabi oscillations

As discussed in the previous section, the energy eigenvalues of a spin-1/2 particle with gyro-magnetic ratio  $\gamma$  are split by a static magnetic field. The spin of a general state precesses in a way that does not change its expectation value along the magnetic field direction. One way to directly access the energy splitting is through emission or absorption of photons of the right energy. In this section, we will discuss another way. The idea is to supplement the main static component  $B$  of the magnetic field with a time-oscillating component  $\tilde{B}$  in a different direction. This oscillatory component drives transitions between the spin eigenstates defined with respect to the static field direction. As we will see, this effect is maximally enhanced (resonant) if the driving frequency for  $\tilde{B}$  is chosen equal to the Larmor precession frequency for  $B$ .

Specifically, let the magnetic field be

$$\vec{B} = B\hat{z} + \tilde{B}[\cos(\omega t)\hat{x} + \sin(\omega t)\hat{y}]. \quad (9.2.1)$$

The oscillatory component rotates in the  $xy$  plane at the driving frequency  $\omega$ , which can be adjusted independently by the experimentalist. Now define

$$\omega_B = -\gamma B, \quad \Gamma = -\gamma \tilde{B}, \quad (9.2.2)$$

both with units of frequency. Note that  $\omega_B$  is the same as the Larmor precession frequency (see the previous section) for the constant component of the magnetic field. The Hamiltonian matrix in the  $S_z$ -eigenstate basis is

$$H = \frac{\hbar\omega_B}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + \frac{\hbar\Gamma}{2} \begin{pmatrix} 0 & e^{-i\omega t} \\ e^{i\omega t} & 0 \end{pmatrix}. \quad (9.2.3)$$

As an aside, this Hamiltonian can also serve as a model for a variety of other 2-state physical systems, for which the static part of the Hamiltonian has been diagonalized by a choice of basis, and there is an off-diagonal contribution varying harmonically with time. Thus the results below have more general interest and applicability.

Suppose that  $\omega_B > 0$ , so that the ground state (neglecting the oscillating part of  $H$ ) is  $|\downarrow\rangle$ . We start in this ground state, and want to find the probability to find the system in the excited state  $|\uparrow\rangle$ , as a function of time. To accomplish this goal, we will find the time-dependent state

$$|\psi(t)\rangle = c_1(t) |\uparrow\rangle + c_2(t) |\downarrow\rangle \quad (9.2.4)$$

that solves the Schrödinger equation for the full  $H$ , subject to the initial condition

$$c_1(0) = 0, \quad c_2(0) = 1. \quad (9.2.5)$$

In matrix form, the time-dependent Schrödinger equation is

$$i\hbar \frac{d}{dt} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \frac{\hbar}{2} \begin{pmatrix} \omega_B & \Gamma e^{-i\omega t} \\ \Gamma e^{i\omega t} & -\omega_B \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}. \quad (9.2.6)$$

It is helpful to define new coefficients  $a_1(t)$  and  $a_2(t)$ , by

$$c_1 = e^{-i\omega t/2} a_1, \quad c_2 = e^{i\omega t/2} a_2. \quad (9.2.7)$$

This is convenient because the resulting coupled first-order differential equations for  $a_1$  and  $a_2$  have coefficients with no explicit time dependence:

$$i\dot{a}_1 + (\omega - \omega_B)a_1/2 - \Gamma a_2/2 = 0, \quad (9.2.8)$$

$$i\dot{a}_2 - (\omega - \omega_B)a_2/2 - \Gamma a_1/2 = 0. \quad (9.2.9)$$

Mindful of the initial conditions in eq. (9.2.5), we now try for a harmonic solution of the form

$$a_1 = b_1 \sin(\Omega t), \quad (9.2.10)$$

$$a_2 = \cos(\Omega t) + b_2 \sin(\Omega t), \quad (9.2.11)$$




Figure 9.2.1: The probability to find a spin-1/2 system in the higher-energy state due to magnetic spin resonance, as a function of time, according to eq. (9.2.16). The peak probability for the spin to flip is maximized near unity for the resonant condition that the driving frequency  $\omega$  is equal to the Larmor precession frequency  $\omega_B = -\gamma B$ , where  $\gamma$  is the gyromagnetic ratio of the particle and  $B$  is the constant part of the magnetic field.

where  $\Omega$  and  $b_1$  and  $b_2$  are constants to be determined. Plugging this guess in, we can require the vanishing of the coefficients of  $\cos(\Omega t)$  and  $\sin(\Omega t)$  separately in each differential equation. Success is found provided that

$$\Omega^2 = [\Gamma^2 + (\omega - \omega_B)^2] / 4 \quad (9.2.12)$$

and

$$b_1 = -i\Gamma/2\Omega, \quad b_2 = i(\omega_B - \omega)/2\Omega. \quad (9.2.13)$$

Putting everything together, the coefficients in  $|\psi(t)\rangle$  are

$$c_1 = -i\frac{\Gamma}{2\Omega} \sin(\Omega t) e^{-i\omega t/2}, \quad (9.2.14)$$

$$c_2 = \left[ \cos(\Omega t) + i\frac{\omega_B - \omega}{2\Omega} \sin(\Omega t) \right] e^{i\omega t/2}. \quad (9.2.15)$$

The probability to find the spin in the excited state is therefore,

$$\mathcal{P}(\uparrow) = |\langle \uparrow | \psi(t) \rangle|^2 = |c_1|^2 = \frac{\Gamma^2}{\Gamma^2 + (\omega - \omega_B)^2} \sin^2(\Omega t). \quad (9.2.16)$$

This shows resonance behavior, for if the driving frequency  $\omega$  is close to the Larmor precession frequency  $\omega_B$ , the system periodically oscillates into the higher state with probability close to 1, as depicted in Figure 9.2.1. These **Rabi oscillations** (named after Isidor Rabi, who both measured and explained them theoretically) occur with frequency  $2\Omega$ , which near resonance is close in magnitude to  $|\Gamma| \approx |\gamma \tilde{B}|$ , where  $\tilde{B}$  is the amplitude of the oscillating magnetic field. Away from the resonance, the probability for a spin flip is non-zero, but smaller, and the frequency of the observed Rabi oscillations is larger.

In real-world applications, the driving magnetic field is rarely of the form we assumed in eq. (9.2.1), because it is not so easy to make a substantial magnetic field whose direction rotates at a high frequency. Instead, an oscillating magnetic field along a fixed direction, like  $\tilde{B} \cos(\omega t) \hat{x}$ , is typically used for the driving field. This is more difficult to solve exactly, but has essentially the same behavior near resonance. The reason is that we can decompose it as

$$\tilde{B} \cos(\omega t) \hat{x} = \frac{1}{2} \tilde{B} [\cos(\omega t) \hat{x} + \sin(\omega t) \hat{y}] + \frac{1}{2} \tilde{B} [\cos(\omega t) \hat{x} - \sin(\omega t) \hat{y}], \quad (9.2.17)$$

which is a superposition of two rotating fields with angular frequencies  $+\omega$  and  $-\omega$ . When one of these is resonant (close to  $\omega_B$ ), the other is far from resonant and therefore has a small effect.

This phenomenon has many applications, not the least of which is **magnetic resonance imaging**, commonly used in medical settings to safely produce maps of tissue using the magnetic moments of hydrogen nuclei (protons) affected in different signature ways by the fields of neighboring atoms in different molecules. In other applications, **nuclear magnetic resonance** is made more complicated and interesting by the fact that the nuclear spin  $s_N$  need not be  $1/2$ , in which case the number of states involved in the resonance is not 2, but  $2s_N + 1$ .

### 9.3 Landau levels for a charged particle moving in a constant uniform magnetic field

Classically, a charged particle in a constant uniform magnetic field undergoes cyclotron motion, following a helical trajectory. This problem is exactly solvable in quantum mechanics, too, and we will now study it in detail.

First, let us gain some insight by reviewing the classical solution. To be concrete, let us take the particle to be an electron with charge  $q = -e$  and mass  $m$ , and in this section we orient the coordinate system so that the magnetic field points in the negative  $z$  direction,

$$\vec{B} = -B \hat{z}. \quad (9.3.1)$$

Then the Lorentz force law, eq. (4.3.12), amounts to a first-order differential equation for the velocity vector  $\vec{v} = d\vec{r}/dt$  of the electron,

$$\frac{d\vec{v}}{dt} = \omega_c \vec{v} \times \hat{z}, \quad (9.3.2)$$

or  $\ddot{x} = \omega_c \dot{y}$  and  $\ddot{y} = -\omega_c \dot{x}$  and  $\ddot{z} = 0$  in rectangular coordinates, where

$$\omega_c = \frac{eB}{mc} \quad (9.3.3)$$

is called the **cyclotron angular frequency**.<sup>†</sup> Since the force has no component along  $\hat{z}$ , the velocity in that direction is constant. The general solution for the classical position of the electron is a helix, which we take to have radius  $R$  and center  $(x_c, y_c)$  in the  $xy$  plane,

$$x(t) = x_c + R \sin(\omega_c t + \phi_0), \quad (9.3.4)$$

$$y(t) = y_c + R \cos(\omega_c t + \phi_0), \quad (9.3.5)$$

$$z(t) = z_0 + v_z t, \quad (9.3.6)$$

where  $\phi_0$ ,  $z_0$ , and  $v_z$  are the other constants needed to specify the initial conditions. It is important that the cyclotron frequency is independent of all 6 of these parameters. Since the classical motion is harmonic with angular frequency  $\omega_c$ , it should not be too surprising that we will be able to solve the corresponding quantum mechanical problem using the algebraic technology that we developed for harmonic oscillators in section 7.3, and that the energy quantum for orbital excitations for a free electron in a magnetic field will turn out to be

$$\hbar\omega_c = 1.15768 \times 10^{-4} \text{ eV} \left( \frac{B}{\text{Tesla}} \right). \quad (9.3.7)$$

This is approximately twice the product of the magnetic field and the Bohr magneton defined in eq. (4.3.26).

The expression for the Hamiltonian depends on the choice of gauge for the vector potential. One convenient choice maintains manifest rotational symmetry about the  $\hat{z}$  axis,

$$\vec{A} = \frac{1}{2}B(y\hat{x} - x\hat{y}) = -\frac{1}{2}Br\hat{\phi}, \quad (9.3.8)$$

where  $(r, \phi, z)$  are the cylindrical coordinates. From the general form of eq. (4.3.28), the total Hamiltonian operator is then

$$H_{\text{total}} = \frac{1}{2m} (\Pi_x^2 + \Pi_y^2 + P_z^2) - e\Phi(z) + \frac{g_e e}{2mc} \vec{B} \cdot \vec{S}, \quad (9.3.9)$$

where we have included spin and allowed for the possibility of a  $z$ -dependent electric potential, and the kinetic momentum operators

$$\Pi_x = P_x + \frac{eB}{2c}Y, \quad \Pi_y = P_y - \frac{eB}{2c}X \quad (9.3.10)$$

obey the commutation relation

$$[\Pi_x, \Pi_y] = i\frac{e\hbar}{c}B. \quad (9.3.11)$$

---

<sup>†</sup>For an isolated electron, the cyclotron frequency is nearly the same as the Larmor frequency  $\omega_B = (g_e/2)\omega_c$  defined in the context of spin precession in section 9.1. However, for electrons that are not isolated,  $m$  in eq. (9.3.3) should be interpreted as the effective mass of the electron moving in its environment, which can differ greatly from the nominal  $0.511 \text{ MeV}/c^2$ . This can give a very different numerical value for  $\omega_c$  in practical applications involving electrons in materials, sometimes modifying eq. (9.3.7) dramatically.

Since  $\vec{S}$  and  $P_z$  and  $\Phi(z)$  commute with  $\Pi_x$  and  $\Pi_y$ , it makes sense to separate the spin and  $z$ -dependent parts of the Hamiltonian as  $H_{\text{total}} = H + H_s + H_z$ , where our primary interest is

$$H = \frac{1}{2m} (\Pi_x^2 + \Pi_y^2), \quad (9.3.12)$$

the Hamiltonian for the 2-dimensional motion in the  $xy$  plane, and the other parts are

$$H_s = -\frac{g_e e B}{2mc} S_z, \quad H_z = \frac{P_z^2}{2m} - e\Phi(z). \quad (9.3.13)$$

The eigenvalues of  $H_s$  are just  $E_s = \mp g_e \hbar e B / 4mc$ , for spin-up states with  $S_z = \hbar/2$  and spin-down states with  $S_z = -\hbar/2$ . Thus, each spin-up state is lower than its counterpart spin-down state by an energy splitting  $g_e \hbar e B / 2mc \approx \hbar\omega_c$ . We now assume that the eigenvalues  $E_z$  of  $H_z$  have also been found. For example, if  $\Phi = 0$ , the  $H_z$  eigenstates are plane waves with momentum  $p_z$  along the  $z$ -direction and  $E_z = p_z^2 / 2m$ . In the opposite extreme, it might be that the potential  $\Phi(z)$  effectively confines the electron to the vicinity of a plane of constant  $z$ , by requiring a large energy gap for excitations along the  $z$  direction. In any case, we can concentrate on the remaining 2-dimensional problem with Hamiltonian  $H$ , to which the constants  $E_s$  and  $E_z$  can be added to find the total energy.

Since  $H$  is quadratic in two Hermitian operators with a constant commutator, we are inspired to follow the logical path of the harmonic oscillator by defining raising and lowering operators

$$a^\dagger = \frac{-i}{\sqrt{2\hbar\omega_c m}} (\Pi_x - i\Pi_y), \quad a = \frac{i}{\sqrt{2\hbar\omega_c m}} (\Pi_x + i\Pi_y). \quad (9.3.14)$$

You can check that they obey the usual ladder operator commutation relation

$$[a, a^\dagger] = 1, \quad (9.3.15)$$

and that the Hamiltonian has the simple form

$$H = \hbar\omega_c (a^\dagger a + 1/2). \quad (9.3.16)$$

So, the energy levels are the same as for a 1-dimensional harmonic oscillator with frequency  $\omega_c$ ,

$$E_n = \hbar\omega_c (n + 1/2), \quad (9.3.17)$$

where we will soon verify that  $n = 0, 1, 2, \dots$ . These are called the **Landau levels** for a charged particle in a magnetic field, after Lev Landau. The energy spacing between them grows linearly with the applied magnetic field.

At this point, one might naively suppose that the Landau level states obey  $a|n\rangle = \sqrt{n}|n-1\rangle$  and  $a^\dagger|n\rangle = \sqrt{n+1}|n+1\rangle$ . However, one must take into account the fact that in the present 2-dimensional problem the states at each level  $n$  are not unique, unlike the 1-dimensional harmonic



oscillator. In fact, each Landau level is infinitely degenerate. This corresponds to the classical property that the energy is determined by the radius  $R$  of the helix, but one can always translate it so that it is centered about any desired point in the  $xy$  plane, without changing the energy.

To understand the quantum version of this degeneracy, let us first define two operators corresponding to the center of the classical trajectory. Solving eqs. (9.3.4) and (9.3.5) for  $(x_c, y_c)$  in terms of  $x(t)$  and  $y(t)$  and their time derivatives  $\dot{x} = \pi_x/m$  and  $\dot{y} = \pi_y/m$ , and then promoting these objects to operators, we define

$$X_c = X + \frac{1}{m\omega_c}\Pi_y, \quad Y_c = Y - \frac{1}{m\omega_c}\Pi_x. \quad (9.3.18)$$

You can check that these are both constants of the motion, meaning that  $[H, X_c] = [H, Y_c] = 0$ . However, they do not commute with each other, as one finds

$$[X_c, Y_c] = -i\frac{\hbar c}{eB}. \quad (9.3.19)$$

Therefore, a CSCO can include either one of them (or any linear combination), but not both. For example, a valid CSCO choice would be  $H$  and  $X_c$ . In that case, the degeneracy of the orthobasis kets with fixed  $n$  is uncountably infinite, and corresponds to the eigenvalue of  $X_c$ .

However, let us make another choice, which we will see involves  $X_c$  and  $Y_c$  in a different way. Since the problem has symmetry under rotations about the  $z$  axis, we know (and can explicitly check) that the angular momentum component

$$L_z = XP_y - YP_x \quad (9.3.20)$$

commutes with  $H$ . Therefore, we take the CSCO to consist of  $H$  and  $L_z$ , and denote the corresponding orthonormal energy eigenstate basis elements as  $|n, \lambda\rangle$ , where

$$H|n, \lambda\rangle = \hbar\omega_c(n + 1/2)|n, \lambda\rangle, \quad (9.3.21)$$

$$L_z|n, \lambda\rangle = \hbar\lambda|n, \lambda\rangle. \quad (9.3.22)$$

From the general theory of rotations, in section 5.3, we know that  $\lambda$  must be an integer, but we will soon find that in the present problem it must satisfy the further restriction of being bounded from below, but not from above. This asymmetry is related to the sign of the magnetic field in eq. (9.3.1). In this basis, the infinite degeneracy of the Landau levels is countable, since  $L_z$  has discrete eigenvalues, unlike  $X_c$ .

From the operator definitions, you can check that

$$[H, a] = -\hbar\omega_c a, \quad [L_z, a] = \hbar a, \quad (9.3.23)$$

$$[H, a^\dagger] = \hbar\omega_c a^\dagger, \quad [L_z, a^\dagger] = -\hbar a^\dagger. \quad (9.3.24)$$

Acting on  $|n, \lambda\rangle$  with eq. (9.3.23), we find that  $a|n, \lambda\rangle$  must be an eigenstate of  $H$  and  $L_z$  with eigenvalues  $\hbar\omega_c(n-1/2)$  and  $\hbar(\lambda+1)$ , respectively. It follows that  $a$  not only lowers  $n$  by 1, but must simultaneously raise  $\lambda$  by 1, so  $a|n, \lambda\rangle = c|n-1, \lambda+1\rangle$ , for some normalization constant  $c$ . Using orthonormality of the basis kets, the norm of the right side is  $|c|^2$ , while the norm of the left side is  $\langle n, \lambda|a^\dagger a|n, \lambda\rangle = n$ , where we have made use of eq. (9.3.16) and the definition of  $n$  in eq. (9.3.21). Therefore, we must have  $c = \sqrt{n}$ , where an arbitrary phase convention has been chosen. Doing a similar computation for  $a^\dagger|n, \lambda\rangle$ , we arrive at

$$a^\dagger|n, \lambda\rangle = \sqrt{n+1}|n+1, \lambda-1\rangle, \quad (9.3.25)$$

$$a|n, \lambda\rangle = \sqrt{n}|n-1, \lambda+1\rangle. \quad (9.3.26)$$

Equation (9.3.26) shows that  $n$  must be a non-negative integer, just as in the case of the ordinary harmonic oscillator, because acting with  $a$  on any of the states  $|0, \lambda\rangle$  results in the null ket. The infinitely degenerate set of states with  $n = 0$  and varying  $\lambda$  is called the **lowest Landau level**.

Acting with  $a$  or  $a^\dagger$  changes both the energy level and the angular momentum. We will now construct raising and lowering operators for the angular momentum that leave the energy fixed. One clue to this construction is the fact, already noted, that  $X_c$  and  $Y_c$  commute with  $H$ . A second clue is that  $X_c$  and  $Y_c$  have a constant commutator with each other. We are therefore inspired to define a second pair of raising and lowering operators,

$$b^\dagger = \frac{1}{\sqrt{2}\ell_B}(X_c + iY_c), \quad b = \frac{1}{\sqrt{2}\ell_B}(X_c - iY_c), \quad (9.3.27)$$

where we have defined the **magnetic length**,

$$\ell_B = \sqrt{\frac{\hbar c}{eB}} = 2.56556 \times 10^{-8} \text{ meters} \left( \frac{\text{Tesla}}{B} \right)^{1/2}. \quad (9.3.28)$$

These operators obey the usual ladder commutation relation

$$[b, b^\dagger] = 1, \quad (9.3.29)$$

and are independent in the sense that they commute with the  $a, a^\dagger$  operators,

$$[a, b] = [a, b^\dagger] = [a^\dagger, b] = [a^\dagger, b^\dagger] = 0. \quad (9.3.30)$$

Furthermore, you are invited to check from the operator definitions that

$$b^\dagger b = \frac{1}{\hbar\omega_c}H + \frac{1}{\hbar}L_z - 1/2, \quad (9.3.31)$$

and that the counterparts of eqs. (9.3.23) and (9.3.24) are

$$[H, b] = 0, \quad [L_z, b] = -\hbar b, \quad (9.3.32)$$

$$[H, b^\dagger] = 0, \quad [L_z, b^\dagger] = \hbar b^\dagger. \quad (9.3.33)$$

With this information, it is now a short exercise, very similar to the derivation of eqs. (9.3.25) and (9.3.26), to check that (up to the usual arbitrary phase choice),

$$b^\dagger |n, \lambda\rangle = \sqrt{\lambda + n + 1} |n, \lambda + 1\rangle, \quad (9.3.34)$$

$$b |n, \lambda\rangle = \sqrt{\lambda + n} |n, \lambda - 1\rangle. \quad (9.3.35)$$

This shows how the infinite degeneracy of each Landau level  $n$  arises, as the operators  $b^\dagger$  and  $b$  change the angular momentum within each level. However, eq. (9.3.35) also shows that  $\lambda$  is restricted to be not less than  $-n$ , as lowering it further would result in the null ket.

The complete set of eigenstates of  $H$  and  $L_z$  can now be constructed, from the state with  $n = \lambda = 0$ , as

$$|n, \lambda\rangle = \frac{(b^\dagger)^{\lambda+n} (a^\dagger)^n}{\sqrt{(\lambda+n)!} \sqrt{n!}} |0, 0\rangle, \quad (9.3.36)$$

where  $n = 0, 1, 2, \dots$  is a non-negative integer, and  $\lambda = -n, -n+1, -n+2, \dots$  so that  $\lambda + n$  is also a non-negative integer. This is a complete orthobasis for the two-dimensional degrees of freedom of a charged particle in a magnetic field.

To construct the corresponding wavefunctions  $\psi_{n,\lambda}(x, y) = \langle x, y | n, \lambda \rangle$ , it is convenient to define dimensionless complex coordinates  $z, z^*$  by

$$z = \frac{x + iy}{\sqrt{2}\ell_B} = \frac{re^{i\phi}}{\sqrt{2}\ell_B}, \quad z^* = \frac{x - iy}{\sqrt{2}\ell_B} = \frac{re^{-i\phi}}{\sqrt{2}\ell_B}, \quad (9.3.37)$$

with partial derivatives defined by

$$\frac{\partial}{\partial z} = \frac{\ell_B}{\sqrt{2}} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right), \quad \frac{\partial}{\partial z^*} = \frac{\ell_B}{\sqrt{2}} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right), \quad (9.3.38)$$

so that

$$\frac{\partial z}{\partial z} = \frac{\partial z^*}{\partial z^*} = 1, \quad \frac{\partial z^*}{\partial z} = \frac{\partial z}{\partial z^*} = 0. \quad (9.3.39)$$

Then the position representations of the ladder operators are simply

$$a = \frac{\partial}{\partial z^*} + z/2, \quad a^\dagger = -\frac{\partial}{\partial z} + z^*/2, \quad (9.3.40)$$

$$b = \frac{\partial}{\partial z} + z^*/2, \quad b^\dagger = -\frac{\partial}{\partial z^*} + z/2. \quad (9.3.41)$$

Now, an eigenstate of  $L_z$  with eigenvalue  $\hbar\lambda$  must have wavefunction proportional to  $e^{i\lambda\phi}$ , as we found on general grounds in section 5.3. Therefore,  $\psi_{0,0}$  cannot depend on  $\phi$ , and so must depend on position only through the combination  $zz^*$ . Writing  $\psi_{0,0} = f(zz^*)$ , and noting that it must be annihilated by  $a$  (and also by  $b$ ), we find  $f'(zz^*) + f(zz^*)/2 = 0$ , which has the solution

$f(zz^*) = ce^{-zz^*/2}$ , with a normalization constant  $c$ . The other states in the lowest Landau level ( $n = 0$ ) are found by acting repeatedly with  $b^\dagger$ . Applying eq. (9.3.41)  $\lambda$  times, this gives the elegant result

$$\psi_{0,\lambda} = \frac{c}{\sqrt{\lambda!}} z^\lambda e^{-zz^*/2}. \quad (9.3.42)$$

Rewriting this in cylindrical coordinates, and requiring  $\int_0^{2\pi} d\phi \int_0^\infty dr r |\psi_{0,0}|^2 = 1$  to fix  $c$ , we obtain the normalized wavefunctions

$$\psi_{0,\lambda} = \frac{1}{\sqrt{2\pi\lambda!} \ell_B} \left( \frac{r e^{i\phi}}{\sqrt{2}\ell_B} \right)^\lambda \exp(-r^2/4\ell_B^2). \quad (9.3.43)$$

The wavefunctions for the higher Landau levels are more complicated, but can now be obtained by acting with  $(a^\dagger)^n/\sqrt{n!}$ .

The wavefunction  $\psi_{0,0}$  is sharply peaked at the origin  $r = 0$ , which is arbitrarily chosen. For nonzero  $\lambda$ , the probability density  $|\psi_{0,\lambda}|^2$  instead has a ring-like profile, vanishing at the origin, and maximum at  $r = R_\lambda = \sqrt{2\lambda}\ell_B$ . For very large angular momentum  $\lambda$ , the radius of the ring  $R_\lambda$  is much larger than its width (defined as the range of  $r$  near  $R_\lambda$  for which the wavefunction is appreciable), which is always comparable to  $\ell_B$ , independent of  $\lambda$ .

If we now suppose that the electron is confined to a disk of radius  $R$  centered at the origin in the  $xy$  plane, then the infinite degeneracy of each Landau level is reduced to a finite degeneracy. Intuitively, valid states must have  $R_\lambda \lesssim R$ , implying  $\lambda \lesssim R^2/2\ell_B^2$ , in order for the wavefunction ring to “fit” inside the disk. This is not exact, because confining the electrons to  $r < R$  introduces a boundary condition that modifies the stationary state wavefunctions in a complicated way. However, in a macroscopic disk with  $R \gg \ell_B$ , only the tiny fraction of states with  $R - R_\lambda$  comparable to  $\ell_B$  will be distorted significantly. For the purposes of counting the number of states in a macroscopic area, this makes almost no difference. So, we estimate that the number  $N$  of states in the lowest Landau level should be (one more than) the maximum allowed  $\lambda$ ,

$$N \approx \lambda_{\max} \approx \frac{R^2}{2\ell_B^2} = \frac{e\Phi_m}{2\pi\hbar c} = \frac{\Phi_m}{\Phi_0}, \quad (9.3.44)$$

where  $\Phi_m = \pi R^2 B$  is the magnetic flux through the disk, and

$$\Phi_0 = 2\pi\hbar c/e = 4.13567 \times 10^{-11} \text{ Tesla} \cdot \text{cm}^2 \quad (9.3.45)$$

is the natural quantum unit<sup>‡</sup> of magnetic flux. Thus, by varying the applied magnetic field, one can dial the number of electrons that can fit in the lowest Landau level, and in each higher energy

---

<sup>‡</sup>In the context of superconductivity, a conventional definition of  $\Phi_0$  is used that is half as big, corresponding to a (Cooper pair of electrons) charge that is twice as large.

level. This affects the conductivity properties of materials, and in particular is an important ingredient in the quantum Hall effect.

It is instructive to treat the same problem in a different gauge. The magnetic field  $\vec{B} = -B\hat{z}$  can also be obtained as  $\vec{\nabla} \times \vec{A}$  where we now choose

$$\vec{A} = -Bx\hat{y}. \quad (9.3.46)$$

This vector potential is invariant under translations in the  $y$  direction, rather than maintaining manifest invariance under rotations about the  $z$  axis. The Hamiltonian in this gauge is

$$H = \frac{1}{2m}P_x^2 + \frac{1}{2m}\left(P_y - \frac{eB}{c}X\right)^2. \quad (9.3.47)$$

Since this version of  $H$  commutes with  $P_y$  due to our gauge choice, we use them as our CSCO, and look for an orthobasis of common eigenstates  $|n, k\rangle$  that satisfy

$$H|n, k\rangle = \hbar\omega_c(n + 1/2)|n, k\rangle, \quad (n = 0, 1, 2, \dots), \quad (9.3.48)$$

$$P_y|n, k\rangle = \hbar k|n, k\rangle, \quad (k = \text{real}). \quad (9.3.49)$$

Here we have used the fact that we already know the energy eigenvalues, since they are measurable and so cannot depend on our choice of gauge. The infinite degeneracy in the Landau levels is now represented by the freedom to choose any  $P_y$  eigenvalue  $\hbar k$ , independently of  $n$ . Substituting  $P_y = \hbar k$ , and defining

$$x_0 = \hbar k / m\omega_c = k\ell_B^2, \quad (9.3.50)$$

the Hamiltonian becomes

$$H = \frac{1}{2m}P_x^2 + \frac{1}{2}m\omega_c^2(X - x_0)^2. \quad (9.3.51)$$

This is simply a 1-dimensional harmonic oscillator with angular frequency  $\omega_c$  describing motion in the  $x$  direction, and displaced by the constant distance  $x_0$  depending on the  $y$  momentum eigenvalue. The orthobasis position representation wavefunctions are, up to normalization,

$$\psi_{n,k}(x, y) = \langle x, y | n, k \rangle = \psi_n(x - x_0)e^{iky}, \quad (9.3.52)$$

where  $\psi_n(x)$  are the harmonic oscillator wavefunctions of section 7.2, with  $\omega = \omega_c$  and length scale  $b = \ell_B$ . Thus our new choice of gauge achieves a clean separation of  $x$  and  $y$  wavefunctions. The more general lesson is that different gauge choices are useful for highlighting different aspects of a problem.

Suppose now that our electron is confined to a rectangular area,  $0 < x < d_x$  and  $0 < y < d_y$ , with macroscopic lengths  $d_x$  and  $d_y$ . We could try to impose a boundary condition that the

wavefunction vanishes outside of this rectangle, but then it would become difficult to solve the  $H, P_y$  eigenvalue equations exactly. Fortunately, for many purposes we can still use the solutions found above, to a good approximation. For the vast majority of states in the lowest Landau levels, the boundary conditions at  $x = 0$  and  $x = d_x$  have almost no effect because the support of the wavefunctions is limited to a distance comparable to  $\ell_B$ . For example, the lowest Landau level ( $n = 0$ ) states have wavefunctions (up to normalization)

$$\psi_{0,k}(x, y) = \exp[-(x - x_0)^2/2\ell_B^2] e^{iky}. \quad (9.3.53)$$

Because of the exponential suppression, the distortion of this wavefunction resulting from the existence of the  $x = 0$  and  $x = d_x$  boundaries is very small except for the tiny fraction of states for which  $x_0$  or  $d_x - x_0$  is comparable to  $\ell_B$ . This argument does not apply for the  $y = 0$  and  $y = d_y$  boundaries, because the wavefunction factor  $e^{iky}$  has constant magnitude. However, we can use the trick of imposing periodic boundary conditions in the  $y$  direction, so that  $y = 0$  is identified with  $y = d_y$ . Since  $d_y$  is a macroscopic length, the microscopic physics should not depend very much on whether we identify the two distant sides, or more generally on what boundary conditions we impose, as long as they are consistent. The point is that imposing periodic boundary conditions in  $y$  is particularly easy to do in a consistent way for the wavefunctions proportional to  $e^{iky}$  (and in particular is much easier than trying to impose  $\psi = 0$  at  $y = 0$  and  $y = d_y$ ). It just restricts the allowed eigenvalues to

$$k = 2\pi n_y/d_y, \quad (n_y = \text{integer}). \quad (9.3.54)$$

We can now repeat the estimate of the number of states per unit area in the lowest Landau level. To a good approximation, the valid states will be the ones for which  $x_0 = \ell_B^2 2\pi n_y/d_y$  fits between 0 and  $d_x$ . This means that the allowed  $n_y$  are the integers in the range  $0 < n_y < d_x d_y / 2\pi \ell_B^2$ , resulting in a degeneracy for the lowest Landau level of

$$N \approx \frac{eB d_x d_y}{2\pi \hbar c} = \frac{\Phi_m}{\Phi_0}. \quad (9.3.55)$$

This agrees exactly with our previous result of eq. (9.3.44) obtained using the rotationally symmetric gauge choice corresponding to the  $H, L_z$  orthobasis.

## 9.4 Exercises

**Exercise 9.1.** Show that the spin expectation value precession result of eq. (9.1.22) obeys Ehrenfest's Theorem, eq. (3.4.16), with  $\vec{S}$  playing the role of  $A$ .

**Exercise 9.2.** A spin-1/2 particle with gyromagnetic ratio  $\gamma$  is in a uniform but time-varying magnetic field  $\vec{B} = B(t)\hat{z}$  so that the Hamiltonian is  $H = -\gamma \vec{B} \cdot \vec{S}$ . At time  $t = 0$ , the

spin of the particle along the direction  $\hat{n} = (\hat{x} + \hat{z})/\sqrt{2}$  is measured to be  $+\hbar/2$ . You should express your answers in terms of the definite integral  $I(t) = \int_0^t dt' B(t')$ . [Hint: in this problem the Hamiltonian is not constant in time, so you need to solve a differential equation for the components of the state vector. That's where the definite integral comes in.]

- (a) If the  $\hat{z}$  component of the spin is measured at time  $t > 0$ , what is the probability that the result will be  $S_z = -\hbar/2$ ?
- (b) If, instead, the  $\hat{y}$  component of the spin is measured at time  $t > 0$ , what is the probability that the result will be  $S_y = +\hbar/2$ ?
- (c) Find the expectation value of  $S_x$  as a function of time  $t > 0$ .

**Exercise 9.3.** For the Rabi oscillation solution of eqs. (9.2.14) and (9.2.15), find the expectation value of the spin operator,  $\langle \vec{S} \rangle$ , as a function of time. Simplify the result as much as possible, and show that it obeys Ehrenfest's Theorem, eq. (3.4.16) with  $\vec{S}$  playing the role of  $A$ .

**Exercise 9.4.** Consider the magnetic spin resonance system treated in section 9.2, but now assume that the system is initially in an eigenstate of  $S_x$  with eigenvalue  $\hbar/2$ , so that the boundary condition for the differential equations (9.2.8) and (9.2.9) is  $a_1(0) = a_2(0) = 1/\sqrt{2}$ . Find the Rabi oscillation solution for  $a_1(t)$  and  $a_2(t)$  with the same  $\Omega^2 = [\Gamma^2 + (\omega - \omega_B)^2]/4$ . Show that the probability to find the system in the state  $|\uparrow\rangle$  is  $\mathcal{P}(\uparrow) = 1/2 + n \sin^2(\Omega t)$ , where  $n$  is a quantity that you will find. What happens to this probability at resonance ( $\omega = \omega_B$ )? Under what conditions can  $|n| = 1/2$ , so that the probability periodically reaches 0 and 1?

**Exercise 9.5.** Prove each of eqs. (9.3.29)-(9.3.35) from the operator definitions given earlier.

**Exercise 9.6.** Find the  $n = 1$  and  $n = 2$  Landau level state wavefunctions  $\psi_{1,\lambda}$  and  $\psi_{2,\lambda}$  in the  $H, L_z$  eigenstate basis. Write your answers in terms of  $z$  and  $z^*$  defined in eq. (9.3.37).

**Exercise 9.7.** Consider an electron moving in the  $xy$  plane in the presence of uniform constant fields  $\vec{B} = -B\hat{z}$  and  $\vec{E} = E\hat{x}$ . Use the same gauge choice as in eqs. (9.3.46) and (9.3.47). The presence of the electric field just adds a term  $eEX$  to the Hamiltonian.

- (a) Show that there are stationary states with energies  $E_{n,k} = \hbar\omega_c(n + 1/2) + c_1 kE/B + c_2 E^2/B^2$ , where  $c_1$  and  $c_2$  are constants that you will determine, with corresponding wavefunctions  $\psi_{n,k}(x, y) = \psi_n(x - x'_0)e^{iky}$  where  $\psi_n$  are the standard 1-d harmonic oscillator wavefunctions, and  $x'_0$  is another constant that you will find.
- (b) A wavepacket superposition of states with nearly the same  $k$  will have a velocity in the  $y$  direction, with dispersion because the energy depends on the wavenumber. Find the group velocity  $v_g = (1/\hbar)\partial E_{n,k}/\partial k$ . (This is equal to the time-averaged velocity for the cycloid motion of the corresponding classical problem.)

## 10 Examples with spherical symmetry

### 10.1 Stationary states with spherical symmetry

Consider a Hamiltonian of the form

$$H = \frac{P^2}{2\mu} + V(R). \quad (10.1.1)$$

In this chapter, we will write  $\mu$  for the mass of the particle, to avoid confusion with the eigenvalue  $\hbar m$  of  $L_z$ , and  $R$  is the operator corresponding to the classical spherical coordinate  $r$ . Thus  $V(R)$  is a spherically symmetric potential, with no dependence on  $\theta$  or  $\phi$ . Likewise, the momentum squared operator

$$P^2 = P_x^2 + P_y^2 + P_z^2 = -\hbar^2 \nabla^2 \quad (10.1.2)$$

is rotationally invariant, so there is no preferred direction associated with the Hamiltonian. It follows that  $H$ ,  $L^2$ , and  $L_z$  are compatible observables, and we can look for an orthobasis of common eigenstates

$$|E, l, m\rangle, \quad (10.1.3)$$

where  $E$  is the energy of the state, the eigenvalue of the Hamiltonian. Because of eq. (8.7.5), these are also parity eigenstates, with eigenvalue  $(-1)^l$ .

In the following, we will work in the position wavefunction representation with

$$\psi_{E,l,m}(r, \theta, \phi) = \langle r, \theta, \phi | E, l, m \rangle = R_{E,l}(r) Y_l^m(\theta, \phi), \quad (10.1.4)$$

where  $R_{E,l}(r)$  is a radial wavefunction. Note that  $R_{E,l}(r)$  will depend on both  $E$  and  $l$  in general, but it will not depend on  $m$ , because the operators  $L_+$  and  $L_-$  raise and lower  $m$  without changing  $l$  or the radial dependence of the wavefunction. The time-independent Schrödinger equation in this basis is

$$\left[ -\frac{\hbar^2 \nabla^2}{2\mu} + V(r) \right] \psi_{E,l,m}(r, \theta, \phi) = E \psi_{E,l,m}(r, \theta, \phi). \quad (10.1.5)$$

Now, using eq. (8.6.12), this becomes

$$-\frac{\hbar^2}{2\mu} \left[ \frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} - \frac{l(l+1)}{r^2} \right] R_{E,l}(r) + V(r) R_{E,l}(r) = E R_{E,l}(r), \quad (10.1.6)$$

where the spherical harmonics have been factored out.

In many cases, we will find that the radial wavefunction  $R_{E,l}(r)$  has a power-law behavior  $r^p$  near the origin, for an integer  $p$ . By requiring the total probability to be finite, it is clear that



$p$  cannot be less than  $-1$ ; otherwise, the normalization integral  $\int dr r^2 |R_{E,l}(r)|^2$  would diverge even for a finite volume. But in almost all cases, one can make the even stronger statement that  $R_{E,l}(r)$  must be finite as  $r \rightarrow 0$ . This is because  $\nabla^2(1/r) = -4\pi\delta(\vec{r})$ , so that  $p = -1$  would necessarily imply the presence of a delta function contribution to the potential  $V(r)$ . In the examples to follow, we will often make use of the finiteness of the radial wavefunction.

It is often useful to define a function

$$U_{E,l}(r) = rR_{E,l}(r), \quad (10.1.7)$$

in which one power of the radial coordinate has been factored out. The eigenvalue differential equation then becomes

$$\left[ -\frac{\hbar^2}{2\mu} \frac{d^2}{dr^2} + \frac{\hbar^2 l(l+1)}{2\mu r^2} + V(r) \right] U_{E,l} = EU_{E,l}, \quad (10.1.8)$$

with the nice feature that there is no term with a single  $r$  derivative; that is the reason for sometimes using  $U_{E,l}$  rather than  $R_{E,l}$ .

Indeed, eq. (10.1.8) for  $U_{E,l}$  is very similar to the one-dimensional time-independent Schrödinger equation (6.3.3) with  $x$  replaced by  $r$ . However, there are two important differences. First, the domain of the independent variable is now limited to non-negative values,

$$0 \leq r < \infty, \quad (10.1.9)$$

unlike the one-dimensional Schrödinger equation. Second, the potential has effectively been modified to

$$V_{\text{eff}}(r) = V(r) + \frac{\hbar^2 l(l+1)}{2\mu r^2}, \quad (10.1.10)$$

where the second term is a repulsive centrifugal contribution that blows up at  $r = 0$ , and therefore makes the wavefunction vanish there unless  $l = 0$ . In fact, if  $V(r)$  is finite at  $r = 0$ , one finds from eq. (10.1.8) that for  $l \neq 0$ ,  $R_{E,l}$  must scale like  $r^l$  at small  $r$ .

With these caveats, the problem of stationary states for a particle in a spherically symmetric potential has been reduced to a particular type of one-dimensional problem. The same results that we derived in section 6.3 [see the discussion surrounding eqs. (6.3.6)–(6.3.8)] for matching wavefunctions at special points holds here as for the one-dimensional problems. In particular, the radial wavefunction is always continuous, and if the potential is finite (not necessarily continuous) at a point  $r = r_0$ , then its first derivative with respect to  $r$  is continuous there.

## 10.2 Free particle in spherical coordinates

We have already discussed the position and momentum eigenstates and wavefunctions for a free particle moving in three dimensions, in section 2.8. For position eigenstates, the CSCO used was

$(X, Y, Z)$ , while for momentum eigenstates, it was  $(P_x, P_y, P_z)$ . Now we will discuss the same physical problem using the CSCO  $(H, L^2, L_z)$  with eigenvalues  $(E, \hbar^2 l(l+1), \hbar m)$ , starting from eq. (10.1.8) with  $V(r) = 0$ . We will solve the time-independent Schrödinger equation for fixed  $(E, l, m)$  in the position wavefunction representation with coordinates  $(r, \theta, \phi)$ , where the domain may consist of only a sub-volume of the whole space. The resulting solutions can then be applied, using matching of the wavefunction, to problems in which a particle moves freely only in that sub-volume. They can also be applied to scattering problems in three dimensions.

To simplify things, define

$$k^2 = 2\mu E/\hbar^2, \quad \rho = kr. \quad (10.2.1)$$

From the discussion in section 2.8, we already know that the allowed eigenvalues  $E$  are positive, so that  $k$  is a real number with units of  $1/[\text{length}]$ . Thus  $\rho$  is a radial coordinate re-scaled to make it dimensionless. In terms of these, eq. (10.1.8) becomes

$$\left( \frac{d^2}{d\rho^2} + 1 - \frac{l(l+1)}{\rho^2} \right) U_l(\rho) = 0. \quad (10.2.2)$$

Note that this equation and the forms of its solutions  $U_l(\rho)$  do not depend on  $E$ , because the dependence has been hidden in  $\rho$ .

Let us start with the case  $l = 0$ . Then eq. (10.2.2) is a familiar equation, with the familiar solutions  $U_0(\rho) = \sin(\rho)$  or  $\cos(\rho)$ . Of these, the first solution  $U_0 = \sin(\rho)$  gives a constant for  $R(r)$  as  $r \rightarrow 0$ . However, the second solution  $U_0 = \cos(\rho)$  approaches a constant as  $r \rightarrow 0$ . This renders it physically unacceptable if the origin is included in the spatial region under consideration, because then  $R(r) \sim 1/r$ , which is divergent and would require a delta function potential  $V(r)$  at the origin, since  $\nabla^2(1/r) = -4\pi\delta(\vec{r})$ . Nevertheless, it is acceptable if we are solving for the free-particle wavefunction only in a region that does not include the origin. We therefore have two solutions for  $L^2 = 0$ , labeled  $A$  and  $B$ ,

$$U_0^A(kr) = \sin(kr), \quad U_0^B(kr) = -\cos(kr), \quad (10.2.3)$$

where the minus sign is a phase choice for later convenience, and the  $B$  solution is understood to be acceptable if, and only if, the origin is excluded. These solutions for  $l = 0$  can be used as seeds to find solutions for  $l > 0$ , as we will now see.

Writing the solution for general  $l$  in the form

$$U_l(\rho) = \rho^{l+1} f_l(\rho), \quad (10.2.4)$$

the differential equation (10.2.2) becomes

$$\left( \frac{d^2}{d\rho^2} + \frac{2(l+1)}{\rho} \frac{d}{d\rho} + 1 \right) f_l(\rho) = 0. \quad (10.2.5)$$

Now we note that if  $f_l(\rho)$  satisfies the differential equation for  $l$ , then

$$f_{l+1}(\rho) \equiv -\frac{1}{\rho} \frac{df_l}{d\rho} \quad (10.2.6)$$

will satisfy it for  $l+1$ , as can be proved by computing  $d^2 f_{l+1}/d\rho^2$  and  $df_{l+1}/d\rho$  and plugging into eq. (10.2.5). So, for each of our two  $l=0$  solutions from eq. (10.2.3),

$$f_0^A(\rho) = \frac{\sin \rho}{\rho}, \quad f_0^B(\rho) = -\frac{\cos \rho}{\rho}, \quad (10.2.7)$$

by recursion we will have a solution

$$f_l = \left( -\frac{1}{\rho} \frac{d}{d\rho} \right)^l f_0. \quad (10.2.8)$$

Therefore, for each value of  $l$ , we have two linearly independent solutions

$$R_l^A(\rho) = \rho^l f_l^A = \rho^l \left( -\frac{1}{\rho} \frac{d}{d\rho} \right)^l \left( \frac{\sin \rho}{\rho} \right) \equiv j_l(\rho), \quad (10.2.9)$$

$$R_l^B(\rho) = \rho^l f_l^B = \rho^l \left( -\frac{1}{\rho} \frac{d}{d\rho} \right)^l \left( -\frac{\cos \rho}{\rho} \right) \equiv n_l(\rho). \quad (10.2.10)$$

The functions  $j_l(\rho)$  are called the **spherical Bessel functions**, and  $n_l(\rho)$  are called the **spherical Neumann functions**. The lowest few are

$$j_0(\rho) = \frac{\sin \rho}{\rho}, \quad n_0(\rho) = -\frac{\cos \rho}{\rho}, \quad (10.2.11)$$

$$j_1(\rho) = \frac{\sin \rho}{\rho^2} - \frac{\cos \rho}{\rho}, \quad n_1(\rho) = -\frac{\cos \rho}{\rho^2} - \frac{\sin \rho}{\rho}, \quad (10.2.12)$$

$$j_2(\rho) = \left( \frac{3}{\rho^3} - \frac{1}{\rho} \right) \sin \rho - \frac{3}{\rho^2} \cos \rho, \quad n_2(\rho) = \left( -\frac{3}{\rho^3} + \frac{1}{\rho} \right) \cos \rho - \frac{3}{\rho^2} \sin \rho. \quad (10.2.13)$$

For small  $\rho$ , they can be shown to behave like

$$j_l(\rho) \sim \frac{\rho^l}{(2l+1)!!}, \quad (10.2.14)$$

$$n_l(\rho) \sim -\frac{(2l-1)!!}{\rho^{l+1}}, \quad (10.2.15)$$

where the double factorial notation means

$$(2l+1)!! = (2l+1)(2l-1) \cdots (1) = (2l+1)!/(2^l l!), \quad (10.2.16)$$

with the special value  $(-1)!! = 1$ . Thus the  $j_l$  solutions are well-behaved at the origin, but the  $n_l$  solutions are not. For large  $\rho$ , they both fall off like  $1/\rho$  multiplied by oscillating functions,

$$j_l(\rho) \sim \frac{1}{\rho} \sin(\rho - l\pi/2), \quad (10.2.17)$$

$$n_l(\rho) \sim -\frac{1}{\rho} \cos(\rho - l\pi/2). \quad (10.2.18)$$

The functions  $n_l(\rho)$  are not needed for the case of a completely free particle (one that has no potential everywhere including  $r = 0$ ). However, they are useful in situations where we piece together the free-particle solution(s) in region(s) not including the origin to other solutions that do include the origin, as for example in section 10.4.

For example, suppose we have a potential that is spherically symmetric and piece-wise constant within radial intervals, so that

$$V(r) = \begin{cases} V_0 & (0 \leq r < a_1), \\ V_1 & (a_1 < r < a_2), \\ \dots & \dots \\ V_N & (a_N < r < \infty). \end{cases} \quad (10.2.19)$$

In each of these regions, the constant potential  $V_n$  can be absorbed into the constant energy, so the possible stationary-state solution wavefunctions are

$$\left( A_l^{(n)} j_l(k_n r) + B_l^{(n)} n_l(k_n r) \right) Y_l^m(\theta, \phi), \quad \text{for } (a_n < r < a_{n+1}), \quad (10.2.20)$$

for some constants  $A_l^{(n)}$  and  $B_l^{(n)}$ , but now with

$$k_n = \sqrt{2\mu(E - V_n)}/\hbar. \quad (10.2.21)$$

The coefficients  $A_l^{(n)}$  and  $B_l^{(n)}$  can be determined by matching the wavefunctions, and their first derivative with respect to  $r$ , at each of the points  $a_n$ . This procedure also simultaneously determines the energy eigenvalue  $E$ . In the region  $0 \leq r < a_1$ , the Neumann function solutions are not allowed, so one must have  $B_l^{(0)} = 0$ , but in all other regions  $B_l^{(n)}$  is allowed to be non-zero. Note that there is a degeneracy  $2l + 1$  for each of these stationary states, due to the fact that the energy does not depend on  $m = -l, \dots, l$ . One can also match such wavefunctions to regions in which the potential is something more complicated.

Different linear combinations of  $j_l$  and  $n_l$  are useful in certain kinds of problems. The **spherical Hankel functions** defined by

$$h_l^{(1)}(\rho) = j_l(\rho) + in_l(\rho) = -i\rho^l \left( -\frac{1}{\rho} \frac{d}{d\rho} \right)^l \left( \frac{e^{i\rho}}{\rho} \right), \quad (10.2.22)$$

$$h_l^{(2)}(\rho) = j_l(\rho) - in_l(\rho) = i\rho^l \left( -\frac{1}{\rho} \frac{d}{d\rho} \right)^l \left( \frac{e^{-i\rho}}{\rho} \right), \quad (10.2.23)$$

correspond to outgoing and ingoing spherical waves, respectively. To understand this, note that from eqs. (10.2.17) and (10.2.18) their behavior for large  $\rho = kr$  is

$$h_l^{(1)}(\rho) \approx \frac{1}{\rho} e^{i[\rho - \pi(l+1)/2]} = i^{-l} e^{i\rho}/\rho, \quad (10.2.24)$$

$$h_l^{(2)}(\rho) \approx \frac{1}{\rho} e^{-i[\rho + \pi(l+1)/2]} = i^l e^{-i\rho}/\rho, \quad (10.2.25)$$

so that the time dependence of the corresponding wavefunctions is

$$\langle \vec{r} | \psi^{(1)}(t) \rangle \propto e^{-itE/\hbar} e^{ikr}/r, \quad (10.2.26)$$

$$\langle \vec{r} | \psi^{(2)}(t) \rangle \propto e^{-itE/\hbar} e^{-ikr}/r. \quad (10.2.27)$$

These maintain constant phase at  $r = (E/\hbar k)t + \text{constant}$  and  $r = -(E/\hbar k)t + \text{constant}$ , respectively. (The phase velocity is therefore  $\omega/k = E/p = \hbar k/2m$ , but recall from the discussion leading to eq. (6.2.11) that wavepacket superpositions will have a group velocity twice this,  $\hbar k/m$ , which is the classical value.)

For problems in which  $E < V_n$  in at least one region,  $k$  will be imaginary in that region. Then the solutions will involve  $j_l(kr)$  and  $n_l(kr)$  or  $h_l^{(1)}(kr)$  and  $h_l^{(2)}(kr)$ , where now  $k = i\sqrt{2\mu(V_n - E)}/\hbar$  is a pure imaginary number. In many such cases, a more convenient basis of solutions is provided by the **modified spherical Bessel functions**,

$$i_l(\rho) = j_l(i\rho)/i^l = \rho^l \left( \frac{1}{\rho} \frac{d}{d\rho} \right)^l \left( \frac{\sinh \rho}{\rho} \right), \quad (10.2.28)$$

$$k_l(\rho) = -i^l h_l^{(1)}(i\rho) = \rho^l \left( -\frac{1}{\rho} \frac{d}{d\rho} \right)^l \left( \frac{e^{-\rho}}{\rho} \right), \quad (10.2.29)$$

where now

$$\rho = \kappa r, \quad \kappa = k/i = \sqrt{2\mu(V_n - E)}/\hbar. \quad (10.2.30)$$

The function  $i_l(\rho)$  is well-defined and useful for regions that include  $r = 0$ , while the function  $k_l(\rho)$  is useful for regions that extend to  $r = \infty$ .

For the rest of this section, we consider the  $(H, L^2, L_z)$  orthobasis eigenstate wavefunctions for the completely free particle whose domain includes all space, including the origin. They are

$$\langle \vec{r} | k, l, m \rangle = \psi_{k,l,m}(\vec{r}) = R_{k,l}(r) Y_l^m(\theta, \phi), \quad (E = \hbar^2 k^2 / 2\mu), \quad (10.2.31)$$

where the radial wavefunctions  $R_{k,l}(r) = A_{k,l} j_l(kr)$  contain a normalization constant  $A_{k,l}$  to be fixed. To do so, we can use the orthonormality relations for the spherical Bessel functions,<sup>†</sup>

$$\int_0^\infty dr r^2 j_l(kr) j_l(k'r) = \frac{\pi}{2k^2} \delta(k - k'), \quad (10.2.32)$$

for each  $l$ . It follows that if we choose  $A_{k,l} = \sqrt{2/\pi}$ , so that the radial wavefunctions are

$$R_{k,l}(r) = \sqrt{\frac{2}{\pi}} j_l(kr), \quad (10.2.33)$$

---

<sup>†</sup>Equation 10.2.32 is technically ill-defined, because the integration does not converge, even for  $k \neq k'$ . It should therefore be viewed as a formal relation, to be used within expressions where  $k$  or  $k'$  is integrated over. This is similar to the technically ill-defined nature of the Fourier integral for the delta function in eq. (2.2.20).

then we have the orthonormality relations,

$$\int dr r^2 R_{k',l}(r) R_{k,l}(r) = \frac{1}{k^2} \delta(k - k'), \quad (10.2.34)$$

and for the full eigenstates and their wavefunctions,

$$\langle k', l', m' | k, l, m \rangle = \int d^3 \vec{r} \psi_{k',l',m'}(\vec{r})^* \psi_{k,l,m}(\vec{r}) = \frac{1}{k^2} \delta(k - k') \delta_{l'l'} \delta_{mm'}. \quad (10.2.35)$$

With our choice of normalization in eqs. (10.2.33)–(10.2.35), we also have the completeness of the orthobasis in terms of an integral over  $k$ ,

$$\int_0^\infty dk k^2 \sum_{l=0}^\infty \sum_{m=-l}^l |k, l, m\rangle \langle k, l, m| = I, \quad (10.2.36)$$

from which follows

$$\int_0^\infty dk k^2 \sum_{l=0}^\infty \sum_{m=-l}^l \psi_{k,l,m}(\vec{r}')^* \psi_{k,l,m}(\vec{r}) = \delta^{(3)}(\vec{r} - \vec{r}'). \quad (10.2.37)$$

A perfectly good alternative normalization choice would be to express the same results in terms of “energy normalization” kets

$$|E, l, m\rangle = \frac{\sqrt{\mu k}}{\hbar} |k, l, m\rangle, \quad (10.2.38)$$

and the corresponding radial wavefunctions

$$R_{E,l}(r) = \sqrt{\frac{2\mu k}{\pi \hbar^2}} j_l(kr). \quad (10.2.39)$$

If one uses  $|E, l, m\rangle$ ,  $R_{E,l}(r)$ , and  $\psi_{E,l,m}(\vec{r})$ , then the Dirac orthonormality and completeness relations will have the same form as eqs. (10.2.34)–(10.2.37), but with  $\delta(E - E')$  replacing  $\delta(k - k')/k^2$  and  $\int_0^\infty dE$  replacing  $\int_0^\infty dk k^2$ . This follows from  $\delta(k - k') = \delta(E - E') \hbar^2 k / \mu$ .

Let us now consider the relation between the plane wavefunctions associated with momentum eigenstates  $|\vec{p}\rangle$  and the spherical waves associated with the  $(H, L^2, L_z)$  eigenstates  $|k, l, m\rangle$ . Using the completeness relation, we have

$$\langle \vec{r} | \vec{p} \rangle = \int_0^\infty dk k^2 \sum_{l=0}^\infty \sum_{m=-l}^l \langle \vec{r} | k, l, m \rangle \langle k, l, m | \vec{p} \rangle. \quad (10.2.40)$$

Now, both  $|\vec{p}\rangle$  and  $|k, l, m\rangle$  are eigenstates of  $H$  with energy eigenvalues  $E = |\vec{p}|^2/2\mu$  and  $\hbar^2 k^2/2\mu$ , respectively. Therefore, Theorem 2.6.5 says that the last inner product in eq. (10.2.40) must vanish unless  $|\vec{p}| = \hbar k$ , and so it must be of the form

$$\langle k, l, m | \vec{p} \rangle = \frac{1}{(2\pi \hbar)^{3/2}} \delta(k - |\vec{p}|/\hbar) \sqrt{\frac{\pi}{2}} C_{l,m}(\theta_{\vec{p}}, \phi_{\vec{p}}), \quad (10.2.41)$$

for some functions  $C_{l,m}$  that depend only on  $l$ ,  $m$ , and the angular coordinates of the vector  $\vec{p}$ . (The  $C_{l,m}$  cannot depend on the magnitude  $k$ , because they are dimensionless and there is no other dimensionful quantity on which they could depend.) Since  $k = |\vec{p}|/\hbar$  is enforced by the delta function, we can define  $\vec{k} = \vec{p}/\hbar$  and use eqs. (2.8.62) and (10.2.31) to find that eq. (10.2.40) reads

$$\frac{e^{i\vec{k}\cdot\vec{r}}}{(2\pi\hbar)^{3/2}} = \frac{1}{(2\pi\hbar)^{3/2}} \sum_{l=0}^{\infty} \sum_{m=-l}^l C_{l,m}(\theta_{\vec{k}}, \phi_{\vec{k}}) j_l(kr) Y_l^m(\theta, \phi). \quad (10.2.42)$$

The claim is that the coefficient functions turn out to be

$$C_{l,m}(\theta_{\vec{k}}, \phi_{\vec{k}}) = 4\pi i^l Y_l^m(\theta_{\vec{k}}, \phi_{\vec{k}})^*, \quad (10.2.43)$$

so that we have the remarkable identity relating plane waves to spherical waves,

$$e^{i\vec{k}\cdot\vec{r}} = \sum_{l=0}^{\infty} \sum_{m=-l}^l 4\pi i^l Y_l^m(\theta_{\vec{k}}, \phi_{\vec{k}})^* j_l(kr) Y_l^m(\theta, \phi). \quad (10.2.44)$$

Note that  $(\theta_{\vec{k}}, \phi_{\vec{k}})$  are the spherical coordinate angles for the vector  $\vec{k}$ , while  $(\theta, \phi)$  are the angles for the vector  $\vec{r}$  in the same coordinate system. The interpretation of eq. (10.2.44) is that a plane wave with momentum  $\vec{p} = \hbar\vec{k}$  consists of a superposition of spherical waves with all allowed (quantized) values of orbital angular momentum.

To prove the claimed eq. (10.2.44), we first consider the case that  $\vec{k} = k\hat{z}$ . Then  $\theta_{\vec{k}} = 0$ , and  $\vec{k} \cdot \vec{r} = kr \cos \theta$ , so  $e^{i\vec{k}\cdot\vec{r}} = e^{ikr \cos \theta}$ . Because this does not depend on  $\phi$  at all, its expansion in terms of spherical harmonics will only include the  $m = 0$  functions  $Y_l^0(\theta, \phi)$ , which are proportional to the ordinary Legendre polynomials  $P_l(\cos \theta)$ . So, in this case the expansion in eq. (10.2.44) has the simpler form

$$e^{ikr \cos \theta} = \sum_{l=0}^{\infty} c_l j_l(kr) P_l(\cos \theta), \quad (10.2.45)$$

where  $c_l$  are some coefficients that we need to determine. Now we can multiply both sides of eq. (10.2.45) by  $P_{l'}(\cos \theta)$ , and integrate with respect to  $\cos \theta$ , making use of the identity

$$\int_{-1}^1 du P_l(u) P_{l'}(u) = \frac{2}{2l+1} \delta_{ll'}. \quad (10.2.46)$$

The result, after relabeling  $l' \rightarrow l$ , is

$$c_l j_l(kr) = \frac{2l+1}{2} \int_{-1}^1 d(\cos \theta) P_l(\cos \theta) e^{ikr \cos \theta}. \quad (10.2.47)$$

This must hold for all  $r$ , so we can consider the leading behavior as  $r \rightarrow 0$  in particular. Using eq. (10.2.14), this gives

$$c_l \frac{(kr)^l}{(2l+1)!!} + \mathcal{O}(r^{l+1}) = \frac{2l+1}{2} \int_{-1}^1 du P_l(u) e^{ikru}. \quad (10.2.48)$$

This shows that the right side evidently must be proportional to  $r^l$  in the limit of small  $r$ . Expanding  $e^{ikru}$  in a power series in  $r$ , using  $e^{ikru} = \sum_{n=0}^{\infty} (ikru)^n / n!$ , this implies the identities

$$\int_{-1}^1 du u^n P_l(u) = 0 \quad (\text{for integer } n < l), \quad (10.2.49)$$

and the result we need (from the  $n = l$  term),

$$c_l \frac{k^l}{(2l+1)!!} = \frac{2l+1}{2} \frac{(ik)^l}{l!} \int_{-1}^1 du u^l P_l(u). \quad (10.2.50)$$

Using eq. (8.6.44) for the Legendre polynomials, one can obtain the integral

$$\int_{-1}^1 du u^l P_l(u) = \frac{2(l!)}{(2l+1)!!}, \quad (10.2.51)$$

by integrating by parts  $l$  times. Thus, eq. (10.2.50) simplifies to

$$c_l = (2l+1) i^l, \quad (10.2.52)$$

so that

$$e^{ikr \cos \theta} = \sum_{l=0}^{\infty} (2l+1) i^l j_l(kr) P_l(\cos \theta). \quad (10.2.53)$$

Now for the case of  $\vec{k}$  in an arbitrary direction, substitute  $kr \cos \theta \rightarrow \vec{k} \cdot \vec{r}$  in the preceding, or

$$\cos \theta \rightarrow \hat{k} \cdot \hat{r}. \quad (10.2.54)$$

Using the spherical harmonic addition identity, eq. (8.6.71), one finally arrives at eq. (10.2.44).

### 10.3 Particle confined to a sphere

As an application of the results of the previous section, consider a particle of mass  $\mu$  that is confined within a sphere of radius  $a$ , but is otherwise free, so that the potential is

$$V(r) = \begin{cases} 0 & (r < a), \\ \infty & (r \geq a). \end{cases} \quad (10.3.1)$$

The stationary-state wavefunctions are

$$\psi_{E,l,m} = \begin{cases} C_{E,l,m} j_l(kr) Y_l^m(\theta, \phi) & (r < a), \\ 0 & (r \geq a), \end{cases} \quad (10.3.2)$$






Figure 10.3.1: The solutions of the transcendental equation  $X = \tan X$  for positive  $X = ka$  provide the eigenvalues for  $k$  for the  $l = 1$  states of a particle confined to a ball of radius  $a$ . This graph shows the lowest three solutions  $X \approx \pi(1.4303, 2.4590, 3.4709) = (4.4934, 7.7253, 10.9041)$ , obtained as the intersections of  $Y = \tan X$  with the line  $Y = X$ .

where  $C_{E,l,m}$  are normalization constants and  $E = \hbar^2 k^2 / 2\mu$ . Continuity of the wavefunction at  $r = a$  requires that

$$j_l(ka) = 0, \quad (10.3.3)$$

and this boundary condition determines the allowed quantized energy levels  $E_{n,l}$ . Let us see how this works for  $l = 0, 1, 2$ .

For  $l = 0$ , eq. (10.3.3) becomes simply  $\sin(ka)/ka = 0$ , so  $ka = n\pi$ , where  $n$  is a positive integer. The energies are therefore

$$E_{n,0} = \frac{\hbar^2 k^2}{2\mu} = \frac{\hbar^2 \pi^2 n^2}{2\mu a^2} \quad (10.3.4)$$

for  $l = m = 0$ , with corresponding wavefunctions

$$\psi_{n,0,0} = C_{n,0,0} \frac{\sin(n\pi r/a)}{n\pi r/a}. \quad (10.3.5)$$

Note that these wavefunctions approach a non-zero constant at  $r = 0$ . The number of zeros of the radial wavefunction, including the one at  $r = a$ , is  $n$ . The constant can be fixed by requiring the unit normalization condition  $1 = 4\pi \int_0^a dr r^2 |\psi_{n,0,0}|^2$ , which yields  $C_{n,0,0} = n\sqrt{\pi/2a^3}$ .

For  $l = 1$ , the boundary condition (10.3.3) reads

$$\tan(ka) = ka, \quad (10.3.6)$$

which is a transcendental equation that can be understood graphically and then solved numerically for  $X = ka$  as shown in Figure 10.3.1. The lowest three energy solutions have  $ka/\pi \approx (1.4303, 2.4590, 3.4709, \dots)$ , so

$$E_{n,1} \approx \frac{\hbar^2 \pi^2}{2\mu a^2} (2.0458, 6.0468, 12.0471, \dots) \quad \text{for } n = (2, 3, 4, \dots), \quad (10.3.7)$$

where the label  $n$  is again the number of zeros of the radial wavefunction. In the  $l = 1$  case,  $r = 0$  and  $r = a$  are always both zeros, so  $n \geq 2$ . Each of the  $l = 1$  energy levels has degeneracy 3, because the quantum number  $m = -1, 0, 1$  does not affect the energy.

For  $l = 2$ , the boundary condition (10.3.3) becomes

$$\tan(ka) = \frac{ka}{1 - (ka)^2/3}, \quad (10.3.8)$$

which again is transcendental, but whose solutions can again be found numerically, with the results  $ka/\pi \approx (1.8346, 2.8950, 3.9225, \dots)$ , so

$$E_{n,2} \approx \frac{\hbar^2 \pi^2}{2\mu a^2} (3.3656, 8.3812, 15.3861, \dots) \quad \text{for } n = (2, 3, 4, \dots), \quad (10.3.9)$$

where again the label  $n$  is the number of zeros of the radial wavefunction  $j_2(kr)$ . Each of these energy levels has degeneracy 5, corresponding to  $m = -2, -1, 0, 1, 2$ .

The energy eigenvalues for higher angular momentum quantum number  $l$  can be numerically solved for in a similar way, and have increasingly higher energies. The degeneracy of each energy level  $E_{n,l}$  is  $2l + 1$ , corresponding to the allowed values of the  $L_z$  eigenvalue  $\hbar m$ . The lowest few energy levels, with  $E_{n,l} < 10\hbar^2\pi^2/\mu a^2$  and  $l \leq 7$ , are depicted in Fig. 10.3.2.




Figure 10.3.2: The lowest energy levels, in units of the ground-state energy  $E_{1,0} = \hbar^2\pi^2/2\mu a^2$ , for a particle of mass  $\mu$  confined to a sphere of radius  $a$ , labeled by the angular momentum quantum number  $l$ . All energy eigenvalues with  $E_{n,l} < 10\hbar^2\pi^2/\mu a^2$  and  $l \leq 7$  are shown.

## 10.4 Particle in a spherical potential well

Consider a particle of mass  $\mu$  in a spherical potential well of radius  $a$  and finite depth  $V_0$ ,

$$V(r) = \begin{cases} -V_0 & (r < a), \\ 0 & (r \geq a). \end{cases} \quad (10.4.1)$$

We will consider bound state solutions, which have  $E < 0$ . As discussed in section 10.2, the stationary-state solution for the region  $r < a$  involves the ordinary spherical Bessel function  $j_l(kr)$ , but now with

$$k = \sqrt{2\mu(E + V_0)}/\hbar. \quad (10.4.2)$$

The Neumann functions  $n_l(\rho)$  are not well-behaved at  $\rho = 0$  and so do not appear for  $r < a$ . The solution for  $r > a$  uses the modified spherical Bessel function  $k_l(\kappa r)$  [see eq. (10.2.29)] with

$$\kappa = \sqrt{-2\mu E}/\hbar, \quad (10.4.3)$$

because  $k_l(\rho)$  is the linear combination that is well-behaved at  $\rho = \infty$ . The stationary-state wavefunctions are therefore

$$\psi_{E,l,m}(\vec{r}) = \begin{cases} A j_l(kr) Y_l^m(\theta, \phi) & (r \leq a), \\ B k_l(\kappa r) Y_l^m(\theta, \phi) & (r \geq a), \end{cases} \quad (10.4.4)$$

where  $A$  and  $B$  are normalization constants, and

$$k^2 + \kappa^2 = 2\mu V_0/\hbar^2 \quad (10.4.5)$$

from eqs. (10.4.2) and (10.4.3).

At  $r = a$ , the wavefunction and its first derivative with respect to  $r$  are continuous, since the potential is finite there. This gives

$$A j_l(ka) = B k_l(\kappa a), \quad (10.4.6)$$

$$kA j_l'(ka) = \kappa B k_l'(\kappa a). \quad (10.4.7)$$

Taking the ratio of these to eliminate  $A$  and  $B$  yields

$$\frac{k j_l'(ka)}{j_l(ka)} = \frac{\kappa k_l'(\kappa a)}{k_l(\kappa a)}, \quad (10.4.8)$$

which is a transcendental equation that can be used together with eq. (10.4.5) to solve for the allowed eigenvalues  $k, \kappa$ , and thus  $E$ . As always for a spherically symmetric potential, the allowed energies depend on  $l$ , but not  $m$ . Either equation (10.4.6) or (10.4.7) then also allows for the ratio  $A/B$  to be found for each  $E, l$ . The remaining unknown corresponds to the

overall magnitude of the constants  $A$  and  $B$ , which can be fixed by the normalization of the wavefunction.

For example, if  $l = 0$ , eq. (10.4.8) yields

$$-k \cot(ka) = \kappa. \quad (10.4.9)$$

Writing dimensionless variables  $X = ka$  and  $Y = \kappa a$ , eqs. (10.4.5) and (10.4.9) give

$$X^2 + Y^2 = 2\mu V_0 a^2 / \hbar^2, \quad (10.4.10)$$

$$-X \cot X = Y. \quad (10.4.11)$$

Fortunately, these are exactly the same equations we encountered for the odd-parity solutions of the one-dimensional square well problem, with  $L \rightarrow 2a$ . [Compare to eqs. (6.5.12) and (6.5.18).] The same graphical and numerical analysis therefore applies. In particular, if we label the  $l = 0$  stationary states by  $n = 1, 2, 3, \dots$ , then the condition for the bound state  $|\psi_n\rangle$  to exist is

$$\frac{\mu V_0 a^2}{\hbar^2} > \frac{\pi^2}{2} (n - 1/2)^2. \quad (10.4.12)$$

Recall that in the case of a particle in a one-dimensional square well, there is always at least one bound state, but for a sufficiently shallow potential well only the even-parity ground state exists as a bound state. For the three-dimensional spherical well, since only the analogs of the odd-parity one-dimensional square-well states exist, eq. (10.4.12) tells us that the existence of a bound state requires the potential to be sufficiently deep,  $V_0 > \pi^2 \hbar^2 / 8\mu a^2$ . The more general lesson is that three-dimensional potentials may have no bound states if they are not sufficiently attractive.

## 10.5 Isotropic three-dimensional harmonic oscillator

In section 7.5, we have already encountered the isotropic three-dimensional harmonic oscillator as a special case of the general anisotropic version, and obtained its stationary-state energy levels and degeneracies. We did this using a CSCO of  $(H_x, H_y, H_z)$ , the individual Hamiltonians for excitations in the  $x$ ,  $y$ , and  $z$  directions. In this section, we will solve the problem again, this time with a CSCO consisting of  $(H, L^2, L_z)$ . This is possible because, in the isotropic case,

$$H = \frac{P^2}{2\mu} + \frac{1}{2} \mu \omega^2 R^2 \quad (10.5.1)$$

is invariant under all rotations and therefore commutes with  $L^2$  and  $L_z$ . Note that we are now using  $\mu$  as the symbol for the mass of the particle.

Before proceeding, let us consider what sort of answer we expect, given the results of section 7.5. At an energy level  $E = \hbar\omega(n + 3/2)$  with  $n = n_x + n_y + n_z$ , the wavefunctions must be linear combinations of wavefunctions of the form

$$\langle x, y, z | n_x, n_y, n_z \rangle \propto H_{n_x}(x/b) H_{n_y}(y/b) H_{n_z}(z/b) e^{-(x^2+y^2+z^2)/2b^2}, \quad (10.5.2)$$

where the length scale in the problem is

$$b = \sqrt{\hbar/\mu\omega}, \quad (10.5.3)$$

and the  $H_{n_a}$  are the Hermite polynomials. After translating to spherical coordinates, such a linear combination that is an eigenstate of  $L^2$  and  $L_z$  must be of the form

$$\langle r, \theta, \phi | n, l, m \rangle = Y_l^m(\theta, \phi) R_{n,l}(r), \quad (10.5.4)$$

where

$$R_{n,l}(r) = (\text{polynomial of degree } n \text{ in } r) e^{-r^2/2b^2}. \quad (10.5.5)$$

Our goal is to solve for the functions  $R_{n,l}(r)$ .

In section 10.1, we found the differential equation for  $U_{n,l} = rR_{n,l}$ , which in the present case becomes

$$\left[ -\frac{\hbar^2}{2\mu} \frac{d^2}{dr^2} + \frac{1}{2} \mu \omega^2 r^2 + \frac{\hbar^2 l(l+1)}{2\mu r^2} - E \right] U_{n,l} = 0. \quad (10.5.6)$$

Inspired by eq. (10.5.5), we define dimensionless quantities  $x$  and  $y(x)$  by

$$x = r/b, \quad U_{n,l} = y(x) e^{-x^2/2}. \quad (10.5.7)$$

When plugged into eq. (10.5.6), this gives

$$y'' - 2xy' + [\mathcal{E} - 1 - l(l+1)/x^2] y = 0, \quad (10.5.8)$$

where

$$\mathcal{E} = 2\mu b^2 E / \hbar^2 = 2E / \hbar\omega \quad (10.5.9)$$

is a dimensionless combination proportional to the energy eigenvalue. We already know from eq. (7.5.13) that the allowed eigenvalues are  $\mathcal{E} = 2n + 3$  for non-negative integers  $n$ , and from eq. (10.5.5) that  $y(x)$  is a polynomial, but in the following derivation we will proceed as if these facts were not known.

We now try a series solution for  $y(x)$ , of the form

$$y = x^q \sum_{p=0}^{\infty} c_p x^p. \quad (10.5.10)$$

Here  $q$  and the  $c_p$  are constants, with  $c_0 \neq 0$  by definition. (Otherwise, we would adjust the value of  $q$  to make it so.) Now we prepare to plug in to eq. (10.5.8) by computing

$$y'' = \sum_{p=0}^{\infty} c_p (p+q)(p+q-1) x^{p+q-2}, \quad (10.5.11)$$

$$y/x^2 = \sum_{p=0}^{\infty} c_p x^{p+q-2}, \quad (10.5.12)$$

$$xy' = \sum_{p=0}^{\infty} c_p (p+q) x^{p+q} = \sum_{p=0}^{\infty} c_{p-2} (p+q-2) x^{p+q-2}, \quad (10.5.13)$$

$$y = \sum_{p=0}^{\infty} c_{p-2} x^{p+q-2}, \quad (10.5.14)$$

where in the last equality of each of eqs. (10.5.13) and (10.5.14) we have used the trick of relabeling  $p \rightarrow p-2$  and defining  $c_{-2} = c_{-1} = 0$ . The motivation behind this relabeling trick is that now all of the summands have the same powers of  $x$ , and so eq. (10.5.8) becomes

$$\sum_{p=0}^{\infty} x^{p+q-2} \left\{ c_p [(p+q)(p+q-1) - l(l+1)] - c_{p-2} [2(p+q-2) + 1 - \mathcal{E}] \right\} = 0. \quad (10.5.15)$$

For this equation to be satisfied for all  $x$ , each coefficient of a given power of  $x$  must vanish, so the quantity in braces must vanish for each  $p$ .

From the first term  $p = 0$ , using  $c_{-2} = 0$  we find

$$c_0 [q(q-1) - l(l+1)] = 0. \quad (10.5.16)$$

Since  $c_0 \neq 0$ , the possible solutions are  $q = l+1$  and  $q = -l$ . However, the latter can be rejected on physical grounds, since it would imply that  $y \sim x^{-l}$  for small  $x$ , which would mean  $R_{n,l} \sim 1/r^{l+1}$  for small  $r$ , and the wavefunction would not be finite at the point  $r = 0$ . Therefore,

$$q = l+1, \quad (10.5.17)$$

which implies that  $R_{n,l} \sim r^l$  for small  $r$ . This suppression for small  $r$  is consistent with the existence of the repulsive  $\hbar^2 l(l+1)/2\mu r^2$  centrifugal barrier contribution to the effective potential for non-zero angular momentum, see eq. (10.1.10) or eq. (10.5.6).

The second term  $p = 1$  gives us

$$c_1 [q(q+1) - l(l+1)] = 0, \quad (10.5.18)$$

because  $c_{-1} = 0$ . Now, plugging in  $q = l + 1$ , this becomes  $2(l + 1)c_1 = 0$ . Since our knowledge of the angular momentum eigenvalue problem tells us that  $l + 1$  cannot vanish, we must have  $c_1 = 0$ .

For all larger  $p$ , the vanishing of eq. (10.5.15) implies (after using  $q = l + 1$ ) that

$$c_p p(p + 2l + 1) = c_{p-2} (2p + 2l - 1 - \mathcal{E}). \quad (10.5.19)$$

Since  $c_1 = 0$ , it follows that  $c_p = 0$  for all odd  $p$ . Defining  $C_j = c_{2j}$ , we have

$$y = x^{l+1} \sum_{j=0}^{\infty} C_j x^{2j}, \quad (10.5.20)$$

with, from eq. (10.5.19) by taking  $p = 2(j + 1)$ , the recurrence relation

$$C_{j+1} = \frac{4j + 2l + 3 - \mathcal{E}}{2(j + 1)(2j + 2l + 3)} C_j. \quad (10.5.21)$$

For large  $j$ , we have  $C_j \approx C_{j-1}/j$ , so if the series does not terminate, it would behave asymptotically like  $y \sim x^{l+1} \sum_j (x^2)^j / j! \sim x^{l+1} e^{x^2}$ , which would imply that  $R_{n,l} \sim r^l e^{-r^2/2b^2} e^{r^2/b^2} \sim r^l e^{r^2/2b^2}$ . As in the case of the one-dimensional harmonic oscillator, such a solution that blows up exponentially as  $r \rightarrow \infty$  is unphysical, since it cannot be normalized.

We can therefore conclude that the series must terminate, and  $y(x)$  is actually a polynomial, in agreement with eq. (10.5.5). Then the recurrence relation eq. (10.5.21) implies that physically valid solutions for  $y(x)$  must have  $C_{k+1} = 0$  for some integer  $k \geq 0$ , and so satisfy

$$\mathcal{E} = 4k + 2l + 3, \quad (10.5.22)$$

or, using eq. (10.5.9),

$$E = \hbar\omega(2k + l + 3/2). \quad (10.5.23)$$

The integer  $n = n_x + n_y + n_z$  must then be equal to  $2k + l$ , and the energy eigenkets  $|E, l, m\rangle$  are linear combinations of the kets  $|n_x, n_y, n_z\rangle$ . An interesting feature of these results is that, unlike the particle-in-a-sphere and particle-in-a-spherical-well examples of sections 10.3 and 10.4, here the degeneracies in the energy levels are not entirely due to the rotational invariance of the problem. For a given  $l$ , there are  $2l + 1$  degenerate states with  $m = -l, \dots, l$  that have the same energy. But, there are also “accidental” energy degeneracies between states with different angular momenta, for example  $k = 0, l = 2$  and  $k = 1, l = 0$ .

Summarizing what we now know about the three-dimensional isotropic harmonic oscillator stationary-state wavefunctions with definite  $(L^2, L_z)$ ,

$$\psi_{k,l,m}(r, \theta, \phi) = AR_{k,l}(r) Y_l^m(\theta, \phi), \quad (10.5.24)$$

where  $A$  is a normalization constant. The radial wavefunctions are of the form

$$R_{k,l}(r) = (r/b)^l P_{k,l}(r^2/b^2) e^{-r^2/2b^2}, \quad (10.5.25)$$

where, with  $z = x^2$ ,

$$P_{k,l}(z) = \sum_{j=0}^k C_j z^j. \quad (10.5.26)$$

Using eq. (10.5.22), the recurrence relation (10.5.21) becomes

$$C_{j+1} = \frac{2(j-k)}{(j+1)(2j+2l-3)} C_j. \quad (10.5.27)$$

For the lowest few values  $k = 0, 1, 2, 3$ , we have (leaving the coefficient  $C_0$  as an arbitrary normalization for the moment):

$$P_{0,l}(z) = C_0, \quad (10.5.28)$$

$$P_{1,l}(z) = C_0 \left( 1 - \frac{2}{2l+3} z \right), \quad (10.5.29)$$

$$P_{2,l}(z) = C_0 \left( 1 - \frac{4}{2l+3} z + \frac{4}{(2l+3)(2l+5)} z^2 \right), \quad (10.5.30)$$

$$P_{3,l}(z) = C_0 \left( 1 - \frac{6}{2l+3} z + \frac{12}{(2l+3)(2l+5)} z^2 - \frac{8}{(2l+3)(2l+5)(2l+7)} z^3 \right). \quad (10.5.31)$$

These polynomials are proportional to **associated Laguerre polynomials**, for which there are unfortunately at least three different notational conventions in common use. Adopting here the definition of the associated Laguerre polynomial  $L_k^\alpha(z)$  used by Mathematica,<sup>†</sup>

$$P_{k,l}(z) = L_k^{l+1/2}(z), \quad (10.5.32)$$

corresponding to the choice

$$C_0 = \frac{(2l+2k+1)!!}{2^k k! (2l+1)!!} = \frac{(2l+2k+1)! l!}{2^{2k} (2l+1)! (l+k)! k!}. \quad (10.5.33)$$

More generally, the conventional normalization is such that, at  $z = 0$  and for any  $\alpha$ ,

$$L_k^\alpha(0) = \frac{\Gamma(k+\alpha+1)}{k! \Gamma(\alpha+1)}. \quad (10.5.34)$$

Here, the **Gamma function** is defined by

$$\Gamma(z) = \int_0^\infty dt t^{z-1} e^{-t}, \quad (10.5.35)$$

---

<sup>†</sup>The relations between the definition of the associated Laguerre polynomials used here and in some other sources is  $\Gamma(N+\alpha+1)L_N^\alpha(z)_{\text{here, Mathematica}} = L_N^\alpha(z)_{\text{some other books}} = (-1)^\alpha L_{N+\alpha}^\alpha(z)_{\text{still other books}}$ .



and satisfies  $\Gamma(z) = (z-1)\Gamma(z-1)$ . When  $z$  is an integer, then  $z! = \Gamma(z+1)$ . Another useful value is  $\Gamma(1/2) = \sqrt{\pi}$ .

An explicit, general form for the associated Laguerre polynomials is

$$L_N^\alpha(z) = \frac{1}{N!} z^{-\alpha} e^z \frac{d^N}{dz^N} (e^{-z} z^{N+\alpha}). \quad (10.5.36)$$

Note that the lower index  $N$  is always an integer equal to the degree of the polynomial, but the raised index  $\alpha$  need not be, as in the present application where  $\alpha = l + 1/2$ . They satisfy the differential equation

$$\left[ z \frac{d^2}{dz^2} + (1 - z + \alpha) \frac{d}{dz} + N \right] L_N^\alpha(z) = 0, \quad (10.5.37)$$

and the orthogonality relation

$$\int_0^\infty dz e^{-z} z^\alpha L_N^\alpha(z) L_{N'}^\alpha(z) = \frac{\Gamma(N + \alpha + 1)}{N!} \delta_{NN'}. \quad (10.5.38)$$

The degree  $N$  is also equal to the number of zeros (in the present case  $N = k$ ), which all occur for positive real  $z$ .

Putting everything together, the wavefunctions for the stationary states of the three-dimensional isotropic harmonic oscillator with energies  $E = \hbar\omega(2k + l + 3/2)$  are

$$\psi_{k,l,m}(r, \theta, \phi) = \sqrt{\frac{2(k!)}{b^3 \Gamma(k + l + 3/2)}} \left(\frac{r}{b}\right)^l L_k^{l+1/2}(r^2/b^2) e^{-r^2/2b^2} Y_l^m(\theta, \phi). \quad (10.5.39)$$

The normalization factor  $A$  in eq. (10.5.24) has been chosen, with the aid of eq. (10.5.38), so that orthonormality holds,

$$\langle k', l', m' | k, l, m \rangle = \int_0^\infty dr r^2 \int d\Omega [\psi_{k',l',m'}(r, \theta, \phi)]^* \psi_{k,l,m}(r, \theta, \phi) = \delta_{kk'} \delta_{ll'} \delta_{mm'}. \quad (10.5.40)$$

## 10.6 Exercises

**Exercise 10.1.** Consider a particle of mass  $\mu$  in a spherically symmetric potential  $V(R)$  in a stationary bound state with energy  $E$  and zero angular momentum, so that the wavefunction  $\psi(r)$  is only a function of  $r$ . The quantity  $|\psi(0)|^2$  is useful for evaluating “contact” quantities that depend on the probability for the particle to be found at the potential source, the origin. (An example is the Darwin term of the hydrogen atom fine structure.) Show that in general

$$|\psi(0)|^2 = \frac{\mu}{2\pi\hbar^2} \langle V'(R) \rangle, \quad (10.6.1)$$

with the standard wavefunction normalization such that  $4\pi \int_0^\infty dr r^2 |\psi(r)|^2 = 1$ . [Hint: multiply the Schrödinger equation  $-\frac{\hbar^2}{2\mu} (\psi'' + \frac{2}{r}\psi') + [V(r) - E] \psi = 0$  by  $\psi'(r)$ , and integrate by parts.] Check this result in the special cases of the  $l = 0$  states of a particle confined to the interior of a sphere, and the ground state of the isotropic harmonic oscillator.

**Exercise 10.2.** Use the spherical Bessel and Neumann function forms in eqs. (10.2.11)-(10.2.13) to verify the large-distance forms of eqs. (10.2.14) and (10.2.15) for  $l = 0, 1, 2$ .

**Exercise 10.3.** Consider a particle of mass  $\mu$  trapped inside a ball of radius  $b$  that has a hard core of radius  $a$ , so that the potential in spherical coordinates is

$$V(r) = \begin{cases} \infty & (\text{for } r < a \text{ and for } r > b). \\ 0 & (\text{for } a < r < b) \end{cases} \quad (10.6.2)$$

This means that the eigenstates of  $H$ ,  $L^2$ , and  $L_z$  have wavefunctions of the form  $\Psi_{E,l,m}(r, \theta, \phi) = [A_jl(kr) + Bn_l(kr)]Y_l^m(\theta, \phi)$  for the region  $a < r < b$ .

(a) Find all of the allowed energy eigenstates and eigenvalues for  $l = 0$ . [Hint: use boundary conditions to solve for the ratio  $B/A$  twice, and require the two expressions to be equal.]

(b) For the case  $l = 1$ , find a transcendental equation whose solutions will yield the energy eigenvalues. Put your equation into the form  $\tan[k(b - a)] = \{\text{an expression not involving sines or cosines}\}$ . [Hint: first put the equation into a form that is polynomial in  $ka$ ,  $kb$ , and their sines and cosines; then use trigonometric identities for  $\sin(kb - ka)$  and  $\cos(kb - ka)$ .]

(c) For the special case  $l = 1$  and  $b = 2a$ , write your transcendental equation in the form  $\tan X = X/(1 + NX^2)$ , where  $X = ka$  and  $N$  is a certain integer that you will discover. Solve for  $X$  numerically to at least 3 digits of accuracy, and obtain the lowest energy for  $l = 1$ . How does it compare to the lowest energy for  $l = 0$  that you found in part (a)?

**Exercise 10.4.** Consider the isotropic 3-d harmonic oscillator problem, with potential  $V(x, y, z) = \frac{1}{2}\mu\omega^2(x^2 + y^2 + z^2)$ . The Hamiltonian  $H$  can be written as the sum of  $H_x = \hbar\omega(a_x^\dagger a_x + 1/2)$ ,  $H_y = \hbar\omega(a_y^\dagger a_y + 1/2)$ , and  $H_z = \hbar\omega(a_z^\dagger a_z + 1/2)$ , which form a CSCO with corresponding orthonormal eigenbasis  $|n_x, n_y, n_z\rangle$ . Another choice of CSCO is  $H$ ,  $L^2$ , and  $L_z$ , with corresponding eigenbasis  $|n, l, m\rangle'$ , where  $n = n_x + n_y + n_z$ . (The ' distinguishes the two types of orthobasis elements, since they both have three integer labels.)

(a) Show that the angular momentum component operators  $L_x$ ,  $L_y$ , and  $L_z$  can be written as

$$L_b = i\hbar\epsilon_{bcd}a_c a_d^\dagger. \quad (10.6.3)$$

(b) Construct the operator  $L^2$  in terms of the ladder operators. You should write the answer in “normal-ordered” form, which means that the commutation relations have been used to ensure that no creation operator appears to the right of an annihilation operator, as

$$L^2 = \hbar^2 [N_1(a_x^{\dagger 2}a_y^2 + a_x^{\dagger 2}a_z^2 + a_y^{\dagger 2}a_x^2 + a_y^{\dagger 2}a_z^2 + a_z^{\dagger 2}a_x^2 + a_z^{\dagger 2}a_y^2) + N_2(a_x^\dagger a_y^\dagger a_x a_y + a_x^\dagger a_z^\dagger a_x a_z + a_y^\dagger a_z^\dagger a_y a_z) + N_3(a_x^\dagger a_x + a_y^\dagger a_y + a_z^\dagger a_z)], \quad (10.6.4)$$

where  $N_1$ ,  $N_2$ , and  $N_3$  are certain integers that you will discover. Note that  $L_x$ ,  $L_y$ ,  $L_z$ , and  $L^2$  all give 0 acting on the ground state with  $n = 0$ .

- (c) For the subspace of states  $|1, 0, 0\rangle$ ,  $|0, 1, 0\rangle$ , and  $|0, 0, 1\rangle$  with  $n = 1$ , find the action of  $L^2$  and  $L_z$ . What are their matrix representations in that basis? Find the eigenvalues and eigenvectors of  $L_z$  within this subspace.
- (d) For the subspace of states with  $n = 2$ , find the action of  $L^2$  on each of the  $|n_x, n_y, n_z\rangle$  basis. Using these results, and using the ordering  $|2, 0, 0\rangle$ ,  $|0, 2, 0\rangle$ ,  $|0, 0, 2\rangle$ ,  $|1, 1, 0\rangle$ ,  $|1, 0, 1\rangle$ ,  $|0, 1, 1\rangle$ , find the corresponding  $6 \times 6$  matrix representation for  $L^2$ . Find the eigenvalues and normalized eigenvectors of  $L^2$  for the  $n = 2$  subspace in that basis.
- (e) Compute the action of  $L_z$  on each of the simultaneous eigenvectors of  $H, L^2$  found in the previous part. Within each sub-subspace of fixed  $n = 2$  and fixed  $l$ , find the eigenvalues and eigenvectors of  $L_z$ , and so conclude by writing the six  $|2, l, m\rangle'$  orthobasis states as linear combinations of the six  $|n_x, n_y, n_z\rangle$  eigenstates.

# 11 Coulomb potential and hydrogen-like atoms

## 11.1 Bound states for hydrogen atom

One of the most important exactly solvable problems in quantum mechanics is that of a particle moving in a Coulomb potential. This includes the hydrogen atom, and more generally hydrogen-like ions, which consist of a single electron with mass  $m_e$  and charge  $-e$  [with the normalization as defined in Gaussian cgs metric system units, see eqs. (1.1.1) and (1.1.2)] and a heavy nucleus with mass  $m_N$  and charge  $Ze$ , where  $Z$  is an integer. As discussed in a general context in section 4.2, the problem can be separated into center-of-mass and relative degrees of freedom, where the latter use a reduced mass  $\mu = m_e m_N / (m_e + m_N)$ , which is very close to  $m_e$ . In this chapter, we neglect the small effects of special relativity and electron and nuclear spins, which will be treated in Chapter 17. So, our Hamiltonian is  $H = P^2/2\mu + V(R)$ , where the potential energy in the position representation is

$$V(r) = -Ze^2/r. \quad (11.1.1)$$

The special case  $Z = 1$  is the hydrogen atom. Because the potential is spherically symmetric, we choose the CSCO to be  $(H, L^2, L_z)$ , and look for an orthobasis of eigenstates  $|E, l, m\rangle$  with wavefunctions

$$\psi_{E,l,m}(r, \theta, \phi) = \langle r, \theta, \phi | E, l, m \rangle = R_{E,l}(r) Y_l^m(\theta, \phi). \quad (11.1.2)$$

The goal is to find the energy eigenvalues and the radial wavefunctions  $R_{E,l}(r)$ .

In this section, we will work out the properties of the bound state solutions, for which the energy eigenvalues are discrete and satisfy  $E < V(\infty) = 0$ , and  $|R_{E,l}|$  decreases exponentially as  $r \rightarrow \infty$ . Unlike the case of the three-dimensional harmonic oscillator, there are also unbound energy eigenstates, for which  $E$  is continuous and non-negative, and  $rR_{E,l}$  oscillates with an amplitude approaching a constant for large  $r$ . The unbound state solutions will be found in section 11.2.

The time-independent Schrödinger equation for the radial wavefunction is

$$\left[ -\frac{\hbar^2}{2\mu} \left( \frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} - \frac{l(l+1)}{r^2} \right) - \frac{Ze^2}{r} - E \right] R_{E,l}(r) = 0. \quad (11.1.3)$$

A good first step is to replace  $r$  by a dimensionless variable. Since  $-\hbar^2/2\mu E$  has units of [length]<sup>2</sup> and is a positive number for the bound states, we define a rescaled radial coordinate

$$s = 2r/b, \quad b = \hbar/\sqrt{-2\mu E}. \quad (11.1.4)$$

With this change of variables, eq. (11.1.3) becomes

$$\left( \frac{d^2}{ds^2} + \frac{2}{s} \frac{d}{ds} - \frac{l(l+1)}{s^2} + \frac{n}{s} - \frac{1}{4} \right) R_{E,l} = 0, \quad (11.1.5)$$

where we have introduced

$$n = \frac{Ze^2}{\hbar} \sqrt{\frac{-\mu}{2E}}, \quad (11.1.6)$$

a dimensionless quantity that parameterizes the energy eigenvalue. Note that  $n$  is a real number for  $E < 0$ , and is pure imaginary for  $E > 0$ . We will soon find out [see eq. (11.1.15)] that  $n$  must be a positive integer for a bound-state solution, but we do not know that yet.

The next part of our strategy is to identify, and factor out, the large-distance and short-distance behaviors of  $R_{E,l}$ . For  $s \rightarrow \infty$ , eq. (11.1.5) becomes  $d^2 R/ds^2 \approx R/4$ , which has two linearly independent solutions,  $e^{s/2}$  and  $e^{-s/2}$ . The first of these is unphysical, as it blows up for  $s \rightarrow \infty$  and so is not normalizable. In the opposite limit  $s \rightarrow 0$ , the last two terms of eq. (11.1.5) can be neglected, and there is a power-law solution for  $R$  proportional to  $s^l$ . We therefore write

$$R_{E,l} = s^l e^{-s/2} f(s), \quad (11.1.7)$$

which factors out the leading behavior in the two limits. Plugging this into eq. (11.1.5) gives

$$s \frac{d^2 f}{ds^2} + (2l + 2 - s) \frac{df}{ds} + (n - l - 1) f = 0, \quad (11.1.8)$$

which we must now solve simultaneously for the eigenfunctions  $f(s)$  and the corresponding eigenvalues  $n$ .

Equation (11.1.8) is a special case of a famous differential equation, called the **confluent hypergeometric equation**, which in its traditional general form is

$$x \frac{d^2 F}{dx^2} + (c - x) \frac{dF}{dx} - aF = 0, \quad (11.1.9)$$

where  $a$  and  $c$  are constants. This has a unique (up to a multiplicative constant) solution that is finite as  $x \rightarrow 0$ , the confluent hypergeometric function, which has a series expansion

$$F(a, c, x) = 1 + \frac{a}{c}x + \frac{a(a+1)}{c(c+1)} \frac{x^2}{2!} + \frac{a(a+1)(a+2)}{c(c+1)(c+2)} \frac{x^3}{3!} + \cdots, \quad (11.1.10)$$

as can be verified by direct substitution into the differential equation. This series converges for all finite  $|x|$ , even if  $a$ ,  $c$ , and  $x$  are complex, provided that  $c \neq 0, -1, -2, \dots$ . One can check (either from the series solution, or by direct substitution into the differential equation) that it has an integral representation

$$F(a, c, x) = \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-a)} \int_0^1 dt e^{tx} t^{a-1} (1-t)^{c-a-1}, \quad (11.1.11)$$

provided that  $\text{Re}[a] > \text{Re}[c] > 0$ . For real  $x$  large and positive, it has the asymptotic form

$$F(a, c, x) \approx \frac{\Gamma(c)}{\Gamma(a)} x^{a-c} e^x \quad (\text{large real } x, \text{ and } a \neq 0, -1, -2, \dots). \quad (11.1.12)$$

However, as we are about to see, the special case relevant for our present purposes will arise when  $a$  is a non-positive integer; in that case,  $F(a, c, x)$  is instead a polynomial in  $x$  of degree  $-a$ , because the series in eq. (11.1.10) terminates.

For the present application to the bound states of the hydrogen-like atom, we have  $x = s$  and  $a = l + 1 - n$  and  $c = 2l + 2$ , by comparing eq. (11.1.8) to (11.1.9). Therefore,  $f(s)$  in eq. (11.1.7) is equal to  $F(l+1-n, 2l+2, s)$ , up to normalization, so

$$R_{n,l} = C_{n,l} s^l e^{-s/2} F(l+1-n, 2l+2, s), \quad (11.1.13)$$

Here we have replaced the subscript label  $E$  by the label  $n$ , which contains the same information, and  $C_{n,l}$  is a normalization constant to be chosen later. In the large-distance limit  $s \rightarrow \infty$ , the asymptotic form of eq. (11.1.12) would seem to tell us

$$R_{n,l} \propto s^{-n-1} e^{s/2} \quad (n-l-1 \neq 0, 1, 2, \dots). \quad (11.1.14)$$

However, the exponential factor  $e^{s/2}$  is the form that we already rejected as non-normalizable and therefore physically unacceptable; it would imply that no matter how far from the nucleus you look, the electron must have infinitely larger probability to be farther away. The only way to get a physically sensible bound state is to arrange for eq. (11.1.14) not to apply. In other words, it is necessary that the series solution eq. (11.1.10) for  $F(l+1-n, 2l+2, s)$  terminates, so that instead of being proportional to  $e^s$ , it is actually a polynomial in  $s$  with degree that we will call  $k$ . This implies that

$$n = k + l + 1. \quad (11.1.15)$$

Since  $k$ , being the degree of the polynomial, is a non-negative integer, and  $l + 1$  is always a positive integer,  $n$  must be a positive integer, called the **principal quantum number** of the hydrogen atom bound state.

Inverting eq. (11.1.6), the allowed energy eigenvalues are

$$E_n = - \left( \frac{e^4 \mu}{2\hbar^2} \right) \frac{Z^2}{n^2} = - \left( \frac{e^2}{2a_0} \right) \frac{Z^2}{n^2}, \quad (11.1.16)$$

which depend only on  $n$ , not  $k$  and  $l$  individually. To write the last expression we have defined<sup>†</sup>

---

<sup>†</sup>The definitions of the Bohr radius and Rydberg energy units used here are appropriate for the infinite nuclear mass limit, with  $\mu = m_e$ . An alternative definition uses, instead of the electron mass  $m_e$ , the reduced mass  $\mu$  for the lightest isotope of hydrogen, which is smaller by a factor  $m_p/(m_e + m_p) \approx 0.999453$ . From here on, we ignore the small difference between  $m_e$  and  $\mu$ , which can be restored by replacing  $m_e \rightarrow \mu$  in the obvious way.




Figure 11.1.1: Stationary-state energy levels of the hydrogen atom for  $l \leq 4$ . The bound state levels with  $E_n = -13.6 \text{ eV}/n^2$  for  $n = 1, 2, 3, \dots$  are discrete and have degeneracy  $n^2$ , with  $0 \leq l \leq n - 1$ . For each non-negative integer  $l$ , there are also unbound continuum energy eigenstates with  $E \geq 0$ .

the **Bohr radius** (named after Niels Bohr),

$$a_0 = \frac{\hbar^2}{e^2 m_e} = 5.292 \times 10^{-11} \text{ meters} = \text{Bohr radius.} \quad (11.1.17)$$

The scale of energy is therefore

$$\frac{e^4 m_e}{2\hbar^2} = \frac{e^2}{2a_0} = \frac{\hbar^2}{2m_e a_0^2} = 2.180 \times 10^{-18} \text{ Joules} = 13.606 \text{ eV} = \text{Rydberg}, \quad (11.1.18)$$

named after Johannes Rydberg. The ground state has  $n = 1$  and  $k = l = 0$ . More generally, for each  $l$ , the lowest possible energy is obtained for  $k = 0$ , so that  $n = l + 1$ .

The energy levels of the hydrogen atom are depicted in Fig. 11.1.1. At each bound-state energy level  $n$ , the values of  $l$  that can occur are the integers from 0 to  $n - 1$ , each with degeneracy  $2l + 1$ , which comes from  $m = -l, \dots, l$ . The total degeneracy for each energy level  $E_n$  is therefore

$$g_n = \sum_{l=0}^{n-1} (2l + 1) = n^2. \quad (11.1.19)$$

Because this includes states with different angular momenta, the energy degeneracy is accidental, meaning that it cannot be explained by rotational invariance alone. This was also the case for the three-dimensional isotropic harmonic oscillator. For hydrogen-like atoms, this degeneracy is slightly broken by spin and relativistic effects, as we will discuss in detail in Chapter 17.

The distinct values of  $l$  are given letter codes that go back to the early history of spectroscopic analysis. This **spectroscopic notation** is as follows:

$$\begin{aligned}
l = 0 &\leftrightarrow s \text{ “sharp”}, \\
l = 1 &\leftrightarrow p \text{ “principal”}, \\
l = 2 &\leftrightarrow d \text{ “diffuse”}, \\
l = 3 &\leftrightarrow f \text{ “fundamental”}, \\
l = 4 &\leftrightarrow g \text{ “grotesque”}, \\
l = 5 &\leftrightarrow h \text{ “horrendous”}, \\
&\dots \qquad \dots
\end{aligned} \tag{11.1.20}$$

The letter codes are in alphabetical order  $f, g, h, i, k, \dots$  for  $l = 3, 4, 5, 6, 7, \dots$ , but  $j$  is omitted. (And I’m just kidding about the names “grotesque” and “horrendous”; unlike the first four, I just made those up, and they do not actually have standard names to fit the standard letters.) A supposedly useful mnemonic is “sober **p**hysicists **d**on’t find **g**iraffes **h**iding in **k**itchens”. The hydrogen atom states are often referred to in the notation  $nl$ , but with  $l = 0, 1, 2, 3, 4, 5, \dots$  replaced by the spectroscopic code<sup>‡</sup> letter  $s, p, d, f, g, h, \dots$ , so that the ground state is called  $1s$  and the first excited states are  $2s$  and  $2p$ , and the second excited states are  $3s$ ,  $3p$ , and  $3d$ .

Returning to the radial wavefunction, eq. (11.1.13) tells us that

$$R_{n,l} = C_{n,l} s^l e^{-s/2} F_{n,l}, \tag{11.1.21}$$

where we are now adopting the shorter notation

$$F_{n,l} = F(l+1-n, 2l+2, s), \tag{11.1.22}$$

which is a polynomial of degree  $k = n - l - 1$ . Recalling that  $s = 2r/b$  from eq. (11.1.4), the exponential factor tells us that the spatial support of wavefunctions is set, in terms of the Bohr radius, by the length scale

$$b = na_0/Z. \tag{11.1.23}$$

Equations (11.1.16) and (11.1.23) show that the magnitude of the binding energy scales like  $Z^2$ , while the characteristic size of a given wavefunction’s support scales like  $1/Z$ . In the classical limit  $\hbar \rightarrow 0$ , the Bohr radius goes to 0 and the binding energy of the ground state goes to  $-\infty$ , in accord with the discussion of the classical instability of atoms in section 1.1.

---

<sup>‡</sup>In some other contexts, the letter codes for orbital angular momentum in eq. (11.1.20) are capitalized.



For the lowest few energy states, the polynomials are, from eq. (11.1.10),

$$F_{1,0} = 1, \quad (11.1.24)$$

$$F_{2,1} = 1, \quad F_{2,0} = 1 - \frac{s}{2}, \quad (11.1.25)$$

$$F_{3,2} = 1, \quad F_{3,1} = 1 - \frac{s}{4}, \quad F_{3,0} = 1 - s + \frac{s^2}{6}, \quad (11.1.26)$$

$$F_{4,3} = 1, \quad F_{4,2} = 1 - \frac{s}{6}, \quad F_{4,1} = 1 - \frac{s}{2} + \frac{s^2}{20}, \quad F_{4,0} = 1 - \frac{3s}{2} + \frac{s^2}{2} - \frac{s^3}{24}. \quad (11.1.27)$$

More generally, in terms of the associated Laguerre polynomials  $L_N^\alpha(x)$  already defined explicitly by eq. (10.5.36), it can be shown that

$$F_{n,l} = \frac{(n-l-1)!(2l+1)!}{(n+l)!} L_{n-l-1}^{2l+1}(s). \quad (11.1.28)$$

The lower index on  $L_N^\alpha$  is always equal to the degree of the associated Laguerre polynomial, which is also the number of its zeros, in this case  $N = k - 1$ . Note, however, that in the case of the isotropic three-dimensional harmonic oscillator the upper index  $\alpha$  was always half-integer [see eq. (10.5.32)], while here  $\alpha = 2l + 1$  is always a positive odd integer.

From Theorem 2.6.5, we know that the kets  $|n, l, m\rangle$  and  $|n', l', m'\rangle$  are orthogonal whenever any of  $n, l$ , or  $m$  differ from  $n', l'$ , or  $m'$ , respectively. The orthogonality conditions  $\delta_{ll'}$  and  $\delta_{mm'}$  are already enforced by the proportionality of the wavefunctions to spherical harmonics. It follows that the associated Laguerre polynomials must also satisfy an orthogonality relation of the form (taking  $x = ns = 2Zr/a_0$ ):

$$\int_0^\infty dx x^{2l+2} L_{n-l-1}^{2l+1}(x/n) L_{n'-l-1}^{2l+1}(x/n') e^{-x(1/n+1/n')/2} = \delta_{nn'} B_{nl}, \quad (11.1.29)$$

for  $n, n' = 1, 2, 3, \dots$  and  $l = 0, 1, \dots, \min(n, n') - 1$ . Although not obvious, it can be checked that this is true, with

$$B_{nl} = \frac{2(n+l)! n^{2l+4}}{(n-l-1)!}. \quad (11.1.30)$$

Note that eq. (11.1.29) is a quite different orthogonality relation for associated Laguerre polynomials than the one that was useful for the three-dimensional harmonic oscillator, eq. (10.5.38).

Putting together eqs. (11.1.4), (11.1.21), (11.1.23), and (11.1.28), we obtain the radial wavefunctions of the hydrogen-like atom bound states,

$$R_{n,l}(r) = A_{n,l} e^{-Zr/na_0} \left( \frac{2Zr}{na_0} \right)^l L_{n-l-1}^{2l+1}(2Zr/na_0), \quad (11.1.31)$$

where we have introduced a new normalization constant factor

$$A_{n,l} = \left( \frac{2Z}{a_0} \right)^{3/2} \frac{n^l}{\sqrt{B_{n,l}}} = \left( \frac{Z}{a_0} \right)^{3/2} \frac{2}{n^2} \sqrt{\frac{(n-l-1)!}{(n+l)!}}, \quad (11.1.32)$$

chosen so that

$$\int_0^\infty dr r^2 R_{n,l}(r) R_{n',l}(r) = \delta_{nn'}. \quad (11.1.33)$$

The full wavefunction [obtained by including the  $Y_l^m(\theta, \phi)$  factor according to eq. (11.1.2)] is

$$\psi_{n,l,m}(r, \theta, \phi) = R_{n,l}(r) Y_l^m(\theta, \phi). \quad (11.1.34)$$

An arbitrary choice of phase has been made in  $A_{n,l}$ , as usual. With this sensible choice, the radial wavefunctions  $R_{n,l}$  are all real.

We now have everything necessary to evaluate the wavefunctions for the lowest few energy levels. For the ground state,

$$R_{1,0}(r) = \left(\frac{Z}{a_0}\right)^{3/2} 2e^{-Zr/a_0}, \quad (11.1.35)$$

and for the first excited states,

$$R_{2,0}(r) = \left(\frac{Z}{a_0}\right)^{3/2} \frac{1}{2\sqrt{2}} \left(2 - \frac{Zr}{a_0}\right) e^{-Zr/2a_0}, \quad (11.1.36)$$

$$R_{2,1}(r) = \left(\frac{Z}{a_0}\right)^{3/2} \frac{1}{2\sqrt{6}} \frac{Zr}{a_0} e^{-Zr/2a_0}, \quad (11.1.37)$$

and for the second excited states,

$$R_{3,0}(r) = \left(\frac{Z}{a_0}\right)^{3/2} \frac{2}{81\sqrt{3}} \left[27 - 18\frac{Zr}{a_0} + 2\left(\frac{Zr}{a_0}\right)^2\right] e^{-Zr/3a_0}, \quad (11.1.38)$$

$$R_{3,1}(r) = \left(\frac{Z}{a_0}\right)^{3/2} \frac{2\sqrt{2}}{81\sqrt{3}} \left[6\frac{Zr}{a_0} - \left(\frac{Zr}{a_0}\right)^2\right] e^{-Zr/3a_0}, \quad (11.1.39)$$

$$R_{3,2}(r) = \left(\frac{Z}{a_0}\right)^{3/2} \frac{2\sqrt{2}}{81\sqrt{15}} \left(\frac{Zr}{a_0}\right)^2 e^{-Zr/3a_0}. \quad (11.1.40)$$

These radial wavefunctions, and the corresponding radial probability densities  $r^2|R_{n,l}|^2$ , are graphed in Figure 11.1.2 for  $Z = 1$ . For future reference, we also note that the hydrogen atom radial wavefunction at  $r = 0$  can be evaluated, using eqs. (11.1.28), 11.1.31), and (11.1.32), as

$$R_{n,l}(0) = 2\left(\frac{Z}{na_0}\right)^{3/2} \delta_{l0}. \quad (11.1.41)$$

In particular, it is only non-zero for states with orbital angular momentum  $l = 0$ .

Putting in the spherical harmonics with the Condon–Shortley phase convention, the wavefunctions for the lowest few energy levels are

$$\psi_{1,0,0} = \left(\frac{Z}{a_0}\right)^{3/2} \frac{1}{\sqrt{\pi}} e^{-Zr/a_0}, \quad (11.1.42)$$




Figure 11.1.2: Radial wavefunctions  $R_{n,l}$  in units of  $1/a_0^{3/2}$  (left column) and radial probability densities  $r^2|R_{n,l}|^2$  in units of  $1/a_0$  (right column) for the  $Z = 1$  hydrogen atom stationary states with  $n = 1, 2, 3$ , as functions of  $r/a_0$ . Note the differing vertical scales.

for the ground state, and

$$\psi_{2,0,0} = \left(\frac{Z}{a_0}\right)^{3/2} \frac{1}{4\sqrt{2\pi}} \left(2 - \frac{Zr}{a_0}\right) e^{-Zr/2a_0}, \quad (11.1.43)$$

$$\psi_{2,1,0} = \left(\frac{Z}{a_0}\right)^{3/2} \frac{1}{4\sqrt{2\pi}} \frac{Zr}{a_0} e^{-Zr/2a_0} \cos \theta, \quad (11.1.44)$$

$$\psi_{2,1,\pm 1} = \mp \left(\frac{Z}{a_0}\right)^{3/2} \frac{1}{8\sqrt{\pi}} \frac{Zr}{a_0} e^{-Zr/2a_0} \sin \theta e^{\pm i\phi}, \quad (11.1.45)$$

for the first excited states, and

$$\psi_{3,0,0} = \left(\frac{Z}{a_0}\right)^{3/2} \frac{1}{81\sqrt{3\pi}} \left[27 - 18\frac{Zr}{a_0} + 2\left(\frac{Zr}{a_0}\right)^2\right] e^{-Zr/3a_0}, \quad (11.1.46)$$

$$\psi_{3,1,0} = \left(\frac{Z}{a_0}\right)^{3/2} \frac{\sqrt{2}}{81\sqrt{\pi}} \left[6\frac{Zr}{a_0} - \left(\frac{Zr}{a_0}\right)^2\right] e^{-Zr/3a_0} \cos \theta, \quad (11.1.47)$$

$$\psi_{3,1,\pm 1} = \mp \left(\frac{Z}{a_0}\right)^{3/2} \frac{1}{81\sqrt{\pi}} \left[6\frac{Zr}{a_0} - \left(\frac{Zr}{a_0}\right)^2\right] e^{-Zr/3a_0} \sin \theta e^{\pm i\phi}, \quad (11.1.48)$$

$$\psi_{3,2,0} = \left(\frac{Z}{a_0}\right)^{3/2} \frac{1}{81\sqrt{6\pi}} \left(\frac{Zr}{a_0}\right)^2 e^{-Zr/3a_0} (3\cos^2 \theta - 1), \quad (11.1.49)$$

$$\psi_{3,2,\pm 1} = \mp \left(\frac{Z}{a_0}\right)^{3/2} \frac{1}{81\sqrt{\pi}} \left(\frac{Zr}{a_0}\right)^2 e^{-Zr/3a_0} \sin \theta \cos \theta e^{\pm i\phi}, \quad (11.1.50)$$

$$\psi_{3,2,\pm 2} = \left(\frac{Z}{a_0}\right)^{3/2} \frac{1}{162\sqrt{\pi}} \left(\frac{Zr}{a_0}\right)^2 e^{-Zr/3a_0} \sin^2 \theta e^{\pm 2i\phi}, \quad (11.1.51)$$

for the second excited states.

Let us now work out the expectation value of  $R^p$  in the ground state, for integer  $p$ ,

$$\langle 1, 0, 0 | R^p | 1, 0, 0 \rangle = \int_0^\infty dr r^2 \int d\Omega r^p \left(\frac{Z}{a_0}\right)^3 \frac{1}{\pi} e^{-2Zr/a_0}. \quad (11.1.52)$$

Using  $\int d\Omega = 4\pi$ , this evaluates to

$$\langle 1, 0, 0 | R^p | 1, 0, 0 \rangle = \begin{cases} \frac{1}{2}(p+2)! \left(\frac{a_0}{2Z}\right)^p & (p \geq -2), \\ \infty & (p \leq -3). \end{cases} \quad (11.1.53)$$

In particular, plugging in  $p = 0$  gives  $\langle 1, 0, 0 | 1 | 1, 0, 0 \rangle = 1$  (confirming the correct normalization of the wavefunction), while  $p = 1$  gives

$$\langle 1, 0, 0 | R | 1, 0, 0 \rangle = \langle R \rangle = \frac{3a_0}{2Z}, \quad (11.1.54)$$

and  $p = -1$  gives

$$\langle 1, 0, 0 | 1/R | 0, 0, 1 \rangle = Z/a_0. \quad (11.1.55)$$

(Note that this is not equal to  $1/\langle R \rangle$ .) We can also directly compute, for the ground state,

$$\langle 1, 0, 0 | P^2 | 1, 0, 0 \rangle = \int_0^\infty dr r^2 \int d\Omega \psi_{1,0,0}^* (-\hbar^2 \nabla^2) \psi_{1,0,0} = \hbar^2 Z^2 / a_0^2. \quad (11.1.56)$$

As a check, the expectation value of the Hamiltonian in the ground state is

$$\langle H \rangle = \langle P^2 \rangle / 2m_e - Ze^2 \langle 1/R \rangle = -Z^2 e^2 / 2a_0 = -Z^2 \text{ Rydberg}, \quad (11.1.57)$$

in agreement with the energy eigenvalue.

Later, we will want more general expressions for the expectation values  $\langle R^p \rangle$  in arbitrary hydrogen-like atom stationary bound states  $|n, l, m\rangle$ , for various integer powers  $p$ . A helpful tool to find these for all integer  $p \geq -1$ , and for all integer  $p \leq -3$  in terms of the single seed case  $p = -2$ , is the **Kramers–Pasternack recurrence formula**, which relates the expectation values for any three consecutive integer powers of the radial coordinate  $R$ . It is

$$\frac{q+1}{n^2} \langle R^q \rangle - (2q+1) \frac{a_0}{Z} \langle R^{q-1} \rangle + \frac{q}{4} [(2l+1)^2 - q^2] \frac{a_0^2}{Z^2} \langle R^{q-2} \rangle = 0, \quad (11.1.58)$$

valid for all  $q > -2l - 1$ .

The proof of eq. (11.1.58) is far from obvious, but goes as follows. Start from the differential equation for the radial wavefunction  $R_{n,l}(s)$  with  $s = 2Zr/na_0$ , as given in eq. (11.1.5). Multiply by  $4s^{q+3}R'_{n,l} + 2(1-q)s^{q+2}R_{n,l}$  (this is perhaps the most non-obvious part), and integrate with respect to  $s$ . Then, expand the integrand and eliminate all derivatives of  $R_{n,l}$  using the integration-by-parts identities<sup>§</sup>

$$\int_0^\infty ds \frac{d}{ds} (s^{q+3} R_{n,l}'^2) = 0, \quad (11.1.59)$$

$$\int_0^\infty ds \frac{d}{ds} (s^{q+2} R_{n,l} R_{n,l}') = 0, \quad (11.1.60)$$

$$\int_0^\infty ds \frac{d}{ds} (s^p R_{n,l}^2) = 0, \quad (11.1.61)$$

with  $p = q+1$ ,  $q+2$ , and  $q+3$  in the last equation. The result is

$$\int_0^\infty ds \{ (q+1)s^{q+2} - 2(2q+1)ns^{q+1} + q[(2l+1)^2 - q^2] s^q \} R_{n,l}^2 = 0. \quad (11.1.62)$$

Finally, using the change-of-integration-variable relation,

$$\langle R^p \rangle = \int_0^\infty dr r^{p+2} [R_{n,l}(r)]^2 = (na_0/2Z)^{p+3} \int_0^\infty ds s^{p+2} [R_{n,l}(s)]^2, \quad (11.1.63)$$

from eqs. (11.1.4), and (11.1.23), we arrive at eq. (11.1.58).

---

<sup>§</sup>The ' denotes a derivative with respect to  $s$ . The boundary terms in eqs. (11.1.59)–(11.1.61) vanish for  $q > -2l - 1$ , because  $R_{n,l}$  scales like  $s^l$  for  $s \rightarrow 0$ , and like  $e^{-s/2}$  for  $s \rightarrow \infty$ .

Using  $q = 0$  in the Kramers–Pasternack formula of eq. (11.1.58), and  $\langle R^0 \rangle = \langle 1 \rangle = 1$ , we immediately find

$$\langle 1/R \rangle = \frac{Z}{n^2 a_0}, \quad (11.1.64)$$

in agreement with the  $n = 1$  special case in eq. (11.1.55). Then, using  $q = 1$ , one obtains

$$\langle R \rangle = \frac{a_0}{2Z} [3n^2 - l(l+1)]. \quad (11.1.65)$$

Using this as a measure of the “size” of the state  $|n, l, m\rangle$ , we note that it does not depend on the magnetic quantum number  $m$ , and

- shrinks with larger nuclear charge  $Z$ ,
- grows with larger  $n$ , for fixed  $l$ ,
- shrinks with larger  $l$ , for fixed  $n$ .

The last two of these features can be compared visually to the probability density distributions in the right column of Figure 11.1.2.

Applying eq. (11.1.58) with  $q = 2$ , we get

$$\langle R^2 \rangle = \frac{a_0^2}{Z^2} \frac{n^2}{2} [5n^2 - 3l(l+1) + 1]. \quad (11.1.66)$$

One can apply eq. (11.1.58) recursively to find  $\langle R^p \rangle$  for any desired positive integer  $p$ . However, for negative  $p$ , one finds an obstacle, that  $\langle 1/R^2 \rangle$  cannot be determined by the recurrence relation alone. In section 15.6, we will use another method to find

$$\langle 1/R^2 \rangle = \frac{Z^2}{a_0^2 n^3 (l+1/2)}. \quad (11.1.67)$$

With this as a seed, all results for  $\langle R^p \rangle$  with  $p \leq -3$  can then be determined by the recurrence relation. In particular, using  $q = -1$  in eq. (11.1.58) yields<sup>¶</sup>

$$\langle 1/R^3 \rangle = \frac{Z^3}{a_0^3 n^3 l(l+1)(l+1/2)}. \quad (11.1.68)$$

The expectation value of  $P^2$  can also be found by relating it to the Hamiltonian,

$$\langle P^2 \rangle = 2m_e (\langle H \rangle + Ze^2 \langle 1/R \rangle) = m_e \left( \frac{e^2}{2a_0} \right) \frac{2Z^2}{n^2} = \frac{\hbar^2 Z^2}{a_0^2 n^2}, \quad (11.1.69)$$

where we have used the known energy eigenvalues and eq. (11.1.64). This result can also be obtained from the Virial Theorem, eq. (3.6.5).

---

<sup>¶</sup>Note that  $\langle 1/R^3 \rangle$  diverges for  $l = 0$ . More generally,  $\langle R^p \rangle = \int_0^\infty dr r^{2+p} |R_{n,l}(r)|^2$  diverges if  $p \leq -2l - 3$ , because  $|R_{n,l}| \propto r^l$  for small  $r$ .

The characteristic size of the hydrogen atom with  $Z = 1$  is set by the Bohr radius  $a_0 = \hbar^2/m_e e^2$ . This could have been estimated by a dimensional analysis construction from the available quantities in the problem, by requiring it to have units of length, and demanding that it vanish in each of the following three extreme limits: (1) the classical limit  $\hbar \rightarrow 0$ , and (2) the limit that the electron mass  $m_e$  is taken very large, and (3) the limit of large  $e^2$  so that the attractive electrical force between the nucleus and the electron is large.

However, if we add the speed of light  $c$  to our toolbox of quantities, then there is a dimensionless quantity that we can form out of the available parameters in the problem, namely the **fine structure constant**,

$$\alpha = \frac{e^2}{\hbar c} \approx 1/137.036. \quad (11.1.70)$$

So, we can construct another significant length scale, the **reduced<sup>||</sup> Compton wavelength** of the electron,

$$\lambda_e = \alpha a_0 = \frac{\hbar}{m_e c} \approx 3.862 \times 10^{-13} \text{ meters}. \quad (11.1.71)$$

Named for Arthur Compton's studies of the scattering of electrons by high-energy photons, this is the (very small) length scale at which virtual electron-positron pair production becomes important for understanding the energy levels of electron bound states. To see this, note that according to Einstein's famous formula relating energy and mass, the amount of energy needed to make such a pair is  $\Delta E = 2m_e c^2$ . For ultra-relativistic particles, energy is related to momentum by  $\Delta E \sim c\Delta p$ , so  $e^-e^+$  pair production becomes important for  $\Delta p > 2m_e c$ . From the uncertainty principle  $(\Delta x)(\Delta p) \geq \hbar/2$ , confinement of the electron to a size  $\Delta x \sim \hbar/4m_e c \sim \lambda_e/4$  will result in large enough fluctuations  $\Delta p$  to cause virtual  $e^-e^+$  pair production to be an issue. Furthermore, if we call  $v = \sqrt{\langle P^2 \rangle}/m_e$  the order of magnitude of the "velocity" of electrons in a hydrogen-like atom state with principal quantum number  $n$ , then

$$v \sim \frac{Z\hbar}{na_0 m_e} = Z\alpha c/n. \quad (11.1.72)$$

These considerations show that the approximation we have made in this section works only because the fine structure constant is small. That is what ensures  $\lambda_e \ll a_0$  and  $v \ll c$ , so that electron-positron pair production and other relativistic corrections to the Hamiltonian can be neglected, to first approximation. However, for atoms with very large  $Z$ , the speeds of atomic electrons increase, and relativistic effects become important.

A related point is that the binding energy of the hydrogen atom is very small compared to  $m_e c^2 = 0.511 \text{ MeV}$ , the rest energy of the electron. In terms of the fine structure constant,

$$1 \text{ Rydberg} = \frac{e^2}{2a_0} = \frac{e^4 m_e}{2\hbar^2} = \frac{1}{2} \alpha^2 m_e c^2. \quad (11.1.73)$$

---

<sup>||</sup>The ordinary Compton wavelength of the electron is defined as  $\lambda_e = h/m_e c = 2\pi\lambda_e$ .

To estimate (very roughly) the size of the relativistic corrections to the energies, we can take the total classical relativistic energy of an electron with momentum  $p$ , and subtract off the rest energy, then expand in small  $p$ ,

$$E_{\text{relativistic}} = \sqrt{m_e^2 c^4 + p^2 c^2} - m_e c^2 = \frac{p^2}{2m_e} - \frac{p^4}{8m_e^3 c^2} + \cdots, \quad (11.1.74)$$

where we have used the binomial expansion  $\sqrt{1+x} = 1 + x/2 - x^2/8 + \cdots$  for small  $x = p^2/m_e^2 c^2$ . The first term is just the nonrelativistic kinetic energy that we have been using in the quantum Hamiltonian. So, we can estimate that relativistic effects should make a difference of order

$$\Delta E_{\text{relativistic}} \sim -\frac{\langle P^2 \rangle^2}{8m_e^3 c^2} = -\frac{\alpha^2 Z^4}{4n^4} \text{ Rydberg}. \quad (11.1.75)$$

However, it turns out that a correct relativistic analysis must take into account spin. We will carry this out in section 17.1, with a final result in eq. (17.1.26). We will do it again in section 27.5, using a manifestly relativistic approach. Our crude estimate in eq. (11.1.75) is parametrically correct in the sense that the fine structure effects indeed modify the binding energies by amounts that are suppressed by  $\alpha^2$  and by powers of  $n$ , but we will see that the numerical details are quite different than the naive estimate of eq. (11.1.75).

When an electron transitions between states of the hydrogen atom, it releases or absorbs a photon with energy equal to the difference in energy levels,  $\Delta E = E_n - E_{n'}$ . This is equal to  $\hbar\omega$  where  $\omega$  is the angular frequency of the photon. Therefore, taking  $Z = 1$  for the remainder of this section,

$$\omega_{n,n'} = \frac{13.6 \text{ eV}}{\hbar} \left( \frac{1}{n'^2} - \frac{1}{n^2} \right), \quad (11.1.76)$$

or equivalently in terms of wavelength,

$$\lambda_{n,n'} = (9.11 \times 10^{-8} \text{ meters}) \frac{n'^2}{1 - n'^2/n^2}. \quad (11.1.77)$$

This formula was found empirically by Rydberg in 1888, generalizing work by Johann Balmer. For transitions between the ground state  $n' = 1$  and the states with  $n \geq 2$ , these spectral lines are called the Lyman series, after Theodore Lyman. They are all in the ultraviolet range,

$$\text{Lyman series: } \omega_{n,1} = (2.067 \times 10^{16} \text{ s}^{-1}) \left( 1 - \frac{1}{n^2} \right), \quad (n = 2, 3, 4, \dots, \infty), \quad (11.1.78)$$

with wavelengths from  $\lambda_{2,1} = 1.216 \times 10^{-7}$  meters to  $\lambda_{\infty,1} = 9.11 \times 10^{-8}$  meters. The latter wavelength, corresponding to a photon emitted when a free electron is captured to the ground state of a hydrogen atom, or absorbed in the process of ionizing a hydrogen atom that was



initially in its ground state, is called the Lyman limit. The spectral line with  $n' = 1$  and  $n = 2$  is called the Lyman-alpha line, and is important in astronomy as an absorption feature when observing distant sources, for example as a tool to learn about intervening gas.

The Balmer series comes from transitions between  $n' = 2$  and all higher-energy states  $n \geq 3$ ,

$$\text{Balmer series: } \omega_{n,2} = (5.168 \times 10^{15} \text{ s}^{-1}) \left(1 - \frac{4}{n^2}\right), \quad (n = 3, 4, 5, \dots, \infty), \quad (11.1.79)$$

with wavelengths ranging from  $\lambda_{3,2} = 6.56 \times 10^{-7}$  meters to  $\lambda_{\infty,2} = 3.97 \times 10^{-7}$  meters. The Balmer lines are particularly interesting because four of them can be seen by the human eye. For  $n = 3$  the Balmer line is red, which is responsible for the reddish color of supernova remnants and star-forming regions, including the Orion nebula. The  $n = 4, 5$ , and  $6$  Balmer lines are greenish-blue, blue, and violet, respectively, while  $n = 7, \dots, \infty$  are in the ultraviolet. The remaining named series are all entirely in the infrared, and are called Paschen (transitions between  $n' = 3$  and  $n \geq 4$ ), Brackett (transitions between  $n' = 4$  and  $n \geq 5$ ), Pfund (transitions between  $n' = 5$  and  $n \geq 6$ ), and Humphreys (transitions between  $n' = 6$  and  $n \geq 7$ ), with higher  $n'$  series not named. There is significant overlap between the infrared series of spectral lines. Within each series, the spectral lines are called  $\alpha, \beta$ , etc. in order of decreasing wavelength.

## 11.2 Unbound states of Coulomb potentials

We now consider the unbound stationary states of the Coulomb problem, with  $E \geq 0$ . These include the ionized states of the hydrogen-like atom, describing an electron that is influenced by the nucleus but not localized near it. The electron's wavefunction can again be split into the product of a radial wavefunction and an angular part consisting of a spherical harmonic,

$$\langle \vec{r} | \Phi_{E,l,m} \rangle = \Phi_{E,l,m}(r, \theta, \phi) = \tilde{R}_{E,l}(r) Y_l^m(\theta, \phi) \quad (E \geq 0). \quad (11.2.1)$$

We write  $\Phi_{E,l,m}$  and  $\tilde{R}_{E,l}$  here to distinguish them from the bound-state wavefunctions  $\psi_{n,l,m}$  and  $R_{n,l}$  of the preceding section. For the unbound states, each  $E \geq 0$  can have any non-negative integer value of  $l$ , as depicted in Fig. 11.1.1.

The differential equation satisfied by the radial wavefunction for unbound states is the same as eq. (11.1.5), with the important differences that the rescaled radial coordinate  $s$  and the quantity  $n$ , defined in eqs. (11.1.4) and (11.1.6) respectively, are imaginary for  $E > 0$ . Accordingly, we make the replacement<sup>†</sup>

$$n \rightarrow \frac{i}{ka_0}, \quad (11.2.2)$$

---

<sup>†</sup>In this section, we set  $\mu = m_e$  and  $Z = 1$  for simplicity, with the understanding that general  $Z$  can always be restored by the replacement  $a_0 \rightarrow a_0/Z$ .

which defines a continuous real wavenumber parameter  $k$  with units of  $1/[\text{length}]$ . Comparing to the definition of  $n$  in eq. (11.1.6) gives

$$ka_0 = \sqrt{\frac{2\hbar^2 E}{m_e e^4}}, \quad (11.2.3)$$

or, equivalently,

$$E = \frac{1}{2}(ka_0)^2 \text{ Rydberg}. \quad (11.2.4)$$

Then, from eq. (11.1.4) the rescaled radial coordinate used in the previous section becomes

$$s = 2ikr. \quad (11.2.5)$$

It follows that, up to normalization, we can obtain the solution in terms of the confluent hypergeometric function defined in eq. (11.1.10), by simply re-using eq. (11.1.13) with the preceding substitutions for  $n$  and  $s$ . Since  $s$  is now imaginary, there is no problem with having an asymptotic behavior like  $e^{s/2}$  for large  $s$ , and therefore no restriction that the confluent hypergeometric function must be a polynomial; that is why there is no discrete quantization of  $k$ . The result is

$$\tilde{R}_{E,l}(r) = C_{k,l} (2kr)^l e^{-ikr} F(l+1 + \frac{i}{ka_0}, 2l+2, 2ikr), \quad (11.2.6)$$

where  $C_{k,l}$  is a normalization constant to be determined, which we will choose to be real.

Despite the appearance of the phase factor  $e^{-ikr}$  and the complex arguments of the confluent hypergeometric function,  $\tilde{R}_{E,l}(r)$  as given in eq. (11.2.6) is real. This follows immediately from a property of the confluent hypergeometric function,

$$F(a, c, z) = e^z F(c - a, c, -z), \quad (11.2.7)$$

which can in turn be proved quickly from the integral representation of eq. (11.1.11) by using the change of integration variable  $t \rightarrow 1 - t$ .

Let us now consider the behavior of the radial wavefunction  $\tilde{R}_{E,l}$  for large  $r$ . This can be done by using the asymptotic form for the confluent hypergeometric function for complex arguments,

$$F(a, c, z) \approx \frac{\Gamma(c)}{\Gamma(c-a)} (-z)^{-a} + \frac{\Gamma(c)}{\Gamma(a)} e^z z^{a-c} \quad (\text{large } |z|), \quad (11.2.8)$$

which generalizes eq. (11.1.12) for real arguments. The  $\Gamma$  function for complex arguments was defined in eq. (10.5.35). Applying this to eq. (11.2.6), the two terms turn out to give contributions that are complex conjugates of each other, as required by the previous paragraph. After some simplification, one finds the asymptotic form

$$\tilde{R}_{E,l}(r) \approx C_{k,l} \frac{(2l+1)! e^{-\pi/2ka_0}}{|\Gamma(l+1 + i/ka_0)|} \frac{1}{kr} \sin\left(kr + \frac{1}{ka_0} \ln(2kr) - \frac{\pi}{2}l - \delta\right) \quad (11.2.9)$$

for large  $kr$ , where

$$\delta = \text{Arg}[\Gamma(l+1+i/ka_0)]. \quad (11.2.10)$$

The radial probability density  $r\tilde{R}_{E,l}(r)$  approaches a sinusoidal oscillation with a constant amplitude at large distances, but with a phase shift that depends logarithmically on  $kr$ .

In order to determine the normalization constant, we first define the orthonormality properties of the states. Since the energy  $E \geq 0$  is continuous, we will use a Dirac orthonormality condition for unbound states  $|E, l, m\rangle$ , along with the ordinary discrete orthonormality of the bound states  $|n, l, m\rangle$  of the previous section. So,

$$\langle E', l', m' | E, l, m \rangle = \delta(E - E') \delta_{ll'} \delta_{mm'}, \quad (11.2.11)$$

$$\langle n', l', m' | E, l, m \rangle = 0, \quad (11.2.12)$$

$$\langle n', l', m' | n, l, m \rangle = \delta_{nn'} \delta_{ll'} \delta_{mm'}. \quad (11.2.13)$$

In terms of the radial wavefunctions for each  $l$ , these become

$$\int_0^\infty dr r^2 \tilde{R}_{E',l}(r) \tilde{R}_{E,l}(r) = \delta(E - E'), \quad (11.2.14)$$

$$\int_0^\infty dr r^2 \tilde{R}_{E',l}(r) R_{n,l}(r) = 0, \quad (11.2.15)$$

$$\int_0^\infty dr r^2 R_{n',l}(r) R_{n,l}(r) = \delta_{nn'}. \quad (11.2.16)$$

Now, for the purposes of normalizing the radial wavefunction in eq. (11.2.9), note that when  $E = E'$ , one need only consider the asymptotic form at very large  $r$ , because contributions to  $\int_0^\infty dr r^2 |\tilde{R}_{E,l}(r)|^2$  from any finite range in  $r$  contribute only an infinitesimal fraction of the total. In the very large  $r$  limit, the logarithmic variation in the phase shift can be neglected, and the normalization problem is the same as for the simpler wavefunctions

$$f_k(r) = A \frac{\sin(kr + \beta)}{kr}, \quad (11.2.17)$$

for some phase shift  $\beta$ , where

$$A = C_{k,l} \frac{(2l+1)! e^{-\pi/2ka_0}}{|\Gamma(l+1+i/ka_0)|}. \quad (11.2.18)$$

The norm of this wavefunction is infinite, but integrating over a finite range gives

$$\int_0^D dr r^2 f_{k'}(r) f_k(r) = \frac{A^2}{2kk'} \left\{ \frac{\sin[(k-k')D]}{k-k'} + \frac{\sin[2\beta] - \sin[2\beta + (k+k')D]}{k+k'} \right\}. \quad (11.2.19)$$

In the formal limit  $D \rightarrow \infty$ , one can interpret the first term in the braces as a delta function distribution [see eq. (2.2.22)], while the remaining part remains bounded for all  $k$  and  $k'$ , and

vanishes for an infinite number of choices  $D = 2\pi q/(k + k')$ , for arbitrarily large integers  $q$ . This allows us to interpret, when inserted in any expression in which  $k$  or  $k'$  is integrated over,

$$\int_0^\infty dr r^2 f_{k'}(r) f_k(r) = A^2 \frac{\pi}{2k^2} \delta(k - k') = A^2 \frac{\pi \hbar^2}{4m_e k} \delta(E - E'). \quad (11.2.20)$$

Thus we can adopt the energy normalization for Dirac orthonormality, by taking  $A^2 = 4m_e k/\pi \hbar^2$ , allowing us to solve eq. (11.2.18) for  $C_{k,l}$ . Using this in eq. (11.2.6), the final result for the unbound energy eigenstate state radial wavefunction is

$$\tilde{R}_{E,l}(r) = \frac{2}{\hbar} \sqrt{\frac{m_e k}{\pi}} \frac{|\Gamma(l+1+i/ka_0)| e^{\pi/2ka_0}}{(2l+1)!} (2kr)^l e^{-ikr} F(l+1+\frac{i}{ka_0}, 2l+2, 2ikr), \quad (11.2.21)$$

where  $k$  is related to  $E$  by eq. (11.2.3) or eq. (11.2.4). Although this is not the simplest result one might have hoped for, we again remark that at least it is real, despite naive appearances.

The completeness relation corresponding to the energy eigenstates of eqs. (11.2.11)-(11.2.13) contains both a sum over bound states and an integral over unbound states:

$$\sum_{n=1}^{\infty} \sum_{l=0}^{n-1} \sum_{m=-l}^l |n, l, m\rangle \langle n, l, m| + \int_0^\infty dE \sum_{l=0}^{\infty} \sum_{m=-l}^l |E, l, m\rangle \langle E, l, m| = I. \quad (11.2.22)$$

This combination of ordinary and Dirac orthonormality and completeness will be crucial in the evaluation of the ground state energy of the hydrogen atom in an electric field (the quadratic Stark effect), in section 15.7.

For a general potential  $V(r) = -Ze^2/r$ , the results above can be obtained by making the replacement  $a_0 \rightarrow a_0/Z$  everywhere in the preceding discussion. This includes the case of a repulsive potential with  $Z < 0$ , but with one qualitative difference: the bound states  $|n, l, m\rangle$  do not exist, and the orthonormality and completeness relations therefore include only the unbound states with continuous positive  $E$ .

### 11.3 Exercises

**Exercise 11.1.** In this problem, we will reconstruct the Coulomb potential from the functional form of its ground-state wavefunction. Suppose that a particle in a spherical potential  $V(r)$  has a stationary state wavefunction  $\psi_E(r, \theta, \phi) = Ne^{-r/b}$ , where  $N$  and  $b$  are constants.

(a) Assume that  $V(r)$  vanishes as  $r \rightarrow \infty$ . Use this to find the energy eigenvalue  $E$ , by matching leading terms in Schrödinger's equation as  $r \rightarrow \infty$ .

(b) Now that you have found  $E$ , consider finite  $r$  and find the potential  $V(r)$ .

**Exercise 11.2.** For the ground state of the hydrogen atom, what would be the maximum allowed classical distance of the electron from the proton, if the energy is 1 Rydberg? Compute the probability for the electron to be found farther away from the proton than this distance.

**Exercise 11.3.** For the ground state of the hydrogen atom, find the uncertainties of the rectangular coordinate  $Z$  and its momentum component  $P_z$ . Check that the uncertainty principle is satisfied by your answers.

**Exercise 11.4.** For the ground state and the first excited states of the hydrogen atom, find all of the matrix elements of the rectangular coordinate operators for every pair of such states,  $\langle n', l', m' | X | n, l, m \rangle$  and  $\langle n', l', m' | Y | n, l, m \rangle$  and  $\langle n', l', m' | Z | n, l, m \rangle$ , with  $n = 1, 2$  and  $n' = 1, 2$ . (Hint: many of them are zero.)

**Exercise 11.5.** Use the Kramers–Pasternack formula (11.1.58) to derive the expectation values  $\langle n, l, m | R^p | n, l, m \rangle$  for all integers  $-3 \leq p \leq 3$ . As seeds, you may use the obvious fact  $\langle 1 \rangle = 1$ , and also  $\langle 1/R^2 \rangle = 2/(a_0^2 n^3 (2l + 1))$ , which will be found in section 15.6 using another trick.

**Exercise 11.6.** (a) For the ground state of the hydrogen atom, by taking the Fourier transform as in eq. (2.8.64), show that the momentum wavefunction is

$$\tilde{\psi}(\vec{p}) = \frac{C}{(p^2 + \hbar^2/a_0^2)^N}, \quad (11.3.1)$$

where  $N$  is an integer and  $C$  is a normalization factor, both of which you will discover. [Hint: since the ground state is spherically symmetric, the momentum wavefunction cannot depend on the direction of  $\vec{p}$ . So, for the purposes of computing the integral, you can take  $\vec{p} = p\hat{z}$ .]

(b) Use your result from the previous part to compute  $\langle P^2 \rangle$  and  $\langle (P^2)^2 \rangle$ . The first of these should agree with the  $n = 1$ ,  $Z = 1$  special case of eq. (11.1.69).

(c) Taking the result of part (a) at face value (ignoring the reality of relativity), compute numerically the probability for the magnitude of the electron's momentum to exceed  $m_e c = \hbar/\alpha a_0$ , where  $m_e$  is the electron's mass.

**Exercise 11.7.** A particle of mass  $\mu$  moves in the potential  $V(R) = -a/R + \hbar^2 b/2\mu R^2$ , where  $a$  and  $b$  are positive constants. Consider stationary-state wavefunctions of the form  $\psi(\vec{r}) = R(r)Y_l^m(\theta, \phi)$ . Find the bound-state wavefunctions and energy eigenvalues for orbital angular momentum  $l$ . (Hint: note that the radial effective potential has the same form as for the hydrogen atom, but with a modification of the angular momentum contribution.)

## 12 Addition of angular momenta

### 12.1 Statement of the problem

In quantum mechanical problems, we often deal with two or more distinct angular momenta. The sum of two independent angular momentum operators,

$$\vec{J} = \vec{J}_1 + \vec{J}_2, \quad (12.1.1)$$

is also an angular momentum operator. This expresses the fact that if all components of  $\vec{J}_1$  commute with all components of  $\vec{J}_2$ , and if the components of  $\vec{J}_1$  and  $\vec{J}_2$  each satisfy the commutator algebra (5.3.33), then so will the components of  $\vec{J}$ .

Since  $J_1^2$ ,  $J_2^2$ ,  $J_{1z}$ , and  $J_{2z}$  are compatible operators, we can find an orthobasis of common eigenkets for them, labeled

$$|j_1 j_2 m_1 m_2\rangle \equiv |j_1 m_1\rangle \otimes |j_2 m_2\rangle, \quad (12.1.2)$$

with eigenvalues  $\hbar^2 j_1(j_1+1)$ ,  $\hbar^2 j_2(j_2+1)$ ,  $\hbar m_1$ , and  $\hbar m_2$ , respectively. We call this the **product orthobasis** for two angular momenta, because its elements consist of the tensor products of elements of the orthobases for the individual angular momenta. Here we have suppressed any degeneracy labels, which might correspond to different radial wavefunctions, for example. For fixed  $j_1$  and  $j_2$ , there are  $2j_1 + 1$  allowed values of  $m_1$ , and  $2j_2 + 1$  allowed values of  $m_2$ ,

$$m_1 = j_1, j_1 - 1, \dots, -j_1 + 1, -j_1, \quad (12.1.3)$$

$$m_2 = j_2, j_2 - 1, \dots, -j_2 + 1, -j_2. \quad (12.1.4)$$

So, there are  $(2j_1 + 1)(2j_2 + 1)$  orthobasis kets of the form eq. (12.1.2) for fixed  $j_1$  and  $j_2$ .

Another set of compatible operators is  $J_1^2$ ,  $J_2^2$ ,  $J^2$ , and  $J_z$ . They also have a set of common eigenkets that form a different orthobasis, whose kets we can write as

$$|j_1 j_2 j m\rangle. \quad (12.1.5)$$

This is called the **total angular momentum orthobasis**. As we will see in the following sections, for fixed  $j_1$  and  $j_2$ , the possible values of  $j$  range from a maximum of  $j_1 + j_2$  to a minimum of  $|j_1 - j_2|$ , with integer increments,

$$j = j_1 + j_2, j_1 + j_2 - 1, \dots, |j_1 - j_2| + 1, |j_1 - j_2|. \quad (12.1.6)$$

Intuitively, the extremes for  $j$  correspond to the two angular momenta aligned in the same and in opposite directions, respectively. Then, for each  $j$ , there are  $2j + 1$  allowed values

$$m = j, j - 1, \dots, -j + 1, -j. \quad (12.1.7)$$

As a check, the number of total angular momentum basis states for fixed  $j_1$  and  $j_2$  is

$$\sum_{j=|j_1-j_2|}^{j_1+j_2} (2j+1) = (2j_1+1)(2j_2+1), \quad (12.1.8)$$

matching the result for the number of product basis kets.

There is a potentially annoying problem of notation to be addressed here, because both orthobases are labeled by four numbers. In many cases, this will not cause confusion, but what if it does? Our solution to this problem will be to use a colon to separate the last two (magnetic quantum number) labels for the product angular momentum orthobasis whenever there is a chance of confusion. All other labels in both the product and total angular momentum bases are separated by no punctuation, or by a comma when it is typographically convenient (or just suits our mood). Also, there will be many occasions in which the labels  $j_1$  and  $j_2$  that are common to both orthobases are fixed, and understood from context. In that case, we can suppress those labels and call the total angular momentum orthobasis kets  $|jm\rangle$  or  $|j, m\rangle$ , and the product orthobasis kets  $|m_1 m_2\rangle$  or  $|m_1, m_2\rangle$ , but use the notation  $|m_1 : m_2\rangle$  if there is a significant chance of confusion with the total angular momentum orthobasis.

A frequently encountered situation is that we might have to evaluate matrix elements involving the dot product of two angular momentum operators. If we are working in the product basis, the most straightforward way to do this is to write the dot product in terms of raising and lowering operators,

$$\vec{J}_1 \cdot \vec{J}_2 = \frac{1}{2} (J_{1+} J_{2-} + J_{1-} J_{2+}) + J_{1z} J_{2z}, \quad (12.1.9)$$

which follows from eq. (8.1.10). Then each of  $J_{1+}$ ,  $J_{1-}$ , and  $J_{1z}$  can be evaluated by their actions on the  $|j_1 m_1\rangle$  component using the rules of eqs. (8.1.7), (8.1.22), (8.1.24), and similarly for the actions of  $J_{2+}$ ,  $J_{2-}$ , and  $J_{2z}$  on the  $|j_2 m_2\rangle$  part. However, it is often more convenient to use instead the total angular momentum basis. This is because one can use  $J^2 = (\vec{J}_1 + \vec{J}_2)^2 = J_1^2 + J_2^2 + 2\vec{J}_1 \cdot \vec{J}_2$  to write

$$\vec{J}_1 \cdot \vec{J}_2 = \frac{1}{2} (J^2 - J_1^2 - J_2^2). \quad (12.1.10)$$

Now, when acting on the total angular momentum basis kets  $|j_1 j_2 j m\rangle$ , this operator just evaluates to a number,

$$\vec{J}_1 \cdot \vec{J}_2 = \frac{\hbar^2}{2} [j(j+1) - j_1(j_1+1) - j_2(j_2+1)], \quad (12.1.11)$$

independent of  $m$ . Solving problems in quantum mechanics is often the art of turning operators into numbers, and using the correct choice of basis can make things easier. The specific lesson

here is that the appearance of a dot product of angular momenta should immediately suggest using the total angular momentum basis.

Because they are both bases, each element of the total angular momentum orthobasis must be a linear combination of the product orthobasis kets, and vice versa. In general, the problem of **addition of angular momenta** is to evaluate the coefficients appearing in these linear combinations.

We will start with two useful and common special cases before taking on the most general case. First, in section 12.2, we consider the case of two spins  $s_1 = s_2 = 1/2$ , for two particles with orbital angular momenta absent or disregarded. Next, in section 12.3, we will take up the example of a single particle with arbitrary orbital angular momentum  $l$  and spin  $s = 1/2$ , which can be combined to form the total angular momentum of the particle. In section 12.4 we will discuss the general case of addition of two arbitrary angular momenta. These results can be applied recursively to combine any number of angular momenta.

## 12.2 Addition of two spins

Consider two spins with  $s_1 = s_2 = 1/2$ . Spin magnitudes are always fixed, so we suppress those labels, and write the four product orthobasis kets as

$$|\uparrow\uparrow\rangle, \quad |\uparrow\downarrow\rangle, \quad |\downarrow\uparrow\rangle, \quad |\downarrow\downarrow\rangle. \quad (12.2.1)$$

Here the first  $\uparrow$  or  $\downarrow$  label on each ket stands for the eigenvalue  $\hbar/2$  or  $-\hbar/2$  of  $S_{1z}$ , and the second label similarly stands for the eigenvalue of  $S_{2z}$ . We say that the individual spins in this basis are either “up” or “down” relative to our choice of the  $\hat{z}$  direction. Now, define the total spin operator by

$$\vec{S} = \vec{S}_1 + \vec{S}_2. \quad (12.2.2)$$

The product orthobasis kets are also eigenkets of  $S_z$ , since

$$S_z |m_1 m_2\rangle = S_{1z} |m_1 m_2\rangle + S_{2z} |m_1 m_2\rangle = \hbar(m_1 + m_2) |m_1 m_2\rangle, \quad (12.2.3)$$

so that  $S_z |\uparrow\uparrow\rangle = \hbar |\uparrow\uparrow\rangle$ , and  $S_z |\uparrow\downarrow\rangle = S_z |\downarrow\uparrow\rangle = 0$ , and  $S_z |\downarrow\downarrow\rangle = -\hbar |\downarrow\downarrow\rangle$ . Choosing a representation in which

$$|\uparrow\uparrow\rangle \leftrightarrow \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad |\uparrow\downarrow\rangle \leftrightarrow \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \quad |\downarrow\uparrow\rangle \leftrightarrow \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \quad |\downarrow\downarrow\rangle \leftrightarrow \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \quad (12.2.4)$$



we have the matrix representation

$$S_z \leftrightarrow \hbar \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}. \quad (12.2.5)$$

However, the product orthobasis kets are not all eigenstates of  $S^2$ . The operation of  $S^2$  on them can be obtained from

$$\begin{aligned} S^2 &= (\vec{S}_1 + \vec{S}_2) \cdot (\vec{S}_1 + \vec{S}_2) = S_1^2 + S_2^2 + 2\vec{S}_1 \cdot \vec{S}_2 \\ &= \frac{3}{2}\hbar^2 + S_{1+}S_{2-} + S_{1-}S_{2+} + 2S_{1z}S_{2z}, \end{aligned} \quad (12.2.6)$$

where in the last equality we have taken advantage of the fact that all of the kets in the state space under consideration are eigenkets of both  $S_1^2$  and  $S_2^2$  with eigenvalues  $3\hbar^2/4$ , and then applied eq. (8.1.10) to write the result in terms of the angular momentum raising and lowering operators. Now, recall from eqs. (8.1.22) and (8.1.24) that, for any  $j = 1/2$  system,

$$J_+ |\uparrow\rangle = 0, \quad J_+ |\downarrow\rangle = \hbar |\uparrow\rangle, \quad (12.2.7)$$

$$J_- |\downarrow\rangle = 0, \quad J_- |\uparrow\rangle = \hbar |\downarrow\rangle. \quad (12.2.8)$$

Applying these for each of  $J = S_1$  and  $S_2$ , we obtain from eq. (12.2.6),

$$S^2 |\uparrow\uparrow\rangle = 2\hbar^2 |\uparrow\uparrow\rangle, \quad (12.2.9)$$

$$S^2 |\uparrow\downarrow\rangle = \hbar^2 |\uparrow\downarrow\rangle + \hbar^2 |\downarrow\uparrow\rangle, \quad (12.2.10)$$

$$S^2 |\downarrow\uparrow\rangle = \hbar^2 |\uparrow\downarrow\rangle + \hbar^2 |\downarrow\uparrow\rangle, \quad (12.2.11)$$

$$S^2 |\downarrow\downarrow\rangle = 2\hbar^2 |\downarrow\downarrow\rangle. \quad (12.2.12)$$

In matrix representation form, this reads

$$S^2 \leftrightarrow \hbar^2 \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix}. \quad (12.2.13)$$

This matrix has eigenvalues  $2\hbar^2$ ,  $2\hbar^2$ ,  $2\hbar^2$ , and 0, which correspond to  $S^2 = \hbar^2 s(s+1)$  with  $s = 1$  and  $s = 0$ . The addition of two spins with  $s_1 = s_2 = 1/2$  thus gives a triplet of states  $|1, m\rangle$  with total spin  $s = 1$  and  $m = -1, 0, 1$ , plus a singlet state with total spin  $s = 0$ , denoted  $|0, 0\rangle$ . By finding the eigenkets corresponding to the eigenvalue pairs  $(s, m)$ , we can summarize

the information about the addition of two spin-1/2 systems as

$$\begin{array}{ll} \text{total spin basis } |s, m\rangle & \text{product basis } |m_1 m_2\rangle \\ |1, 1\rangle & = |\uparrow\uparrow\rangle, \end{array} \quad (12.2.14)$$

$$|1, 0\rangle = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle), \quad (12.2.15)$$

$$|1, -1\rangle = |\downarrow\downarrow\rangle, \quad (12.2.16)$$

$$|0, 0\rangle = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle). \quad (12.2.17)$$

Both orthobases have 4 members, and we can of course invert the relationship for the  $m = 0$  states, to find

$$|\uparrow\downarrow\rangle = \frac{1}{\sqrt{2}}(|1, 0\rangle + |0, 0\rangle), \quad (12.2.18)$$

$$|\downarrow\uparrow\rangle = \frac{1}{\sqrt{2}}(|1, 0\rangle - |0, 0\rangle). \quad (12.2.19)$$

As a way of expressing the fact that the tensor product of two spin-1/2 systems gives a sum of angular momenta 0 and 1, we write

$$\frac{1}{2} \otimes \frac{1}{2} = 0_A \oplus 1_S, \quad (12.2.20)$$

Here, the  $\otimes$  represents the addition of angular momenta, while the  $\oplus$  indicates the combinations of total angular momenta. The subscripts  $A$  and  $S$  in this notation are a reminder that the  $s = 0$  singlet state is antisymmetric under exchange of the two spins, while each of the  $s = 1$ ,  $m = 1, 0, -1$  triplet states is symmetric, as can be seen in eqs. (12.2.14)-(12.2.17).

An important practical question is: which orthobasis should we use? The answer depends on the problem under consideration. As a general rule, it is convenient to use a basis in which the operators most important to us are diagonal. For example, suppose that we have a Hamiltonian of the form

$$H = b_1 S_{1z} + b_2 S_{2z}, \quad (12.2.21)$$

which could occur if both spins are interacting with a magnetic field, but not with each other. This Hamiltonian is diagonal in the product basis, where the kets are already eigenvalues of  $S_{1z}$  and  $S_{2z}$ , but it is diagonal in the total angular momentum basis only if  $b_1 = b_2$ . On the other hand, suppose that our Hamiltonian is of the form

$$H = a \vec{S}_1 \cdot \vec{S}_2, \quad (12.2.22)$$

which comes from the magnetic moments of the particles interacting with each other. In that case, we can use the standard trick of writing

$$\vec{S}_1 \cdot \vec{S}_2 = \frac{1}{2}(S^2 - S_1^2 - S_2^2). \quad (12.2.23)$$

The right side shows that this operator is diagonal in the total spin basis, where it evaluates to

$$\vec{S}_1 \cdot \vec{S}_2 = \frac{\hbar^2}{2} [s(s+1) - 3/2], \quad (12.2.24)$$

with  $s = 1$  when acting on the triplet states and  $s = 0$  for the singlet state. In other cases, a judgment can be made about which is the most convenient basis, often based on which part of the Hamiltonian, or some other observable of interest, is the most important.

As a classic example, the **hyperfine** splitting of the hydrogen atom ground state arises from the interaction of the electron's spin with the spin of the proton. The hyperfine Hamiltonian  $H_{\text{hf}}$  has exactly the form of eq. (12.2.22), with a constant  $a$  that is positive, and very small compared to the Rydberg energy scale divided by  $\hbar^2$ . Before taking into account fine or hyperfine effects, the hydrogen atom energy eigenstates can be given in the product spin basis labeled as

$$| \underbrace{n, l, m_l}_{\text{orbital}}, \overbrace{s_e, m_{s_e}}^{\text{electron spin}}, \underbrace{s_p, m_{s_p}}_{\text{proton spin}} \rangle = | n, l, m_l, m_{s_e}, m_{s_p} \rangle, \quad (12.2.25)$$

or we can use the total spin basis,

$$| n, l, m_l, s_e, s_p, s, m_s \rangle = | n, l, m_l, s, m_s \rangle. \quad (12.2.26)$$

In both cases, the individual spin labels  $s_e, s_p = 1/2$  are completely fixed and therefore can be suppressed, as indicated. Our present interest is the effect on the ground state with  $n = 1$  and  $l = 0$ . Since  $H_{\text{orbital}} = P^2/2m_e - e^2/R$  trivially commutes with

$$H_{\text{hf}} = a \vec{S}_e \cdot \vec{S}_p, \quad (12.2.27)$$

they have an orthobasis of common eigenstates. The eigenvalues of  $H_{\text{hf}}$  are found immediately in the total spin basis, using eq. (12.2.24), as

$$E_{\text{hf}} = \begin{cases} a\hbar^2/4 & (s = 1), \\ -3a\hbar^2/4 & (s = 0), \end{cases} \quad (12.2.28)$$

for the triplet and singlet total spin states, respectively. The energy splitting between these states is therefore  $a\hbar^2$ , which for the ground state of the hydrogen atom is, numerically,

$$\Delta E_{\text{hf}} = a\hbar^2 = 5.87 \times 10^{-6} \text{ eV}, \quad (12.2.29)$$

corresponding to a wavelength  $\lambda = 2\pi c/a\hbar = 0.211$  meters. This is the famous 21 centimeter line of radio astronomy. The rate for transitions between the  $s = 1$  and  $s = 0$  states turns out to be highly suppressed (for reasons to be discussed in section 22.5), but space is big and mostly cold, so it is a very useful observational tool in astrophysics and experimental cosmology.

There is a special consideration when the two spin-1/2 particles are identical. The **Pauli exclusion principle** says that two identical particles with half-integer spin cannot be in the same quantum state, and more generally that the quantum state describing two identical particles must be antisymmetric under their exchange. Such particles are called **fermions**, after Enrico Fermi, while particles with integer spin are called **bosons** after Satyendra Nath Bose. For example, suppose that the orbital wavefunctions of two spin-1/2 fermions are  $\psi_a(\vec{r})$  and  $\psi_b(\vec{r})$ , corresponding to single-particle kets (neglecting spins for the moment)  $|\psi_a\rangle$  and  $|\psi_b\rangle$ . From these, one can form symmetric and antisymmetric combinations

$$|\psi_a, \psi_b\rangle_S = \frac{1}{\sqrt{2}}(|\psi_a, \psi_b\rangle + |\psi_b, \psi_a\rangle), \quad (12.2.30)$$

$$|\psi_a, \psi_b\rangle_A = \frac{1}{\sqrt{2}}(|\psi_a, \psi_b\rangle - |\psi_b, \psi_a\rangle), \quad (12.2.31)$$

where, on the right-hand sides, the first and second entries in each ket correspond to particle labels 1 and 2, respectively. Then the allowed states constructed from the tensor product of the orbital and total spin states must be either symmetric in orbital kets and antisymmetric in spin kets, or vice versa,

$$|\psi_a, \psi_b\rangle_S \otimes \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) \quad (s = 0), \quad (12.2.32)$$

$$|\psi_a, \psi_b\rangle_A \otimes |\uparrow\uparrow\rangle \quad (s = 1, m_s = 1), \quad (12.2.33)$$

$$|\psi_a, \psi_b\rangle_A \otimes \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle) \quad (s = 1, m_s = 0), \quad (12.2.34)$$

$$|\psi_a, \psi_b\rangle_A \otimes |\downarrow\downarrow\rangle \quad (s = 1, m_s = -1). \quad (12.2.35)$$

Other kets, such as  $|\psi_a, \psi_b\rangle_S \otimes |\uparrow\uparrow\rangle$ , do not exist as physical states. For example, the ground state of helium, with two electrons, has an orbital part that is symmetric under interchange of the positions of the two electrons. The spin state is therefore the antisymmetric total spin  $s = 0$  combination.

## 12.3 Addition of orbital angular momentum and spin

In this section, we consider the combination of two angular momenta  $j_1$  and  $j_2$ , with  $j_1$  arbitrary and  $j_2 = 1/2$ . This applies, for example, to the case where we are combining the orbital  $\vec{L}$  and intrinsic  $\vec{S}$  angular momenta for a single spin-1/2 particle such as the electron. We will frame our discussion in that context.

As we saw in section 8.2, the spin states of an electron can be associated with a two-component spinor, with the spin operator  $\vec{S}$  represented by  $2 \times 2$  matrices  $\frac{\hbar}{2}\vec{\sigma}$ . In the position wavefunction spinor representation corresponding to the CSCO  $(\vec{R}, S^2, S_z)$ , the orbital angular

momentum operators  $\vec{L}$  and  $L^2$  are represented by differential operators proportional to the unit matrix in the spin sector, for example [compare eqs. (8.6.7) and (8.6.8)]

$$L_z \leftrightarrow -i\hbar \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \frac{\partial}{\partial \phi}, \quad L_+ \leftrightarrow \hbar e^{i\phi} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \left( \frac{\partial}{\partial \theta} + i \frac{\cos \theta}{\sin \theta} \frac{\partial}{\partial \phi} \right), \quad (12.3.1)$$

etc. However, it is often more useful in problems with spherical symmetry to replace  $\vec{R}$  in the CSCO with the radial coordinate operator  $R$  and the operators  $L^2$  and  $L_z$ , so that all of the angular momentum operators are represented by matrices rather than differential operators on wavefunctions.

The angular momentum product orthobasis consists of common eigenkets of the compatible operators  $(L^2, S^2, L_z, S_z)$ . The corresponding basis kets can be labeled  $|l s m_l m_s\rangle$ , where we have suppressed one or more degeneracy labels (which might correspond to the radial wavefunction). Then, also suppressing the labels  $l, s$ , the product basis states are

$$|m_l, \uparrow\rangle \quad \text{and} \quad |m_l, \downarrow\rangle, \quad (12.3.2)$$

where  $m_s = 1/2$  and  $-1/2$  are represented by  $\uparrow$  and  $\downarrow$ , respectively. Using  $\vec{J} = \vec{L} + \vec{S}$ , we have another choice of compatible operators,  $(L^2, S^2, J^2, J_z)$ , associated to the total angular momentum orthobasis eigenkets

$$|l s j m\rangle = |j, m\rangle. \quad (12.3.3)$$

Our goal is to express the kets of eq. (12.3.3) as linear combinations of the product basis kets (12.3.2), for each  $l$ .

If  $l = 0$ , then things are very easy; the product basis kets  $|0, \uparrow\rangle$  and  $|0, \downarrow\rangle$  are already eigenstates of  $J^2$  with eigenvalue  $3\hbar^2/4$ , and of  $J_z$  with eigenvalues  $\pm\hbar/2$ , respectively. Therefore, the total angular momentum basis kets have  $j = 1/2$  and  $m = \pm 1/2$ , and are  $|\frac{1}{2}, \frac{1}{2}\rangle = |0, \uparrow\rangle$  and  $|\frac{1}{2}, -\frac{1}{2}\rangle = |0, \downarrow\rangle$ .

For  $l \neq 0$ , we begin with some preliminary counting, in order to know what to expect. There are  $(2l+1)2$  product basis kets  $|m_l, \uparrow\rangle$  and  $|m_l, \downarrow\rangle$ . They are all eigenkets of  $J_z = L_z + S_z$ , and the largest eigenvalue of  $J_z$  is  $\hbar m = \hbar(l+1/2)$ . This implies that there must be, in the total angular momentum basis, a multiplet with  $j = l+1/2$ , which will have  $2(l+1/2)+1 = 2l+2$  basis elements. Exactly one of those will have  $m = l-1/2$ , but we also know that there are two linearly independent states with that eigenvalue in the product basis, namely  $|l, \downarrow\rangle$  and  $|l-1, \uparrow\rangle$ . Therefore, there must also be a multiplet with  $j = l-1/2$ , which has  $2(l-1/2)+1 = 2l$  orthobasis members. Since we have accounted for all  $4l+2$  linearly independent kets, we have established that the tensor product of angular momentum  $l$  with angular momentum  $1/2$  must

consist of states with total angular momenta  $j = l + 1/2$  and  $l - 1/2$ , and no others. In notation similar to eq. (12.2.20), this is expressed as

$$l \otimes \frac{1}{2} = (l - \frac{1}{2}) \oplus (l + \frac{1}{2}). \quad (12.3.4)$$

Unlike eq. (12.2.20), there are no  $A$  or  $S$  subscripts here, because one cannot define antisymmetrization or symmetrization with respect to angular momenta that are not the same.

To construct a complete map between the total and product orthobases, let us start with the state  $|m_l, m_s\rangle = |l, \uparrow\rangle$ . Acting on this with  $J_z$  gives

$$J_z |l, \uparrow\rangle = (L_z + S_z) |l, \uparrow\rangle = \hbar(l + 1/2) |l, \uparrow\rangle. \quad (12.3.5)$$

This is the unique eigenstate of  $J^2$  and  $J_z$  with quantum numbers  $j = l + 1/2$  and  $m = l + 1/2$ . So, up to a phase that we can set arbitrarily,

$$|l + 1/2, l + 1/2\rangle = |l, \uparrow\rangle. \quad (12.3.6)$$

Now we can use this to construct all of the other states with  $j = l + 1/2$ , by acting repeatedly with the lowering operator  $J_- = L_- + S_-$ . From eq. (8.1.24),

$$J_- |l + 1/2, l + 1/2\rangle = \hbar\sqrt{2l + 1} |l + 1/2, l - 1/2\rangle, \quad (12.3.7)$$

so we get

$$|l + 1/2, l - 1/2\rangle = \frac{1}{\hbar\sqrt{2l + 1}} (S_- + L_-) |l, \uparrow\rangle, \quad (12.3.8)$$

or, using eq. (8.1.24) again to evaluate the action of each of  $S_-$  and  $L_-$ ,

$$|l + 1/2, l - 1/2\rangle = \frac{1}{\sqrt{2l + 1}} (|l, \downarrow\rangle + \sqrt{2l} |l - 1, \uparrow\rangle). \quad (12.3.9)$$

This is our second total angular momentum orthobasis eigenket. Acting with  $J_-$  on it gives

$$|l + 1/2, l - 3/2\rangle = \frac{1}{\sqrt{2l + 1}} (\sqrt{2} |l - 1, \downarrow\rangle + \sqrt{2l - 1} |l - 2, \uparrow\rangle). \quad (12.3.10)$$

Using the same strategy of applying  $J_-$ , by induction we obtain all of the  $j = l + 1/2$  kets,

$$|l + 1/2, m\rangle = \frac{1}{\sqrt{2l + 1}} (\sqrt{l - m + 1/2} |m + 1/2, \downarrow\rangle + \sqrt{l + m + 1/2} |m - 1/2, \uparrow\rangle) \quad (12.3.11)$$

for all  $m = -l - 1/2, \dots, l + 1/2$ .

Having found all the states with total angular momentum  $j = l + 1/2$ , now we find the states with  $j = l - 1/2$ . Each state with  $j = l - 1/2$  must be a linear combination of the form

$$|l - 1/2, m\rangle = a |m + 1/2, \downarrow\rangle + b |m - 1/2, \uparrow\rangle. \quad (12.3.12)$$

We know this because the product orthobasis kets on the right side are the only two that have the correct eigenvalue  $m$  of  $J_z/\hbar$ . Now, we can appeal to Theorem 2.6.5, which assures us that, since  $J^2$  is Hermitian, the state  $|l-1/2, m\rangle$  must be orthogonal to the state  $|l+1/2, m\rangle$ . Carrying out the inner product of eq. (12.3.11) with eq. (12.3.12), we learn that

$$a\sqrt{l-m+1/2} = -b\sqrt{l+m+1/2}. \quad (12.3.13)$$

As an arbitrary phase convention, we choose  $a$  real and positive. Requiring unit normalization of the ket then uniquely determines that for  $j = l - 1/2$ ,

$$|l-1/2, m\rangle = \frac{1}{\sqrt{2l+1}} \left( \sqrt{l+m+1/2} |m+1/2, \downarrow\rangle - \sqrt{l-m+1/2} |m-1/2, \uparrow\rangle \right), \quad (12.3.14)$$

for  $m = -l+1/2, \dots, l-1/2$ .

That concludes our task. To recap, in eqs. (12.3.11) and (12.3.14), we have obtained the  $J^2, J_z$  orthobasis kets as linear combinations of the product orthobasis kets.

As the simplest non-trivial concrete example, which we will have occasion to use later (at the end of section 17.3), consider the addition of an angular momentum  $l = 1$  to a spin-1/2:

$$1 \otimes \frac{1}{2} = \frac{3}{2} \oplus \frac{1}{2}. \quad (12.3.15)$$

Applying eq. (12.3.11), we have the  $j = 3/2$  total angular momentum basis states

$$|\frac{3}{2}, \frac{3}{2}\rangle = |1, \uparrow\rangle, \quad (12.3.16)$$

$$|\frac{3}{2}, \frac{1}{2}\rangle = \sqrt{\frac{1}{3}} |1, \downarrow\rangle + \sqrt{\frac{2}{3}} |0, \uparrow\rangle, \quad (12.3.17)$$

$$|\frac{3}{2}, -\frac{1}{2}\rangle = \sqrt{\frac{2}{3}} |0, \downarrow\rangle + \sqrt{\frac{1}{3}} |-1, \uparrow\rangle, \quad (12.3.18)$$

$$|\frac{3}{2}, -\frac{3}{2}\rangle = |-1, \downarrow\rangle, \quad (12.3.19)$$

and applying eq. (12.3.14) we get the  $j = 1/2$  total angular momentum basis states

$$|\frac{1}{2}, \frac{1}{2}\rangle = \sqrt{\frac{2}{3}} |1, \downarrow\rangle - \sqrt{\frac{1}{3}} |0, \uparrow\rangle, \quad (12.3.20)$$

$$|\frac{1}{2}, -\frac{1}{2}\rangle = \sqrt{\frac{1}{3}} |0, \downarrow\rangle - \sqrt{\frac{2}{3}} |-1, \uparrow\rangle. \quad (12.3.21)$$

## 12.4 The general case and Clebsch–Gordan coefficients

Now consider the general case of addition of angular momenta  $\vec{J}_1$  and  $\vec{J}_2$ . Let us apply reasoning similar to the counting that led to eq. (12.3.4). The largest eigenvalue of  $J_z = J_{1z} + J_{2z}$  is  $\hbar(j_1 + j_2)$ , and there is only one such state,  $|j_1 : j_2\rangle$ , so there must be exactly one total angular momentum multiplet with  $j = j_1 + j_2$ . (In this section, product angular momentum basis kets

will always be distinguished by use of a colon.) The subspace with  $J_z$  eigenvalue  $\hbar(j_1 + j_2 - 1)$  has dimension 2, spanned by the two states in the product basis  $|j_1 - 1 : j_2\rangle$  and  $|j_1 : j_2 - 1\rangle$ . One linear combination of these is found in the  $j = j_1 + j_2$  multiplet, so to contain the other there must also be exactly one total angular momentum multiplet with  $j = j_1 + j_2 - 1$ . Similarly, the subspace with  $J_z$  eigenvalue  $\hbar(j_1 + j_2 - 2)$  has dimension 3, spanned by the product basis kets  $|j_1 - 2 : j_2\rangle$  and  $|j_1 - 1 : j_2 - 1\rangle$  and  $|j_1 : j_2 - 2\rangle$ . Two linear combinations of these will occur in the  $j = j_1 + j_2$  and  $j = j_1 + j_2 - 1$  multiplets that we already know about, so there must also be exactly one total angular momentum multiplet with  $j = j_1 + j_2 - 2$ .

Continuing in this way, one finds that the tensor product of a multiplet with  $J_1^2$  eigenvalue  $\hbar^2 j_1(j_1 + 1)$  and a multiplet with  $J_2^2$  eigenvalue  $\hbar^2 j_2(j_2 + 1)$  must consist of a sum of multiplets with  $J^2 = \hbar^2 j(j + 1)$ , with  $j$  taking on the values from  $|j_1 - j_2|$  to  $j_1 + j_2$ , with integer increments,

$$j_1 \otimes j_2 = |j_1 - j_2| \oplus \cdots \oplus (j_1 + j_2). \quad (12.4.1)$$

Each of the total angular momentum basis kets  $|j_1 j_2 j, m\rangle$  is a linear combination of the product basis kets  $|j_1 j_2 m_1 : m_2\rangle$ . By the completeness of the latter, we can write

$$|j_1 j_2 j, m\rangle = \sum_{m_1=-j_1}^{j_1} \sum_{m_2=-j_2}^{j_2} |j_1 j_2 m_1 : m_2\rangle \langle j_1 j_2 m_1 : m_2 | j_1 j_2 j, m\rangle \quad (12.4.2)$$

$$= \sum_{m_1=-j_1}^{j_1} \sum_{m_2=-j_2}^{j_2} |j_1 j_2 m_1 : m_2\rangle C_{m_1 m_2 m}^{j_1 j_2 j} \quad (12.4.3)$$

where the inner products

$$C_{m_1 m_2 m}^{j_1 j_2 j} \equiv \langle j_1 j_2 m_1 : m_2 | j_1 j_2 j, m\rangle \quad (12.4.4)$$

are known as **Clebsch–Gordan coefficients**, after mathematicians Alfred Clebsch and Paul Gordan. Various different notations and conventions for them appear in the literature. We will usually use the  $C$  notation for these inner products, as a way of saving space, with commas inserted between the superscripts or subscripts when it helps to make the meaning clear.

The Clebsch–Gordan coefficients obey selection rules; they can be non-zero only if

$$|j_1 - j_2| \leq j \leq j_1 + j_2, \quad (\text{triangle condition}), \quad (12.4.5)$$

$$j_1 + j_2 - j \text{ is an integer}, \quad (12.4.6)$$

$$m = m_1 + m_2. \quad (12.4.7)$$

The first two of these simply restate eq. (12.4.1), and the last follows immediately from evaluating the matrix element of  $J_z = J_{1z} + J_{2z}$  between  $\langle j_1 j_2 m_1 : m_2 |$  and  $|j_1 j_2 j, m\rangle$ .



In general, the Clebsch–Gordan coefficients are subject to ambiguities, because the normalized states  $|j_1 j_2 j, m\rangle$  are determined only up to a phase. These phase ambiguities are resolved here by adopting a convention that  $C_{m_1 m_2 m}^{j_1 j_2 j}$  is real and positive when  $m_1 = j_1$  and  $m = j$ . (Note that this convention gives a special role to the first of the angular momenta being combined.) The coefficient vanishes unless  $m_2 = j - j_1$ , so we can specify the convention as

$$\text{phase convention: } C_{j_1, j-j_1, j}^{j_1 j_2 j} \text{ is real and positive.} \quad (12.4.8)$$

Then, because the actions of  $J_-$  and  $J_{1-}$  and  $J_{2-}$  each only involve real coefficients, as given by eq. (8.1.24), we find, by acting on eq. (12.4.3) repeatedly with  $J_-$  on the left side and its equivalent  $J_{1-} + J_{2-}$  on the right side, the convenient and useful result that all of the Clebsch–Gordan coefficients will be real.

One can also use completeness with respect to the orthobasis elements  $|j_1 j_2 j, m\rangle$ , to obtain the inverse relation to eq. (12.4.3), which gives each product orthobasis ket as a linear combination of the total angular momentum orthobasis kets,

$$|j_1 j_2 m_1 : m_2\rangle = \sum_{j=|j_1-j_2|}^{j_1+j_2} \sum_{m=-j}^j |j_1 j_2 j, m\rangle \langle j_1 j_2 j, m | j_1 j_2 m_1 : m_2\rangle \quad (12.4.9)$$

$$= \sum_{j=|j_1-j_2|}^{j_1+j_2} \sum_{m=-j}^j |j_1 j_2 j, m\rangle C_{m_1 m_2 m}^{j_1 j_2 j}. \quad (12.4.10)$$

The matrix elements on the right are actually the complex conjugates of the Clebsch–Gordan coefficients, but in our phase convention, all of them are real anyway.

As eqs. (12.4.3) and (12.4.10) demonstrate, results for total angular momentum orthobasis states in terms of product states, or vice versa, can always be expressed in terms of Clebsch–Gordan coefficients. For example, the content of the results for  $l \otimes \frac{1}{2} = (l - \frac{1}{2}) \oplus (l + \frac{1}{2})$  that we found in eqs. (12.3.11) and (12.3.14) can be expressed, writing  $m = m_l \pm 1/2$  as appropriate, as

$$C_{m_l, -\frac{1}{2}, m_l - \frac{1}{2}}^{l, \frac{1}{2}, l + \frac{1}{2}} = \sqrt{\frac{l+1-m_l}{2l+1}}, \quad C_{m_l, \frac{1}{2}, m_l + \frac{1}{2}}^{l, \frac{1}{2}, l + \frac{1}{2}} = \sqrt{\frac{l+1+m_l}{2l+1}}, \quad (12.4.11)$$

$$C_{m_l, -\frac{1}{2}, m_l - \frac{1}{2}}^{l, \frac{1}{2}, l - \frac{1}{2}} = \sqrt{\frac{l+m_l}{2l+1}}, \quad C_{m_l, \frac{1}{2}, m_l + \frac{1}{2}}^{l, \frac{1}{2}, l - \frac{1}{2}} = -\sqrt{\frac{l-m_l}{2l+1}}, \quad (12.4.12)$$

from which the results in eqs. (12.3.16)–(12.3.21) follow as the special case with  $l = 1$ . Although we claimed to be interested in the case that  $l$  was an orbital angular quantum number, there is no difference in the arithmetic for any angular momentum, and  $l$  can even be taken to be a half integer in these formulas.

As another useful example, consider the angular momentum addition problem  $j \otimes 1 = (j-1) \oplus j \oplus (j+1)$ . By a similar strategy, to be formalized below in the general case, the

relevant non-zero Clebsch–Gordan coefficients are found to be

$$C_{m, \pm 1, m \pm 1}^{j, 1, j+1} = \sqrt{\frac{(j+1 \pm m)(j+2 \pm m)}{2(j+1)(2j+1)}}, \quad C_{m, 0, m}^{j, 1, j+1} = \sqrt{\frac{(j+1-m)(j+1+m)}{(j+1)(2j+1)}}, \quad (12.4.13)$$

$$C_{m, \pm 1, m \pm 1}^{j, 1, j} = \mp \sqrt{\frac{(j+1 \pm m)(j \mp m)}{2j(j+1)}}, \quad C_{m, 0, m}^{j, 1, j} = \sqrt{\frac{m}{j(j+1)}}, \quad (12.4.14)$$

$$C_{m, \pm 1, m \pm 1}^{j, 1, j-1} = \sqrt{\frac{(j-1 \mp m)(j \mp m)}{2j(2j+1)}}, \quad C_{m, 0, m}^{j, 1, j-1} = -\sqrt{\frac{(j+m)(j-m)}{j(2j+1)}}. \quad (12.4.15)$$

One can easily find published or online tables of Clebsch–Gordan coefficients, and software implementations of them, but it is a good idea to read the fine print to make sure of the phase convention being used.

We now point out some important identities satisfied by Clebsch–Gordan coefficients in general. First, by using eqs. (8.1.22) and (8.1.24) to evaluate<sup>†</sup>

$$\langle m_1 : m_2 | J_- | j, m \rangle = \langle m_1 : m_2 | (J_{1-} + J_{2-}) | j, m \rangle = (\langle j, m | (J_{1+} + J_{2+}) | m_1 : m_2 \rangle)^*, \quad (12.4.16)$$

one finds

$$\begin{aligned} \sqrt{j(j+1) - m(m-1)} C_{m_1, m_2, m-1}^{j_1 j_2 j} &= \sqrt{j_1(j_1+1) - m_1(m_1+1)} C_{m_1+1, m_2, m}^{j_1 j_2 j} \\ &+ \sqrt{j_2(j_2+1) - m_2(m_2+1)} C_{m_1, m_2+1, m}^{j_1 j_2 j}, \end{aligned} \quad (12.4.17)$$

Similarly, from evaluating matrix elements of  $J_+$ ,

$$\begin{aligned} \sqrt{j(j+1) - m(m+1)} C_{m_1, m_2, m+1}^{j_1 j_2 j} &= \sqrt{j_1(j_1+1) - m_1(m_1-1)} C_{m_1-1, m_2, m}^{j_1 j_2 j} \\ &+ \sqrt{j_2(j_2+1) - m_2(m_2-1)} C_{m_1, m_2-1, m}^{j_1 j_2 j}. \end{aligned} \quad (12.4.18)$$

From the total angular momentum orthonormality relations, using completeness of the product basis and the reality of the Clebsch–Gordan coefficients in our chosen convention, we also obtain

$$\sum_{m_1=-j_1}^{j_1} \sum_{m_2=-j_2}^{j_2} C_{m_1 m_2 m}^{j_1 j_2 j} C_{m_1 m_2 m'}^{j_1 j_2 j'} = \delta_{j j'} \delta_{m m'}. \quad (12.4.19)$$

Similarly, from the product basis orthonormality relations,

$$\sum_{j=|j_1-j_2|}^{j_1+j_2} \sum_{m=-j}^j C_{m_1 m_2 m}^{j_1 j_2 j} C_{m'_1 m'_2 m}^{j_1 j_2 j} = \delta_{m_1 m'_1} \delta_{m_2 m'_2}, \quad (12.4.20)$$

which follows from completeness of the total angular momentum orthobasis.

---

<sup>†</sup>In the rest of this section, the labels  $j_1, j_2$  on bras and kets are always the same, and so are omitted.

Consider eq. (12.4.18) in the special case  $m = j$ , so that the left side vanishes. This gives  $C_{m_1-1, m_2, j}^{j_1 j_2 j} = - \left( \sqrt{j_2(j_2+1) - m_2(m_2-1)} / \sqrt{j_1(j_1+1) - m_1(m_1-1)} \right) C_{m_1, m_2-1, j}^{j_1 j_2 j}$ . This relation can be used to lower  $m_1$  and raise  $m_2$  in unit steps, starting from  $m_1 = j_1$ , giving a sign flip each time. Therefore, in our convention

$$\text{sign} \left( C_{m_1, j-m_1, j}^{j_1 j_2 j} \right) = (-1)^{j_1-m_1}. \quad (12.4.21)$$

In particular, taking  $m_1 = j - j_2$ , we must have

$$\text{sign} \left( C_{j-j_2, j_2, j}^{j_1 j_2 j} \right) = (-1)^{j_1+j_2-j}. \quad (12.4.22)$$

Meanwhile, directly from our convention in eq. (12.4.18), we also have

$$\text{sign} \left( C_{j_2, j-j_2, j}^{j_2 j_1 j} \right) = 1. \quad (12.4.23)$$

Now, the states  $|j_1 j_2 j, j\rangle$  and  $|j_2 j_1 j, j\rangle$  obtained by interchanging the roles of  $\vec{J}_1$  and  $\vec{J}_2$  are really the same state physically, so they can only differ by a phase, which in our convention must be simply a sign.<sup>‡</sup> Comparing the previous two equations, we see that this sign must be  $|j_1 j_2 j, j\rangle = (-1)^{j_1+j_2-j} |j_2 j_1 j, j\rangle$ . Now we can act on this equation repeatedly with  $J_-$  to get the relation between the more general states  $|j_1 j_2 j, m\rangle$  and  $|j_2 j_1 j, m\rangle$ . Doing so cannot change the relative sign, because the operator  $J_- = J_{1-} + J_{2-}$  does not know or care which of the two angular momenta we chose to be the first one, so

$$|j_1 j_2 j, m\rangle = (-1)^{j_1+j_2-j} |j_2 j_1 j, m\rangle. \quad (12.4.24)$$

We therefore arrive at the rule for Clebsch–Gordan coefficients when we interchange the two angular momenta being added,

$$C_{m_1 m_2 m}^{j_1 j_2 j} = (-1)^{j_1+j_2-j} C_{m_2 m_1 m}^{j_2 j_1 j}. \quad (12.4.25)$$

Although we used a convention choice in the intermediate steps of deriving it, eq. (12.4.25) is actually independent of the convention choice. A similar sort of strategy (with details omitted here) can be used to show that if one simultaneously flips the signs of  $m_1$ ,  $m_2$ , and  $m$ , then the Clebsch–Gordan coefficients must also satisfy

$$C_{m_1 m_2 m}^{j_1 j_2 j} = (-1)^{j_1+j_2-j} C_{-m_1, -m_2, -m}^{j_1 j_2 j} \quad (12.4.26)$$

in our phase convention.

---

<sup>‡</sup>Note that we are implicitly defining the *product* orthobasis kets to be invariant under interchange of the roles of the two angular momenta,  $|j_1 j_2 m_1 : m_2\rangle = |j_2 j_1 m_2 : m_1\rangle = |j_1, m_1\rangle \otimes |j_2, m_2\rangle$ .

The following recipe can be used to construct the total angular momentum orthobasis states, and thus the Clebsch–Gordan coefficients, in a general case. Assume that  $j_1 \geq j_2$ . (Otherwise, one can simply reverse the roles of  $j_1$  and  $j_2$ .) Start with the state with the highest possible  $m$ , which is  $m = j_1 + j_2 = j$ . Since there is only one product orthobasis state with  $J_{1z} + J_{2z}$  eigenvalue equal to  $\hbar(j_1 + j_2)$ , we have, using our phase convention choice,

$$|j_1 + j_2, j_1 + j_2\rangle = |j_1 : j_2\rangle. \quad (12.4.27)$$

Now we follow the strategy of repeatedly acting with the lowering operator  $J_- = J_{1-} + J_{2-}$  to find new states  $|j_1 + j_2, m\rangle$ . Using eq. (8.1.24) gives

$$J_- |j_1 + j_2, j_1 + j_2\rangle = \hbar\sqrt{2(j_1 + j_2)} |j_1 + j_2, j_1 + j_2 - 1\rangle, \quad (12.4.28)$$

which also can be evaluated as

$$J_{1-} |j_1 : j_2\rangle + J_{2-} |j_1 : j_2\rangle = \hbar\sqrt{2j_1} |j_1 - 1 : j_2\rangle + \hbar\sqrt{2j_2} |j_1 : j_2 - 1\rangle. \quad (12.4.29)$$

Therefore,

$$|j_1 + j_2, j_1 + j_2 - 1\rangle = \frac{1}{\sqrt{j_1 + j_2}} \left( \sqrt{j_2} |j_1 : j_2 - 1\rangle + \sqrt{j_1} |j_1 - 1 : j_2\rangle \right). \quad (12.4.30)$$

[The special cases  $j_1 = j_2 = 1/2$  and  $j_1 = l, j_2 = 1/2$  were previously found in eqs. (12.2.15) and (12.3.9), respectively.] Continuing to act with  $J_-$ , we similarly find all of the orthobasis kets  $|j_1 + j_2, m\rangle$ . Eventually we will obtain  $m = -j_1 - j_2$ , which is then annihilated by  $J_-$ .

Next, we proceed to obtain the states that have total  $j = j_1 + j_2 - 1$ . We start by writing the most general possible expression for the state with the largest possible  $m$ ,

$$|j_1 + j_2 - 1, j_1 + j_2 - 1\rangle = a |j_1 : j_2 - 1\rangle + b |j_1 - 1 : j_2\rangle. \quad (12.4.31)$$

The coefficients  $a$  and  $b$  can be uniquely identified by requiring that this ket is orthogonal to  $|j_1 + j_2, j_1 + j_2 - 1\rangle$ , that the ket is normalized so that  $|a|^2 + |b|^2 = 1$ , and that  $a$  is real and positive according to our phase convention. This gives

$$|j_1 + j_2 - 1, j_1 + j_2 - 1\rangle = \frac{1}{\sqrt{j_1 + j_2}} \left( \sqrt{j_1} |j_1 : j_2 - 1\rangle - \sqrt{j_2} |j_1 - 1 : j_2\rangle \right). \quad (12.4.32)$$

[Again, we had previously derived this in the special cases  $j_1 = j_2 = 1/2$  and  $j_1 = l, j_2 = 1/2$ , in eqs. (12.2.17) and (12.3.14), respectively.] Now, we again apply the operator  $J_-$  repeatedly, to obtain all of the other states  $|j_1 + j_2 - 1, m\rangle$ .

Continuing on our vaunted quest, we proceed to the states with  $j = j_1 + j_2 - 2$ , starting again with the maximum value of  $m$ . This must be some linear combination of the three product orthobasis kets with  $m = m_1 + m_2 = j_1 + j_2 - 2$ , which we can write as

$$|j_1 + j_2 - 2, j_1 + j_2 - 2\rangle = a |j_1 : j_2 - 2\rangle + b |j_1 - 1 : j_2 - 1\rangle + c |j_1 - 2 : j_2\rangle. \quad (12.4.33)$$

The three unknown coefficients  $a$ ,  $b$ ,  $c$  can be obtained by demanding that this be orthogonal to both of the kets  $|j_1 + j_2, j_1 + j_2 - 2\rangle$  and  $|j_1 + j_2 - 1, j_1 + j_2 - 2\rangle$ , as required by Theorem 2.6.5 (since all three are eigenvectors of  $J^2$  with different eigenvalues), and that the ket is normalized, so  $|a|^2 + |b|^2 + |c|^2 = 1$ , and that  $a$  is real and positive according to our phase convention. The remaining states  $|j_1 + j_2 - 2, m\rangle$  are then obtained by acting repeatedly with  $J_-$ .

The preceding illustrates the general procedure, which is shown schematically in Figure 12.4.1. Each column represents the repeated action of  $J_-$  on the state with the highest  $m$  for a given  $j$ . After completing each column, we move to the next column by first writing a candidate ket  $|j, j\rangle$  as a linear combination of the product orthobasis kets that have  $m_1 + m_2 = m = j$ . The coefficients are determined by requiring orthogonality to all of the previously obtained total angular momentum orthobasis kets with that value of  $m$  and total angular momentum  $j + 1$  or larger. Demanding that the ket be normalized and obey our phase convention, the state is then uniquely determined, and the rest of the states  $|j, m\rangle$  follow by applying  $J_-$  repeatedly. The




Figure 12.4.1: A plan to compute the total angular momentum orthobasis kets  $|j_1 j_2 j, m\rangle$ , abbreviated here as  $|j, m\rangle$ , in terms of product orthobasis kets, when two angular momenta  $j_1$  and  $j_2$  are combined. The process starts at the upper left with  $j = m = j_1 + j_2$ . Each column has fixed  $j$ , and is constructed from top to bottom using  $J_-$ . Then the next column is started by constructing the state with maximum  $m$ , by requiring it to be orthogonal to all of the previously found kets with that same  $m$ . The process ends after the rightmost column with  $j = |j_1 - j_2|$  is finished, with  $m = -|j_1 - j_2|$ . This process provides the Clebsch–Gordan coefficients  $\langle m_1 : m_2 | j, m \rangle = \langle j_1 j_2 m_1 : m_2 | j_1 j_2 j, m \rangle = C_{m_1 m_2 m}^{j_1 j_2 j}$ .

process iterates by moving to the next column with  $j$  lower by 1. Eventually, we will finish the last column of orthobasis kets, which will have  $j = |j_1 - j_2|$ , and there will be none more.

Consider the special case of adding two angular momenta  $j_1 = 1$  and  $j_2 = 1$ . The resulting allowed values of  $j$  are 0, 1, and 2. By following the procedure summarized in Figure 12.4.1, one finds the  $j = 2$  states

$$|2, 2\rangle = |1: 1\rangle, \quad (12.4.34)$$

$$|2, 1\rangle = (|1: 0\rangle + |0: 1\rangle)/\sqrt{2}, \quad (12.4.35)$$

$$|2, 0\rangle = (|1: -1\rangle + 2|0: 0\rangle + |-1: 1\rangle)/\sqrt{6}, \quad (12.4.36)$$

$$|2, -1\rangle = (|0: -1\rangle + |-1: 0\rangle)/\sqrt{2}, \quad (12.4.37)$$

$$|2, -2\rangle = |-1: -1\rangle, \quad (12.4.38)$$

which are all symmetric under  $m_1 \leftrightarrow m_2$ , followed by the  $j = 1$  states

$$|1, 1\rangle = (|1: 0\rangle - |0: 1\rangle)/\sqrt{2}, \quad (12.4.39)$$

$$|1, 0\rangle = (|1: -1\rangle - |-1: 1\rangle)/\sqrt{2}, \quad (12.4.40)$$

$$|1, -1\rangle = (|0: -1\rangle - |-1: 0\rangle)/\sqrt{2}, \quad (12.4.41)$$

which are each antisymmetric under the same exchange, and finally the  $j = 0$  state

$$|0, 0\rangle = (|1: -1\rangle - |0: 0\rangle + |-1: 1\rangle)/\sqrt{3}, \quad (12.4.42)$$

which is symmetric. [These results can also be checked using the general Clebsch–Gordan coefficients for  $j \otimes 1$  in eqs. (12.4.13)-(12.4.15).] To summarize what has been learned, one writes

$$1 \otimes 1 = 2_S \oplus 1_A \oplus 0_S, \quad (12.4.43)$$

where the  $S$  and  $A$  subscripts indicate the symmetry or antisymmetry of the total angular momentum orthobasis kets under exchange of the two product basis eigenvalues  $m_1$  and  $m_2$ .

More generally, for the combination of two equal angular momenta  $j_1 = j_2 = j$ , the symmetry and antisymmetry properties for exchange of  $m_1$  and  $m_2$  are summarized by

$$j \otimes j = (2j)_S \oplus (2j-1)_A \oplus (2j-2)_S \oplus \cdots \oplus 0, \quad (12.4.44)$$

with alternating  $S$  and  $A$ , so that the singlet on the right is symmetric if  $j$  is an integer and antisymmetric if  $j$  is a half-integer. These symmetry and antisymmetry properties follow immediately from eq. (12.4.25).

The simplest nontrivial example that cannot be obtained from eqs. (12.4.11)-(12.4.12) or from (12.4.13)-(12.4.15) is  $j_1 = j_2 = 3/2$ . Call the resulting total angular momentum quantum number  $J$ . Then you can show that the  $J = 3, m_J = 3$  and  $J = 3, m_J = 0$  and  $J = 0, m_J = 0$  states are, in terms of product basis states  $|m_1 : m_2\rangle$ ,

$$|3, 3\rangle = \left|\frac{3}{2} : \frac{3}{2}\right\rangle, \quad (12.4.45)$$

$$|3, 0\rangle = \frac{1}{\sqrt{20}} \left( \left|\frac{3}{2} : -\frac{3}{2}\right\rangle + \left|-\frac{3}{2} : \frac{3}{2}\right\rangle + 3\left|\frac{1}{2} : -\frac{1}{2}\right\rangle + 3\left|-\frac{1}{2} : \frac{1}{2}\right\rangle \right), \quad (12.4.46)$$

$$|0, 0\rangle = \frac{1}{2} \left( \left|\frac{3}{2} : -\frac{3}{2}\right\rangle - \left|-\frac{3}{2} : \frac{3}{2}\right\rangle - \left|\frac{1}{2} : -\frac{1}{2}\right\rangle + \left|-\frac{1}{2} : \frac{1}{2}\right\rangle \right), \quad (12.4.47)$$

respectively. It is left as Exercise 12.7 to work out the other total angular momentum orthobasis states  $|J, m_J\rangle$  in this example.

One can also combine three (or more) angular momenta, to obtain an orthobasis of kets that are eigenstates of the observables  $J^2$  and  $J_z$  for the total angular momentum  $\vec{J} = \vec{J}_1 + \vec{J}_2 + \vec{J}_3$ . This can be done recursively, by first combining  $\vec{J}_1$  and  $\vec{J}_2$  into an angular momentum operator  $\vec{J}_{12}$ , and then combining the results with  $\vec{J}_3$ . For example, combining three spin-1/2 angular momenta results in

$$\frac{1}{2} \otimes \frac{1}{2} \otimes \frac{1}{2} = (1 \oplus 0) \otimes \frac{1}{2} = (1 \otimes \frac{1}{2}) \oplus (0 \otimes \frac{1}{2}) = (3/2)_S \oplus (1/2)_a \oplus (1/2)_b. \quad (12.4.48)$$

The subscript  $S$  on the  $j = 3/2$  multiplet indicates that the states in it are symmetric under exchange of any two of the three individual spins, as we will soon check. There are also two  $j = 1/2$  multiplets in the result, distinguished by degeneracy labels  $a$  and  $b$ . To check the multiplicities of states, we note that on the left side of eq. (12.4.48) there are  $2^3 = 8$  product orthobasis states, since each individual spin has 2 values of  $m_s = \pm 1/2$ . On the right, the multiplicities of the total angular momentum states are  $4 + 2 + 2 = 8$ . To explicitly construct the total angular momentum orthobasis kets, one can first use eqs. (12.2.14)–(12.2.17) to construct the  $J_{12}^2, J_{12z}$  eigenstate kets. Combining the resulting  $j_{12} = 1$  states with the third spin, one finds, by using eq. (12.3.11), that the  $j = 3/2$  states are

$$\left|\frac{3}{2}, \frac{3}{2}\right\rangle = |\uparrow\uparrow\uparrow\rangle, \quad (12.4.49)$$

$$\left|\frac{3}{2}, \frac{1}{2}\right\rangle = \frac{1}{\sqrt{3}} (|\uparrow\uparrow\downarrow\rangle + |\uparrow\downarrow\uparrow\rangle + |\downarrow\uparrow\uparrow\rangle), \quad (12.4.50)$$

$$\left|\frac{3}{2}, -\frac{1}{2}\right\rangle = \frac{1}{\sqrt{3}} (|\uparrow\downarrow\downarrow\rangle + |\downarrow\uparrow\downarrow\rangle + |\downarrow\downarrow\uparrow\rangle), \quad (12.4.51)$$

$$\left|\frac{3}{2}, -\frac{3}{2}\right\rangle = |\downarrow\downarrow\downarrow\rangle, \quad (12.4.52)$$

and, by using eq. (12.3.14), the  $j = 1/2$  states

$$|\frac{1}{2}, \frac{1}{2}, a\rangle = \frac{1}{\sqrt{6}}(2|\uparrow\uparrow\downarrow\rangle - |\uparrow\downarrow\uparrow\rangle - |\downarrow\uparrow\uparrow\rangle), \quad (12.4.53)$$

$$|\frac{1}{2}, -\frac{1}{2}, a\rangle = \frac{1}{\sqrt{6}}(|\uparrow\downarrow\downarrow\rangle + |\downarrow\uparrow\downarrow\rangle - 2|\downarrow\downarrow\uparrow\rangle). \quad (12.4.54)$$

From combining the  $j_{12} = 0$  states from eq. (12.2.17) with the third spin, we have the additional  $j = 1/2$  total angular momentum basis states

$$|\frac{1}{2}, \frac{1}{2}, b\rangle = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\uparrow\rangle - |\downarrow\uparrow\uparrow\rangle), \quad (12.4.55)$$

$$|\frac{1}{2}, -\frac{1}{2}, b\rangle = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\downarrow\rangle - |\downarrow\uparrow\downarrow\rangle). \quad (12.4.56)$$

The explicit forms for the  $j = 3/2$  states show that they are indeed each symmetric under exchange of any two of the three individual spins. For the  $j = 1/2$  states, the ones labeled  $a$  are symmetric, and the ones labeled  $b$  are antisymmetric, under exchange of the first two spins. However, these  $j = 1/2$  states do not have a complete symmetry or antisymmetry under exchange of every pair of spins. Also note that there is no totally antisymmetric combination of three or more spin  $1/2$  states, simply because in the product basis there are only two values  $\uparrow$  and  $\downarrow$  available, so that every product orthobasis ket is symmetric under interchange of at least one pair of spins.

Similarly, combining four spin- $1/2$  angular momenta gives

$$\frac{1}{2} \otimes \frac{1}{2} \otimes \frac{1}{2} \otimes \frac{1}{2} = (1 \oplus 0) \otimes (1 \oplus 0) = 2_S \oplus 1 \oplus 1 \oplus 1 \oplus 0 \oplus 0. \quad (12.4.57)$$

Here the counting is that the number of product orthobasis states on the left is  $2^4 = 16$ , while the counting of multiplicities of total angular momentum orthobasis states on the right is  $5 + 3 + 3 + 3 + 1 + 1 = 16$ . Only the total spin 2 multiplet is totally symmetric, and there is no totally antisymmetric multiplet. The other multiplets on the right side of eq. (12.4.57) have mixed symmetry properties under exchange of the spins.

In the case of  $N$  electrons in an atom, the angular momentum eigenstates are often given in spectroscopic notation, defined as follows. First, combine all of the individual spin operators  $\vec{S}_i$  to obtain the total spin angular momentum operator  $\vec{S} = \sum_i \vec{S}_i$ . The eigenvalues of the operator  $S^2 = \vec{S} \cdot \vec{S}$  are then denoted by  $\hbar^2 S(S+1)$  where  $S$  is a number.<sup>§</sup> Clearly, if  $N$  is even, then the number  $S$  must be an integer with  $0 \leq S \leq N/2$ . If  $N$  is odd, then the number  $S$  must be half-integer, with  $1/2 \leq S \leq N/2$ . Next, combine all of the individual orbital angular momenta operators  $\vec{L}_i$  to obtain the total orbital angular momentum operator  $\vec{L} = \sum_i \vec{L}_i$ . The

---

<sup>§</sup>Following a common but potentially confusing notation, in the case of combined angular momenta, capital letters are often used both for the names of the operators as well as the corresponding quantum numbers.



eigenvalues of the operator  $L^2 = \vec{L} \cdot \vec{L}$  are likewise denoted  $\hbar^2 L(L+1)$ , where the quantum number  $L$  is always an integer. Finally, the operators  $\vec{S}$  and  $\vec{L}$  are combined to form the total angular momentum operator  $\vec{J}$ , and the operators  $J^2$  and  $J_z$  have eigenvalues denoted  $\hbar^2 J(J+1)$  and  $\hbar m_J$ , respectively. The traditional notation for a multiplet of common eigenstates of the observables  $S^2$ ,  $L^2$ , and  $J^2$  is then

$$^{2S+1}L_J, \quad (12.4.58)$$

where  $S$ ,  $L$ , and  $J$  are the quantum numbers, but with  $L$  replaced by the capital letter code  $S$ ,  $P$ ,  $D$ ,  $F$ ,  $G$ ,  $\dots$ , according to whether the number  $L$  is  $0, 1, 2, 3, 4, \dots$ , as indicated in eq. (11.1.20). The degeneracy, or multiplicity, of each group of states denoted by  $^{2S+1}L_J$  is  $2J+1$ , since the eigenvalue  $m_J$  can take on the values  $-J, -J+1, \dots, J-1, J$ .

For example, a single electron always has  $S = 1/2$ . For  $L = 0$ , it has  $J = 1/2$ , and for  $L \geq 1$  it can have  $J = L \pm 1/2$ , as we saw in section 12.3. So, the list of total angular momentum multiplets for a single electron in an atom is

$$^2S_{1/2}, ^2P_{1/2}, ^2P_{3/2}, ^2D_{3/2}, ^2D_{5/2}, ^2F_{5/2}, ^2F_{7/2}, ^2G_{7/2}, ^2G_{9/2}, \dots \quad (12.4.59)$$

For two electrons, the possible total spin quantum numbers are  $S = 0$  and  $1$ , as we saw in section 12.2. The orbital angular momenta  $\vec{L}_1$  and  $\vec{L}_2$  with quantum numbers  $l_1$  and  $l_2$  can be combined into  $L = |l_1 - l_2|, \dots, l_1 + l_2$ . Therefore, the possible total angular momentum eigenstates for two electrons are

$$^1S_0, ^1P_1, ^1D_2, ^1F_3, ^1G_4, \dots \quad (12.4.60)$$

for  $S = 0$ , and

$$^3S_1, ^3P_0, ^3P_1, ^3P_2, ^3D_1, ^3D_2, ^3D_3, ^3F_2, ^3F_3, ^3F_4, \dots \quad (12.4.61)$$

for  $S = 1$ . However, as we will discuss in more detail in section 18.1, Fermi–Dirac statistics requires that the total state must be antisymmetric under exchange of the two electrons. In the special case that the electrons have the same radial wavefunction and the same orbital angular momentum quantum number  $l_1 = l_2$ , then one can use eq. (12.4.44) to see that if  $L$  is odd (an antisymmetric position wavefunction) then only the symmetric spin combination  $S = 1$  is allowed, and if  $L$  is even (a symmetric position wavefunction), then only the antisymmetric spin combination  $S = 0$  is allowed.

## 12.5 How spherical harmonics combine

The spherical harmonics are the position representations of the orbital angular momentum eigenstates of  $L^2$  and  $L_z$ . In this section, we will learn how the addition of angular momentum technology is realized in terms of the spherical harmonics.

Let us start by considering fixed  $l_1$ ,  $l_2$ ,  $l$ , and  $m$ . We then define the following function of  $\theta, \phi$ , as a linear combination of products of spherical harmonics weighted by the Clebsch–Gordan coefficients defined by eq. (12.4.4),

$$\Psi_l^m(\theta, \phi) \equiv \sum_{m_1, m_2} Y_{l_1}^{m_1}(\theta, \phi) Y_{l_2}^{m_2}(\theta, \phi) C_{m_1 m_2 m}^{l_1 l_2 l}, \quad (12.5.1)$$

where the sums are over all  $m_1$  and  $m_2$  such that the Clebsch–Gordan coefficient does not vanish. Our goal is to find another simple expression for  $\Psi_l^m(\theta, \phi)$ . We claim that, as its labeling suggests, this wavefunction is an eigenstate of the differential operators  $L^2$  and  $L_z$ , with eigenvalues  $\hbar^2 l(l+1)$  and  $\hbar m$ , respectively, and it is therefore proportional to  $Y_l^m(\theta, \phi)$ .

To prove this claim, we will check the action of the differential operators  $L_z$ ,  $L_+$ , and  $L_-$  on it. First, by applying eq. (8.6.27), we find

$$L_z \Psi_l^m = \sum_{m_1, m_2} (\hbar m_1 + \hbar m_2) Y_{l_1}^{m_1} Y_{l_2}^{m_2} C_{m_1 m_2 m}^{l_1 l_2 l}. \quad (12.5.2)$$

The Clebsch–Gordan coefficient enforces that all non-zero contributions have  $m_1 + m_2 = m$ , so

$$L_z \Psi_l^m = \hbar m \Psi_l^m. \quad (12.5.3)$$

Next, applying eq. (8.6.29) we find

$$\begin{aligned} L_- \Psi_l^m &= \hbar \sum_{m_1, m_2} \left[ \sqrt{l_1(l_1+1) - m_1(m_1-1)} Y_{l_1}^{m_1-1} Y_{l_2}^{m_2} \right. \\ &\quad \left. + \sqrt{l_2(l_2+1) - m_2(m_2-1)} Y_{l_1}^{m_1} Y_{l_2}^{m_2-1} \right] C_{m_1 m_2 m}^{l_1 l_2 l}. \end{aligned} \quad (12.5.4)$$

Since we are summing over all  $m_1$  and  $m_2$ , we can use the trick of relabeling  $m_1 \rightarrow m_1 + 1$  in the first term and  $m_2 \rightarrow m_2 + 1$  in the second term, to obtain

$$\begin{aligned} L_- \Psi_l^m &= \hbar \sum_{m_1, m_2} \left[ \sqrt{l_1(l_1+1) - m_1(m_1+1)} C_{m_1+1, m_2, m}^{l_1 l_2 l} \right. \\ &\quad \left. + \sqrt{l_2(l_2+1) - m_2(m_2+1)} C_{m_1, m_2+1, m}^{l_1 l_2 l} \right] Y_{l_1}^{m_1} Y_{l_2}^{m_2}. \end{aligned} \quad (12.5.5)$$

Now the Clebsch–Gordan recurrence relation eq. (12.4.17) turns this into

$$L_- \Psi_l^m = \hbar \sum_{m_1, m_2} \sqrt{l(l+1) - m(m-1)} C_{m_1, m_2, m-1}^{l_1 l_2 l} Y_{l_1}^{m_1} Y_{l_2}^{m_2}, \quad (12.5.6)$$

or, using the definition of eq. (12.5.1), simply,

$$L_- \Psi_l^m = \hbar \sqrt{l(l+1) - m(m-1)} \Psi_l^{m-1}. \quad (12.5.7)$$

In an exactly analogous way, one finds

$$L_+ \Psi_l^m = \hbar \sqrt{l(l+1) - m(m+1)} \Psi_l^{m+1}. \quad (12.5.8)$$

Taken together, eqs. (12.5.3), (12.5.7) and (12.5.8), combined with  $L^2 = (L_+ L_- + L_- L_+)/2 + L_z^2$  from eq. (8.1.15), show that this wavefunction also obeys

$$L^2 \Psi_l^m = \hbar^2 l(l+1) \Psi_l^m. \quad (12.5.9)$$

Since the spherical harmonic  $Y_l^m$  is the unique  $(\theta, \phi)$  function with  $L^2$  eigenvalue  $\hbar^2 l(l+1)$  and  $L_z$  eigenvalue  $\hbar m$ , eqs. (12.5.3) and (12.5.9) show that  $\Psi_l^m$  must be proportional to it, and

$$\Psi_l^m(\theta, \phi) = c_{l_1 l_2 l m} Y_l^m(\theta, \phi) \quad (12.5.10)$$

as claimed, for some constant  $c_{l_1 l_2 l m}$  (independent of  $\theta, \phi$ ).

Furthermore, by acting with  $L_-$  on both sides of the previous equation, we obtain

$$L_- \Psi_l^m = c_{l_1 l_2 l m} L_- Y_l^m = c_{l_1 l_2 l m} \hbar \sqrt{l(l+1) - m(m-1)} Y_l^{m-1}, \quad (12.5.11)$$

but also from eq. (12.5.7) this is equal to

$$L_- \Psi_l^m = \hbar \sqrt{l(l+1) - m(m-1)} c_{l_1 l_2 l, m-1} Y_l^{m-1}. \quad (12.5.12)$$

Comparing these informs us that  $c_{l_1 l_2 l m} = c_{l_1 l_2 l, m-1}$  for all  $-l+1 \leq m \leq l$ , so the constant of proportionality does not actually depend on  $m$ , and we therefore drop that label.

All of the preceding just served to prove that we can write

$$\sum_{m_1, m_2} Y_{l_1}^{m_1}(\theta, \phi) Y_{l_2}^{m_2}(\theta, \phi) C_{m_1 m_2 m}^{l_1 l_2 l} = c_{l_1 l_2 l} Y_l^m(\theta, \phi) \quad (12.5.13)$$

for some proportionality constant  $c_{l_1 l_2 l}$ . Our remaining task is to identify this constant. Fortunately, there is a trick to do this very easily; just consider the special case  $\theta = 0$ . Using the fact found in eq. (8.6.53),

$$Y_l^m(0, \phi) = \delta_{m0} \sqrt{\frac{2l+1}{4\pi}}, \quad (12.5.14)$$

the double sum in eq. (12.5.13) collapses to a single term with  $m_1 = m_2 = 0$ , and it reads

$$\sqrt{\frac{2l_1+1}{4\pi}} \sqrt{\frac{2l_2+1}{4\pi}} C_{000}^{l_1 l_2 l} = c_{l_1 l_2 l} \sqrt{\frac{2l+1}{4\pi}}. \quad (12.5.15)$$

Solving this for  $c_{l_1 l_2 l}$ , eq. (12.5.13) becomes

$$\sum_{m_1, m_2} Y_{l_1}^{m_1}(\theta, \phi) Y_{l_2}^{m_2}(\theta, \phi) C_{m_1 m_2 m}^{l_1 l_2 l} = \sqrt{\frac{(2l_1+1)(2l_2+1)}{4\pi(2l+1)}} C_{000}^{l_1 l_2 l} Y_l^m(\theta, \phi). \quad (12.5.16)$$

This accomplishes our goal.

To derive a related interesting and useful identity, start with eq. (12.5.16), multiply by  $C_{m'_1 m'_2 m}^{l_1 l_2 l}$ , then sum over all  $l, m$ , then apply the orthogonality of the Clebsch–Gordan coefficients of eq. (12.4.20) on the left side, and finally relabel  $m'_1 \rightarrow m_1$  and  $m'_2 \rightarrow m_2$ . The result is

$$Y_{l_1}^{m_1}(\theta, \phi) Y_{l_2}^{m_2}(\theta, \phi) = \sum_{l=|l_1-l_2|}^{l_1+l_2} \sqrt{\frac{(2l_1+1)(2l_2+1)}{4\pi(2l+1)}} C_{000}^{l_1 l_2 l} C_{m_1 m_2 m}^{l_1 l_2 l} Y_l^m(\theta, \phi), \quad (12.5.17)$$

where  $m = m_1 + m_2$  on the right side. The product of any two spherical harmonics is thus an appropriately weighted sum over the spherical harmonics selected by the corresponding addition of angular momenta.

Another closely related identity follows. Start from eq. (12.5.17), multiply by  $Y_{l'}^{m'}(\theta, \phi)^*$ , integrate over the angular coordinates, then apply the orthonormality of the spherical harmonics using eq. (8.6.31) on the right side, and finally relabel  $l' \rightarrow l$  and  $m' \rightarrow m$ . The result is

$$\int d\Omega [Y_l^m(\theta, \phi)]^* Y_{l_1}^{m_1}(\theta, \phi) Y_{l_2}^{m_2}(\theta, \phi) = \sqrt{\frac{(2l_1+1)(2l_2+1)}{4\pi(2l+1)}} C_{000}^{l_1 l_2 l} C_{m_1 m_2 m}^{l_1 l_2 l}. \quad (12.5.18)$$

This is the **Wigner–Eckart formula for spherical harmonics**. It is a special case of the general Wigner–Eckart theorem (a statement about matrix elements of operators, and applicable not just to orbital angular momentum) to be derived in section 13.3. Remarkably, the integral only depends on the magnetic quantum numbers  $m_1, m_2$ , and  $m$  through the coefficient  $C_{m_1 m_2 m}^{l_1 l_2 l}$ , so the Clebsch–Gordan selection rules (12.4.5)–(12.4.7) also govern which integrals of this type can be non-zero. This formula can be used to evaluate the integral of any product of three spherical harmonics (with or without complex conjugates), simply by making use of eq. (8.6.51).

## 12.6 Exercises

**Exercise 12.1.** Two angular momentum operators  $\vec{L}$  and  $\vec{S}$  each satisfy the commutation algebra of the form in eq. (5.3.33). Use this to show that all three components of  $\vec{J} = \vec{L} + \vec{S}$  commute with  $\vec{L} \cdot \vec{S}$ .

**Exercise 12.2.** An atomic electron is in a state with orbital angular momentum  $l = 2$ , with  $L_z$  and  $S_z$  eigenvalues 0 and  $\hbar/2$ , respectively. If its total angular momentum squared is measured to be  $J^2 = \hbar^2 j(j+1)$ , what are the possible results for  $j$ , and their probabilities?

**Exercise 12.3.** Two particles both have spin 1/2, and one of them is in a  $p$ -wave orbital angular momentum state while the other is in a  $d$ -wave orbital angular momentum state. If  $\vec{J}$  is the total angular momentum operator, what are the allowed eigenvalues of  $J^2$ , and their degeneracies? [Hint: start by figuring out the total number of basis states by multiplying

together the individual angular momentum multiplicities. You should find 60, so that should be the sum of the degeneracies.]

**Exercise 12.4.** Consider a quantum system with two independent spin-1/2 operators,  $\vec{S}_1$  and  $\vec{S}_2$ , so that the state space is spanned by an orthobasis of  $S_{1z}$  and  $S_{2z}$  eigenstates  $|\uparrow\uparrow\rangle$ ,  $|\uparrow\downarrow\rangle$ ,  $|\downarrow\uparrow\rangle$ , and  $|\downarrow\downarrow\rangle$ . In each ket, the first entry labels states with  $S_{1z}$  eigenvalue  $\pm\hbar/2$ , and the second entry labels states with  $S_{2z}$  eigenvalue  $\pm\hbar/2$ . At time  $t = 0$ , the system is in the state

$$|\psi(0)\rangle = \frac{1}{2}|\uparrow\uparrow\rangle + \frac{1}{2}|\uparrow\downarrow\rangle + \frac{1}{\sqrt{2}}|\downarrow\downarrow\rangle. \quad (12.6.1)$$

(a) At time  $t = 0$ , we simultaneously measure  $S^2$  and  $S_z$ , where  $\vec{S} = \vec{S}_1 + \vec{S}_2$  is the total spin operator. What are the possible outcomes, and their probabilities?

(b) Suppose that instead of the above measurements, we let the system evolve until time  $t$ , with the Hamiltonian  $H = \omega_1 S_{1z} + \omega_2 S_{2z}$ , where  $\omega_1$  and  $\omega_2$  are constants. What is the state at time  $t$ ? If we measure  $S_{1z}$  at time  $t$ , what are the possible outcomes, and their probabilities? Use these results to find the expectation value  $\langle S_{1z} \rangle$  as a function of time.

(c) Now suppose that instead the Hamiltonian of the system is  $H = a\vec{S}_1 \cdot \vec{S}_2$ , where  $a$  is a constant. What is the state at time  $t$ , and what are the possible outcomes and probabilities for a measurement of  $S_{1z}$ ? Use these results to find  $\langle S_{1z} \rangle$  as a function of time. [Hint: to find the state at time  $t$ , it is best to use the total angular momentum basis, in which  $H$  is diagonal.]

**Exercise 12.5.** Consider the hyperfine splitting of the  $n = 1$  level of the hydrogen atom in the presence of a constant external magnetic field  $\vec{B} = B\hat{z}$ . Since we are only treating the ground state, there is no orbital angular momentum  $\vec{L}$ . Let the electron and proton spin operators be  $\vec{S}$  and  $\vec{I}$  respectively, and let their total be  $\vec{J} = \vec{S} + \vec{I}$ . The Hamiltonian is

$$H = \frac{E_\gamma}{\hbar^2} \vec{S} \cdot \vec{I} + \frac{2\mu_B}{\hbar} \vec{B} \cdot \vec{S}, \quad (12.6.2)$$

where  $\mu_B$  is the Bohr magneton and  $E_\gamma$  is the energy of the 21.4 cm line. (There is also an interaction term  $-(g_p\mu_N/\hbar)\vec{B} \cdot \vec{I}$ , but it is neglected because  $\mu_N \ll \mu_B$ .)

(a) Evaluate each of the two operators  $\vec{S} \cdot \vec{I}$  and  $\vec{B} \cdot \vec{S}$  acting on each of the product basis states labeled by the eigenvalues of  $S_z$  and  $I_z$ , and find the matrix representation of  $H$  in that basis.

(b) Repeat part (a), but this time use the total angular momentum basis labeled by the eigenvalues of  $J^2$  and  $J_z$ .

(c) In the limit that  $B$  is so large that  $E_\gamma$  can be neglected, find the energy eigenvalues, and the corresponding energy eigenstates in the product orthobasis.

(d) In the limit that  $B$  is so small that it can be neglected, find the energy eigenvalues and the corresponding energy eigenstates in the total angular momentum basis.

- (e) Find the energy eigenvalues for general  $B$ . Expand them to show agreement with the special limits obtained in parts (c) and (d). (You do not need to find the energy eigenstates.)
- (f) Graph the energy eigenvalues in units of  $\mu\text{eV}$  as a function of  $0 < B < 1200$  gauss. Label the energies near  $B = 0$  with  $j = 0$  and  $j = 1$ , and the energies for large  $B$  with  $m_S$  and  $m_I$ .

**Exercise 12.6.** The electron and its antiparticle, the positron, can form bound states before they annihilate, called positronium. The positronium states with lowest energy and zero orbital angular momenta have a Hamiltonian

$$H = \frac{a}{\hbar^2} \vec{S}_e \cdot \vec{S}_p + \frac{b}{\hbar} (S_{ez} - S_{pz}), \quad (12.6.3)$$

where  $\vec{S}_e$  and  $\vec{S}_p$  are the spin operators for the electron and positron respectively, and  $a = 8.41 \times 10^{-4}$  eV, and  $b = 2\hbar e B / m_e c$  where  $B$  is an external magnetic field in the  $\hat{z}$  direction.

- (a) Working in the total spin basis of eigenstates of  $S^2$  and  $S_z$  where  $\vec{S} = \vec{S}_e + \vec{S}_p$ , find the Hamiltonian as a  $4 \times 4$  matrix in terms of  $a$  and  $b$ , and find its eigenvalues and eigenvectors.
- (b) The  $^1S_0$  singlet (parapositronium) state decays rather quickly by annihilation to two photons, with a lifetime of about  $1.25 \times 10^{-10}$  seconds, while the  $^3S_1$  triplet (orthopositronium) states are relatively stable, with a lifetime of about  $1.42 \times 10^{-7}$  seconds. (Assume that these statements do not depend on the applied magnetic field.) Estimate the magnetic field  $B$  required to make the ground state lifetime longer by 10%.

**Exercise 12.7.** For the addition of two angular momenta  $j_1 = j_2 = 3/2$ , find all of the total angular momentum states in terms of the product basis states, completing eqs. (12.4.45)-(12.4.47). Summarize the results as a list of the nonzero Clebsch–Gordan coefficients.

**Exercise 12.8.** General expressions for the Clebsch–Gordan coefficients for the addition of angular momenta  $l$  and 1 were given in eqs. (12.4.13)-(12.4.15). Use the results to write each of  $Y_l^m(\theta, \phi) \cos \theta$  and  $Y_l^m(\theta, \phi) \sin \theta \cos \phi$  and  $Y_l^m(\theta, \phi) \sin \theta \sin \phi$  as linear combinations of other spherical harmonics, with constant coefficients.

## 13 Tensor operators and useful rules they obey

### 13.1 Irreducible tensor operators

Consider a finite spatial rotation generated by a unitary operator, so that a state ket  $|\psi\rangle$  is transformed to  $|\psi'\rangle = U(\vec{\alpha})|\psi\rangle$ , as discussed in section 5.3. Recall, from the discussion surrounding eqs. (2.5.26)–(2.5.31), that for any operator  $A$  we can consider the corresponding rotated version defined by  $A' = U(\vec{\alpha})AU(\vec{\alpha})^\dagger$ , so that any matrix element  $\langle\chi|A|\psi\rangle$  is equal to the matrix element of the rotated operator for the rotated states,  $\langle\chi'|A'|\psi'\rangle$ . We will now study operators that transform under rotations in a simple and useful way, specifically, as irreducible representations of the rotation group.

An **irreducible tensor operator** (also known as a **spherical tensor operator**) of order  $j$  is a collection of  $2j + 1$  component operators labeled by an index  $m$ ,

$$T_m^{(j)}, \quad (m = -j, \dots, j), \quad (13.1.1)$$

which, by definition, are required to obey the following transformation rule for every unitary rotation operator  $U(\vec{\alpha})$ ,

$$U(\vec{\alpha}) T_m^{(j)} U(\vec{\alpha})^\dagger = \sum_{m'=-j}^j T_{m'}^{(j)} D_{m'm}^{(j)}(\vec{\alpha}), \quad (13.1.2)$$

with the Wigner rotation matrices  $D^{(j)}(\vec{\alpha})$  as defined in eq. (8.5.2). Using the form of  $U(\vec{\alpha})$  for an infinitesimal rotation in terms of the total angular momentum operator  $\vec{J}$  in eq. (8.5.1), and the actions of  $J_z$ ,  $J_+$ , and  $J_-$  on angular momentum eigenstates as given in eqs. (8.1.7), (8.1.22), and (8.1.24), which are reproduced here for convenience,

$$J_z |j, m\rangle = \hbar m |j, m\rangle, \quad (13.1.3)$$

$$J_+ |j, m\rangle = \hbar \sqrt{j(j+1) - m(m+1)} |j, m+1\rangle, \quad (13.1.4)$$

$$J_- |j, m\rangle = \hbar \sqrt{j(j+1) - m(m-1)} |j, m-1\rangle, \quad (13.1.5)$$

it is left to Exercise 13.1 to show that the components of an irreducible tensor operator must obey the commutation relations

$$[J_z, T_m^{(j)}] = \hbar m T_m^{(j)}, \quad (13.1.6)$$

$$[J_+, T_m^{(j)}] = \hbar \sqrt{j(j+1) - m(m+1)} T_{m+1}^{(j)}, \quad (13.1.7)$$

$$[J_-, T_m^{(j)}] = \hbar \sqrt{j(j+1) - m(m-1)} T_{m-1}^{(j)}. \quad (13.1.8)$$

Equations (13.1.6)–(13.1.8) serve as an equivalent necessary and sufficient requirement for  $T_m^{(j)}$  to form an irreducible tensor operator. Their similarity to eqs. (13.1.3)–(13.1.5) suggests that

the operators  $T_m^{(j)}$  “carry” angular momentum as indicated by the quantum number labels  $j$  and  $m$ . Indeed this intuition is valid, as we will explain in the following, with the most precise statement to be given in eq. (13.3.9).

The simplest type of irreducible tensor operator is obtained for  $j = 0$  with a single component, and is called a **scalar operator**. A scalar operator  $S$  obeys  $U(\vec{\alpha}) S U(\vec{\alpha})^\dagger = S$ , or equivalently it commutes with the total angular momentum operators  $J_z$ ,  $J_+$ ,  $J_-$ , and therefore also with  $J_x$  and  $J_y$  and  $J^2$ . Consider a state  $|\alpha, j, m\rangle$  that is an eigenstate of  $J^2$  and  $J_z$  with eigenvalues  $\hbar^2 j(j+1)$  and  $\hbar m$ , with  $\alpha$  a degeneracy label. It follows that the state  $S|\alpha, j, m\rangle$  is also an eigenstate of  $J^2$  and  $J_z$  with the same eigenvalues. In that sense, the scalar operator  $S$  carries no total angular momentum.

Irreducible tensor operators with  $j = 1/2$ , or more generally with half-integer  $j$ , are called **spinor operators**. However, these cannot be observables, because they are necessarily double-valued; under a rotation by  $2\pi$  they acquire a minus sign, just as we saw for matrix representations of rotations of states with half-integer angular momentum in eqs. (8.5.16) and (8.5.19). Observables can be constructed by taking the product of an even number of spinor operators, using Theorem 13.1.1 below. We will not discuss them further here, although they do play a very important role in quantum field theories with fermions.

An irreducible tensor operator with order  $j = 1$  and three components  $m = -1, 0, 1$  is called a **vector operator**  $V_m^{(1)}$ . These can be used to define the familiar rectangular  $x, y, z$  components of  $\vec{V} = \hat{x}V_x + \hat{y}V_y + \hat{z}V_z$ , as follows:

$$V_x = (V_{-1}^{(1)} - V_1^{(1)})/\sqrt{2}, \quad (13.1.9)$$

$$V_y = i(V_{-1}^{(1)} + V_1^{(1)})/\sqrt{2}, \quad (13.1.10)$$

$$V_z = V_0^{(1)}. \quad (13.1.11)$$

These can be shown, using eqs. (13.1.6)–(13.1.8) with  $j = 1$ , to obey the commutation relations

$$[J_a, V_b] = i\hbar\epsilon_{abc}V_c, \quad (a, b, c = x, y, z), \quad (13.1.12)$$

which is often taken as the defining property of a vector operator in rectangular coordinates. For example, the rectangular coordinate position operators found in  $\vec{R} = \hat{x}X + \hat{y}Y + \hat{z}Z$  can be expressed in terms of components of an irreducible tensor operator  $R^{(1)}$ , as

$$R_0^{(1)} = Z, \quad (13.1.13)$$

$$R_1^{(1)} = -(X + iY)/\sqrt{2}, \quad (13.1.14)$$

$$R_{-1}^{(1)} = (X - iY)/\sqrt{2}, \quad (13.1.15)$$

and it is left as an exercise to check that the requisite commutation relations (13.1.6)–(13.1.8) are indeed satisfied. Besides the position operator, other examples of vector operators are the



momentum operator  $\vec{P}$ , the orbital angular momentum  $\vec{L}$ , the spin  $\vec{S}$ , and the total angular momentum operator itself,  $\vec{J} = \vec{L} + \vec{S}$ . Note that the angular momentum raising and lowering operators defined by eq. (8.1.8) are actually given in the irreducible tensor component formalism by  $J_+ = -\sqrt{2}J_1^{(1)}$  and  $J_- = \sqrt{2}J_{-1}^{(1)}$ , respectively, while  $J_z = J_0^{(1)}$ .

Given any two irreducible tensor operators, one can multiply them to construct others. The way to do this is specified by the following theorem.

**Theorem 13.1.1. (Product rules for tensor operators)** *If  $T^{(j_1)}$  and  $S^{(j_2)}$  are irreducible tensor operators of orders  $j_1$  and  $j_2$  respectively, and  $C_{m_1 m_2 m}^{j_1 j_2 j}$  are the Clebsch–Gordan coefficients defined in eq. (12.4.4), then*

$$W_m^{(j)} = \sum_{m_1} \sum_{m_2} T_{m_1}^{(j_1)} S_{m_2}^{(j_2)} C_{m_1 m_2 m}^{j_1 j_2 j} \quad (m = -j, \dots, j) \quad (13.1.16)$$

*is an irreducible tensor operator of order  $j$ . Furthermore,*

$$T_{m_1}^{(j_1)} S_{m_2}^{(j_2)} = \sum_j \sum_m W_m^{(j)} C_{m_1 m_2 m}^{j_1 j_2 j}. \quad (13.1.17)$$

The proof is left as Exercise 13.3.

As an example, consider two vector operators  $V^{(1)}$  and  $U^{(1)}$ . Then, using the results of eqs. (12.4.34)–(12.4.42) to extract the necessary Clebsch–Gordan coefficients for  $1 \otimes 1 = 0 \oplus 1 \oplus 2$ , we apply eq. (13.1.16) to construct the following three tensor operators, of orders 0, 1, and 2:

$$W_0^{(0)} = \left( V_1^{(1)} U_{-1}^{(1)} + V_{-1}^{(1)} U_1^{(1)} - V_0^{(1)} U_0^{(1)} \right) / \sqrt{3}, \quad (13.1.18)$$

and

$$W_{\pm 1}^{(1)} = \pm \left( V_{\pm 1}^{(1)} U_0^{(1)} - V_0^{(1)} U_{\pm 1}^{(1)} \right) / \sqrt{2}, \quad (13.1.19)$$

$$W_0^{(1)} = \left( V_1^{(1)} U_{-1}^{(1)} - V_{-1}^{(1)} U_1^{(1)} \right) / \sqrt{2}, \quad (13.1.20)$$

and

$$W_{\pm 2}^{(2)} = V_{\pm 1}^{(1)} U_{\pm 1}^{(1)}, \quad (13.1.21)$$

$$W_{\pm 1}^{(2)} = \left( V_{\pm 1}^{(1)} U_0^{(1)} + V_0^{(1)} U_{\pm 1}^{(1)} \right) / \sqrt{2}, \quad (13.1.22)$$

$$W_0^{(2)} = \left( V_1^{(1)} U_{-1}^{(1)} + V_{-1}^{(1)} U_1^{(1)} + 2V_0^{(1)} U_0^{(1)} \right) / \sqrt{6}. \quad (13.1.23)$$

Now,  $W_0^{(0)}$  is a scalar operator, equal to  $-\vec{V} \cdot \vec{U} / \sqrt{3}$ . It is also not hard to check that  $W_m^{(1)}$  forms a vector operator, and using eqs. (13.1.9)–(13.1.11), the corresponding rectangular components

are found to be those of  $i(\vec{V} \times \vec{U})/\sqrt{2}$ . The  $j = 2$  tensor operator can be rewritten, in terms of the rectangular components of  $\vec{V}$  and  $\vec{U}$ , as

$$W_{\pm 2}^{(2)} = (V_x \pm iV_y)(U_x \pm iU_y)/2, \quad (13.1.24)$$

$$W_{\pm 1}^{(2)} = [(\mp V_x - iV_y)U_z + V_z(\mp U_x - iU_y)]/2, \quad (13.1.25)$$

$$W_0^{(2)} = (2V_zU_z - V_xU_x - V_yU_y)/\sqrt{6}. \quad (13.1.26)$$

A different basis for these  $j = 2$  operators consists of the five operators

$$\begin{aligned} V_xU_y + V_yU_x, & \quad V_xU_z + V_zU_x, & \quad V_yU_z + V_zU_y, \\ V_xU_x - V_yU_y, & \quad 2V_zU_z - V_xU_x - V_yU_y, \end{aligned} \quad (13.1.27)$$

which can also be recognized as a basis for the entries of the traceless symmetric  $3 \times 3$  matrix

$$\mathcal{W}_{ab} = (V_aU_b + V_bU_a)/2 - \delta_{ab}\vec{V} \cdot \vec{U}/3, \quad (a, b = x, y, z). \quad (13.1.28)$$

Therefore,  $W_m^{(2)}$  and  $\mathcal{W}_{ab}$  are the same operators in different guises. In terms of rectangular components, the product of  $V_a$  and  $U_b$  decomposes to a linear combination of the irreducible tensor operators with  $j = 0, 1, 2$ , as

$$V_aU_b = \delta_{ab}\vec{V} \cdot \vec{U}/3 + \epsilon_{abc}(\vec{V} \times \vec{U})_c/2 + \mathcal{W}_{ab}. \quad (13.1.29)$$

This can be checked using  $\epsilon_{abc}(\vec{V} \times \vec{U})_c = V_aU_b - V_bU_a$ . Thus  $V_aU_b$  is an example of a **reducible tensor operator**.<sup>†</sup>

As a further specialization, we can take  $\vec{V} = \vec{U} = \vec{R}$ , the position operator. The resulting  $j = 2$  spherical tensor is called the **quadrupole moment operator**,

$$Q_{\pm 2}^{(2)} = (X \pm iY)^2/2, \quad (13.1.30)$$

$$Q_{\pm 1}^{(2)} = (\mp X - iY)Z, \quad (13.1.31)$$

$$Q_0^{(2)} = (2Z^2 - X^2 - Y^2)/\sqrt{6}. \quad (13.1.32)$$

This five-component operator has the same content as the five rectangular components of the traceless symmetric tensor

$$\mathcal{Q}_{ab} = R_aR_b - \delta_{ab}R^2/3, \quad (13.1.33)$$

in the sense that each is comprised of (complex) linear combinations of the other. This operator will appear in our study of absorption and emission of electromagnetic radiation, in section 22.5.

---

<sup>†</sup>A **Cartesian tensor operator**  $T_{a_1, \dots, a_n}$  with  $n \geq 2$  indices  $a_i = x, y, z$  is generically (if no special symmetry, antisymmetry, or trace conditions are imposed) reducible, in the sense that its entries can be written as linear combinations of irreducible tensor operators with more than one  $j$ .

It is straightforward to check that the position representation of the quadrupole moment operator can be expressed in terms of spherical harmonics as

$$Q_m^{(2)} = \sqrt{\frac{8\pi}{15}} r^2 Y_2^m(\theta, \phi) \quad (13.1.34)$$

This is not an accident, as any  $2l+1$ -component operator with position representation of the form  $f(r)Y_l^m(\theta, \phi)$  can be shown to satisfy the requirements (13.1.6)-(13.1.8) for a tensor operator of order  $l$ . In particular, consider the **multipole moment operator**

$$T_m^{(l)} = r^l Y_l^m(\theta, \phi). \quad (13.1.35)$$

As we have just seen, the quadrupole moment operator is the special case  $l = 2$  (up to a normalization). The dipole moment operator with  $l = 1$  is equivalent to the position vector operator, as in eqs. (13.1.13)-(13.1.15). The octopole moment operator has  $l = 3$ , etc. To justify the terminology, recall that for a classical charge distribution  $\rho(\vec{r})$ , the electrostatic potential is

$$\Phi(\vec{r}) = \int d^3\vec{r}' \frac{\rho(\vec{r}')}{|\vec{r} - \vec{r}'|}. \quad (13.1.36)$$

Applying the expansion formula (8.6.82), this can be rewritten as

$$\Phi(\vec{r}) = \sum_{l=0}^{\infty} \frac{4\pi}{2l+1} \sum_{m=-l}^l q_{lm}^* \frac{Y_l^m(\theta, \phi)}{r^{l+1}}, \quad (13.1.37)$$

where

$$q_{lm} = \int d^3\vec{r}' \rho(\vec{r}') T_m^{(l)}(r', \theta', \phi'). \quad (13.1.38)$$

Thus, for a particle of charge  $q$  and wavefunction  $\psi(\vec{r})$ , the  $q_{lm}$  can be obtained by integrating  $T_m^{(l)}$  weighted by the charge probability distribution  $\rho(\vec{r}) = q|\psi(\vec{r})|^2$ . In bra-ket form,

$$q_{lm} = q \langle \psi | T_m^{(l)} | \psi \rangle, \quad (13.1.39)$$

the expectation value of the multipole moment operator.

## 13.2 Selection rules for scalar and vector operators and the Landé projection formula

As remarked in the previous section, scalar operators do not change the total angular momentum quantum numbers of the states they act on. Furthermore, their matrix elements between total angular momentum ( $J^2, J_z$ ) eigenstates do not depend on the magnetic quantum number. Both properties are incorporated in the following:

**Theorem 13.2.1. (Scalar operator selection rule)** *If  $S$  is a scalar operator, and  $|\alpha jm\rangle$  are eigenstates of the total angular momentum operators  $J^2$  and  $J_z$ , with additional labels  $\alpha, \beta, \dots$ , then*

$$\langle \beta j' m' | S | \alpha j m \rangle = \delta_{j'j} \delta_{m'm} \langle \beta j || S || \alpha j \rangle, \quad (13.2.1)$$

where  $\langle \beta j || S || \alpha j \rangle$  does not depend on  $m$ .

As terminology, the object  $\langle \beta j || S || \alpha j \rangle$  is called a **reduced matrix element**. This is not a matrix element in the usual sense, but rather just a symbol denoting a number that depends on  $\alpha, \beta$ , and  $j$  and on the choice of the operator  $S$ , but not on  $m$  or  $m'$ .

To prove Theorem 13.2.1, consider  $\langle \beta j' m' | [J_z, S] | \alpha j m \rangle$ , which is 0 from the definition of a scalar operator. Evaluating each  $J_z$  acting on the bra or ket to which it is adjacent, this immediately tells us that  $(\hbar m' - \hbar m) \langle \beta j' m' | S | \alpha j m \rangle = 0$ , so the matrix element in question can only be non-zero if  $m' = m$ . Similarly, the matrix element of  $[J^2, S] = 0$  tells us that  $[\hbar^2 j'(j' + 1) - \hbar^2 j(j + 1)] \langle \beta j' m' | S | \alpha j m \rangle = 0$ , so the matrix element can only be non-zero if  $j' = j$ , since the other root  $j' = -j - 1$  is impossible because  $j$  and  $j'$  are both non-negative. It remains to show that the matrix element is independent of  $m$ . This follows from

$$\langle \beta, j, m+1 | S | \alpha, j, m+1 \rangle = \langle \beta, j, m | S | \alpha, j, m \rangle, \quad (13.2.2)$$

which can be obtained by using the following facts:  $|\alpha, j, m+1\rangle$  is proportional to  $J_+ |\alpha, j, m\rangle$ ; and  $\langle \beta, j, m |$  is proportional to  $\langle \beta, j, m+1 | J_+$  with the same constant of proportionality, namely  $\hbar \sqrt{j(j+1) - m(m+1)}$ ; and  $[J_+, S] = 0$  for a scalar operator.

Theorem 13.2.1 is a powerful selection rule which can be applied, for example, if  $S$  is a part of the Hamiltonian, which is often invariant under rotations and therefore a scalar operator. (Examples of exceptions include a charged particle in the presence of an external electric or magnetic field, which will always pick out a special direction that destroys the rotational symmetry. For example, if there is a constant external electric field  $\vec{E}$ , the Hamiltonian will contain a term proportional to  $\vec{E} \cdot \vec{R}$ , which is not a scalar operator. This is because  $\vec{E}$ , being constant, does not transform as a vector operator in the sense we have defined.) This theorem will even be useful to us in the seemingly trivial case that  $S$  is the identity operator.

We now turn our attention to vector operators, which in this section we will treat in terms of their rectangular coordinate components  $V_a$  for  $a = x, y, z$  with  $\vec{V} = \hat{x}V_x + \hat{y}V_y + \hat{z}V_z$ . By definition, these satisfy

$$[J_a, V_b] = i\hbar \epsilon_{abc} V_c. \quad (13.2.3)$$

One useful observation is that the operator  $\vec{J} \cdot \vec{V} = \vec{V} \cdot \vec{J}$  does not depend on the ordering, since each  $J_a$  commutes with the corresponding component  $V_a$ . Furthermore, it is a scalar operator:

$$[J_a, \vec{J} \cdot \vec{V}] = 0. \quad (13.2.4)$$

One can also check straightforwardly that

$$[J^2, V_a] = i\hbar\epsilon_{abc}(V_b J_c + J_c V_b). \quad (13.2.5)$$

From this, and the identity  $\epsilon_{abc}\epsilon_{dec} = \delta_{ad}\delta_{be} - \delta_{ae}\delta_{bd}$ , one can establish the double commutator relation

$$[J^2, [J^2, V_a]] = 2\hbar^2 (J^2 V_a + V_a J^2 - 2J_a(\vec{J} \cdot \vec{V})). \quad (13.2.6)$$

We now exploit eq. (13.2.6) by evaluating both sides as a matrix element between orthobasis states  $\langle\beta, j', m'|\$  and  $|\alpha, j, m\rangle$ , where  $\alpha$  and  $\beta$  are degeneracy labels. Each instance of  $J^2$  can be immediately evaluated as either  $\hbar^2 j(j+1)$  or  $\hbar^2 j'(j'+1)$ , leading to

$$\begin{aligned} \hbar^4 ([j(j+1) - j'(j'+1)]^2 - 2[j(j+1) + j'(j'+1)]) \langle\beta, j', m'|V_a|\alpha, j, m\rangle = \\ -4\hbar^2 \langle\beta, j', m'|J_a(\vec{J} \cdot \vec{V})|\alpha, j, m\rangle. \end{aligned} \quad (13.2.7)$$

The matrix element on the right can be separated, by using completeness, to get

$$\langle\beta, j', m'|J_a(\vec{J} \cdot \vec{V})|\alpha, j, m\rangle = \sum_{\gamma, j'', m''} \langle\beta, j', m'|J_a|\gamma, j'', m''\rangle \langle\gamma, j'', m''|\vec{J} \cdot \vec{V}|\alpha, j, m\rangle. \quad (13.2.8)$$

Here, the first matrix element vanishes unless  $\gamma = \beta$  and  $j'' = j'$ , while according to Theorem 13.2.1 the second matrix element vanishes unless  $j'' = j$  and  $m'' = m$ . Therefore, only one term in the completeness sum contributes, and eq. (13.2.7) simplifies to a remarkably useful formula,

$$\begin{aligned} \hbar^2(j + j' + 2)(j + j')[(j - j')^2 - 1] \langle\beta, j', m'|V_a|\alpha, j, m\rangle = \\ -4\delta_{jj'} \langle j, m'|J_a|j, m\rangle \langle\beta, j, m|\vec{J} \cdot \vec{V}|\alpha, j, m\rangle, \end{aligned} \quad (13.2.9)$$

where the irrelevant label  $\beta$  has been removed from the first matrix element on the right.

One important application of eq. (13.2.9) is the special case  $j' = j$ . Then it immediately reduces to the following result due to Alfred Landé:

**Theorem 13.2.2. (Vector operator projection rule)** *If an operator  $\vec{V}$  transforms as a vector with respect to rotations generated by the angular momentum operator  $\vec{J}$ , then the Landé projection formula holds:*

$$\langle\beta, j, m'|\vec{V}|\alpha, j, m\rangle = \langle j, m'|\vec{J}|j, m\rangle \frac{\langle\beta, j|\vec{J} \cdot \vec{V}|\alpha, j\rangle}{\hbar^2 j(j+1)} \quad (13.2.10)$$

for  $j \neq 0$ , where the reduced matrix element

$$\langle \beta, j \| \vec{J} \cdot \vec{V} \| \alpha, j \rangle \equiv \langle \beta, j, m | \vec{J} \cdot \vec{V} | \alpha, j, m \rangle \quad (13.2.11)$$

does not depend on  $m$  or  $m'$ . For the special case  $j = 0$ ,

$$\langle \beta, 0, 0 | \vec{V} | \alpha, 0, 0 \rangle = 0. \quad (13.2.12)$$

Because  $\vec{J} \cdot \vec{V}$  is a scalar operator, Theorem 13.2.1 reminds us that the right side of eq. (13.2.11) does not actually depend on  $m$ , as claimed. So, in applications, we can pick any convenient value for  $m$  to do the evaluation of  $\langle \beta, j \| \vec{J} \cdot \vec{V} \| \alpha, j \rangle$ . (The most convenient choice is usually either  $m = j$  or  $m = 0$ .) Of course, if  $j = 0$  then eq. (13.2.10) fails, but in that case  $j = m = m' = 0$ , and the vanishing of  $\langle \beta, 0, 0 | \vec{V} | \alpha, 0, 0 \rangle$  follows from spherical symmetry. The  $j = 0$  states do not have any special direction along which the matrix element of a vector operator could point.

Intuitively, the Landé projection formula says that, within each subspace of fixed  $j$ , all vector operators are proportional to each other, and in particular to the angular momentum operator. As a check, if we take  $\vec{V} = \vec{J}$ , then  $\vec{J} \cdot \vec{V} = J^2$ , and eq. (13.2.10) is satisfied. The reason for the word “projection” in the name is that it implies that fixed- $j$  matrix elements of  $\vec{V}$  are equal to those of  $\vec{J}(\vec{J} \cdot \vec{V})/J^2$ , the formal geometric projection of  $\vec{V}$  onto the angular momentum operator direction. The projection formula is a particularly useful special case of the Wigner–Eckart theorem discussed in the next section.

Another important application of eq. (13.2.9) occurs if we take  $\vec{V} = \vec{R}$ , the position operator. As a notable example, this will be useful for evaluating absorption and emission of electromagnetic radiation in the electric dipole approximation (in section 22.4). For this reason, the conditions on the matrix elements of the components of  $\vec{R}$  that we are about to derive are often called the **dipole selection rules**.

When considering matrix elements of  $\vec{R}$ , we can take  $\vec{J}$  to be just the orbital angular momentum operator  $\vec{L}$ , since that is what generates rotations for  $\vec{R}$ ; the spin is not relevant. Further,  $\vec{L} \cdot \vec{R} = (\vec{R} \times \vec{P}) \cdot \vec{R} = 0$ , so the right side of eq. (13.2.9) vanishes, and it gives simply

$$\hbar^2(l + l' + 2)(l + l')[l(l - l')^2 - 1] \langle \beta, l', m' | \vec{R} | \alpha, l, m \rangle = 0. \quad (13.2.13)$$

Therefore, for the matrix element to be non-zero, we need

$$(l + l' + 2)(l + l')[l(l - l')^2 - 1] = 0. \quad (13.2.14)$$

Now,  $l + l' + 2$  cannot vanish because  $l$  and  $l'$  are always non-negative. Also, the condition  $l + l' = 0$  can only be satisfied if  $l = l' = 0$ , but then  $\langle \beta, l', m' | \vec{R} | \alpha, l, m \rangle$  can only be  $\langle \beta, 0, 0 | \vec{R} | \alpha, 0, 0 \rangle$ , which must vanish anyway due to the spherical symmetry of the state wavefunctions. From

eq. (13.2.14), the only other possibility for a non-vanishing matrix element is  $(l-l')^2 = 1$ . Thus, we have the first dipole selection rule,

$$\langle \beta, l', m' | \vec{R} | \alpha, l, m \rangle \neq 0 \quad \text{requires } l' - l = \pm 1. \quad (13.2.15)$$

Since this holds for the whole vector  $\vec{R}$ , it is also true for the matrix elements of each of its components,  $X$ ,  $Y$ , and  $Z$ . The remaining dipole selection rules are

$$\langle \beta, l', m' | Z | \alpha, l, m \rangle \neq 0 \quad \text{requires } m' = m, \quad (13.2.16)$$

$$\langle \beta, l', m' | (X + iY) | \alpha, l, m \rangle \neq 0 \quad \text{requires } m' = m + 1, \quad (13.2.17)$$

$$\langle \beta, l', m' | (X - iY) | \alpha, l, m \rangle \neq 0 \quad \text{requires } m' = m - 1. \quad (13.2.18)$$

To prove these, we can suppress the labels  $\alpha, \beta$ , which play no role. Taking the matrix element of the commutation relation  $[L_z, Z] = 0$  gives  $\langle l', m' | [L_z, Z] | l, m \rangle = (\hbar m' - \hbar m) \langle l', m' | Z | l, m \rangle = 0$ . This shows that if  $\langle l', m' | Z | l, m \rangle$  is to be non-zero, we need  $m' = m$ . A simple alternative proof uses the  $\phi$ -dependence of the position representations of the spherical harmonics and  $X$ ,  $Y$ , and  $Z$ . We have  $\langle l', m' | Z | l, m \rangle \propto \int_0^{2\pi} d\phi (e^{-im'\phi})(1)(e^{im\phi}) \propto \delta_{mm'}$ , and similarly,  $\langle l', m' | (X \pm iY) | l, m \rangle \propto \int_0^{2\pi} d\phi (e^{-im'\phi})(e^{\pm i\phi})(e^{im\phi}) \propto \delta_{m', m \pm 1}$ . Equations (13.2.17) and (13.2.18) also imply that the corresponding matrix elements of  $X$  and  $Y$  vanish unless  $m' = m \pm 1$ .

As a last application of eq. (13.2.9), consider the selection rules for the matrix elements  $\langle \beta, j, m' | \vec{V} | \alpha, j, m \rangle$  of a general vector operator. The only difference compared to the special case of  $\vec{R}$  that led to eq. (13.2.15) is that now we must admit the possibility that the right side of eq. (13.2.9) does not vanish for  $j = j'$ . Therefore, the **general vector operator selection rules** are

$$\langle \beta, j', m' | \vec{V} | \alpha, j, m \rangle \neq 0 \quad \text{requires } j' - j = 0, \pm 1. \quad (13.2.19)$$

$$\langle \beta, j', m' | V_z | \alpha, j, m \rangle \neq 0 \quad \text{requires } m' = m, \quad (13.2.20)$$

$$\langle \beta, j', m' | (V_x + iV_y) | \alpha, j, m \rangle \neq 0 \quad \text{requires } m' = m + 1, \quad (13.2.21)$$

$$\langle \beta, j', m' | (V_x - iV_y) | \alpha, j, m \rangle \neq 0 \quad \text{requires } m' = m - 1. \quad (13.2.22)$$

The last three can be obtained by using  $[J_z, V_z] = 0$  and  $[J_+, V_+] = 0$  and  $[J_-, V_-] = 0$ .

### 13.3 The Wigner–Eckart Theorem and selection rules for tensor operators

In this section we will state and prove the **Wigner–Eckart Theorem** for matrix elements of irreducible tensor operators. Developed independently by Wigner and Carl Eckart, this is a generalization of three simpler results that we have already given: eq. (12.5.18) for spherical

harmonics, Theorem 13.2.1 for scalar operators, and Theorem 13.2.2 for vector operators. The reader may wish to contemplate the parallels between the following derivations and those used in section 12.5 for spherical harmonics.

Consider an irreducible tensor operator  $T_{m_1}^{(j_1)}$  with  $m_1 = -j_1, \dots, j_1$ , and a multiplet of angular momentum eigenstates  $|\alpha j_2 m_2\rangle$  with  $m_2 = -j_2, \dots, j_2$ , where  $\alpha$  is a degeneracy label. Now we construct the states

$$|\gamma j m\rangle = \sum_{m_1, m_2} T_{m_1}^{(j_1)} |\alpha j_2 m_2\rangle C_{m_1 m_2 m}^{j_1 j_2 j}, \quad (13.3.1)$$

for  $j = |j_1 - j_2|, \dots, j_1 + j_2$  and  $m = -j, \dots, j$ , where  $C_{m_1 m_2 m}^{j_1 j_2 j}$  are Clebsch–Gordan coefficients as defined by eq. (12.4.4). We claim that, as the labeling suggests, eq. (13.3.1) is an eigenstate of  $J^2$  and  $J_z$  with eigenvalues  $\hbar^2 j(j+1)$  and  $\hbar m$ , respectively. (It is not necessarily normalized to unity.) To prove this, it suffices to show that

$$J_z |\gamma j m\rangle = \hbar m |\gamma j m\rangle, \quad (13.3.2)$$

$$J_- |\gamma j m\rangle = \hbar \sqrt{j(j+1) - m(m-1)} |\gamma, j, m-1\rangle, \quad (13.3.3)$$

$$J_+ |\gamma j m\rangle = \hbar \sqrt{j(j+1) - m(m+1)} |\gamma, j, m+1\rangle, \quad (13.3.4)$$

as then the result for  $J^2 = (J_+ J_- + J_- J_+)/2 + J_z^2$  follows.

To prove eq. (13.3.2), we use eq. (13.1.6) to obtain

$$J_z |\gamma j m\rangle = \sum_{m_1, m_2} T_{m_1}^{(j_1)} (\hbar m_1 + J_z) |\alpha j_2 m_2\rangle C_{m_1 m_2 m}^{j_1 j_2 j}. \quad (13.3.5)$$

Then, use  $J_z |\alpha j_2 m_2\rangle = \hbar m_2 |\alpha j_2 m_2\rangle$ , and eq. (13.3.2) follows by noting that the Clebsch–Gordan coefficient enforces that only  $m_1 + m_2 = m$  contributes in the sum.

To prove eq. (13.3.3), we similarly use eq. (13.1.8) and then eq. (13.1.5) to find

$$\begin{aligned} J_- |\gamma j m\rangle = \hbar \sum_{m_1, m_2} C_{m_1 m_2 m}^{j_1 j_2 j} & \left( \sqrt{j_1(j_1+1) - m_1(m_1-1)} T_{m_1-1}^{(j_1)} |\alpha j_2 m_2\rangle \right. \\ & \left. + \sqrt{j_2(j_2+1) - m_2(m_2-1)} T_{m_1}^{(j_1)} |\alpha, j_2, m_2-1\rangle \right). \end{aligned} \quad (13.3.6)$$

Now, since we are summing over all  $m_1$  and  $m_2$ , we can use the trick of relabeling the indices according to  $m_1 \rightarrow m_1 + 1$  in the first term and  $m_2 \rightarrow m_2 + 1$  in the second term. The result is

$$J_- |\gamma j m\rangle = \hbar \sum_{m_1, m_2} T_{m_1}^{(j_1)} |\alpha j_2 m_2\rangle \left( \sqrt{j_1(j_1+1) - m_1(m_1+1)} C_{m_1+1, m_2, m}^{j_1 j_2 j} \right. \quad (13.3.7)$$

$$\left. + \sqrt{j_2(j_2+1) - m_2(m_2+1)} C_{m_1, m_2+1, m}^{j_1 j_2 j} \right). \quad (13.3.8)$$

Equation (13.3.3) then follows immediately from using the Clebsch–Gordan recurrence relation eq. (12.4.17). The proof of eq. (13.3.4) is completely analogous.



Having established that the states  $|\gamma jm\rangle$  defined by eq. (13.3.1) are really eigenstates of  $J^2$  and  $J_z$ , we can now invert the relationship by solving for each  $T_{m_1}^{(j_1)} |\alpha j_2 m_2\rangle$  in terms of  $|\gamma jm\rangle$ . Starting from eq. (13.3.1), multiply by  $C_{m'_1 m'_2 m}^{j_1 j_2 j}$ , sum over all  $j$  and  $m$ , and then use the Clebsch–Gordan orthonormality relation eq. (12.4.20) to collapse the  $m_1, m_2$  sums to a single term, and finally relabel  $m'_1 \rightarrow m_1$  and  $m'_2 \rightarrow m_2$ . The result is

$$T_{m_1}^{(j_1)} |\alpha j_2 m_2\rangle = \sum_{j,m} |\gamma jm\rangle C_{m_1 m_2 m}^{j_1 j_2 j}. \quad (13.3.9)$$

This is the precise version of the intuitive statement that  $T_{m_1}^{(j_1)}$  carries angular momentum quantum numbers  $j_1$  and  $m_1$ , and that it imparts them to the states it acts on, yielding a sum of states that occur in the addition of angular momenta  $j_1$  and  $j_2$ .

Now acting on eq. (13.3.9) with  $\langle \beta j' m' |$ , and then relabeling  $j' \rightarrow j$  and  $m' \rightarrow m$ , we obtain

$$\langle \beta jm | T_{m_1}^{(j_1)} |\alpha j_2 m_2\rangle = \langle \beta jm | \gamma jm\rangle C_{m_1 m_2 m}^{j_1 j_2 j}. \quad (13.3.10)$$

By its construction, the state  $|\gamma jm\rangle$  depends on  $\alpha$  and the choice of tensor operator  $T^{(j_1)}$  and  $j$ . However, the matrix element  $\langle \beta jm | \gamma jm\rangle$  clearly does not depend on  $m_1$  or  $m_2$ , and applying Theorem 13.2.1 with  $S$  taken to be the identity operator, we learn that it does not actually depend on  $m$  either. Therefore, we have proved a powerfully general result:

**Theorem 13.3.1. (Wigner–Eckart)** *If  $T^{(j_1)}$  is an irreducible tensor operator, and  $|\alpha j_2 m_2\rangle$  and  $|\beta jm\rangle$  are eigenstates of  $J^2$  and  $J_z$  with eigenvalues as labeled, then†*

$$\langle \beta jm | T_{m_1}^{(j_1)} |\alpha j_2 m_2\rangle = \langle \beta j || T^{(j_1)} || \alpha j_2 \rangle C_{m_1 m_2 m}^{j_1 j_2 j}, \quad (13.3.11)$$

where the reduced matrix element  $\langle \beta j || T^{(j_1)} || \alpha j_2 \rangle$  does not depend on  $m$ ,  $m_1$ , or  $m_2$ .

For  $j_1 = m_1 = 0$ , so that  $T_0^{(0)} = S$  is a scalar operator, we can use the fact that  $C_{0mm'}^{0jj'} = \delta_{jj'} \delta_{mm'}$  to recover Theorem 13.2.1 as a special case of the Wigner–Eckart Theorem. This is hardly surprising, since we just used the former in the proof of the latter. The Landé projection formula, eq. (13.2.10), can be shown to be equivalent to the special case obtained for vector operators, when  $j_1 = 1$ . The triple spherical harmonic integral relation eq. (12.5.18) is equivalent to the special case in which the generator of rotations  $\vec{J}$  is taken to be the orbital angular momentum  $\vec{L}$ , and the tensor operator is taken to be defined by the position representation  $T_{m_1}^{(l_1)} = Y_{l_1}^{m_1}(\theta, \phi)$ .

One way to use the Wigner–Eckart Theorem is as a labor-saving device. If we know the matrix element on the left side of eq. (13.3.11) for just one instance  $(m_1, m_2, m)$  for which it is

---

†Some sources define the reduced matrix element with other normalization conventions, most commonly by including a factor of  $1/\sqrt{2j+1}$  (and sometimes a different sign) on the right side of eq. (13.3.11).

non-zero, then we can infer the reduced matrix element. Once that is done, all other cases of  $(m_1, m_2, m)$  with the same  $\alpha, \beta, j_1, j_2$ , and  $j$  will also be known.

The Wigner–Eckart Theorem also provides selection rules, since it implies that the matrix element can only be non-zero if the Clebsch–Gordan coefficient  $C_{m_1 m_2 m}^{j_1 j_2 j}$  is non-zero. This means that the same selection rules apply, so that  $\langle \beta j m | T_{m_1}^{(j_1)} | \alpha j_2 m_2 \rangle$  can be non-zero only if

$$j = |j_1 - j_2|, \dots, j_1 + j_2 - 1, j_1 + j_2, \quad (13.3.12)$$

$$m = m_1 + m_2. \quad (13.3.13)$$

These **tensor operator selection rules** generalize the results in eqs. (13.2.19)–(13.2.22) that we obtained for the special case of vector operators.

## 13.4 Exercises

**Exercise 13.1.** Prove the angular momentum commutation relations for tensor operators, eqs. (13.1.6)–(13.1.8), from earlier results.

**Exercise 13.2.** Suppose that  $\vec{V}$  and  $\vec{W}$  are rectangular-coordinate vector operators, obeying eq. (13.1.12). Show that their cross-product  $\vec{V} \times \vec{W}$  is also a vector operator by the same definition, and that their dot product  $\vec{V} \cdot \vec{W}$  is a scalar operator (commutes with  $\vec{J}$ ).

**Exercise 13.3.** Prove Theorem 13.1.1, which allows tensor operators to be constructed from products of other tensor operators.

**Exercise 13.4.** Consider angular momentum eigenstates  $|\alpha, j, m_j\rangle$  with  $J^2$  and  $J_z$  eigenvalues  $\hbar^2 j(j+1)$  and  $\hbar m_j$ , respectively. For what  $j, m_j$  is the matrix element  $\langle \alpha, j, m_j | (X^2 - Y^2) | \beta, 0, 0 \rangle$  possibly non-zero? What about  $\langle \alpha, j, m_j | XY | \beta, 0, 0 \rangle$  and  $\langle \alpha, j, m_j | XZ | \beta, 0, 0 \rangle$ ?

**Exercise 13.5.** Show that for any angular momentum eigenstate  $|\alpha, j, m_j\rangle$ , the expectation values of the components of an irreducible tensor operator  $T_m^{(l)}$  vanish unless  $l \leq 2j$  and  $m = 0$ . This applies in particular to the multipole moment operator defined in eq. (13.1.35). Thus the only non-zero quadrupole moment [as defined in eq. (13.1.39)] for an angular momentum eigenstate is  $q_{20}$ , and only for states with  $j \geq 1$ . A spin-0 or spin-1/2 particle cannot have a quadrupole moment, even if it is composite.

**Exercise 13.6.** The deuteron is a composite particle with charge  $Q = 1$  and spin one, which means that  $q = e$  and  $j = 1$  in its rest frame. Apply the Wigner-Eckart theorem to find the ratios of the electric quadrupole moment operator expectation values  $q_{20}$ , defined in eq. (13.1.39), for the three states with  $m_j = 1, 0, -1$ .

**Exercise 13.7.** Consider a basis of angular momentum eigenstates  $|\alpha, j, m\rangle$  for a particle of unit charge, for fixed  $\alpha$  and  $j$ , and  $m = -j, \dots, j$ . Suppose that the electric dipole and quadrupole moments defined by the expectation values in the highest- $m$  state,

$$p = \langle \alpha, j, j | Z | \alpha, j, j \rangle, \quad (13.4.1)$$

$$q = \langle \alpha, j, j | (2Z^2 - X^2 - Y^2) | \alpha, j, j \rangle / \sqrt{6}, \quad (13.4.2)$$

are known. Use the Wigner–Eckart Theorem to find, in terms of  $p$  and  $q$  and appropriate Clebsch–Gordan coefficients,

- (a) all of the matrix elements  $\langle \alpha, j, m' | A | \alpha, j, m \rangle$  for each of  $A = X$  and  $Y$  and  $Z$ .
- (b) all of the matrix elements  $\langle \alpha, j, m' | A | \alpha, j, m \rangle$  for each of the operators  $A = XY$  and  $XZ$  and  $YZ$  and  $X^2 - Y^2$  and  $(2Z^2 - X^2 - Y^2)/\sqrt{6}$ . For which  $m, m'$  can the results be non-zero?

## 14 Entanglement, the EPR problem, hidden variables, and Bell inequalities

### 14.1 The Einstein–Podolsky–Rosen problem in Bohm’s formulation

Although Einstein played an important role in the early development of quantum mechanics, like many others he was troubled by some of its implications. This was in part due to his discomfort with the reliance of Postulates 4 and 5 on probabilities for the result of a measurement and the subsequent state, rather than completely deterministic time evolution. Even after he grudgingly accepted the possibility of non-determinism, Einstein found the implications of entanglement of quantum states to be difficult to accept, and doubted whether the standard formulation of quantum mechanics should be regarded as a complete framework.

In 1935, Einstein, Boris Podolsky, and Nathan Rosen [*Phys. Rev.* **47**, 777, (1935)] described a thought experiment that highlighted the perceived weirdness associated with measurements involving entangled states. This problem is often called the EPR paradox, but it should be emphasized at the outset that although it may be troubling to humans accustomed to classical ways of thinking, in the end we will find that there is really nothing paradoxical about it. (Indeed, EPR themselves did not call it a “paradox”.) EPR described the problem in terms of measurements of positions and momenta, but David Bohm later distilled the basic ideas into a simpler form in terms of measurements of components of two entangled spins, and that is the version we will now describe.

Consider a system of two spin-1/2 particles, labeled 1 and 2, in a state with the total spin  $\vec{S} = \vec{S}_1 + \vec{S}_2$  equal to 0,

$$|S = 0\rangle = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) = \frac{1}{\sqrt{2}}(|+\hat{z}, -\hat{z}\rangle - |-\hat{z}, +\hat{z}\rangle). \quad (14.1.1)$$

The notation is such that, for example,  $|\uparrow\downarrow\rangle = |+\hat{z}, -\hat{z}\rangle \equiv |+\hat{z}\rangle_1 \otimes |-\hat{z}\rangle_2$  is an eigenstate of both  $S_{1z}$  and  $S_{2z}$ , with eigenvalues  $+\hbar/2$  and  $-\hbar/2$ , respectively. The state of the system is thus represented by a ket in the tensor product of the Hilbert spaces for the two individual particles. In the following, we will purposely avoid the  $\uparrow$  and  $\downarrow$  notations for the spins, because we will want to consider components of the spins other than the  $z$  component. In fact, since the  $S = 0$  state is spherically symmetric, there is nothing special about the  $\hat{z}$  direction, and up to an irrelevant phase we could just as easily write the state in eq. (14.1.1) as

$$|S = 0\rangle = \frac{1}{\sqrt{2}}(|+\hat{n}, -\hat{n}\rangle - |-\hat{n}, +\hat{n}\rangle) \quad (14.1.2)$$

for any convenient unit vector direction  $\hat{n}$ . Importantly, there is no way to write this total spin-0 ket as a single product of kets in the individual spin-1/2 particle Hilbert spaces; no matter the




Figure 14.1.1: The setup for Bohm's version of the EPR problem. Decays of neutral pions at rest produce two spin-1/2 particles moving in opposite directions, in an entangled spin state with total spin 0, and with spatial wavefunctions that have no overlap at late times. Two distant observers Alice and Bob can each measure any component they choose of the spin of the particle that arrives at their respective detector A or B.

choice of bases for the individual spins, it is always a non-trivial linear combination of such products. Thus, it is an example of an entangled state as defined in section 2.9.

The particles also have spatial wavefunctions that we need not write explicitly. The only thing important for us is that they describe localized wavepackets moving away from each other. The two spins, although entangled, are therefore separated by a large distance at late times. This is sometimes called **nonlocal entanglement**.

One way to realize such a state would be an electron-positron pair from the decay of a neutral pion at rest in a suitable reference frame,<sup>†</sup>

$$\pi^0 \rightarrow e^- e^+. \quad (14.1.3)$$

The pion has spin 0, and since we are in its rest frame, the total angular momentum is also 0. We imagine that this experiment will be conducted many times, with two ideal observers Alice and Bob located on opposite sides of the point where the pions decay, and far away, as shown in Figure 14.1.1. They each have a small detector capable of measuring any desired component of the spin of a particle moving through it. Since the detectors are far apart, where the spatial wavefunctions for the two particles have no overlap, Alice can only measure one of the spins, and Bob can only measure the other. Let us choose the labeling of the particles so that Alice always makes measurements on particle 1 (which on an event-by-event basis might be either the electron or the positron), and Bob always on particle 2. They can each measure any component of that spin, or choose to make no measurement. For simplicity, we assume that the Hamiltonian is just that of free particles, so that the spin states have no Hamiltonian time evolution. Throughout this chapter, we will use spin operators with a factor of  $\hbar/2$  extracted, so

$$\vec{\sigma}_k = \frac{2}{\hbar} \vec{S}_k \quad (14.1.4)$$

for particle  $k$ , represented by the Pauli matrices.

---

<sup>†</sup>This is a very rare decay, occurring for less than  $10^{-7}$  of all neutral pions. But this is a thought experiment; we only care that it is possible in principle, not about such practical difficulties.

Now suppose that Alice measures All three components of  $\vec{\sigma}_1$  commute with all three components of  $\vec{\sigma}_2$ , because they operate on distinct one-particle Hilbert spaces.  $\sigma_{1z}$ . The result will be either +1 or -1, with equal probability. If the result was +1, then if Bob measured  $\sigma_{2z}$  the result will necessarily be -1. This follows from the form of the state eq. (14.1.1). We can say that Alice has collapsed the state to  $|+\hat{z}, -\hat{z}\rangle$  by her measurement of  $\sigma_{1z}$ , so that the only possible outcome for Bob is  $\sigma_{2z} = -1$ . Conversely, if Alice instead measured  $\sigma_{1z}$  to be -1, then if Bob measures  $\sigma_{2z}$  the result must be +1. There is a perfect anticorrelation between their measurements.

This perfect anticorrelation in Alice's and Bob's measured results for  $\sigma_{1z}$  and  $\sigma_{2z}$  is not at all surprising. In fact, it is actually much more general, in the sense that it does not even rely on quantum mechanics being correct. It would inevitably be true in any alternative theory, just as long as angular momentum is conserved. There is a similar perfect anticorrelation between the measured electric charges; if Alice sees that particle 1 is an electron, then she can immediately be sure that Bob's particle 2 will be a positron, due to conservation of charge. As an analogy, suppose that Charlie at the central point had taken a pair of shoes and shipped one to Alice and one to Bob, concealed in boxes. Before opening her box, Alice would know nothing about the shoe Bob received, but if she opens her box to find a left shoe, she would immediately gain the information that Bob's box has a right shoe. This has nothing to do with quantum mechanics, of course.

However, spins are much more interesting than electric charges or shoes, and can reveal nontrivial aspects of measurement correlations in quantum mechanics. This is because spins can point in arbitrary directions, and Alice and Bob could choose to make measurements of different spin components. For example, suppose Alice measures  $\sigma_{1x}$  but Bob still measures  $\sigma_{2z}$ . If Alice obtains the result  $\sigma_{1x} = +1$ , then from eq. (14.1.2) with  $\hat{n} = \hat{x}$ , we can say that the total state collapses to  $|+\hat{x}, -\hat{x}\rangle$ . Then, since  $|-\hat{x}\rangle_2 = (|+\hat{z}\rangle_2 - |-\hat{z}\rangle_2)/\sqrt{2}$ , Bob will obtain the results  $\sigma_{2z} = \pm 1$  randomly, and with equal probability. The same random results will occur for Bob's measurement of  $\sigma_{2z}$  if the result of Alice's measurement was instead  $\sigma_{1x} = -1$ , or if she decided to make no measurement at all.

To be more general, suppose that Alice and Bob measure arbitrary spin components  $\hat{a} \cdot \vec{\sigma}_1$  and  $\hat{b} \cdot \vec{\sigma}_2$ , respectively, for unit vectors  $\hat{a}$  and  $\hat{b}$ . Neither can predict with certainty what the other will measure on an event-by-event basis (unless  $\hat{a}$  and  $\hat{b}$  are parallel), but there is a statistical correlation between their measurements, which can be expressed as the expectation value

$$C(\hat{a}, \hat{b}) = \langle \hat{a} \cdot \vec{\sigma}_1 \hat{b} \cdot \vec{\sigma}_2 \rangle \quad (14.1.5)$$

in the state  $|S = 0\rangle$ . A labor-saving trick for evaluating this is to note that when acting on the state  $|S = 0\rangle$ , one can always replace  $\hat{b} \cdot \vec{\sigma}_2$  with  $-\hat{b} \cdot \vec{\sigma}_1$ , followed by use of the Pauli matrix

identity  $(\hat{a} \cdot \vec{\sigma})(\hat{b} \cdot \vec{\sigma}) = \hat{a} \cdot \hat{b} + i(\hat{a} \times \hat{b}) \cdot \vec{\sigma}$  from eq. (8.2.17). Then, since the expectation value of  $\vec{\sigma}_1$  in the  $S = 0$  state vanishes, the prediction of quantum mechanics is

$$C(\hat{a}, \hat{b}) = -\hat{a} \cdot \hat{b}. \quad (14.1.6)$$

This agrees with the special cases already discussed; for  $\hat{a} = \hat{b}$ , the measurements are perfectly anticorrelated, and when  $\hat{a}$  and  $\hat{b}$  are orthogonal there is no correlation. More generally,  $C(\hat{a}, \hat{b})$  is always in the range from  $-1$  (perfect anticorrelation) to  $+1$  (perfect correlation).

The key issue that bothered EPR is that the possible results obtained by Bob would seem to depend on the choices made by Alice, and vice versa, even if they are so far apart from each other that there is no possibility of communication between them when they each choose which spin measurement to make. They could even make their individual decisions about which spin components to measure at random for each individual pion decay after it has occurred and while the electron and positron are already in flight. At the last moment before particle 1 arrives, Alice could decide to measure  $\sigma_{1z}$ . Then, from the results of Alice's measurement alone, she can be immediately certain of the result if Bob measures  $\sigma_{2z}$ , a measurement occurring far away and quite out of her control. Or, she could decide instead to measure  $\sigma_{1x}$ , in which case (regardless of the outcome of her own measurement) she would correctly assign a probability of 50% to the outcome of a measurement by Bob of  $\sigma_{2z}$ . The same follows if she decided to make no measurement at all. This is all despite the fact that, because of the finite speed of light, no signal carrying the news of her last-moment decision could reach Bob before his measurement occurs. Einstein referred to this sort of thing as “spooky action at a distance”, and argued that it pointed to an incompleteness of the standard quantum theory as a description of reality. However, Nature does not care whether we, or even Einstein, find a phenomenon peculiar or spooky. There are two things to check to make sure that there is not a problem with the quantum theory, let alone a true paradox.

The first thing to check is that the predictions of quantum mechanics are really consistent and unambiguous. In the preceding, we have phrased things in terms of Alice's measurement collapsing the state, with implications for what Bob will see. But the situation is symmetrical; we could just as easily describe any pair of measurements in terms of Bob collapsing the state, with implications for what Alice will see. For the case in which Alice measured  $\sigma_{1z} = +1$  and Bob measured  $\sigma_{2z} = -1$ , we could choose to say that Bob has collapsed the total state to  $|+\hat{z}, -\hat{z}\rangle$ , which means that the only possible result for Alice's measurement was, indeed,  $\sigma_{1z} = +1$ . This outcome is fully consistent with the interpretation in which Alice's measurement collapses the state first. A third way is to view the collapse symmetrically, by taking Alice's and Bob's measurements to be a single measurement. (This is possible because all components of  $\vec{\sigma}_1$  and  $\vec{\sigma}_2$  commute with each other.) Again, the state collapses to  $|+\hat{z}, -\hat{z}\rangle$ . For any given outcome of




Figure 14.1.2: If instantaneous communication were possible, then you could send a signal to a relay  $R$  which would be instantaneous (at constant time) in some other reference frame, but would arrive at an earlier time in your own reference frame. The relay could then send a signal, instantaneous in a different reference frame, which could be used to somehow prevent your parents from meeting, preventing your own birth. Fortunately, instantaneous communication is not a consequence of quantum mechanics.

both measurements, quantum mechanics indeed always gives the same result for the collapse of the state regardless of what order we assign to the measurements.<sup>‡</sup> To see this more generally, suppose that Alice measures  $\hat{a} \cdot \vec{\sigma}_1$  to be  $+1$  and Bob measures  $\hat{b} \cdot \vec{\sigma}_2$  to be  $+1$ . As you will verify when you do Exercise 14.1, these three ways of describing the state collapse give the same probabilities for this outcome, and the same final state.

The second thing to check is that quantum mechanics does not provide any instantaneous communication between the distant Alice and Bob, whose measurements are events occurring at a space-like separation in the language of special relativity. In general, an instantaneous (or even faster-than-light) communication between space-like separated points in one inertial reference frame would appear to travel backwards in time in some other inertial frame that is moving with respect to the first. This would imply causality paradoxes, because signals by combinations of actors in different inertial frames can be constructed in such a way that a message could be received on a time-like path before it is sent from a previous point on the same path. For example, you could use instantaneous communications to convince your parents not to meet, thus paradoxically preventing your own birth, as illustrated in Figure 14.1.2.

To see that Alice and Bob's experiments do not actually allow instantaneous communication, we need only note that no matter what each of them chooses to measure, the results of their individual measurements will always appear to be completely random, until they meet to compare them or at least send a conventional signal at a speed not exceeding that of light. Even if Alice and Bob agree to seek maximum anticorrelation by measuring  $\sigma_{1z}$  and  $\sigma_{2z}$  respectively (and even assuming that they do not break their promises!), each will record their data locally as a random sequence of  $+1$  and  $-1$ , containing no information. *After* communicating by conventional means, they can check that their results do have correlations in agreement with quantum mechanics, but the key point is that no information is transmitted in either direction

<sup>‡</sup>This is necessary, because if the measurements of Alice and Bob are space-like separated events, special relativity says that there is no invariant sense of which occurred first. There is an inertial frame in which they are simultaneous, but in some other inertial frames, Alice's measurement was first, and in others Bob's was first.



until that happens.

Why, then, did EPR view the situation as disturbing? The essential issue is that one must not fall into the trap of thinking of the state of a system as always defined only by local information, a viewpoint sometimes called **local realism**. If you try to think about particle 2 only in terms of non-entangled kets such as  $|+\hat{z}\rangle_2$  or  $|-\hat{x}\rangle_2$ , then it might indeed appear paradoxical that Alice's distant measurement of  $S_{1z}$  can affect what Bob measures. However, that way of thinking about the state of particle 2 is explicitly rejected by quantum mechanics, which insists that we take into account the full nonlocal entanglement in order to get correct and consistent predictions. The quantum state describes both particles at the same time, not one particle or the other. The real value of the EPR problem is that it forces us to recognize this fundamental truth about how quantum mechanics works.

One sometimes sees grand pronouncements, inspired by the EPR problem, along the lines of “quantum mechanics is nonlocal”. However, one must be careful about what this implies, because the word “nonlocal” has several completely different meanings in physics. For example, it is possible to define a Hamiltonian that has nonlocal dynamics; that is what quantum field theorists usually mean when they talk about the possibility of nonlocality. That is certainly not the case in the preceding discussion, as we took  $H = 0$  for the spin degrees of freedom, and even the  $H$  describing the time evolution of the spatial wavefunctions was (implicitly) just that of free particle propagation. The entanglement of the state occurred due to perfectly local processes. The rejection of local realism in favor of nonlocal entanglement of states does not imply that we need to accept a nonlocal Hamiltonian, for which there is absolutely no experimental evidence.

## 14.2 Hidden variables and Bell's inequality

Having learned in the previous section that standard quantum mechanics rejects local realism in favor of allowing nonlocal entanglement, it must be recognized that this is a falsifiable hypothesis to be experimentally tested against alternatives. There is a general class of alternatives called **hidden variables** theories, which attempt to incorporate local realism in a way consistent with experiment. In the remaining sections of this chapter, we will consider local hidden variables theories, see how they make predictions that are incompatible with quantum mechanics, and understand why experiments show that they cannot be correct.

Hidden variables are supposed to be quantities that we cannot directly measure or control, but are necessary to completely characterize the state of a system. They could be part of some completely deterministic proposed alternative to quantum mechanics, or could be an extra part of a framework that otherwise looks like conventional quantum mechanics. In the latter version, it is not just a ket  $|\psi\rangle$  that describes the state of the system, but rather the state ket

together with the hidden variable(s), which we will generically call  $\lambda$ . They are supposed to be distributed in a way that is either truly random, or just seems to be random because of our lack of knowledge of them. For any given system, the distribution of the hidden variables can be described by a probability density function  $\mu(\lambda)$  with the properties

$$\int d\lambda \mu(\lambda) = 1, \quad (14.2.1)$$

$$\mu(\lambda) \geq 0, \quad (14.2.2)$$

so that  $\mu(\lambda_0) d\lambda$  is the probability that  $\lambda$  will be found between  $\lambda_0$  and  $\lambda_0 + d\lambda$ . The function  $\mu(\lambda)$  might depend on the particulars of the system and its history.

The presence of  $\lambda$  is supposed to be responsible for the illusion of random outcomes of measurements. For some observable  $A$ , and a state fully characterized by non-hidden properties  $\psi$  and hidden variables  $\lambda$ , the hypothesis is that the outcome of a measurement is not random, but rather determined as some function

$$\text{measured value of } A = f_{A,\psi}(\lambda). \quad (14.2.3)$$

The function  $f_{A,\psi}(\lambda)$  can be chosen so that as  $\lambda$  runs over all possible values, it returns all allowed values predicted by quantum mechanics for the measurement of  $A$  in the state  $\psi$ . Because we have no way of knowing what  $\lambda$  is, the results of individual measurements appear random. So, the mean value of many measurements of  $A$  for apparently identical states  $\psi$  will be

$$\overline{A} = \int d\lambda \mu(\lambda) f_{A,\psi}(\lambda). \quad (14.2.4)$$

This is the counterpart of the expectation value  $\langle A \rangle = \langle \psi | A | \psi \rangle$  in standard quantum theory.

For example, let us consider how this would work for a system with two spin-1/2's combined to have total spin  $S = 0$  in an entangled state coming from  $\pi^0$  decay, as considered in section 14.1. To be general, consider the measurements by Alice and Bob of arbitrary components of the spins of particle 1 and 2, say  $\hat{a} \cdot \vec{\sigma}_1$  and  $\hat{b} \cdot \vec{\sigma}_2$ , where  $\hat{a}$  and  $\hat{b}$  are unit vectors. Then the outcomes for Alice's and Bob's measurements are respectively determined by some functions

$$\hat{a} \cdot \vec{\sigma}_1 = f_A(\hat{a}, \lambda) = \pm 1, \quad (14.2.5)$$

$$\hat{b} \cdot \vec{\sigma}_2 = f_B(\hat{b}, \lambda) = \pm 1. \quad (14.2.6)$$

To incorporate local realism, the value of  $\lambda$  is assumed to be the same for both particles 1 and 2, and is fixed at the moment that the parent pion decays. Thus, in each event,  $\lambda$  has an unknown, but specific, value, which already determines what the ensuing measurement of any component of the spins will be, through the functions  $f_A$  and  $f_B$ .

In the case of spins, the functions determining the measurement outcomes must satisfy some constraints. First, a measured value for  $\hat{n} \cdot \vec{\sigma}_k$  is the same as the negative of a measured value for  $-\hat{n} \cdot \vec{\sigma}_k$ , so  $f_A(\hat{n}, \lambda) = -f_A(-\hat{n}, \lambda)$  and  $f_B(\hat{n}, \lambda) = -f_B(-\hat{n}, \lambda)$ , for any  $\hat{n}$ . Also, to be consistent with angular momentum conservation, it must be true that whenever  $\hat{a} = \hat{b} = \hat{n}$ , the measurement results are perfectly anticorrelated in the case of the EPR setup. Thus,

$$f_B(\hat{n}, \lambda) = -f_A(\hat{n}, \lambda) \quad (14.2.7)$$

so that the measurements are consistent with total spin  $S = 0$ .

We can now write down the hidden-variables prediction for the statistical correlation of Alice's and Bob's measurements. It is given by the product of the possible measurement results, integrated over the hidden variables weighted by their probability density,

$$C(\hat{a}, \hat{b}) = \int d\lambda \mu(\lambda) f_A(\hat{a}, \lambda) f_B(\hat{b}, \lambda). \quad (14.2.8)$$

This hidden variables result should be contrasted with the prediction of quantum mechanics that we found in eq. (14.1.6), which was  $C(\hat{a}, \hat{b})_{\text{QM}} = -\hat{a} \cdot \hat{b}$ .

The hidden variables result for  $C(\hat{a}, \hat{b})$  is clearly less specific than the quantum mechanics prediction, because we have not committed to a particular form for the functions  $\mu$ ,  $f_A$ , and  $f_B$ . Nevertheless, it is possible to draw some specific conclusions. In the case that  $\hat{b} = \hat{a}$ , the hidden variables prediction is

$$\begin{aligned} C(\hat{a}, \hat{a}) &= \int d\lambda \mu(\lambda) f_A(\hat{a}, \lambda) f_B(\hat{a}, \lambda) = - \int d\lambda \mu(\lambda) f_A(\hat{a}, \lambda)^2 = - \int d\lambda \mu(\lambda) \\ &= -1, \end{aligned} \quad (14.2.9)$$

where we used eq. (14.2.7) to get the second equality, then the fact that  $f_A(\hat{a}, \lambda) = \pm 1$ , and finally eq. (14.2.1). In this special case, the hidden variables prediction does agree with that of quantum mechanics. This had to be true, simply because in our hidden variables theory we built in the perfect anticorrelation required by angular momentum conservation, through eq. (14.2.7).

However, for more general  $\hat{a}$  and  $\hat{b}$ , it is not immediately clear whether the hidden variables predictions for  $C(\hat{a}, \hat{b})$  can be made to always agree with the predictions of quantum mechanics. In 1964, John S. Bell<sup>†</sup> answered the question by discovering that the correlations predicted by hidden variables theories must satisfy an inequality that is clearly violated by the correlations predicted by quantum mechanics. Surprisingly, this is true for *any* choices of the functions  $\mu(\lambda)$ ,  $f_A(\hat{a}, \lambda)$ , and  $f_A(\hat{b}, \lambda)$  in the hidden variables theory. This provides a way for experiment to decisively settle the question of whether the general hidden variables idea or standard quantum mechanics is true.

---

<sup>†</sup>*Physics*, **1**, 195, (1964), reprinted in J.S. Bell, *Speakable and Unsayable in Quantum Mechanics*, (1987).

Bell's inequality is remarkably simple to derive. We start with

$$C(\hat{a}, \hat{b}) - C(\hat{a}, \hat{c}) = \int d\lambda \mu(\lambda) \left[ f_A(\hat{a}, \lambda) f_B(\hat{b}, \lambda) - f_A(\hat{a}, \lambda) f_B(\hat{c}, \lambda) \right] \quad (14.2.10)$$

$$= - \int d\lambda \mu(\lambda) \left[ f_A(\hat{a}, \lambda) f_A(\hat{b}, \lambda) - f_A(\hat{a}, \lambda) f_A(\hat{c}, \lambda) \right], \quad (14.2.11)$$

where we have used eq. (14.2.7) to replace  $f_B$  in favor of  $f_A$ . Multiply the second term on the right side by  $1 = [f_A(\hat{b}, \lambda)]^2$ , and rearrange to get

$$C(\hat{a}, \hat{b}) - C(\hat{a}, \hat{c}) = - \int d\lambda \mu(\lambda) \left[ 1 - f_A(\hat{b}, \lambda) f_A(\hat{c}, \lambda) \right] f_A(\hat{a}, \lambda) f_A(\hat{b}, \lambda). \quad (14.2.12)$$

Now, take the absolute value of both sides, and use the general fact that the absolute value of any integral is always less than or equal to the integral of the magnitude of the integrand:

$$\left| \int d\lambda F(\lambda) \right| \leq \int d\lambda |F(\lambda)|. \quad (14.2.13)$$

Let  $F(\lambda)$  be the integrand on the right side of eq. (14.2.12). Then since  $f_A(\hat{a}, \lambda) f_A(\hat{b}, \lambda)$  is always equal to  $\pm 1$ , and both  $\mu(\lambda)$  and  $1 - f_A(\hat{b}, \lambda) f_A(\hat{c}, \lambda)$  are always non-negative, we have  $|F(\lambda)| = \mu(\lambda) \left[ 1 - f_A(\hat{b}, \lambda) f_A(\hat{c}, \lambda) \right]$ . Therefore we obtain

$$|C(\hat{a}, \hat{b}) - C(\hat{a}, \hat{c})| \leq \int d\lambda \mu(\lambda) \left[ 1 - f_A(\hat{b}, \lambda) f_A(\hat{c}, \lambda) \right] \quad (14.2.14)$$

or, using eqs. (14.2.1) and (14.2.7) and (14.2.8) again,

$$|C(\hat{a}, \hat{b}) - C(\hat{a}, \hat{c})| \leq 1 + C(\hat{b}, \hat{c}). \quad (14.2.15)$$

This is Bell's original inequality governing correlations of spin-1/2 measurements in a total spin  $S = 0$  state in hidden variables theories.

To see that Bell's inequality is predicted to be violated by quantum mechanics, consider what happens if we substitute the quantum prediction of the correlation from eq. (14.1.6) into eq. (14.2.15),

$$|\hat{a} \cdot \hat{b} - \hat{a} \cdot \hat{c}| \leq 1 - \hat{b} \cdot \hat{c}. \quad (14.2.16)$$

This reduces the question to a purely geometric claim about three arbitrary unit vectors. A single example will be sufficient. Take  $\hat{a} = \hat{x}$ ,  $\hat{b} = \hat{y}$ , and  $\hat{c} = (\hat{x} + \hat{y})/\sqrt{2}$ . The left side of eq. (14.2.16) is  $1/\sqrt{2}$ , while the right side is  $1 - 1/\sqrt{2}$ , which is less, so the inequality is violated. Thus, quantum mechanics makes a specific prediction that cannot be satisfied in any local hidden variables theory.

### 14.3 Quantum mechanics vs. hidden variables, without inequalities

There are various other Bell-type inequalities, which show that quantum mechanics and hidden variables theories can be distinguished by measuring statistical correlations. In this section we will discuss another way, which is theoretically even sharper because it does not rely on inequalities or statistics, although in practical terms it is somewhat harder to realize experimentally. The result in this example is that quantum mechanics and hidden variables theories make predictions that disagree, not just for correlations on a statistical basis, but for certain measurements on an event-by-event basis with 100% probability.

Consider the following entangled state involving three spin-1/2 particles,<sup>†</sup>

$$|\psi\rangle = \frac{1}{\sqrt{2}} \left( |+\hat{z}, +\hat{z}, +\hat{z}\rangle - |-\hat{z}, -\hat{z}, -\hat{z}\rangle \right). \quad (14.3.1)$$

We assume that the three particles, labeled 1, 2, and 3, start from a central point and are spatially separated without disturbing the spins. The particles are observed by Alice, Bob, and Charlie at three remote sites. It is agreed that each of them will independently measure a random choice of either the  $x$  or  $y$  component of the spin, so that Alice always measures either  $\sigma_{1x}$  or  $\sigma_{1y}$ , while Bob always measures  $\sigma_{2x}$  or  $\sigma_{2y}$ , and Charlie always measures  $\sigma_{3x}$  or  $\sigma_{3y}$ . The experiment is repeated many times, always with the same state  $|\psi\rangle$ , and then the observers meet to compare their results.

Let us first analyze the situation according to the standard rules of quantum mechanics. The state  $|\psi\rangle$  is not an eigenstate of any of the individual spin operators  $\sigma_{1x}$ ,  $\sigma_{1y}$ ,  $\sigma_{2x}$ ,  $\sigma_{2y}$ ,  $\sigma_{3x}$ , or  $\sigma_{3y}$ , and each of the individual measurements of Alice, Bob, and Charlie will have a random result  $\pm 1$  with equal probability. However, you can check that  $|\psi\rangle$  has the remarkable property that it is an eigenstate of all four of the operators of the form

$$\Omega_{abc} = \sigma_{1a}\sigma_{2b}\sigma_{3c}, \quad (14.3.2)$$

where  $(a, b, c) = (x, x, x)$ ,  $(x, y, y)$ ,  $(y, x, y)$ , and  $(y, y, x)$ . For the operator  $\Omega_{xxx}$  the eigenvalue is  $-1$ , and for each of the other three operators  $\Omega_{xyy}$ ,  $\Omega_{yxy}$ , and  $\Omega_{yyx}$ , the eigenvalue is  $+1$ .

Before meeting, Alice, Bob, and Charlie have not learned anything except that their own individual measurement results appear completely random. However, consider what they find when they finally get together to compare their results. They decide to first look only at the results for cases in which two of them had chosen to measure the  $y$  component of the spin, and the other had measured the  $x$  component of the spin. In those cases, we can equivalently say

---

<sup>†</sup>This three-spin state was proposed by N. David Mermin *Am. J. Phys.* **58**, 731, (1990), following a similar four-spin state given by Daniel Greenberger, Michael Horne, and Anton Zeilinger, in *Bell's Theorem, Quantum Theory, and Conceptions of the Universe* (1989). Examples of this type are commonly called **GHZ states**.

that collectively they had measured one of the observables  $\Omega_{xyy}$  or  $\Omega_{yxy}$  or  $\Omega_{yyx}$ , and because

$$\Omega_{xyy} |\psi\rangle = \Omega_{yxy} |\psi\rangle = \Omega_{yyx} |\psi\rangle = |\psi\rangle, \quad (14.3.3)$$

they must find that the product of their measured results was definitely +1, every time. (Sometimes all three of them will have measured the result to be +1, and sometimes two of them will have obtained -1 and the other will have obtained +1.) On the other hand, in the cases where all three had chosen to measure the  $x$  component of the spin, we can equivalently say that their choices amounted to measuring  $\Omega_{xxx}$ . Now, since

$$\Omega_{xxx} |\psi\rangle = -|\psi\rangle, \quad (14.3.4)$$

it must be that the product of their three measurements is -1. (Sometimes they will all get the result -1, and sometimes two will get +1 and the other will get -1.)

So far, we have analyzed the experiment assuming the predictions of quantum mechanics. Now let us try to explain these results in terms of a general local hidden variables theory, in which the measurement of  $\sigma_{1x}$  is supposed to be determined to be some  $f_A(\hat{x}, \lambda) = \pm 1$  and the measurement of  $\sigma_{2y}$  is  $f_B(\hat{y}, \lambda) = \pm 1$ , etc. The hidden variable(s)  $\lambda$  are supposed to be different for each repetition of the experiment, giving the illusion of randomness. Then, the observed results for  $\Omega_{xyy}, \Omega_{yxy}, \Omega_{yyx}$  would imply that, respectively,

$$f_A(\hat{x}, \lambda) f_B(\hat{y}, \lambda) f_C(\hat{y}, \lambda) = 1, \quad (14.3.5)$$

$$f_A(\hat{y}, \lambda) f_B(\hat{x}, \lambda) f_C(\hat{y}, \lambda) = 1, \quad (14.3.6)$$

$$f_A(\hat{y}, \lambda) f_B(\hat{y}, \lambda) f_C(\hat{x}, \lambda) = 1, \quad (14.3.7)$$

for every  $\lambda$ . Now, multiplying these three equations together, and using  $f_A(\hat{y}, \lambda)^2 = f_B(\hat{y}, \lambda)^2 = f_C(\hat{y}, \lambda)^2 = 1$ , we discover that

$$f_A(\hat{x}, \lambda) f_B(\hat{x}, \lambda) f_C(\hat{x}, \lambda) = 1. \quad (14.3.8)$$

This tells us that in those cases where they had all measured the  $x$  component of spin, the product of their measurements should have been +1, with 100% certainty, according to the hidden variables theory. This is in direct contradiction to the quantum mechanics prediction.

To recapitulate: if we stipulate that the measurement of  $\Omega_{xyy}$ ,  $\Omega_{yxy}$ , or  $\Omega_{yyx}$  always gives +1, then, according to any local hidden variables theory, measurement of  $\Omega_{xxx}$  must always give +1 as well. However, the prediction of quantum mechanics, for the given state  $|\psi\rangle$ , is that  $\Omega_{xxx}$  always gives -1. The hidden variables and quantum mechanics hypotheses make contradictory predictions. One of them must be wrong. The experimental verdict will be revealed at the end of the next section.

## 14.4 Aspect's experiments and the demise of local hidden variables

In practice, the most decisive real-world experimental tests, many first conducted by Alain Aspect and collaborators, involve polarizations of photons in entangled states, rather than spin-1/2 systems. In a series of increasingly sensitive experiments, the Bell-type inequalities of local hidden variables theories have been put to the test and found to be clearly violated, and in just the way predicted by standard quantum mechanics. In this section, we will describe a Bell-type inequality for hidden variables, and the corresponding quantum prediction, as tested by A. Aspect, J. Dalibard, G. Roger, *Phys. Rev. Lett.* **49**, 1804 (1982), referred to in the following as the ADR experiment.

Consider observables  $A(\hat{a})$  and  $B(\hat{b})$  that can only take on the values  $\pm 1$ , and depend on unit vectors  $\hat{a}$  and  $\hat{b}$ , and are associated with spatially separated detectors. These generalize the spin-1/2 observables  $\hat{a} \cdot \vec{\sigma}_1$  and  $\hat{b} \cdot \vec{\sigma}_2$  of Alice and Bob in the previous two sections. The experiment can change  $\hat{a}$  and  $\hat{b}$  independently at any time. According to the hidden variables proposal, the measured values are supposed to be determined by some functions

$$A(\hat{a}) = f_A(\hat{a}, \lambda) = \pm 1, \quad (14.4.1)$$

$$B(\hat{b}) = f_B(\hat{b}, \lambda) = \pm 1. \quad (14.4.2)$$

The hidden variables  $\lambda$  are assumed to be local, implying that they are the same for  $A$  and  $B$ , since they will have originated from a single location and event. Their probability density distribution is some non-negative function  $\mu(\lambda)$ , just as in eqs. (14.2.1). The correlation between measurement results  $C(\hat{a}, \hat{b})$  is again given by the formula eq. (14.2.8). However, since the observables are not necessarily components of angular momenta (and will not be in the ADR experiment), we will not assume or use eq. (14.2.7). This means that the Bell inequality we are about to derive applies more generally than the original one.

Consider the combination of correlations

$$\begin{aligned} C(\hat{a}, \hat{b}) - C(\hat{a}, \hat{b}') + C(\hat{a}', \hat{b}) + C(\hat{a}', \hat{b}') &= \int d\lambda \mu(\lambda) \left[ f_A(\hat{a}, \lambda) f_B(\hat{b}, \lambda) - f_A(\hat{a}, \lambda) f_B(\hat{b}', \lambda) \right. \\ &\quad \left. + f_A(\hat{a}', \lambda) f_B(\hat{b}, \lambda) + f_A(\hat{a}', \lambda) f_B(\hat{b}', \lambda) \right], \end{aligned} \quad (14.4.3)$$

for arbitrary unit vectors  $\hat{a}$  and  $\hat{a}'$  associated with observable  $A$ , and  $\hat{b}$  and  $\hat{b}'$  associated with observable  $B$ . Since each of the four terms is bounded by  $-1$  and  $+1$ , in any conceivable theory the left side must be in the range from  $-4$  to  $4$ , inclusive. However, in hidden variables theories, a stronger statement holds. This is because a simple brute-force enumeration of all of the possible results for the contents of the square brackets on the right side shows that it can only be  $2$  or  $-2$ . (See Table 14.4.3.) Using eq. (14.2.1), this implies that the magnitude of the integral on the

$f_A(\hat{a}, \lambda)$	$f_A(\hat{a}', \lambda)$	$f_B(\hat{b}, \lambda)$	$f_B(\hat{b}', \lambda)$	[result]
1	1	1	1	2
1	1	1	-1	2
1	1	-1	1	-2
1	-1	1	1	-2
1	1	-1	-1	-2
1	-1	1	-1	2
1	-1	-1	1	-2
1	-1	-1	-1	2

Table 14.4.3: Possible measurement results pertaining to the ADR experiment, as dictated by eqs. (14.4.1)–(14.4.3), in a theory of local hidden variables. In addition to the eight possibilities shown, there are eight more in which all measurements have the opposite signs. The [result] in the last column is the quantity inside the square brackets on the right side of eq. (14.4.3). It is 2 or -2 in all cases.

right side of eq. (14.4.3) is bounded by 2, and so we have another Bell-type result, the CHSH inequality, named for John Clauser, Michael Horne, Abner Shimony, and Richard Holt,

$$|C(\hat{a}, \hat{b}) - C(\hat{a}, \hat{b}') + C(\hat{a}', \hat{b}) + C(\hat{a}', \hat{b}')| \leq 2, \quad (14.4.4)$$

which must be obeyed in any local hidden variables theory.

In the ADR experiment, an excited state of a calcium atom with total angular momentum  $J = 0$  decays to an intermediate state with  $J = 1$  by emitting one photon in an electric dipole transition, and then to another  $J = 0$  state by emitting a second photon in another electric dipole transition. The photons travel in opposite directions to distant detectors  $A$  and  $B$ , which by a choice of coordinates are located on the positive and negative  $z$  axis, respectively. Both photons are emitted with circular polarization, with angular momenta  $S_{1z} = \pm\hbar$  and  $S_{2z} = \mp\hbar$ , in a total spin  $S = 0$  state by angular momentum conservation. Thus, if one photon is in a  $L$ -circular polarization state, then the other also has  $L$ -circular polarization, because both its momentum direction and its spin are opposite to the first. Similarly, the two photons could both have  $R$ -circular polarization.

However, the detectors  $A$  and  $B$  measure the linear polarizations of the photons, not circular polarizations, as follows. A linear polarizer sends each photon into one photomultiplier if the linear polarization is along a selected direction, and into a different one if the linear polarization is in the orthogonal direction. (The allowed polarizations are always orthogonal to the photon momentum direction.) If detector  $A$  measures the linear polarization along the chosen direction  $\hat{a}$ , then it records  $A(\hat{a}) = +1$ , and if it instead detects the polarization perpendicular to  $\hat{a}$ , then it records  $A(\hat{a}) = -1$ . Similarly, if  $B$  sees the linear polarization along the chosen direction  $\hat{b}$ , then it records  $B(\hat{b}) = +1$ , and if it detects the polarization perpendicular to  $\hat{b}$ , then it records



$B(\hat{b}) = -1$ . The experiment is designed so that detector  $A$  can switch between two choices  $\hat{a}$  and  $\hat{a}'$ , and detector  $B$  can switch between  $\hat{b}$  and  $\hat{b}'$ . The switching is done randomly and quickly, with times shorter than the light travel time between the two detectors. This avoids possible biasing of the results by some hypothetical communication mechanism that might relate the choices made by the detectors.

Let us now work out the quantum mechanics prediction for the correlations of measurements of  $A(\hat{a})$  and  $B(\hat{b})$ . As already noted, the photons are produced in a state where the circular polarizations are the same, but could be either both  $L$  or both  $R$ . Since the atomic transitions have equal amplitudes for these two possibilities due to spherical symmetry, we can write

$$|S = 0\rangle = \frac{1}{\sqrt{2}}(|L, L\rangle + |R, R\rangle). \quad (14.4.5)$$

where the first and second ket labels refer to the photons that are seen by detectors  $A$  and  $B$ , respectively. The single-photon orthobasis states obey  $\langle R|R\rangle = \langle L|L\rangle = 1$  and  $\langle L|R\rangle = \langle R|L\rangle = 0$ . To make contact with the measurements made by the experiment, it is convenient to rewrite the state using a linear polarization orthobasis. Our chosen coordinates imply that the momenta of the photons that reach detectors  $A$  and  $B$  point in the directions  $\hat{k}_A = \hat{z}$  and  $\hat{k}_B = -\hat{z}$ . Then the transformations from the circular polarization orthobases to the linear polarization orthobases are

$$|L\rangle_A = \frac{1}{\sqrt{2}}(|\hat{x}\rangle_A - i|\hat{y}\rangle_A), \quad (14.4.6)$$

$$|R\rangle_A = \frac{1}{\sqrt{2}}(|\hat{x}\rangle_A + i|\hat{y}\rangle_A), \quad (14.4.7)$$

$$|L\rangle_B = \frac{1}{\sqrt{2}}(|\hat{x}\rangle_B + i|\hat{y}\rangle_B), \quad (14.4.8)$$

$$|R\rangle_B = \frac{1}{\sqrt{2}}(|\hat{x}\rangle_B - i|\hat{y}\rangle_B), \quad (14.4.9)$$

where  $|\hat{x}\rangle$  and  $|\hat{y}\rangle$  are states with linear polarizations along the  $x$  and  $y$  axes respectively, and  $\langle \hat{x}|\hat{x}\rangle = \langle \hat{y}|\hat{y}\rangle = 1$  and  $\langle \hat{x}|\hat{y}\rangle = \langle \hat{y}|\hat{x}\rangle = 0$  in each case. Using eqs. (14.4.6)–(14.4.9) in eq. (14.4.5),

$$|S = 0\rangle = \frac{1}{\sqrt{2}}(|\hat{x}, \hat{x}\rangle + |\hat{y}, \hat{y}\rangle). \quad (14.4.10)$$

Note that this state has even parity, because it is invariant under  $(\hat{x}, \hat{y}) \rightarrow (-\hat{x}, -\hat{y})$ , in agreement with the fact that the transition involves two even-parity atomic states.

A state  $|\hat{n}\rangle$  describing a photon with linear polarization  $\hat{n}$  has, up to an arbitrary phase,

$$\langle \hat{n}|\hat{x}\rangle = \hat{n}_x, \quad \langle \hat{n}|\hat{y}\rangle = \hat{n}_y. \quad (14.4.11)$$

Therefore, if we use  $|\hat{a}, \hat{b}\rangle$  to denote the state in which the linear polarizations are  $\hat{a}$  and  $\hat{b}$ , then

$$\langle \hat{a}, \hat{b}|S = 0\rangle = \frac{1}{\sqrt{2}}(\hat{a}_x \hat{b}_x + \hat{a}_y \hat{b}_y), \quad (14.4.12)$$




Figure 14.4.1: The configuration for the unit vectors  $\hat{a}$ ,  $\hat{a}'$ ,  $\hat{b}$ , and  $\hat{b}'$  that provides the maximum violation of the Bell-type inequality eq. (14.4.4) in the ADR photon polarization experiment, using the quantum mechanics prediction of eq. (14.4.15). The three angles labeled  $\theta$  are all equal to  $\pi/8$ , so that  $\theta_{ab} = \theta_{a'b} = \theta_{a'b'} = \pi/8$ , and  $\theta_{ab'} = 3\pi/8$ .

again up to an irrelevant phase. So, the probability for the linear polarizations to be observed as  $A(\hat{a}) = 1$ ,  $B(\hat{b}) = 1$  is

$$\frac{1}{2}(\hat{a} \cdot \hat{b})^2 = \frac{1}{2} \cos^2 \theta_{ab}, \quad (14.4.13)$$

where  $\theta_{ab}$  is the angle between  $\hat{a}$  and  $\hat{b}$ . The same result is obtained for the probability to observe the polarizations both orthogonal to  $\hat{a}$  and  $\hat{b}$ , so that  $A(\hat{a}) = -1$ ,  $B(\hat{b}) = -1$ . The probabilities to observe  $A(\hat{a}) = 1$ ,  $B(\hat{b}) = -1$  and  $A(\hat{a}) = -1$ ,  $B(\hat{b}) = 1$  are both

$$\frac{1}{2}|\hat{a} \times \hat{b}|^2 = \frac{1}{2} \sin^2 \theta_{ab}. \quad (14.4.14)$$

Weighting each of the outcomes for  $A(\hat{a})B(\hat{b})$  by these respective probabilities, we arrive at

$$\begin{aligned} C(\hat{a}, \hat{b}) &= \langle A(\hat{a})B(\hat{b}) \rangle = \frac{1}{2} \cos^2 \theta_{ab} + \frac{1}{2} \cos^2 \theta_{ab} - \frac{1}{2} \sin^2 \theta_{ab} - \frac{1}{2} \sin^2 \theta_{ab} \\ &= \cos(2\theta_{ab}). \end{aligned} \quad (14.4.15)$$

This is the quantum mechanics prediction for the correlation.

The local hidden variables Bell inequality, eq. (14.4.4), is incompatible with the quantum mechanics result of eq. (14.4.15) for many choices of  $\hat{a}$ ,  $\hat{a}'$ ,  $\hat{b}$ , and  $\hat{b}'$ . (Due to the geometry of the experiment and the transverse polarization of photons, these four unit vectors must all lie in the  $xy$  plane.) The most extreme violation of the inequality, used in the actual ADR experiment, occurs if one chooses the unit vectors as depicted in Figure 14.4.1, with

$$C(\hat{a}, \hat{b}) = C(\hat{a}', \hat{b}) = C(\hat{a}', \hat{b}') = \cos(\pi/4) = \frac{1}{\sqrt{2}}, \quad (14.4.16)$$

$$C(\hat{a}, \hat{b}') = \cos(3\pi/4) = -\frac{1}{\sqrt{2}}. \quad (14.4.17)$$

The (idealized) quantum mechanics prediction for that particular geometry is therefore

$$|C(\hat{a}, \hat{b}) - C(\hat{a}, \hat{b}') + C(\hat{a}', \hat{b}) + C(\hat{a}', \hat{b}')| = 2\sqrt{2} \approx 2.828. \quad (14.4.18)$$

The expected quantum mechanics prediction for the ADR experiment as performed was slightly lower at  $2.70 \pm 0.05$ , due to experimental non-ideal realities. This was in excellent agreement

with the actual experimental result  $2.697 \pm 0.015$  reported by ADR. This experimental result is also completely incompatible with the upper bound of 2 from eq. (14.4.4) in all theories with local hidden variables.

Further experiments have sharpened the result that local hidden variables cannot explain observed phenomena, and nonlocal entanglement of states as predicted by quantum mechanics is both consistent and supported by the experimental evidence. In particular, these experiments address a perverse loophole in Bell's original argument that we have not mentioned yet. It is natural to expect that the detector settings made in each trial (in the Aspect experiment case, the choice of  $\hat{a}$  or  $\hat{a}'$ , and the choice of  $\hat{b}$  or  $\hat{b}'$ ) should not be correlated in any special way with the hidden variables  $\lambda$  of the decaying particle. But what if they are, because of some influence in their shared past? This could mean that event statistics are somehow biased by the detector setting choices in just such a very special way as to allow a hidden variables theory to not only violate Bell-type inequalities but closely mimic the standard quantum mechanics predictions, a possibility sometimes called **superdeterminism**. Experiments of various types have taken great pains to choose the detector settings randomly in very different (and often quite convoluted) ways to discount this already implausible conspiracy theory scenario. Furthermore, the striking perfect-correlation quantum mechanics prediction for a state of the GHZ type discussed in section 14.3 (but with photon polarizations rather than spins) has been confirmed within experimental uncertainties by J.-W. Pan, D. Bouwmeester, M. Daniell, H. Weinfurter and A. Zeilinger, "Experimental test of quantum nonlocality in three-photon Greenberger–Horne–Zeilinger entanglement", *Nature* 403, 515 (2000). The experimental conclusion is that "local realism" is not real.

Since local hidden variables theories do not agree with experiment, one might entertain the possibility of "nonlocal hidden variables", which could allow interactions between space-like separated points. This is hard to rule out in general because it could predict anything, including causality violation; one might as well explain experiments by invoking sorcery. Furthermore, nonlocal hidden variables abandon the original motivation of local realism, and in any case are a cure for a disease that does not exist.

## 14.5 Exercises

**Exercise 14.1.** Consider the nonlocally entangled EPR state  $|S = 0\rangle$  of section 14.1. Suppose that Alice and Bob measure  $\hat{a} \cdot \vec{\sigma}_1$  and  $\hat{b} \cdot \vec{\sigma}_2$ , respectively. Let  $\theta$  be the angle between  $\hat{a}$  and  $\hat{b}$ . The four possible outcomes of their measurements are denoted  $(+1, +1)$ ,  $(+1, -1)$ ,  $(-1, +1)$ , and  $(-1, -1)$ .

(a) Suppose Alice's measurement was first. For the outcome  $\hat{a} \cdot \vec{\sigma}_1 = +1$ , the associated projection

operator can be written as

$$P_{\hat{a} \cdot \vec{\sigma}_1 = +1} = |\hat{a}, \hat{n}\rangle \langle \hat{a}, \hat{n}| + |\hat{a}, -\hat{n}\rangle \langle \hat{a}, -\hat{n}| \quad (14.5.1)$$

for any choice of unit vector  $\hat{n}$ . It is convenient to choose  $\hat{n} = \hat{a}$ , both in eq. (14.5.1) and in the expression for the starting state  $|S = 0\rangle$  in eq.(14.1.2). Apply Postulates 4 and 5 to show that the probability of this outcome is  $1/2$ , and that the normalized state after the measurement is  $|\hat{a}, -\hat{a}\rangle$ . Now consider Bob's subsequent measurement. The projection operator associated with Bob's outcome  $\hat{b} \cdot \vec{\sigma}_2 = +1$  is

$$P_{\hat{b} \cdot \vec{\sigma}_2 = +1} = |\hat{n}, \hat{b}\rangle \langle \hat{n}, \hat{b}| + |-\hat{n}, \hat{b}\rangle \langle -\hat{n}, \hat{b}|, \quad (14.5.2)$$

where again  $\hat{n}$  is any unit vector. It is again convenient to choose  $\hat{n} = \hat{a}$ . Apply Postulates 4 and 5 again to show that the probability for the outcome  $(+1, +1)$  is  $\mathcal{P}(+1, +1) = \frac{1}{2} \sin^2(\theta/2)$ , and that the state after the measurements is  $|\hat{a}, \hat{b}\rangle$ .

(b) Repeat part (a), but this time take Bob's measurement first. Show that the final state and the probability for the outcome  $(+1, +1)$  are the same as found in part (a).

(c) Repeat parts (a) and (b), but this time assuming the measurements are simultaneous, using a projection operator

$$P_{\hat{a} \cdot \vec{\sigma}_1 = +1, \hat{b} \cdot \vec{\sigma}_2 = +1} = |\hat{a}, \hat{b}\rangle \langle \hat{a}, \hat{b}|. \quad (14.5.3)$$

Check that the results agree with those found in parts (a) and (b).

(d) Repeat the analysis of parts (a), (b), and (c) for each of the other three outcomes  $(+1, -1)$ ,  $(-1, +1)$ , and  $(-1, -1)$ . As a check, your four probabilities should add to 1. As a further check, use your four probabilities to obtain the expectation value  $\langle \hat{a} \cdot \vec{\sigma}_1 \hat{b} \cdot \vec{\sigma}_2 \rangle$  given in eq. (14.1.6).

**Exercise 14.2.** Show that Mermin's 3-spin GKZ-type state in eq. (14.3.1) satisfies the eigenvalue equations (14.3.3) and (14.3.4) for the operators  $\Omega_{xyy}$ ,  $\Omega_{yxy}$ ,  $\Omega_{yyx}$  and  $\Omega_{xxx}$  defined in terms of the individual spin components in eq. (14.3.2).

## 15 Stationary-state perturbation theory

### 15.1 Perturbative expansion for energy eigenstates

Given a Hamiltonian  $H$  that does not depend on time, we would like to solve the time-independent Schrödinger equation,

$$H|\psi_n\rangle = E_n|\psi_n\rangle, \quad (15.1.1)$$

as an eigenvalue problem for the stationary states  $|\psi_n\rangle$  and their energies  $E_n$ . Unfortunately, in many cases this is too hard to do exactly. However, suppose that it is possible to split the Hamiltonian into two parts,

$$H = H_0 + \lambda W, \quad (15.1.2)$$

where  $H_0$  is a simpler Hamiltonian for which we already know how to solve the eigenvalue problem, and  $\lambda$  is a small parameter, so that  $W$  can be treated as a perturbation. Then we can hope to find an approximate solution for the full Hamiltonian  $H$ , by expanding in  $\lambda$ . We will now work out how to do this. This is called **time-independent perturbation theory** or **stationary-state perturbation theory**.

To begin, assume that we have found the complete set of eigenstates  $|n\rangle$  and eigenvalues  $\mathcal{E}_n$  for  $H_0$ , so that

$$H_0|n\rangle = \mathcal{E}_n|n\rangle. \quad (15.1.3)$$

In general, the states  $|n\rangle$  for a given  $\mathcal{E}_n$  might be degenerate, but for now we assume they are not. For each unperturbed stationary state  $|n\rangle$  and energy  $\mathcal{E}_n$ , we want to solve for the corresponding  $|\psi_n\rangle$  and energy  $E_n$ . The connection between the unperturbed and perturbed stationary-state solutions is assumed to be smoothly continuous in  $\lambda$  in order for the following strategy to work. In particular, there should be a one-to-one correspondence between the known  $|n\rangle$  and the desired  $|\psi_n\rangle$  for all  $\lambda$ , as long as it is not too large.

The perturbation theory strategy is to write the unknown full eigenstates and energy eigenvalues each as a power series in  $\lambda$ ,

$$|\psi_n\rangle = |\psi_n^{(0)}\rangle + \lambda|\psi_n^{(1)}\rangle + \lambda^2|\psi_n^{(2)}\rangle + \lambda^3|\psi_n^{(3)}\rangle + \cdots, \quad (15.1.4)$$

$$E_n = E_n^{(0)} + \lambda E_n^{(1)} + \lambda^2 E_n^{(2)} + \lambda^3 E_n^{(3)} + \cdots. \quad (15.1.5)$$

Now,  $\lambda = 0$  must recover the unperturbed solutions, so

$$|\psi_n^{(0)}\rangle = |n\rangle, \quad (15.1.6)$$

$$E_n^{(0)} = \mathcal{E}_n. \quad (15.1.7)$$

The unperturbed stationary states are a complete orthobasis, with

$$\sum_n |n\rangle \langle n| = I, \quad (15.1.8)$$

$$\langle n'|n\rangle = \delta_{nn'}. \quad (15.1.9)$$

This is always true, because  $H_0$  is assumed to be an observable.

We could also choose unit normalization for the eigenkets  $|\psi_n\rangle$ , but we won't. It is much more convenient to choose the following normalization condition for them instead:

$$\langle n|\psi_n\rangle = 1 \quad (\text{for each } n), \quad (15.1.10)$$

which will result in  $\langle \psi_n|\psi_n\rangle \neq 1$ . After obtaining the solutions for  $|\psi_n\rangle$ , we can re-normalize them later, as a last step. Plugging eq. (15.1.4) into eq. (15.1.10), we have

$$\langle n| (|n\rangle + \lambda|\psi_n^{(1)}\rangle + \lambda^2|\psi_n^{(2)}\rangle + \cdots) = 1. \quad (15.1.11)$$

Since  $\langle n|n\rangle = 1$ , in order to have a solution for general  $\lambda$ , it is necessary and sufficient that the coefficient of  $\lambda^k$  in eq. (15.1.11) vanishes for each  $k \geq 1$ , which implies

$$\langle n|\psi_n^{(k)}\rangle = 0 \quad (\text{for } k = 1, 2, 3, \dots). \quad (15.1.12)$$

Thus, the normalization choice eq. (15.1.10) ensures that all of the corrections to each state are orthogonal to the corresponding unperturbed state. This is extremely important, and will be used repeatedly in the following.

The time-independent Schrödinger equation (15.1.1) now reads

$$\begin{aligned} (H_0 + \lambda W) (|n\rangle + \lambda|\psi_n^{(1)}\rangle + \lambda^2|\psi_n^{(2)}\rangle + \cdots) = \\ (\mathcal{E}_n + \lambda E_n^{(1)} + \lambda^2 E_n^{(2)} + \cdots) (|n\rangle + \lambda|\psi_n^{(1)}\rangle + \lambda^2|\psi_n^{(2)}\rangle + \cdots). \end{aligned} \quad (15.1.13)$$

Expanding both sides, and matching the coefficient of each power  $\lambda^k$ , we get

$$\lambda^0: \quad H_0|n\rangle = \mathcal{E}_n|n\rangle, \quad (15.1.14)$$

$$\lambda^1: \quad H_0|\psi_n^{(1)}\rangle + W|n\rangle = \mathcal{E}_n|\psi_n^{(1)}\rangle + E_n^{(1)}|n\rangle, \quad (15.1.15)$$

$$\lambda^2: \quad H_0|\psi_n^{(2)}\rangle + W|\psi_n^{(1)}\rangle = \mathcal{E}_n|\psi_n^{(2)}\rangle + E_n^{(1)}|\psi_n^{(1)}\rangle + E_n^{(2)}|n\rangle, \quad (15.1.16)$$

and, in general,

$$\lambda^k: \quad H_0|\psi_n^{(k)}\rangle + W|\psi_n^{(k-1)}\rangle = \mathcal{E}_n|\psi_n^{(k)}\rangle + \sum_{j=1}^{k-1} E_n^{(j)}|\psi_n^{(k-j)}\rangle + E_n^{(k)}|n\rangle. \quad (15.1.17)$$

Equation (15.1.14) is, of course, just a repetition of eq. (15.1.3).

Consider first the effect of the perturbation at order  $\lambda^1$ . Taking the inner product of eq. (15.1.15) with  $\langle n|$  gives

$$\langle n|H_0|\psi_n^{(1)}\rangle + \langle n|W|n\rangle = \mathcal{E}_n\langle n|\psi_n^{(1)}\rangle + E_n^{(1)}\langle n|n\rangle. \quad (15.1.18)$$

Now using  $\langle n|H_0 = \mathcal{E}_n\langle n|$  on the first term, we see that it and the first term on the right side are equal, and both vanish due to eq. (15.1.12). Therefore, using the orthonormality condition eq. (15.1.9) on the last term, eq. (15.1.18) simplifies to

$$E_n^{(1)} = \langle n|W|n\rangle, \quad (15.1.19)$$

so that

$$E_n = \mathcal{E}_n + \lambda\langle n|W|n\rangle + \dots. \quad (15.1.20)$$

To first order in the expansion in  $\lambda$ , the energy shift in the state  $|n\rangle$  is simply the expectation value of the Hamiltonian perturbation. This is the most important and commonly used result from stationary-state perturbation theory.

To find the first-order correction to the energy eigenkets, we note that completeness of the  $H_0$  eigenstates allows us to write

$$|\psi_n^{(1)}\rangle = \sum_{m \neq n} |m\rangle \langle m|\psi_n^{(1)}\rangle, \quad (15.1.21)$$

where we have made good use of eq. (15.1.12) to dispense with the  $m = n$  term. We now need to find the coefficients  $\langle m|\psi_n^{(1)}\rangle$  for  $m \neq n$ . To do so, take the inner product of  $\langle m|$  acting on eq. (15.1.15),

$$\langle m|H_0|\psi_n^{(1)}\rangle + \langle m|W|n\rangle = \mathcal{E}_n\langle m|\psi_n^{(1)}\rangle + E_n^{(1)}\langle m|n\rangle. \quad (15.1.22)$$

The last term vanishes due to the orthonormality of the unperturbed stationary states eq. (15.1.9), and the first term on the left can be simplified using  $\langle m|H_0 = \mathcal{E}_m\langle m|$ . Therefore,

$$(\mathcal{E}_n - \mathcal{E}_m)\langle m|\psi_n^{(1)}\rangle = \langle m|W|n\rangle \quad (m \neq n). \quad (15.1.23)$$

Now, assuming that there are no degeneracies, so that  $\mathcal{E}_m \neq \mathcal{E}_n$  for  $m \neq n$ , we can solve for  $\langle m|\psi_n^{(1)}\rangle$ , and eq. (15.1.21) becomes

$$|\psi_n^{(1)}\rangle = \sum_{m \neq n} |m\rangle \frac{\langle m|W|n\rangle}{\mathcal{E}_n - \mathcal{E}_m}. \quad (15.1.24)$$

We have thus obtained the perturbed energy eigenstates at first order in the expansion in  $\lambda$ ,

$$|\psi_n\rangle = |n\rangle + \lambda \sum_{m \neq n} |m\rangle \frac{\langle m|W|n\rangle}{\mathcal{E}_n - \mathcal{E}_m} + \dots. \quad (15.1.25)$$

We can now see why the normalization condition eq. (15.1.10) was so useful and important; it gave us eq. (15.1.12), which in turn allowed us to exclude  $m = n$  from the sum, which avoided a term with a disastrous vanishing denominator<sup>†</sup> in  $|\psi_n^{(1)}\rangle$ , and in similar expressions to follow.

Having solved the perturbation theory problem at order  $\lambda^1$ , we can continue iteratively to obtain the perturbed energies and their eigenstates at any desired order in  $\lambda$ . We will now find the necessary recurrence relations in general. At order  $\lambda^k$ , the inner product of  $\langle n|$  with eq. (15.1.17) is

$$\langle n|H_0|\psi_n^{(k)}\rangle + \langle n|W|\psi_n^{(k-1)}\rangle = \mathcal{E}_n \langle n|\psi_n^{(k)}\rangle + \sum_{j=1}^{k-1} E_n^{(j)} \langle n|\psi_n^{(k-j)}\rangle + E_n^{(k)}. \quad (15.1.26)$$

The first term on the left vanishes because of eq. (15.1.12), after pulling out  $H_0 = \mathcal{E}_n$ . On the right side of the equality, using eq. (15.1.12) again ensures that all of the terms vanish except the last one, where we have used  $\langle n|n\rangle = 1$ . So, we find the remarkable result

$$E_n^{(k)} = \langle n|W|\psi_n^{(k-1)}\rangle, \quad (15.1.27)$$

that the  $\lambda^k$  correction to each energy eigenvalue is always obtained in terms of the  $\lambda^{k-1}$  correction to the corresponding ket.

Also, using completeness, we can write

$$|\psi_n^{(k)}\rangle = \sum_{m \neq n} |m\rangle \langle m|\psi_n^{(k)}\rangle, \quad (15.1.28)$$

where once again eq. (15.1.12) has been used to eliminate the  $m = n$  term. To find the coefficients  $\langle m|\psi_n^{(k)}\rangle$  for  $m \neq n$ , we take  $\langle m|$  acting on eq. (15.1.17), which gives

$$(\mathcal{E}_n - \mathcal{E}_m) \langle m|\psi_n^{(k)}\rangle = \langle m|W|\psi_n^{(k-1)}\rangle - \sum_{j=1}^{k-1} E_n^{(j)} \langle m|\psi_n^{(k-j)}\rangle, \quad (15.1.29)$$

where  $\langle m|n\rangle = 0$  has been used to eliminate the term proportional to  $E_n^{(k)}$ . The right-hand side only involves expressions from orders less than  $k$ . Therefore, eqs. (15.1.27), (15.1.28), and (15.1.29) can be used to solve iteratively for  $E_n^{(k)}$  and then  $|\psi_n^{(k)}\rangle$  at each successive level  $k$ .

Let us apply the preceding for  $k = 2$ . Evaluating eq. (15.1.27) using eq. (15.1.24) gives

$$E_n^{(2)} = \sum_{m \neq n} \frac{|\langle m|W|n\rangle|^2}{\mathcal{E}_n - \mathcal{E}_m}. \quad (15.1.30)$$

We have thus obtained the second-order corrections to the energies. Also, eq. (15.1.29) gives

$$\langle m|\psi_n^{(2)}\rangle = \frac{1}{\mathcal{E}_n - \mathcal{E}_m} \left( \langle m|W|\psi_n^{(1)}\rangle - E_n^{(1)} \langle m|\psi_n^{(1)}\rangle \right), \quad (15.1.31)$$

---

<sup>†</sup>Of course, the denominator is still a disaster if the unperturbed states have energy degeneracies, so that  $\mathcal{E}_m = \mathcal{E}_n$  for some  $m \neq n$ . This issue is addressed by “degenerate perturbation theory”, in section 15.6.



or, using eqs. (15.1.19) and (15.1.24),

$$\langle m|\psi_n^{(2)}\rangle = \sum_{p \neq n} \frac{\langle m|W|p\rangle\langle p|W|n\rangle}{(\mathcal{E}_n - \mathcal{E}_p)(\mathcal{E}_n - \mathcal{E}_m)} - \frac{\langle m|W|n\rangle\langle n|W|n\rangle}{(\mathcal{E}_n - \mathcal{E}_m)^2} \quad (15.1.32)$$

for  $n \neq m$ . Summarizing the results to order  $\lambda^2$ :

$$E_n = \mathcal{E}_n + \lambda\langle n|W|n\rangle + \lambda^2 \sum_{m \neq n} \frac{|\langle m|W|n\rangle|^2}{\mathcal{E}_n - \mathcal{E}_m} + \dots, \quad (15.1.33)$$

and

$$\begin{aligned} |\psi_n\rangle &= |n\rangle + \lambda \sum_{m \neq n} |m\rangle \frac{\langle m|W|n\rangle}{\mathcal{E}_n - \mathcal{E}_m} \\ &+ \lambda^2 \sum_{m \neq n} |m\rangle \left( \sum_{p \neq n} \frac{\langle m|W|p\rangle\langle p|W|n\rangle}{(\mathcal{E}_n - \mathcal{E}_p)(\mathcal{E}_n - \mathcal{E}_m)} - \frac{\langle m|W|n\rangle\langle n|W|n\rangle}{(\mathcal{E}_n - \mathcal{E}_m)^2} \right) + \dots \end{aligned} \quad (15.1.34)$$

Don't forget that, by design, this ket  $|\psi_n\rangle$  does not have unit norm, but now it can be renormalized as  $|\psi_n\rangle/\sqrt{\langle\psi_n|\psi_n\rangle}$ , if desired. In doing so, it often makes sense to expand in  $\lambda$  and drop the terms of order  $\lambda^3$  or higher. From eq. (15.1.34), we find using the orthonormality of the unperturbed kets that

$$\langle\psi_n|\psi_n\rangle = 1 + \lambda^2 \sum_{m \neq n} \frac{|\langle m|W|n\rangle|^2}{(\mathcal{E}_n - \mathcal{E}_m)^2} + \dots, \quad (15.1.35)$$

so that, working consistently to second order in  $\lambda$ , the normalized eigenstate can be obtained by just multiplying the first term  $|n\rangle$  in eq. (15.1.34) by

$$1/\sqrt{\langle\psi_n|\psi_n\rangle} = 1 - \frac{\lambda^2}{2} \sum_{m \neq n} \frac{|\langle m|W|n\rangle|^2}{(\mathcal{E}_n - \mathcal{E}_m)^2} + \dots \quad (15.1.36)$$

It is also worth noting that, using eq. (15.1.27), the energy eigenvalue to all orders in  $\lambda$  is

$$E_n = \mathcal{E}_n + \sum_{k=1}^{\infty} \lambda^k \langle n|W|\psi_n^{(k-1)}\rangle = \mathcal{E}_n + \lambda \langle n|W \sum_{k=1}^{\infty} \lambda^{k-1} |\psi_n^{(k-1)}\rangle, \quad (15.1.37)$$

which can be rewritten

$$E_n = \mathcal{E}_n + \lambda \langle n|W|\psi_n\rangle. \quad (15.1.38)$$

The all-orders energy  $E_n$  is therefore known exactly in terms of the all-orders state ket  $|\psi_n\rangle$ . Of course, the latter may well be only known partially as a perturbative expansion.

So far, we have used the parameter  $\lambda$  as a way of keeping track of the order of perturbation theory. In any given term, the power of  $\lambda$  is the same as the number of matrix elements of

$W$ , so we can simply absorb  $\lambda$  into the definition of  $W$  without losing any information. (The convergence of perturbation theory still relies on the assumption that the properties of  $W$  allow it to be treated as a small correction. For example,  $W$  may contain one or more small dimensionless parameters that effectively play the role of  $\lambda$ .) Equivalently, taking  $\lambda = 1$ , we have

$$E_n = \mathcal{E}_n + E_n^{(1)} + E_n^{(2)} + E_n^{(3)} + \cdots, \quad (15.1.39)$$

$$|\psi_n\rangle = |n\rangle + |\psi_n^{(1)}\rangle + |\psi_n^{(2)}\rangle + |\psi_n^{(3)}\rangle + \cdots. \quad (15.1.40)$$

The preceding results can be expressed more compactly with the following simplifying notation,

$$W_{mn} = \langle m|W|n\rangle, \quad (15.1.41)$$

$$\mathcal{E}_{nm} = \mathcal{E}_n - \mathcal{E}_m. \quad (15.1.42)$$

Then, the energy eigenvalue corrections are summarized as

$$E_n^{(1)} = W_{nn}, \quad (15.1.43)$$

$$E_n^{(2)} = \sum_{m \neq n} \frac{|W_{mn}|^2}{\mathcal{E}_{nm}}, \quad (15.1.44)$$

$$E_n^{(3)} = \sum_{m \neq n} \sum_{p \neq n} \frac{W_{nm}W_{mp}W_{pn}}{\mathcal{E}_{np}\mathcal{E}_{nm}} - W_{nn} \sum_{m \neq n} \frac{|W_{mn}|^2}{\mathcal{E}_{nm}^2}, \quad (15.1.45)$$

etc., and for the (un-normalized) energy eigenstates,

$$|\psi_n^{(1)}\rangle = \sum_{m \neq n} |m\rangle \frac{W_{mn}}{\mathcal{E}_{nm}}, \quad (15.1.46)$$

$$|\psi_n^{(2)}\rangle = \sum_{m \neq n} |m\rangle \left( \sum_{p \neq n} \frac{W_{mp}W_{pn}}{\mathcal{E}_{np}\mathcal{E}_{nm}} - \frac{W_{nn}W_{mn}}{\mathcal{E}_{nm}^2} \right), \quad (15.1.47)$$

etc. All higher order corrections likewise involve powers of matrix elements of  $W$  in the numerators, and powers of unperturbed energy differences  $\mathcal{E}_{nm}$ 's in the denominators. This means that the perturbative expansion, as we have constructed it here, may fail to converge for a state  $|n\rangle$  if there is some other state  $|m\rangle$  such that  $\mathcal{E}_{nm}$  is zero, or small compared to the matrix elements of  $W$ . To treat such cases, we will need to use degenerate perturbation theory (section 15.6), or almost-degenerate perturbation theory (section 15.8), respectively.

## 15.2 Simple examples of perturbation theory

Consider the one-dimensional harmonic oscillator with a linear perturbation:

$$H_0 = \frac{P^2}{2m} + \frac{1}{2}m\omega^2 X^2, \quad W = -fX. \quad (15.2.1)$$

Here  $f$  is a constant force, which plays the role of the expansion parameter  $\lambda$  in the discussion of section 15.1. Before solving this as a problem in perturbation theory for small  $f$ , we recall that it is special in that the stationary states can actually be found exactly. As already noted at the end of section 7.4, this is because the total Hamiltonian can be rewritten, by completing the square, as

$$H = \frac{P^2}{2m} + \frac{1}{2}m\omega^2\left(X - \frac{f}{m\omega^2}\right)^2 - \frac{f^2}{2m\omega^2}, \quad (15.2.2)$$

so that after a shift of variables  $X \rightarrow X + f/m\omega^2$ , the new  $H$  will have the same  $m$ ,  $\omega$ , and energy eigenvalues, except that the latter will be offset by a constant  $-f^2/2m\omega^2$ . Thus we have

$$E_n = \hbar\omega(n + 1/2) - f^2/2m\omega^2, \quad (15.2.3)$$

exactly. Furthermore, the corresponding energy eigenstates must be

$$|\psi_n\rangle = T(f/m\omega^2)|n\rangle, \quad (15.2.4)$$

where  $T(a)$  is the translation operator of eq. (5.2.1), and  $|n\rangle$  are the stationary states of  $H_0$ . We can expand this exact result to linear order in  $f$ , using eq. (7.3.4) for the momentum operator in terms of harmonic oscillator creation and annihilation operators, to get

$$T(f/m\omega^2) = I + \frac{f}{\sqrt{2m\hbar\omega^3}}(a^\dagger - a) + \mathcal{O}(f^2), \quad (15.2.5)$$

so that the eigenstates of  $H$  are

$$|\psi_n\rangle = |n\rangle + \frac{f}{\sqrt{2m\hbar\omega^3}}\left(\sqrt{n+1}|n+1\rangle - \sqrt{n}|n-1\rangle\right) + \mathcal{O}(f^2) \quad (15.2.6)$$

in terms of the eigenstates of  $H_0$ .

Now let us use solve the same problem using the technology of perturbation theory that we have developed. The unperturbed energy eigenvalues and eigenstates are  $\mathcal{E}_n = \hbar\omega(n + 1/2)$  and  $|n\rangle$ . The general matrix elements needed for the perturbative expansion are

$$\begin{aligned} W_{n'n} = \langle n'|W|n\rangle &= -f \langle n'|X|n\rangle = -f\sqrt{\frac{\hbar}{2m\omega}} \langle n'|(a^\dagger + a)|n\rangle \\ &= -f\sqrt{\frac{\hbar}{2m\omega}} \left(\sqrt{n+1}\delta_{n',n+1} + \sqrt{n}\delta_{n',n-1}\right). \end{aligned} \quad (15.2.7)$$

Now, since this vanishes for  $n' = n$ , the first-order correction to the energies are all 0. From eqs. (15.1.43) and (15.1.44), we get, through second order,

$$E_n^{(1)} = 0, \quad (15.2.8)$$

$$E_n^{(2)} = \frac{|-f\sqrt{\hbar(n+1)/2m\omega}|^2}{\mathcal{E}_n - \mathcal{E}_{n+1}} + \frac{|-f\sqrt{\hbar n/2m\omega}|^2}{\mathcal{E}_n - \mathcal{E}_{n-1}}, \quad (15.2.9)$$

where only two terms in the sum contributed. Now  $\mathcal{E}_n - \mathcal{E}_{n+1} = -\hbar\omega$  and  $\mathcal{E}_n - \mathcal{E}_{n-1} = \hbar\omega$ , so eq. (15.2.9) simplifies to

$$E_n^{(2)} = -f^2/2m\omega^2, \quad (15.2.10)$$

and we have

$$E_n = \hbar\omega(n + 1/2) - f^2/2m\omega^2 + \dots. \quad (15.2.11)$$

This agrees with the exact result of eq. (15.2.3). In fact, this comparison shows that the possible higher-order terms (indicated by the ellipsis here) must actually conspire to vanish. The first-order perturbative correction to the stationary state  $|n\rangle$ , applying eq. (15.1.46), also only has two terms in the sum,

$$|\psi_n^{(1)}\rangle = -f\sqrt{\frac{\hbar}{2m\omega}} \left( \frac{\sqrt{n+1}|n+1\rangle}{-\hbar\omega} + \frac{\sqrt{n}|n-1\rangle}{\hbar\omega} \right) \quad (15.2.12)$$

$$= \frac{f}{\sqrt{2m\hbar\omega^3}} \left( \sqrt{n+1}|n+1\rangle - \sqrt{n}|n-1\rangle \right). \quad (15.2.13)$$

Again this agrees with the exact result in eq. (15.2.4), after the expansion for linear order in  $f$  as in eq. (15.2.6).

As a second example, less amenable to an easy exact calculation, consider the potential

$$V(x) = \begin{cases} V_0 & (|x| < a/2), \\ 0 & (a/2 < |x| < L/2), \\ \infty & (|x| > L/2), \end{cases} \quad (15.2.14)$$

as in Figure 15.2.1. This is an infinite square well with a perturbation bump in the middle, so we define  $H = H_0 + W$  where, using a hybrid operator/position-representation notation,

$$H_0 = \frac{P^2}{2m} + \begin{cases} 0 & (|x| < L/2), \\ \infty & (|x| > L/2), \end{cases} \quad (15.2.15)$$




Figure 15.2.1: An infinite one-dimensional square well potential with width  $L$ , with a symmetric bump of height  $V_0$  and width  $a$  to be treated as a perturbation.

and the perturbation Hamiltonian is the bump potential

$$W = \begin{cases} V_0 & (|x| < a/2), \\ 0 & (|x| > a/2). \end{cases} \quad (15.2.16)$$

The infinite square well problem defined by  $H_0$  was solved in section 6.4. The wavefunctions  $\langle x|n\rangle = \phi_n(x)$  for the unperturbed  $H_0$  eigenstates  $|n\rangle$  were found to be

$$\phi_n(x) = \sqrt{\frac{2}{L}} \cos(n\pi x/L) \quad (n = 1, 3, 5, \dots), \quad (15.2.17)$$

$$\phi_n(x) = \sqrt{\frac{2}{L}} \sin(n\pi x/L) \quad (n = 2, 4, 6, \dots), \quad (15.2.18)$$

with, in both cases, energies

$$\mathcal{E}_n = \frac{\hbar^2 \pi^2 n^2}{2mL^2}. \quad (15.2.19)$$

Applying eq. (15.1.19) or its equivalent eq. (15.1.43), the leading correction to the energy eigenvalue for the state  $n$  is

$$E_n^{(1)} = V_0 \int_{-a/2}^{a/2} dx |\phi_n(x)|^2. \quad (15.2.20)$$

For the ground state with  $n = 1$ , the energy eigenvalue to first order in  $V_0$  evaluates to

$$E_1 = \frac{\hbar^2 \pi^2}{2mL^2} + V_0 \left( \frac{a}{L} + \frac{1}{\pi} \sin(\pi a/L) \right). \quad (15.2.21)$$

As a check, note that if  $a = L$ , then the perturbation simply adds  $V_0$  to the energy eigenvalue, independent of  $L$ . This is just a constant shift in the Hamiltonian.

For the change in the ground-state wavefunction, we apply eq. (15.1.46). This requires us to find  $W_{k1}/\mathcal{E}_{1k}$ , which for odd  $k \neq 1$  is

$$\frac{\langle k|W|1\rangle}{\mathcal{E}_1 - \mathcal{E}_k} = \frac{2mL^2}{\hbar^2 \pi^2 (1 - k^2)} \frac{2V_0}{L} \int_{-a/2}^{a/2} dx \cos\left(\frac{\pi k x}{L}\right) \cos\left(\frac{\pi x}{L}\right) \quad (15.2.22)$$

$$= \frac{4V_0 m L^2}{\hbar^2 \pi^3 (1 - k^2)} \left[ \frac{\sin(\pi a(k-1)/2L)}{k-1} + \frac{\sin(\pi a(k+1)/2L)}{k+1} \right]. \quad (15.2.23)$$

For even  $k$  we have instead

$$\frac{\langle k|W|1\rangle}{\mathcal{E}_1 - \mathcal{E}_k} = 0, \quad (15.2.24)$$

because of

$$\int_{-a/2}^{a/2} dx \sin\left(\frac{\pi k x}{L}\right) \cos\left(\frac{\pi x}{L}\right) = 0. \quad (15.2.25)$$

This reflects (sorry not sorry about the pun) the parity selection rule of eq. (5.4.16). Since the perturbation Hamiltonian has even parity  $\pi_W = 1$ , and  $\pi_k = (-1)^{k-1}$  for a state  $|k\rangle$ , the parity selection rule tells us that  $\langle k|W|n\rangle$  vanishes if  $k+n$  is odd, and in particular that  $\langle k|W|1\rangle = 0$  for even  $k$ . It follows that the corrected ground-state wavefunction is

$$\psi_1(x) = \sqrt{\frac{2}{L}} \cos\left(\frac{\pi x}{L}\right) + \psi_1^{(1)}(x), \quad (15.2.26)$$

with

$$\psi_1^{(1)}(x) = \sum_{k=3,5,7,\dots} \frac{\langle k|W|1\rangle}{\mathcal{E}_1 - \mathcal{E}_k} \phi_k(x) \quad (15.2.27)$$

The corrections to the energies and wavefunctions of the other stationary states are left as an exercise.

For a third example, let us estimate the effect of the proton size on the energy levels of the hydrogen atom. In doing so, we will treat the proton (not completely realistically) as a ball of uniform charge density with radius  $a_p$ . For all  $r \geq a_p$ , the electric field is the same as if the proton were a point charge. For  $r \leq a_p$ , an elementary application of Gauss' Law in classical electrostatics shows that the electric field rises linearly with  $r$ , so that the potential energy of the electron is

$$V(r) = \begin{cases} -e^2 \left( \frac{3}{2a_p} - \frac{r^2}{2a_p^3} \right), & (r \leq a_p), \\ -\frac{e^2}{r}, & (r \geq a_p), \end{cases} \quad (15.2.28)$$

as shown in Figure 15.2.2. Therefore, compared to our treatment in section 11.1, the perturbation to the Hamiltonian is

$$W = \begin{cases} e^2 \left( \frac{1}{r} - \frac{3}{2a_p} + \frac{r^2}{2a_p^3} \right), & (r \leq a_p), \\ 0, & (r \geq a_p), \end{cases} \quad (15.2.29)$$




Figure 15.2.2: The solid line shows the shape of the potential energy for the electron in the hydrogen atom, with the proton modeled as a sphere of radius  $a_p$  with uniform charge density. The dashed line shows the potential with the proton modeled as a point, as in section 11.1.

in the position representation. So, the first-order correction to the ground state energy is

$$E_{1,0,0}^{(1)} = \langle 1, 0, 0 | W | 1, 0, 0 \rangle \quad (15.2.30)$$

$$= 4\pi \int_0^{a_p} dr r^2 \left( \frac{1}{\sqrt{\pi a_0^3}} e^{-r/a_0} \right)^2 e^2 \left( \frac{1}{r} - \frac{3}{2a_p} + \frac{r^2}{2a_p^3} \right), \quad (15.2.31)$$

where the  $4\pi$  factor comes from the angular integration. Within the radial integral,  $e^{-2r/a_0} \approx 1$  is an excellent approximation, since  $a_p \ll a_0$ . The integration then yields

$$E_{1,0,0}^{(1)} = \frac{2e^2 a_p^2}{5a_0^3} = \frac{4a_p^2}{5a_0^2} \text{Rydberg} \approx 2.1 \times 10^{-10} \text{Rydberg}, \quad (15.2.32)$$

where we have used  $a_p \approx 8.5 \times 10^{-16}$  meters in the last approximation.

In the preceding, we used the small size of the proton compared to the Bohr radius to effectively replace the radial wavefunction by its (very nearly constant) value at  $r = 0$ , with  $|R_{1,0}(r)|^2 \rightarrow |R_{1,0}(0)|^2$ . For a general  $n, l, m$  state, the same procedure will give

$$E_{n,l,m}^{(1)} = |R_{n,l}(0)|^2 \int d\Omega |Y_l^m(\theta, \phi)|^2 \int_0^{a_p} dr r^2 e^2 \left( \frac{1}{r} - \frac{3}{2a_p} + \frac{r^2}{2a_p^3} \right). \quad (15.2.33)$$

Now,  $\int d\Omega |Y_l^m|^2 = 1$ , and  $R_{n,l}(0) = 0$  unless  $l = 0$ , which also implies  $m = 0$ . From eq. (11.1.41), we know that the hydrogen atom radial wavefunction evaluated at  $r = 0$  is  $R_{n,0}(0) = \delta_{l0} 2/(na_0)^{3/2}$ . So, we find

$$E_{n,l,m}^{(1)} = \delta_{l0} \delta_{m0} \frac{4a_p^2}{5a_0^2 n^3} \text{Rydberg}. \quad (15.2.34)$$

Thus the finite proton radius only affects the energies of the states with no angular momentum, and with decreasing importance for larger principal quantum number  $n$ .

The preceding calculation is just an estimate, as the proton is certainly not a ball of uniform charge density. A more sophisticated calculation can be used to define the effective charge radius of the proton, which can then be compared to various experimental determinations of the same quantity. This has recently been the subject of some interest and controversy, because experimental determinations of the proton charge radius inferred from electron scattering seem to be slightly larger than those obtained from measuring the energy levels of muonic hydrogen, in which a muon replaces the electron. The muon is used because its much larger mass gives it a smaller Bohr radius than the electron, leading to a larger effect in eq. (15.2.34) by a factor  $m_\mu^2/m_e^2 \approx 42753$ ; the heavier muon is much more likely to be found inside the proton than an electron, and therefore provides a better probe.

### 15.3 Helium atom ground state from first-order perturbation theory

In this section, we consider the helium atom consisting of two electrons each with mass  $m_e$  and charge  $-e$ , and a heavy nucleus with charge  $Ze = +2e$  at the origin, as pictured in Figure 15.3.1.




Figure 15.3.1: The helium atom consists of two electrons with mass  $m_e$  and charge  $-e$  at positions  $\vec{r}_1$  and  $\vec{r}_2$ , and a very heavy nucleus of charge  $+2e$  at the origin.

The Hamiltonian is<sup>†</sup>

$$H = H_1 + H_2 + H_{12}, \quad (15.3.1)$$

in which, in the position representation,

$$H_1 = -\frac{\hbar^2}{2m_e} \nabla_1^2 - \frac{Ze^2}{r_1}, \quad H_2 = -\frac{\hbar^2}{2m_e} \nabla_2^2 - \frac{Ze^2}{r_2}, \quad (15.3.2)$$

$$H_{12} = \frac{e^2}{|\vec{r}_1 - \vec{r}_2|}, \quad (15.3.3)$$

where  $Z = 2$ , and  $\vec{r}_1$  and  $\vec{r}_2$  describe the position coordinates of the two electrons, and  $\nabla_1^2$  and  $\nabla_2^2$  are the corresponding Laplacians. If the electrons did not interact with each other, then  $H_1$  and  $H_2$  would just be the Hamiltonians for hydrogen-like atoms with  $Z = 2$ . The electron-electron interaction term  $H_{12}$  is the source of complications, and in this section we treat it as the perturbation  $W$ , although it is not particularly small.

The lowest energy eigenstate of the unperturbed Hamiltonian  $H_0 = H_1 + H_2$  is simply the tensor product of two hydrogen-like atomic ground states with  $(n, l, m) = (1, 0, 0)$ ,

$$|(1, 0, 0), (1, 0, 0)\rangle = |1, 0, 0\rangle_1 \otimes |1, 0, 0\rangle_2. \quad (15.3.4)$$

The position wavefunction for the unperturbed state is

$$\psi(\vec{r}_1, \vec{r}_2) = \langle \vec{r}_1 | 1, 0, 0 \rangle_1 \langle \vec{r}_2 | 1, 0, 0 \rangle_2 = \psi_{1,0,0}(\vec{r}_1) \psi_{1,0,0}(\vec{r}_2), \quad (15.3.5)$$

where, as we found in eq. (11.1.42),

$$\psi_{1,0,0}(\vec{r}) = \sqrt{\frac{Z^3}{\pi a_0^3}} e^{-Zr/a_0}. \quad (15.3.6)$$

---

<sup>†</sup>For simplicity, we ignore small effects due to spin interactions and the finite mass of the nucleus. Also, we ignore the fact that the electrons are identical; this issue turns out to not affect the following treatment of the ground state because the electron spins will be oriented oppositely. It does affect the excited states of helium, as we will discuss later, in section 18.3.



The corresponding unperturbed total energy eigenvalue for  $H_1 + H_2$  is, including the contributions of both electrons,

$$\mathcal{E}_0 = 2 \left( -\frac{Z^2 e^2}{2a_0} \right) = -8 \text{ Rydberg} = -108.85 \text{ eV}. \quad (15.3.7)$$

This turns out to be about 30% lower than the experimental value (the minimum energy needed to completely ionize the helium atom). The reason is that the electron-electron Coulomb repulsive potential  $H_{12}$  raises the energy, by an amount that we now estimate.

At first order in perturbation theory, the ground-state energy correction is

$$E^{(1)} = \langle (1, 0, 0), (1, 0, 0) | H_{12} | (1, 0, 0), (1, 0, 0) \rangle \quad (15.3.8)$$

$$= \int d^3 \vec{r}_1 \int d^3 \vec{r}_2 |\psi_{1,0,0}(\vec{r}_1)|^2 |\psi_{1,0,0}(\vec{r}_2)|^2 \frac{e^2}{|\vec{r}_1 - \vec{r}_2|}. \quad (15.3.9)$$

We therefore have

$$E^{(1)} = e^2 \left( \frac{Z^3}{\pi a_0^3} \right)^2 I, \quad (15.3.10)$$

with

$$I = \int_0^\infty dr_1 r_1^2 \int d\Omega_1 \int_0^\infty dr_2 r_2^2 \int d\Omega_2 e^{-2Zr_1/a_0} e^{-2Zr_2/a_0} \frac{1}{|\vec{r}_1 - \vec{r}_2|}. \quad (15.3.11)$$

Our challenge now is to evaluate the integral  $I$ . To do so, recall the identity [rewritten from eq. (8.6.82)]

$$\frac{1}{|\vec{r}_1 - \vec{r}_2|} = \sum_{l=0}^{\infty} \frac{4\pi}{2l+1} \frac{r_{\min}^l}{r_{\max}^{l+1}} \sum_{m=-l}^l Y_l^m(\theta_2, \phi_2)^* Y_l^m(\theta_1, \phi_1), \quad (15.3.12)$$

in which

$$r_{\max} = \max(r_1, r_2), \quad r_{\min} = \min(r_1, r_2). \quad (15.3.13)$$

Using eq. (15.3.12) in the integral  $I$ , we therefore have  $\int d\Omega_1 Y_l^m(\theta_1, \phi_1) = 0$  unless  $l = m = 0$ , and similarly for the  $d\Omega_2$  integral. Since only that one term survives, we can replace

$$\frac{1}{|\vec{r}_1 - \vec{r}_2|} \rightarrow \frac{4\pi}{r_{\max}} Y_0^0(\theta_2, \phi_2)^* Y_0^0(\theta_1, \phi_1) \delta_{l0} = \delta_{l0}/r_{\max}, \quad (15.3.14)$$

eliminating all of the angular dependences of the integrand. So, using  $\int d\Omega_1 = \int d\Omega_2 = 4\pi$ ,

$$I = (4\pi)^2 \int_0^\infty dr_1 r_1^2 \int_0^\infty dr_2 r_2^2 e^{-2Zr_1/a_0} e^{-2Zr_2/a_0} \frac{1}{r_{\max}}. \quad (15.3.15)$$

Separating the  $dr_2$  integral into two parts, according to whether  $r_2 < r_1$  or  $r_2 > r_1$ , now gives

$$I = (4\pi)^2 \int_0^\infty dr_1 \left( \int_0^{r_1} dr_2 r_1 r_2^2 + \int_{r_1}^\infty dr_2 r_1^2 r_2 \right) e^{-2Zr_1/a_0} e^{-2Zr_2/a_0}, \quad (15.3.16)$$

which can be done straightforwardly. The two parts turn out to contribute equally to the total,

$$I = \frac{5\pi^2 a_0^5}{8Z^5}. \quad (15.3.17)$$

Putting this into eq. (15.3.10), the first-order correction to the ground-state energy is

$$E^{(1)} = \frac{5Ze^2}{8a_0}. \quad (15.3.18)$$

Combining this with eq. (15.3.7), and plugging in  $Z = 2$ , the total ground-state energy for helium estimated from first-order perturbation theory is

$$E = \mathcal{E}_0 + E^{(1)} = (-8 + 5/2) \text{ Rydberg} = -74.83 \text{ eV}. \quad (15.3.19)$$

Note that the first-order correction is not suppressed by an adjustable small parameter compared to the unperturbed energy; it is  $5/16$  as large in magnitude. This might cause worry that perhaps the perturbative expansion is not converging fast enough to be trusted. Nevertheless, our final result differs by only about 5.3% from the experimental result of  $-79.01 \text{ eV}$ . In section 16.4, we will use a non-perturbative technique, the variational method, to do even better.

## 15.4 Brillouin–Wigner perturbation theory

The standard organization of stationary-state perturbation theory summarized in eqs. (15.1.39)–(15.1.47) has an alternative, called **Brillouin–Wigner perturbation theory** after Léon Brillouin and Wigner, which sometimes has better numerical convergence behavior. To derive it, we start with the eigenvalue problem in the form

$$(H_0 + W - E_n) |\psi_n\rangle = 0, \quad (15.4.1)$$

using the same notations as in section 15.1, but with  $\lambda = 1$  from the start. Now, act with the unperturbed bra  $\langle m|$ , then use  $\langle m|H_0 = \langle m|\mathcal{E}_m$ , and rearrange the result to get

$$\langle m|\psi_n\rangle = \frac{\langle m|W|\psi_n\rangle}{E_n - \mathcal{E}_m}. \quad (15.4.2)$$

Note the simultaneous presence in the denominator of the full energy eigenvalue  $E_n$  and the unperturbed energy  $\mathcal{E}_m$ . From completeness of the unperturbed states  $|m\rangle$ , we also have

$$|\psi_n\rangle = \sum_m |m\rangle \langle m|\psi_n\rangle = |n\rangle \langle n|\psi_n\rangle + \sum_{m \neq n} |m\rangle \langle m|\psi_n\rangle. \quad (15.4.3)$$

As in section 15.1.47, we choose the normalization condition  $\langle n|\psi_n\rangle = 1$ , and use eq. (15.4.2) in (15.4.3) to obtain

$$|\psi_n\rangle = |n\rangle + \sum_{m \neq n} |m\rangle \frac{\langle m|W|\psi_n\rangle}{E_n - \mathcal{E}_m}. \quad (15.4.4)$$

This has a straightforward iterative solution,

$$\begin{aligned} |\psi_n\rangle &= |n\rangle + \sum_{m \neq n} |m\rangle \frac{1}{E_n - \mathcal{E}_m} \langle m|W|n\rangle \\ &+ \sum_{m \neq n} \sum_{p \neq n} |m\rangle \frac{1}{E_n - \mathcal{E}_m} \langle m|W|p\rangle \frac{1}{E_n - \mathcal{E}_p} \langle p|W|n\rangle + \cdots, \end{aligned} \quad (15.4.5)$$

as can be checked by plugging it in. To find an equation for the corresponding energy eigenvalues, we now use eq. (15.1.38) to obtain

$$\begin{aligned} E_n &= \mathcal{E}_n + \langle n|W|n\rangle + \sum_{m \neq n} \langle n|W|m\rangle \frac{1}{E_n - \mathcal{E}_m} \langle m|W|n\rangle \\ &+ \sum_{m \neq n} \sum_{p \neq n} \langle n|W|m\rangle \frac{1}{E_n - \mathcal{E}_m} \langle m|W|p\rangle \frac{1}{E_n - \mathcal{E}_p} \langle p|W|n\rangle + \cdots. \end{aligned} \quad (15.4.6)$$

Equations (15.4.5) and (15.4.6) summarize Brillouin–Wigner perturbation theory. They have a simpler structure than the standard perturbation theory of section 15.1, but have the disadvantage that they are not yet a complete solution, since the full energy  $E_n$  appears on both sides of eq. (15.4.6).

At first order in Brillouin–Wigner perturbation theory, only the first two terms on the right-hand side of eq. (15.4.6) are included, and the solution for  $E_n$  is the same as in standard perturbation theory. At second order or beyond, the equation for  $E_n$  is a non-linear algebraic equation, and often can only be obtained numerically as an approximation. However, this can be a good thing, because the results so obtained are often more accurate than those found from the standard perturbation theory of section 15.1 at the same order. One way of thinking about this is that, at a given order in perturbation theory, the  $E_n$  appearing in the denominators of Brillouin–Wigner perturbation theory contain more complete information than the  $\mathcal{E}_n$  appearing in the denominators of standard perturbation theory. Once  $E_n$  has been obtained by solving eq.(15.4.6) truncated at a specific order in the expansion, it can be plugged into eq. (15.4.5) to obtain the corresponding energy eigenstate.

## 15.5 Dalgarno–Lewis method for simplifying perturbation theory

The formulas of stationary-state perturbation theory found in section 15.1 for the state, and for the energy beyond first order, contain sums over all of the unperturbed states. For example, the second-order correction to the energy contains the sum

$$E_n^{(2)} = \sum_{m \neq n} \frac{|\langle m|W|n\rangle|^2}{\mathcal{E}_n - \mathcal{E}_m} \quad (15.5.1)$$

While this is a well-defined formal solution, in practice it is sometimes hard to compute such sums directly. It is especially difficult if some of the unperturbed states form a continuum, in

which case the corresponding part of the summation will actually be an integration. Fortunately, there is an elegant method, due to Alexander Dalgarno and John T. Lewis, that allows these sums to be recast into a form that can often be solved exactly. Even when an exact solution cannot be obtained, the **Dalgarno–Lewis method** still allows for a systematic approximation, often by numerically solving a differential equation.

Suppose that, given  $|n\rangle$ , one can find an operator  $A$  that satisfies

$$[A, H_0] |n\rangle = (W - c) |n\rangle, \quad (15.5.2)$$

where  $H_0$  and  $W$  are the unperturbed and perturbation parts of the Hamiltonian, respectively, and  $c$  is a constant. Actually,  $c$  is not arbitrary, because after acting on both sides of eq. (15.5.2) with  $\langle n|$ , the left side vanishes, which implies  $c = \langle n|W|n\rangle = E_n^{(1)}$ . Also, since only the commutator of  $A$  appears in eq. (15.5.2), one can always add a constant multiple of the identity operator to it, to ensure that

$$\langle n|A|n\rangle = 0. \quad (15.5.3)$$

Note that the required operator  $A$  is different for each  $|n\rangle$ . Now, for every other unperturbed orthobasis state  $|m\rangle$ , eq. (15.5.2) implies  $\langle m|W|n\rangle = \langle m|[A, H_0]|n\rangle + c \langle m|n\rangle$ , or, since orthonormality of the  $H_0$  eigenstates eliminates the  $c$  term,

$$\langle m|W|n\rangle = (\mathcal{E}_n - \mathcal{E}_m) \langle m|A|n\rangle. \quad (15.5.4)$$

The Dalgarno–Lewis idea is to use eq. (15.5.4) to cancel the energy denominators in the results of perturbation theory, by expressing matrix elements of  $W$  in terms of matrix elements of  $A$ . So, if we can find  $A$ , the expressions in perturbation theory can then be simplified using the completeness relation for the unperturbed states.

For example, the first-order correction to the state is, from eq. (15.1.24),

$$|\psi_n^{(1)}\rangle = \sum_{m \neq n} |m\rangle \frac{\langle m|W|n\rangle}{\mathcal{E}_n - \mathcal{E}_m} = \sum_{m \neq n} |m\rangle \langle m|A|n\rangle = \sum_m |m\rangle \langle m|A|n\rangle. \quad (15.5.5)$$

The last equality relies on the fact that the additional term with  $m = n$  vanishes, because of eq. (15.5.3). Now, from the completeness relation  $\sum_m |m\rangle \langle m| = I$ , we conclude that the first-order correction to the state  $|n\rangle$  is simply

$$|\psi_n^{(1)}\rangle = A |n\rangle. \quad (15.5.6)$$

In words, the desired operator  $A$  turns the unperturbed state into the first-order correction to the state. Similarly, the second-order correction to the energy is, from eq. (15.5.1),

$$E_n^{(2)} = \sum_{m \neq n} \langle n|W|m\rangle \langle m|A|n\rangle = \sum_m \langle n|W|m\rangle \langle m|A|n\rangle = \langle n|WA|n\rangle, \quad (15.5.7)$$

where we have again used eq. (15.5.3) to account for the vanishing of the additional term with  $m = n$ , and then used completeness. As a check, this can be rewritten as

$$E_n^{(2)} = \langle n | W | \psi_n^{(1)} \rangle, \quad (15.5.8)$$

in accord with eq. (15.1.27) with  $k = 2$ . In a similar way, it is not too hard to show that even the third-order correction to the energy can be written as

$$E_n^{(3)} = \langle \psi_n^{(1)} | W | \psi_n^{(1)} \rangle - \langle \psi_n^{(1)} | \psi_n^{(1)} \rangle \langle n | W | n \rangle, \quad (15.5.9)$$

without energy-difference denominators.

Of course, this wonderful simplification does not come for free. To make use of the Dalgarno–Lewis method, one must either find the operator  $A$ , or equivalently (and more directly) solve for the ket  $A|n\rangle = |\psi_n^{(1)}\rangle$ . Rewriting eq. (15.5.2), the key equation to be solved is

$$(\mathcal{E}_n - H_0) |\psi_n^{(1)}\rangle = W|n\rangle - |n\rangle \langle n | W | n \rangle. \quad (15.5.10)$$

In favorable cases, this can be solved exactly for  $|\psi_n^{(1)}\rangle$  or the corresponding wavefunction; we will see an example of this in section 15.7 when we treat the ground state of the hydrogen atom in an electric field (the quadratic Stark effect). Even when this is not feasible, it is often possible to cast eq. (15.5.10) in the position representation as a differential equation that can be solved approximately, perhaps numerically. This is often faster and simpler than trying to do the sums over all other states  $|m\rangle$  with energy denominators  $\mathcal{E}_n - \mathcal{E}_m$ , especially if there are an infinite number of such states. The great advantage of the Dalgarno–Lewis method is that eq. (15.5.10) only involves the eigenvalues and matrix elements of the unperturbed state  $|n\rangle$  that one is considering. Once it has been solved to find  $|\psi_n^{(1)}\rangle$ , the second and third-order corrections to the energy follow immediately from eqs. (15.5.8) and (15.5.9), respectively.

## 15.6 Degenerate perturbation theory

As noted in section 15.1, a special problem arises when unperturbed states  $|n\rangle$  have degeneracies, because then the energy differences  $\mathcal{E}_{nm} = \mathcal{E}_n - \mathcal{E}_m$  appearing in the denominators of eqs. (15.1.44)–(15.1.47) will vanish. In this section, we show how to deal with this difficulty.

The bad-denominator problem is related to the fact that perturbation theory requires a smooth one-to-one correspondence between the unperturbed states and the perturbed states as the expansion parameter  $\lambda$  varies. Specifically, this is what allowed us to demand that the unperturbed state  $|n\rangle$  and the corrections to it  $|\psi_n^{(k)}\rangle$  are orthogonal, as we did in eq. (15.1.12). This in turn was used to exclude the terms in the sums that would otherwise have had vanishing energy difference denominators. In the degenerate case, the essential problem is that for a given




Figure 15.6.1: Perturbed energies as a function of the expansion parameter  $\lambda$ , for non-degenerate (left) and degenerate (right) perturbation theory. In degenerate perturbation theory, for each of the perturbed states  $|\psi_1\rangle$ ,  $|\psi_2\rangle$ ,  $|\psi_3\rangle$  there is an ambiguity in choosing the corresponding unperturbed states  $|1\rangle$ ,  $|2\rangle$ ,  $|3\rangle$ , since arbitrary linear combinations of them have the same  $H_0$  eigenvalue  $\mathcal{E}_1 = \mathcal{E}_2 = \mathcal{E}_3$ . This ambiguity is resolved (at least in part) by choosing the unperturbed states to diagonalize the perturbation Hamiltonian  $W$  within each subspace of degenerate  $\mathcal{E}_n$ .

perturbed state  $|\psi_n\rangle$  there is an ambiguity in deciding which of the degenerate unperturbed states  $|n\rangle$  corresponds to it and therefore can be excluded from the sums, since arbitrary linear combinations of them have the same  $H_0$  eigenvalue  $\mathcal{E}_n$ . This is illustrated in Figure 15.6.1, which compares the situations for non-degenerate and degenerate unperturbed states.

To resolve the problem, we need to answer the following question: if a group of unperturbed states  $\{|n\rangle\}$  have the same unperturbed energy  $\mathcal{E}_n$ , then which linear combinations of them are the  $|\psi_n^{(0)}\rangle$ ? The problem of avoiding bad (vanishing) denominators in perturbation theory is the same as the problem of choosing the correct linear combinations of unperturbed energy states.

Consider what happens when we return to eq. (15.1.23), rewritten here for convenience,

$$(\mathcal{E}_n - \mathcal{E}_m) \langle m | \psi_n^{(1)} \rangle = \langle m | W | n \rangle \quad (m \neq n), \quad (15.6.1)$$

but now allowing for the possibility that some unperturbed states have degenerate unperturbed energies. Whenever  $\mathcal{E}_m \neq \mathcal{E}_n$ , we can still solve

$$\langle m | \psi_n^{(1)} \rangle = \frac{W_{mn}}{\mathcal{E}_n - \mathcal{E}_m}, \quad (\mathcal{E}_m \neq \mathcal{E}_n). \quad (15.6.2)$$

However, for  $\mathcal{E}_m = \mathcal{E}_n$ , we instead learn only that  $\langle m | W | n \rangle$  must vanish for  $m \neq n$ , in order to have a consistent perturbative solution. It is therefore required that the matrix elements of  $W$  form a diagonal matrix when restricted to each subspace with degenerate unperturbed energies,

$$\langle m | W | n \rangle = \delta_{nm} W_{nn} \quad (\text{for } \mathcal{E}_m = \mathcal{E}_n). \quad (15.6.3)$$

Fortunately, this is always possible to arrange by a suitable choice of the orthobasis of stationary states  $|n\rangle$ . This is assured by Theorem 2.6.6, since  $W$  is Hermitian.

Equation 15.6.3 is a necessary (but not always sufficient, as we will see) requirement on the choice of unperturbed states. The point is that by choosing a “good” basis of unperturbed states in this way, the perturbation Hamiltonian matrix  $W_{mn}$  does not connect pairs of distinct unperturbed states for which the energy-difference denominators would vanish.

In practice, one must solve the eigenvalue problem for  $g_{\mathcal{E}_n} \times g_{\mathcal{E}_n}$  matrices, formed by the matrix elements of the operator  $W$  restricted to each  $g_{\mathcal{E}_n}$ -dimensional degenerate subspace. Once we choose the states  $|n\rangle$  as the resulting orthonormal eigenvectors, then it follows from eq. (15.1.19) that the eigenvalues are the corresponding first-order corrections to the energies,

$$E_n^{(1)} = W_{nn}, \quad (15.6.4)$$

the same result as for the non-degenerate case. Note that it is only necessary to diagonalize  $W$  within the subspaces for degenerate  $\mathcal{E}_n$ . In particular, for each unperturbed state  $|n\rangle$  that is non-degenerate, there is nothing to do;  $W$  restricted to that subspace is simply a  $1 \times 1$  matrix, so  $|n\rangle$  is automatically an eigenvector, with eigenvalue  $W_{nn} = \langle n|W|n\rangle$ .

Often, only the first-order energy corrections are required; then eq. (15.6.4) is sufficient provided that the unperturbed energy basis has been chosen to satisfy eq. (15.6.3), and in that case you can skip the complications of the next few paragraphs, from here to eq. (15.6.15).

If one needs the first-order corrections to the state ket  $|\psi_n^{(1)}\rangle$  and the second-order corrections to the energies  $E_n^{(2)}$ , then one must confront the fact that eq. (15.6.1) provides no information about  $\langle m|\psi_n^{(1)}\rangle$  for  $\mathcal{E}_m = \mathcal{E}_n$ . By completeness,

$$|\psi_n^{(1)}\rangle = \sum'_m |m\rangle \frac{W_{mn}}{\mathcal{E}_{nm}} + \sum''_m |m\rangle \langle m|\psi_n^{(1)}\rangle, \quad (15.6.5)$$

where we have introduced the notations, always for fixed  $n$  = the state being perturbed,

$$\sum'_m = \text{sum over all } m \text{ such that } \mathcal{E}_m \neq \mathcal{E}_n, \quad (15.6.6)$$

$$\sum''_m = \text{sum over all } m \text{ such that } m \neq n \text{ and } \mathcal{E}_m = \mathcal{E}_n, \quad (15.6.7)$$

and used eq. (15.6.2) in the first summation. To solve for the remaining matrix elements  $\langle m|\psi_n^{(1)}\rangle$  appearing in the last summation, one must use the equations governing the perturbative expansion at order  $\lambda^2$ , and possibly beyond.

To see how this goes, consider eq. (15.1.29) with  $k = 2$  and  $\mathcal{E}_m = \mathcal{E}_n$ , which yields

$$E_n^{(1)} \langle m|\psi_n^{(1)}\rangle = \langle m|W|\psi_n^{(1)}\rangle, \quad (\text{for } m \neq n \text{ and } \mathcal{E}_m = \mathcal{E}_n). \quad (15.6.8)$$

After using  $E_n^{(1)} = W_{nn}$ , and eq. (15.6.5) with  $m$  replaced by  $p$ , and defining an operator  $W^{(2)}$  with matrix elements

$$W_{mn}^{(2)} \equiv \sum_p' \frac{W_{mp}W_{pn}}{\mathcal{E}_{np}}, \quad (15.6.9)$$

eq. (15.6.8) becomes

$$W_{nn} \langle m | \psi_n^{(1)} \rangle = W_{mn}^{(2)} + \sum_p'' W_{mp} \langle p | \psi_n^{(1)} \rangle, \quad (\text{for } m \neq n \text{ and } \mathcal{E}_m = \mathcal{E}_n). \quad (15.6.10)$$

In the last summation, we know that  $W_{mp} = \delta_{mp}W_{mm}$ , because of eq. (15.6.3). Therefore, only the single term with  $p = m$  in that sum survives, and eq. (15.6.10) reduces to

$$(W_{nn} - W_{mm}) \langle m | \psi_n^{(1)} \rangle = W_{mn}^{(2)}, \quad (\text{for } m \neq n \text{ and } \mathcal{E}_m = \mathcal{E}_n). \quad (15.6.11)$$

Now, for states that are non-degenerate at first order, so that  $W_{mm} \neq W_{nn}$ , we obtain

$$\langle m | \psi_n^{(1)} \rangle = \frac{W_{mn}^{(2)}}{W_{nn} - W_{mm}} \quad (\text{for } \mathcal{E}_m = \mathcal{E}_n \text{ and } W_{mm} \neq W_{nn}), \quad (15.6.12)$$

but in the opposite case we learn only that

$$W_{mn}^{(2)} = 0, \quad (\text{for } m \neq n \text{ and } \mathcal{E}_m = \mathcal{E}_n \text{ and } W_{mm} = W_{nn}). \quad (15.6.13)$$

This is a second condition that must be imposed on the choice of the unperturbed states, in addition to eq. (15.6.3).

In words, we have found that we must choose the unperturbed kets  $|n\rangle$  so that on subspaces where both  $H_0$  and  $W$  are proportional to the identity, the matrix  $W^{(2)}$  is also diagonal. Fortunately, this is always possible, because  $W^{(2)}$  is a Hermitian matrix when restricted<sup>†</sup> to this subspace. Now, the second-order corrections to the energy eigenvalues are obtained from eq. (15.1.27), and are simply

$$E_n^{(2)} = W_{nn}^{(2)} = \sum_m' \frac{|W_{mn}|^2}{\mathcal{E}_{nm}}, \quad (15.6.14)$$

where it is crucial that the unperturbed states satisfy the conditions given in both eqs. (15.6.3) and (15.6.13), and there is no vanishing denominator problem because of the fact that the sum excludes all terms with  $\mathcal{E}_m = \mathcal{E}_n$ .

There may still be some ambiguity in the matrix elements  $\langle m | \psi_n^{(1)} \rangle$ , if for some  $m \neq n$  one has  $\mathcal{E}_m = \mathcal{E}_n$  and  $W_{mm} = W_{nn}$  and  $W_{mn}^{(2)} = W_{nn}^{(2)}$ , so that the degeneracy is not completely lifted

---

<sup>†</sup>The operator  $W^{(2)}$  is not Hermitian on the whole state space, because  $n$  and  $m$  are not treated symmetrically in the definition eq. (15.6.9). However, it is Hermitian within the subspace of matrix elements with  $\mathcal{E}_m = \mathcal{E}_n$ , for which  $W_{mn}^{(2)*} = W_{nm}^{(2)}$ .



even at second order. In that case, one must use the information from third order, that is,  $k = 3$  in eq. (15.1.17). One then finds a third condition on the choice of unperturbed states  $|n\rangle$ , that the operator  $W^{(3)}$  with matrix elements

$$W_{mn}^{(3)} \equiv \sum_q' \sum_p' \frac{W_{mq}W_{qp}W_{pn}}{\mathcal{E}_{nq}\mathcal{E}_{np}} - W_{nn} \sum_p' \frac{W_{mp}W_{pn}}{\mathcal{E}_{np}^2} \quad (15.6.15)$$

is diagonal on each subspace of states where the operators  $H_0$ ,  $W$ , and  $W^{(2)}$  are proportional to the identity operator. If the degeneracy of unperturbed states is still not removed at third order (that is,  $W^{(3)}$  has degenerate eigenvalues when restricted to that subspace of states), one must continue to  $k = 4$  in eq. (15.1.17), etc. These considerations from  $k \geq 3$  affect the determination of  $|\psi_n^{(1)}\rangle$  (through  $\langle m|\psi_n^{(1)}\rangle$ ) even though they seemingly involve higher orders in perturbation theory, but they do not affect the determination of  $E_n^{(2)}$  given by eq. (15.6.14).

Practical applications of degenerate perturbation theory often only need  $E_n^{(1)}$  as given in eq. (15.6.4), and therefore one only needs to worry about choosing the unperturbed states to satisfy eq. (15.6.3). Examples will appear in sections 15.7 and 15.8 and in Chapter 17.

To conclude this section, we exploit first-order perturbation theory to infer the exact matrix element  $\langle 1/R^2 \rangle$  for the states  $|n, l, m\rangle$  of the hydrogen atom, fulfilling a promise made in section 11.1. Consider the Hamiltonian  $H_\lambda = H_0 + \lambda/R^2$ , where  $H_0 = P^2/2m_e - e^2/R$  is the unperturbed hydrogen atom Hamiltonian, and  $\lambda$  is a small parameter. In this case, the operator  $1/R^2$  commutes with the angular momentum operators  $L^2$  and  $L_z$ , so it is diagonal in the orthobasis of states  $|n, l, m\rangle$  for fixed  $n$  and different  $l, m$ . Therefore  $|n, l, m\rangle$  is a good basis for degenerate first-order perturbation theory in  $\lambda$ , and the first-order correction to the energy eigenvalue for each state is simply proportional to the expectation value that we seek,

$$E_{n,l,m}^{(1)} = \lambda \langle 1/R^2 \rangle_{n,l,m}. \quad (15.6.16)$$

Meanwhile, the exact energy eigenvalues  $E_{\lambda,k,l,m}$  of  $H_\lambda$  are the solutions to the radial wavefunction eigenvalue problem

$$\left[ -\frac{\hbar^2}{2m_e} \left( \frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} - \frac{l(l+1)}{r^2} \right) + \frac{\lambda}{r^2} \right] R_{k,l} = E_{\lambda,k,l,m} R_{k,l}, \quad (15.6.17)$$

where the integer  $k$  is the radial quantum number (the number of zeros of  $R_{k,l}(r)$  for  $0 < r < \infty$ ). Now we can define a quantity  $l'$  by

$$\frac{\hbar^2}{2m_e} l'(l'+1) = \frac{\hbar^2}{2m_e} l(l+1) + \lambda, \quad (15.6.18)$$

or, equivalently,

$$l' = \sqrt{(l+1/2)^2 + 2m_e\lambda/\hbar^2} - 1/2. \quad (15.6.19)$$

Note that  $l'$  need not be an integer, unlike  $l$ . Nevertheless, the analysis of the radial wavefunction eigenvalue problem that led to eq. (11.1.16) goes through, but with  $l$  replaced everywhere by  $l'$ , and  $n$  replaced by  $k + l' + 1$ . In particular, it is crucial that  $k$  is a non-negative integer, in order that the wavefunction is normalizable. The result is

$$E_{\lambda,k,l,m} = -\frac{e^2}{2a_0(k+l'+1)^2}. \quad (15.6.20)$$

Expanding in small  $\lambda$ , for fixed integers  $k$  and  $l$ ,

$$E_\lambda = E_{\lambda=0} + \lambda \left. \frac{dE_\lambda}{d\lambda} \right|_{\lambda=0} + \dots, \quad (15.6.21)$$

and by comparing to eq. (15.6.16), we find

$$\langle 1/R^2 \rangle_{n,l,m} = \frac{d}{d\lambda} \left( \frac{-e^2}{2a_0(k+l'+1)^2} \right) \Big|_{\lambda=0} = \frac{e^2}{a_0(k+l+1)^3} \left. \frac{dl'}{d\lambda} \right|_{\lambda=0}. \quad (15.6.22)$$

From eq. (15.6.19), we have

$$\left. \frac{dl'}{d\lambda} \right|_{\lambda=0} = \frac{m_e}{\hbar^2(l+1/2)}. \quad (15.6.23)$$

Plugging eq. (15.6.23) into eq. (15.6.22), and using  $n = k + l + 1$ , we obtain

$$\langle 1/R^2 \rangle_{n,l,m} = \frac{m_e e^2}{\hbar^2 a_0 n^3 (l+1/2)} = \frac{1}{a_0^2 n^3 (l+1/2)}. \quad (15.6.24)$$

Equation (11.1.67) follows by restoring the general nuclear charge  $Z$ , using the rule  $a_0 \rightarrow a_0/Z$ .

The same sort of perturbation theory trick can be used to check the result we found for  $\langle 1/R \rangle_{n,l,m}$  in eq. (11.1.64). Consider  $H_Z = P^2/2m_e - Ze^2/R$ , the Hamiltonian for general  $Z$ . We rewrite this as  $H_Z = H_0 + W$ , where  $H_0 = P^2/2m_e - e^2/R$  is the  $Z = 1$  Hamiltonian, and  $W = (1 - Z)e^2/R$  is treated as a perturbation by taking  $Z$  to be close to 1. Then, for an eigenstate of  $H_0$  with principal quantum number  $n$ , the first-order energy correction is

$$E_n^{(1)} = (1 - Z)e^2 \langle 1/R \rangle_{n,l,m}, \quad (15.6.25)$$

which is proportional to the expectation value that we seek. (Again we are making use of the fact that the basis  $|n, l, m\rangle$  is a good one for first-order degenerate perturbation theory, in the sense that the perturbation  $1/R$  is diagonal for fixed  $n$  and different  $l, m$ .) On the other hand, the exact energies were found in eq. (11.1.16) to be

$$E_n = -\frac{Z^2 e^2}{2a_0 n^2} = -\frac{e^2}{2a_0 n^2} [1 + 2(Z - 1) + (Z - 1)^2]. \quad (15.6.26)$$

Comparing the terms linear in small  $Z - 1$  in eqs. (15.6.25) and (15.6.26), we must have

$$\langle 1/R \rangle_{n,l,m} = \frac{1}{a_0 n^2}. \quad (15.6.27)$$

This confirms eq. (11.1.64) with  $Z = 1$ .

## 15.7 Hydrogen atom in electric field (Stark effect)

Consider a hydrogen atom in a static external electric field  $\vec{E} = E\hat{z}$ . The corresponding electrostatic potential seen by the electron is  $\Phi(\vec{r}) = -Ez$ , giving a potential energy contribution to the Hamiltonian,

$$W = -e\Phi = eEZ, \quad (15.7.1)$$

which we will treat as a perturbation. The calculation of shifts in the atomic energy levels, known as the Stark effect, provides a nice example of both non-degenerate second-order and degenerate first-order perturbation theory.

For the ground state  $|n, l, m\rangle = |1, 0, 0\rangle$ , the first-order correction to the energy vanishes,

$$E_{1,0,0}^{(1)} = \langle 1, 0, 0 | W | 1, 0, 0 \rangle = eE \langle 1, 0, 0 | Z | 1, 0, 0 \rangle = 0. \quad (15.7.2)$$

This can be understood as a consequence of the parity selection rule, eq. (5.4.16). More generally, the expectation value of  $Z$  in any parity eigenstate is always 0, because  $\Pi Z \Pi = -Z$ , so that  $\pi_Z = -1$ , giving  $\pi_Z \pi_{n,l,m}^2 = -1$ . Another way to see that the matrix element vanishes is the dipole selection rule of eq. (13.2.15).

Since the first-order correction to the ground-state energy is 0, we turn to the second-order correction. Applying the general formula of eq. (15.1.44), and remembering that the unperturbed energies do not depend on  $l$  and  $m$ , we have

$$E_{1,0,0}^{(2)} = e^2 E^2 \sum_{n=2}^{\infty} \sum_{l=0}^{n-1} \sum_{m=-l}^l \frac{|\langle n, l, m | Z | 1, 0, 0 \rangle|^2}{\mathcal{E}_1 - \mathcal{E}_n} + e^2 E^2 \int_0^{\infty} d\mathcal{E} \sum_{l=0}^{\infty} \sum_{m=-l}^l \frac{|\langle \mathcal{E}, l, m | Z | 1, 0, 0 \rangle|^2}{\mathcal{E}_1 - \mathcal{E}}. \quad (15.7.3)$$

Note that this includes integration over the unbound states  $|\mathcal{E}, l, m\rangle$  with continuous unperturbed energies  $\mathcal{E} > 0$  as well as summation over the bound states  $|n, l, m\rangle$  with discrete unperturbed energies  $\mathcal{E}_n = -e^2/2a_0n^2$ , excluding  $n = 1$ . Equation (15.7.3) simplifies considerably if we use the dipole selection rules of eqs. (13.2.15) and (13.2.16), which inform us that only the  $l = 1$  and  $m = 0$  matrix elements are non-zero. Therefore, it can be rewritten as

$$E_{1,0,0}^{(2)} = -e^2 E^2 \left( \sum_{n=2}^{\infty} \frac{|\langle n, 1, 0 | Z | 1, 0, 0 \rangle|^2}{\mathcal{E}_n - \mathcal{E}_1} + \int_0^{\infty} d\mathcal{E} \frac{|\langle \mathcal{E}, 1, 0 | Z | 1, 0, 0 \rangle|^2}{\mathcal{E} - \mathcal{E}_1} \right), \quad (15.7.4)$$

where the overall minus sign appears because the denominators have flipped sign to make them positive. Since both the sum and the integral are clearly positive, eq. (15.7.4) is negative; the presence of the electric field lowers the hydrogen atom energy.

The evaluation of eq. (15.7.4) is not trivial, so we will study it using a series of tricks with instructive value. First, we will find a simple lower bound on its magnitude, then a simple upper

bound, and then attempt a numerical approximation by including only the bound state sum contribution. Finally, we will obtain the exact value of  $E_{1,0,0}^{(2)}$  by exploiting the power of the Dalgarno–Lewis method of section 15.5.

To obtain the lower bound, note that the magnitude of eq. (15.7.4) is certainly greater than that of the  $n = 2$  term alone, since all of the contributions have the same sign. So, using  $\mathcal{E}_2 - \mathcal{E}_1 = 3e^2/8a_0$ ,

$$|E_{1,0,0}^{(2)}| > (n=2 \text{ term only}) = \frac{8}{3}a_0E^2|\langle 2, 1, 0|Z|1, 0, 0\rangle|^2. \quad (15.7.5)$$

The relevant matrix element is

$$\langle 2, 1, 0|Z|1, 0, 0\rangle = \int_0^{2\pi} d\phi \int_{-1}^1 d(\cos\theta) \int_0^\infty dr r^2 (R_{21}Y_1^0)^* r \cos\theta (R_{10}Y_0^0) = \frac{128\sqrt{2}}{243}a_0. \quad (15.7.6)$$

Therefore, we have

$$|E_{1,0,0}^{(2)}| > \frac{2^{18}}{3^{11}}E^2a_0^3 \approx 1.4798E^2a_0^3 \quad (15.7.7)$$

as a lower bound on the energy shift of the ground state due to the external electric field  $E$ .

We also obtain a bound from the other direction, by noting that the denominators in eq. (15.7.4) with  $n \neq 2$  are all larger than they would be if each instance of  $\mathcal{E}_n$  or  $\mathcal{E}$  were replaced by  $\mathcal{E}_2$ . So,

$$|E_{1,0,0}^{(2)}| < \frac{8}{3}a_0E^2 \left( \sum_{n=2}^\infty |\langle n, 1, 0|Z|1, 0, 0\rangle|^2 + \int_0^\infty d\mathcal{E} |\langle \mathcal{E}, 1, 0|Z|1, 0, 0\rangle|^2 \right). \quad (15.7.8)$$

Now that the annoying energy-difference denominators are gone, we can evaluate the whole quantity in parentheses using the completeness identity of eq. (11.2.22), remembering that the missing  $l \neq 1$  and  $m \neq 0$  terms vanish by the dipole selection rules. The result is

$$|E_{1,0,0}^{(2)}| < \frac{8}{3}a_0E^2 \langle 1, 0, 0|Z^2|1, 0, 0\rangle. \quad (15.7.9)$$

Another trick: by rotational symmetry of the ground state, we can replace  $Z^2$  in the expectation value by the average of  $X^2$ ,  $Y^2$ , and  $Z^2$ , which is easier to compute. So

$$\langle 1, 0, 0|Z^2|1, 0, 0\rangle = \frac{1}{3}\langle 1, 0, 0|(X^2 + Y^2 + Z^2)|1, 0, 0\rangle = \frac{1}{3}\langle 1, 0, 0|R^2|1, 0, 0\rangle = a_0^2, \quad (15.7.10)$$

where, at the end, we have used eq. (11.1.53) with  $p = 2$ . Thus we have

$$|E_{1,0,0}^{(2)}| < \frac{8}{3}E^2a_0^3 \approx 2.6667a_0^3E^2 \quad (15.7.11)$$

as an upper bound on the energy shift to complement the lower bound in eq. (15.7.7).

As an attempt to do better than the bounds in eqs. (15.7.7) and (15.7.11), one can add up the exact contributions from all bound states  $n = 2, 3, 4, \dots$  with  $\mathcal{E}_n = -e^2/2a_0n^2$  in eq. (15.7.4). It is possible, but quite non-trivial, to show that

$$e^2 \frac{|\langle n, 1, 0 | Z | 1, 0, 0 \rangle|^2}{\mathcal{E}_n - \mathcal{E}_1} = \frac{(2n)^9 (n-1)^{2n-6}}{3(n+1)^{2n+6}} a_0^3. \quad (15.7.12)$$

The sum over all integers  $n \geq 2$  in eq. (15.7.4) is then found to converge to a numerical value

$$E_{1,0,0}^{(2), \text{partial}} \approx -1.8316 E^2 a_0^3, \quad (15.7.13)$$

which is indeed between the two bounds that we have already obtained. However, this turns out to be numerically far from the true answer; as we are about to discover, the integral over unbound states contributes significantly.

Fortunately, one can obtain the correct result while avoiding having to perform directly the difficult sum plus integration, by instead using the Dalgarno–Lewis method described in section 15.5. To do this, we must first use eq. (15.5.10) to solve for the first-order correction to the ground-state wavefunction,  $\psi_{1,0,0}^{(1)}(\vec{r})$ . In the position representation, this equation is

$$(\mathcal{E}_1 - H_0)\psi_{1,0,0}^{(1)} = eE r \cos \theta \psi_{1,0,0}^{(0)}, \quad (15.7.14)$$

where  $\mathcal{E}_1 = -e^2/2a_0$  and  $\psi_{1,0,0}^{(0)} = e^{-r/a_0}/\sqrt{\pi a_0^3}$  from the results of section 11.1, and we have used the fact that  $W = eEZ$  has vanishing expectation value in the unperturbed ground state. The unperturbed Hamiltonian operator is

$$H_0 = -\frac{\hbar^2}{2m_e} \nabla^2 - \frac{e^2}{r} = -\frac{a_0 e^2}{2} \left[ \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial}{\partial \theta} \right) \right] - \frac{e^2}{r}, \quad (15.7.15)$$

where, in the second equality, we have used the fact that there is no  $\phi$  dependence because of the symmetry of the problem about the  $z$ -axis, and  $\hbar^2/2m_e = a_0 e^2/2$ . Given the form of eq. (15.7.14), it is reasonable to try for a solution of the form

$$\psi_{1,0,0}^{(1)} = f(r) \cos \theta \psi_{1,0,0}^{(0)}, \quad (15.7.16)$$

where  $f(r)$  is to be determined. With this guess, the differential equation (15.7.14) becomes

$$\frac{1}{2} f'' + \left( \frac{1}{r} - \frac{1}{a_0} \right) f' - \frac{1}{r^2} f = \frac{E}{e a_0} r, \quad (15.7.17)$$

in which we are delighted to see that the  $\theta$  dependence has disappeared. Since the inhomogeneous part is linear in  $r$ , we are inspired to make the further guess that  $f$  is a quadratic polynomial in  $r$ , and are rewarded with the simple solution

$$f = -\frac{E}{e} (a_0 + r/2) r. \quad (15.7.18)$$

[This corresponds to the Dalgarno–Lewis operator  $A = -\frac{E}{e}(a_0 + R/2)Z$  as the solution to eqs. (15.5.2) and (15.5.3), although we do not directly need it.]

Having successfully found  $\psi_{1,0,0}^{(1)}$ , the rest is relatively easy. We get, using eq. (15.5.8) in the position representation,

$$E_{1,0,0}^{(2)} = \int_0^{2\pi} d\phi \int_{-1}^1 d(\cos \theta) \int_0^\infty dr r^2 (eEr \cos \theta) \left( -\frac{E}{e}(a_0 + r/2)r \cos \theta \right) |\psi_{1,0,0}^{(0)}|^2, \quad (15.7.19)$$

which evaluates finally to the correct, and surprisingly simple, result

$$E_{1,0,0}^{(2)} = -\frac{9}{4}a_0^3 E^2 = -2.25a_0^3 E^2. \quad (15.7.20)$$

Because the leading correction to the ground state energy comes from second order in perturbation theory, and is therefore quadratic in the applied electric field, this is called the **quadratic Stark effect**.

Let us now see how to connect eq. (15.7.20) to an experimental measurement. In general, for a dipole moment  $\vec{p}$  induced by an external electric field  $\vec{E}$ , the polarizability  $\alpha$  is defined by

$$\vec{p} = \alpha \vec{E}. \quad (15.7.21)$$

As the electric field is increased, the change in the energy of the dipole is

$$dU = -\vec{p} \cdot d\vec{E} = -\alpha E dE, \quad (15.7.22)$$

so that the total energy of the induced dipole is

$$U = -\frac{1}{2}\alpha E^2. \quad (15.7.23)$$

Comparing to eq. (15.7.20), we have  $-\frac{9}{4}a_0^3 E^2 = -\frac{1}{2}\alpha E^2$ , or

$$\alpha = \frac{9}{2}a_0^3 = 6.67 \times 10^{-31} \text{ m}^3, \quad (15.7.24)$$

which agrees well with the experimental value for the polarizability of atomic hydrogen in the ground state.

Let us now turn to the Stark effect for the first excited ( $n = 2$ ) stationary states of the hydrogen atom. The unperturbed states are

$$|n, l, m\rangle = |2, 0, 0\rangle, |2, 1, 0\rangle, |2, 1, 1\rangle, \text{ and } |2, 1, -1\rangle, \quad (15.7.25)$$

which all have the same unperturbed energy  $\mathcal{E}_2 = -e^2/8a_0$ . We therefore need to apply the method of degenerate perturbation theory as outlined in section 15.6. To start, we must choose the unperturbed orthobasis of  $n = 2$  states as the eigenkets of the  $4 \times 4$  matrix

$$W_{(l',m'),(l,m)} = \langle 2, l', m' | eEZ | 2, l, m \rangle. \quad (15.7.26)$$

Using the dipole selection rule of eqs. (13.2.15) and (13.2.16), the only non-zero entries of this matrix come from

$$\langle 2, 1, 0 | Z | 2, 0, 0 \rangle = \langle 2, 0, 0 | Z | 2, 1, 0 \rangle, \quad (15.7.27)$$

which can be evaluated as

$$\langle 2, 1, 0 | Z | 2, 0, 0 \rangle = \int_0^{2\pi} d\phi \int_{-1}^1 d(\cos \theta) \int_0^\infty dr r^2 (R_{21}Y_1^0)^* (r \cos \theta) (R_{20}Y_0^0) \quad (15.7.28)$$

$$= \frac{1}{16} \int_{-1}^1 d(\cos \theta) \cos^2 \theta \int_0^\infty dr \left( \frac{r}{a_0} \right)^4 (2 - r/a_0) e^{-r/a_0} \quad (15.7.29)$$

$$= -3a_0. \quad (15.7.30)$$

So, in the orthobasis of eq. (15.7.25), the perturbation Hamiltonian has the matrix representation

$$W \leftrightarrow -3a_0eE \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad (15.7.31)$$

which has eigenvalues  $-3a_0eE$ ,  $+3a_0eE$ ,  $0$ , and  $0$ . Thus, the energy eigenvalues to first order in the perturbing electric field and the corresponding unperturbed energy states are

$$\begin{array}{cc} \text{Energies} & \text{States} \end{array} \quad (15.7.32)$$

$$-\frac{e^2}{8a_0} - 3a_0eE \quad \frac{1}{\sqrt{2}} (|2, 0, 0\rangle + |2, 1, 0\rangle), \quad (15.7.33)$$

$$-\frac{e^2}{8a_0} + 3a_0eE \quad \frac{1}{\sqrt{2}} (|2, 0, 0\rangle - |2, 1, 0\rangle), \quad (15.7.34)$$

$$-\frac{e^2}{8a_0} \quad |2, 1, 1\rangle, \quad |2, 1, -1\rangle. \quad (15.7.35)$$

Because the perturbed  $n = 2$  energy levels vary linearly with the applied electric field, this is called the **linear Stark effect**.

Now that we have found the leading results in perturbation theory for the  $n = 1$  and  $n = 2$  energy levels, it is time to confess the awful truth that in this example, the perturbative expansion does not converge! The perturbation Hamiltonian  $W = eEZ$  is negative with an arbitrarily large magnitude; it approaches  $-\infty$  for  $z \rightarrow -\infty$ . This implies that in the idealized problem of a hydrogen atom in a uniform electric field filling infinite space, there are no bound states at all. Instead, the exact stationary-state solutions will have only continuous energies, with resonances near the bound state energies that we have obtained. For this reason, the perturbative expansion in powers of the electric field  $E$  actually cannot converge to the supposed atomic bound states.

Fortunately, however, in any real-world version of the Stark effect, the electric field is cut off at some large distance, and the perturbation Hamiltonian remains finite. So, our effort was not at all wasted. In fact, the practically relevant results really are the perturbative ones we found, not the “exact” calculation for unbound states based on an unattainable ideal.

## 15.8 Almost-degenerate perturbation theory

The perturbative expansion can also face problems if the perturbed state is almost, but not quite, degenerate with some other states. This means that for some subset of stationary states  $|n_1\rangle, |n_2\rangle, \dots |n_N\rangle$ , the differences in unperturbed energies  $\mathcal{E}_{n_j} - \mathcal{E}_{n_k}$  are all small compared to the magnitudes of the matrix elements of  $W$ . In this case, the terms in the perturbation expansions for  $E_{n_j}$  and  $|\psi_{n_j}\rangle$  will be finite, but will not become smaller at higher orders, and convergence will fail. To get sensible results we must formulate **almost-degenerate perturbation theory** to evaluate them, as follows.

For inspiration, we write the perturbation Hamiltonian as

$$W = \sum_n \sum_m |n\rangle \langle n| W |m\rangle \langle m|, \quad (15.8.1)$$

by using completeness twice. Consider a projection operator onto the subspace of almost-degenerate states that we wish to consider,

$$P_{\text{deg}} = \sum_k |n_k\rangle \langle n_k|. \quad (15.8.2)$$

Now we define the perturbation Hamiltonian projected onto the almost-degenerate subspace,

$$W_{\text{deg}} = P_{\text{deg}} W P_{\text{deg}}, \quad (15.8.3)$$

which is equivalent to only including the orthobasis elements for the degenerate subspace in the summations in eq. (15.8.1). Note that  $P_{\text{deg}} W_{\text{deg}} P_{\text{deg}} = W_{\text{deg}}$ , since  $P_{\text{deg}}^2 = P_{\text{deg}}$ , as always for projection operators.

The strategy is to now re-divide the full Hamiltonian as

$$H = H'_0 + W', \quad (15.8.4)$$

where

$$H'_0 = H_0 + W_{\text{deg}}, \quad (15.8.5)$$

$$W' = W - W_{\text{deg}}. \quad (15.8.6)$$

One now endeavors to solve the eigenvalue problem for  $H'_0$  exactly, and then treat  $W'$  as the perturbation. In favorable circumstances, for example if the dimension of the almost-degenerate



subspace is small, this can be done exactly. In less favorable cases, one might have to solve the eigenvalue problem for  $H'_0$  in an approximation, before then applying  $W'$  as the perturbation. In any case, let us refer to the unperturbed eigenstates of  $H'_0$  within the almost-degenerate subspace as  $|n'_j\rangle$ , with unperturbed energies  $\mathcal{E}_{n'_j}$ . The point is that all of the  $W'$  matrix elements within the almost-degenerate subspace vanish,

$$\langle n'_j | W' | n'_k \rangle = 0. \quad (15.8.7)$$

This follows because the left side can be rewritten, using  $|n'_k\rangle = P_{\text{deg}}|n'_k\rangle$ , as

$$\langle n'_j | P_{\text{deg}}(W - W_{\text{deg}})P_{\text{deg}} | n'_k \rangle = \langle n'_j | (W_{\text{deg}} - W_{\text{deg}}) | n'_k \rangle = 0. \quad (15.8.8)$$

One can now fearlessly apply the usual non-degenerate perturbation theory results of eqs. (15.1.43)–(15.1.47) with  $H'_0$  as the unperturbed Hamiltonian with eigenstates  $|n'_j\rangle$ , and  $W'$  as the perturbation, because small denominators  $\mathcal{E}_{n'_j} - \mathcal{E}_{n'_k}$  can never appear in the sums.

As a minimal example of all three types of stationary-state perturbation theory (non-degenerate, degenerate, almost-degenerate), consider a system with only 2 states, with

$$H_0 = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}, \quad W = \begin{pmatrix} 0 & c \\ c^* & 0 \end{pmatrix}, \quad (15.8.9)$$

where  $a, b$  are real numbers and  $c$  is complex. Of course,  $2 \times 2$  matrices are easy, so the eigenvalues for  $H = H_0 + W$  can be done exactly in this example, with the results

$$E_{1,2} = \frac{1}{2} \left( a + b \mp \sqrt{(a - b)^2 + 4|c|^2} \right). \quad (15.8.10)$$

If one applies non-degenerate perturbation theory, one readily obtains (assuming that  $b > a$ )

$$E_1 = a - |c|^2/(b - a) + \cdots, \quad E_2 = b + |c|^2/(b - a) + \cdots, \quad (15.8.11)$$

with the perturbation  $c$  contributing quadratically. On the other hand, if  $b = a$ , then the previous expansion fails, but one can apply degenerate perturbation theory, with the result

$$E_1 = a - |c| + \cdots, \quad E_2 = a + |c| + \cdots, \quad (15.8.12)$$

in which the perturbation  $c$  contributes linearly. In this simple example, the application of almost-degenerate perturbation theory is just the exact result, which behaves quadratically with  $|c|$  for small  $|c| \ll |a - b|$ , and transitions to behaving linearly with  $|c|$  for large  $|c| \gg |a - b|$ . These statements are illustrated in Figure 15.8.1. The fact that increasing the off-diagonal perturbation moves the energy eigenvalues apart is called **level repulsion**.




Figure 15.8.1: Perturbation theory results at leading order for energy eigenvalues of a two-state system with Hamiltonian given by eq. (15.8.9), for non-degenerate perturbation theory with  $b - a \gg |c|$  (left), degenerate perturbation theory with  $a = b$  (center), and almost-degenerate perturbation theory with  $a \approx b$  (right). For a two-state system, almost-degenerate perturbation theory is the exact result, by definition. In all three cases, the energy eigenvalues are moved apart by the perturbation, illustrating level repulsion.

As a less minimal example of almost-degenerate perturbation theory, consider a state space of dimension 3, with matrix representations for the unperturbed and perturbation Hamiltonians

$$H_0 = \begin{pmatrix} a - \Delta & 0 & 0 \\ 0 & a + \Delta & 0 \\ 0 & 0 & b \end{pmatrix}, \quad W = \begin{pmatrix} 0 & \delta & \epsilon \\ \delta & 0 & 0 \\ \epsilon & 0 & 0 \end{pmatrix}, \quad (15.8.13)$$

where  $a, b, \Delta, \delta$ , and  $\epsilon$  are all real constants. The exact eigenvalues of  $H = H_0 + W$  can be obtained by solving a cubic equation, but the results are complicated. However, it is worth noting that, because of the theorem in linear algebra that the sum of the eigenvalues of a matrix is equal to the trace, the exact energies must satisfy a simple sum rule,

$$E_1 + E_2 + E_3 = 2a + b, \quad (15.8.14)$$

regardless of the values of the other parameters. Now let us see what perturbation theory says.

First, let us find the energy eigenvalues by applying non-degenerate perturbation theory. Because  $H_0$  is diagonal, the unperturbed stationary states are the kets represented by the unit column vectors

$$|1\rangle = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad |2\rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad |3\rangle = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad (15.8.15)$$

with

$$\mathcal{E}_1 = a - \Delta, \quad \mathcal{E}_2 = a + \Delta, \quad \mathcal{E}_3 = b. \quad (15.8.16)$$

From eq. (15.1.43), the first-order corrections to the energies vanish, because  $W$  has no non-zero diagonal entries. Applying (15.1.44) gives the energy eigenvalues at second order in non-degenerate perturbation theory in  $\delta, \epsilon$ ,

$$E_1 = a - \Delta + \epsilon^2/(a - b - \Delta) - \delta^2/2\Delta, \quad (15.8.17)$$

$$E_2 = a + \Delta + \delta^2/2\Delta, \quad (15.8.18)$$

$$E_3 = b + \epsilon^2/(b - a + \Delta). \quad (15.8.19)$$

As a check, these do satisfy the sum rule eq.(15.8.14).

However, if  $\Delta$  is small compared to  $\delta$ , then the last term in each of eqs. (15.8.17) and (15.8.18) will be large, signaling a breakdown in the perturbative expansion because the unperturbed states  $|1\rangle$  and  $|2\rangle$  are too close in energy. In that case, one must use almost-degenerate perturbation theory. In this example,

$$P_{\text{deg}} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad W_{\text{deg}} = \begin{pmatrix} 0 & \delta & 0 \\ \delta & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (15.8.20)$$

and so

$$H'_0 = \begin{pmatrix} a - \Delta & \delta & 0 \\ \delta & a + \Delta & 0 \\ 0 & 0 & b \end{pmatrix}, \quad W' = \begin{pmatrix} 0 & 0 & \epsilon \\ 0 & 0 & 0 \\ \epsilon & 0 & 0 \end{pmatrix}. \quad (15.8.21)$$

Note that  $W'$  is indeed 0 within the  $2 \times 2$  subspace of almost-degenerate states. It is not too hard to find the unperturbed energies and eigenstates of  $H'_0$ ,

$$\mathcal{E}_{1'} = a - \sqrt{\delta^2 + \Delta^2}, \quad |1'\rangle = \sqrt{\frac{r_+}{2}} |1\rangle - \sqrt{\frac{r_-}{2}} |2\rangle, \quad (15.8.22)$$

$$\mathcal{E}_{2'} = a + \sqrt{\delta^2 + \Delta^2}, \quad |2'\rangle = \sqrt{\frac{r_-}{2}} |1\rangle + \sqrt{\frac{r_+}{2}} |2\rangle, \quad (15.8.23)$$

$$\mathcal{E}_{3'} = b, \quad |3'\rangle = |3\rangle, \quad (15.8.24)$$

where

$$r_{\pm} = 1 \pm \frac{\Delta}{\sqrt{\delta^2 + \Delta^2}}. \quad (15.8.25)$$

We then apply eqs. (15.1.43) and (15.1.44) with  $H'_0$  and  $W'$ , while being careful to note that the matrix representation of  $W'$  in eq. (15.8.21) is in the original basis  $|1\rangle, |2\rangle, |3\rangle$ , so that

$$W' = \epsilon |1\rangle\langle 3| + \epsilon |3\rangle\langle 1|. \quad (15.8.26)$$

The first-order contributions to the energies vanish because  $W'$  is also off-diagonal in the basis  $|1'\rangle, |2'\rangle, |3'\rangle$ . The second-order energy corrections are

$$E_{1'}^{(2)} = |\langle 3'|W'|1'\rangle|^2/(\mathcal{E}_{1'} - \mathcal{E}_{3'}), \quad (15.8.27)$$

$$E_{2'}^{(2)} = |\langle 3'|W'|2'\rangle|^2/(\mathcal{E}_{2'} - \mathcal{E}_{3'}), \quad (15.8.28)$$

$$E_{3'}^{(2)} = |\langle 1'|W'|3'\rangle|^2/(\mathcal{E}_{3'} - \mathcal{E}_{1'}) + |\langle 2'|W'|3'\rangle|^2/(\mathcal{E}_{3'} - \mathcal{E}_{2'}), \quad (15.8.29)$$

resulting in the energy eigenvalues at second-order in almost-degenerate perturbation theory,

$$E_{1'} = a - \sqrt{\delta^2 + \Delta^2} + \frac{\epsilon^2 r_+}{2(a - b - \sqrt{\delta^2 + \Delta^2})}, \quad (15.8.30)$$

$$E_{2'} = a + \sqrt{\delta^2 + \Delta^2} + \frac{\epsilon^2 r_-}{2(a - b + \sqrt{\delta^2 + \Delta^2})}, \quad (15.8.31)$$

$$E_{3'} = b + \frac{\epsilon^2(b - a - \Delta)}{(b - a)^2 - \delta^2 - \Delta^2}, \quad (15.8.32)$$

now with  $\epsilon$  as the sole expansion parameter. Again one can check that the sum rule eq.(15.8.14) is indeed satisfied.

The almost-degenerate perturbation theory expansion in  $\epsilon$  will be a good one as long as  $|\epsilon/(\mathcal{E}_{1'} - \mathcal{E}_{3'})| = |\epsilon/(a - b - \sqrt{\delta^2 + \Delta^2})|$  and  $|\epsilon/(\mathcal{E}_{2'} - \mathcal{E}_{3'})| = |\epsilon/(a - b + \sqrt{\delta^2 + \Delta^2})|$  are both small. Conversely, if they are not, then even this expansion will break down, and almost-degenerate perturbation theory would, unfortunately, consist of solving the original  $3 \times 3$  problem exactly.

## 15.9 Exercises

**Exercise 15.1.** Consider a harmonic oscillator supplemented by a quartic potential term, so that the Hamiltonian is  $H = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 X^2 + \lambda X^4$ . Treat the last term as a small perturbation.

(a) Compute the correction to the energy of the state  $|n\rangle$  at first order in  $\lambda$ . For fixed  $n$ , your answer is good for sufficiently small  $\lambda$ . However, you should find that it grows quadratically with  $n$ , so if  $\lambda$  is fixed, no matter how small it is, the perturbation expansion will break down for sufficiently large  $n$ .

(b) Find the ground state in terms of the unperturbed energy eigenkets, at first order in  $\lambda$ .

**Exercise 15.2.** The Hamiltonian for a rigid rotator in a magnetic field perpendicular to the  $y$  axis is  $H = aL^2 + bL_z + cL_x$ , where  $a$ ,  $b$ , and  $c$  are positive constants, and  $\vec{L}$  is the angular momentum operator.

(a) Write down the exact energy eigenvalues. (This should be quick. Start by writing  $bL_z + cL_x = b'\hat{n} \cdot \vec{L}$ , where  $\hat{n}$  is a unit vector; what is  $b'$ ?)

(b) Now take  $c \ll a, b$ . Find the unperturbed energy eigenvalues with  $c = 0$  for the eigenkets

$|l, m\rangle$  of  $L^2$  and  $L_z$ , and use them to obtain the energy eigenvalues at second order in the perturbation. Compare them to the exact answer from part (a). [Hint: the necessary matrix elements  $\langle l', m' | L_x | l, m \rangle$  can be found in eq. (8.4.4).]

**Exercise 15.3.** A spinless particle of charge  $e$  and mass  $m$  is confined to a cubic box of side  $L$  with center at the origin. A weak uniform electric field  $\vec{E} = E\hat{z}$  is applied, with direction parallel to one of the sides of the box, and the electrostatic potential for the field is taken to be zero at the center of the cubic box.

(a) Write down the unperturbed energy eigenvalues and corresponding normalized wavefunctions, in terms of appropriate quantum numbers. [Hint: for a particle confined to a 1-d box, the wavefunctions were given in eq. (6.4.10) and (6.4.11), for odd and even  $n$ , respectively. However, here you may find it more convenient to use the unified expression  $\varphi_n(x) = \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi x}{L} + \frac{n\pi}{2}\right)$ , which is equivalent up to irrelevant signs.]

(b) Find the change in the ground state energy at second order in the electric field. You should leave your answer in terms of an infinite sum of the form  $\text{SUM} = \sum_{n=2,4,6,\dots} \frac{n^p}{(n^2-1)^q}$ , where  $p$  and  $q$  are certain integers that you will find. Evaluate SUM to 3 significant digits. How does the change in the ground state energy scale with the size of the box  $L$ ?

**Exercise 15.4.** Consider the Stark effect for the  $n = 3$  states of the Hydrogen atom:

(a) For the unperturbed states  $|n, l, m\rangle$  with  $n = 3$ , find all of the non-zero matrix elements  $\langle 3, l', m' | Z | 3, l, m \rangle$ . (Use selection rules, and the fact that  $Z$  is Hermitian, to reduce the labor.)

(b) Include an electric field  $\vec{E} = E\hat{z}$ , treated as a perturbation. Obtain the energy eigenvalue corrections at first order in degenerate perturbation theory. Give the corresponding energy eigenstates to zeroth order in the perturbation, in terms of the unperturbed eigenstates  $|3, l, m\rangle$ .

**Exercise 15.5.** Starting from Brillouin–Wigner perturbation theory at second order in the state eq. (15.4.5) and third order in the energy eq. (15.4.6), derive the corresponding results for standard perturbation theory, given at the end of section 15.1.

**Exercise 15.6.** Consider a 3-state system with Hamiltonian  $H_0 + W$ , where in a certain basis

$$H_0 = \begin{pmatrix} \mathcal{E}_1 & 0 & 0 \\ 0 & \mathcal{E}_2 & 0 \\ 0 & 0 & \mathcal{E}_2 \end{pmatrix}, \quad W = \begin{pmatrix} 0 & a & b \\ a & 0 & c \\ b & c & 0 \end{pmatrix}, \quad (15.9.1)$$

where  $\mathcal{E}_1$  and  $\mathcal{E}_2$  are distinct but of the same order, and  $a, b, c$  are much smaller and real.

(a) Assuming that  $c \neq 0$ , find the energy eigenvalues at second order in degenerate perturbation theory. Recall that this requires going to a new unperturbed basis, chosen so that  $W$  is diagonal within the  $2 \times 2$  subspace corresponding to the degenerate  $H_0$  eigenvalues.

- (b) Now consider  $c = 0$ . Again compute the three energy eigenvalues at second order in degenerate perturbation theory. Note that, as explained in the discussion leading to eq. (15.6.13), since the degeneracy is not lifted at first order, it is necessary to choose a different unperturbed basis, such that the matrix  $W^{(2)}$  defined in eq. (15.6.9) is diagonal within the  $2 \times 2$  subspace corresponding to the degenerate  $H_0$  eigenvalues. (It is not diagonal, or even Hermitian, on the whole  $3 \times 3$  state space.) You should find that the two energy eigenvalues arising from  $\mathcal{E}_2$  are *not* the same as the  $c \rightarrow 0$  limit of part (a).
- (c) Check that the sum of the three energy eigenvalues in both cases is equal to  $\mathcal{E}_1 + 2\mathcal{E}_2$ , which is required because that is the trace of  $H$ . For the case  $c = 0$ , compute the exact eigenvalues of  $H$ , and check that expanding them to second order in the perturbations reproduces the results of part (b).

## 16 The variational method

### 16.1 Estimate and upper bound on the ground state energy

In Chapter 15, we studied how to use perturbation theory to find approximate solutions to the Hamiltonian eigenvalue problem. In this chapter, we will consider the **variational method** (also known as the **Rayleigh–Ritz method**), which applies even in situations for which it is not possible to make a sensible perturbation series expansion. Because the variational method is not based on a perturbative expansion, there is no need to know the exact solution to a simpler unperturbed problem.

The key idea comes from an inequality satisfied by the expectation value of the Hamiltonian in any state  $|\psi\rangle$  of our choice. Although the eigenstates  $|E_n, u_{E_n}\rangle$  of  $H$  and the corresponding energy eigenvalues  $E_n$  and degeneracy labels  $u_{E_n}$  are unknown, we can use completeness to write

$$\langle\psi|H|\psi\rangle = \sum_n \sum_{u_{E_n}} \langle\psi|H|E_n, u_{E_n}\rangle \langle E_n, u_{E_n}|\psi\rangle = \sum_n \sum_{u_{E_n}} E_n \langle\psi|E_n, u_{E_n}\rangle \langle E_n, u_{E_n}|\psi\rangle \quad (16.1.1)$$

$$\geq E_0 \sum_n \sum_{u_{E_n}} \langle\psi|E_n, u_{E_n}\rangle \langle E_n, u_{E_n}|\psi\rangle, \quad (16.1.2)$$

where we used  $E_n \geq E_0$  to get the last line. Now we can again use completeness to rewrite  $\sum_n \sum_{u_{E_n}} \langle\psi|E_n, u_{E_n}\rangle \langle E_n, u_{E_n}|\psi\rangle = \langle\psi|\psi\rangle$ , and obtain the bound

$$\frac{\langle\psi|H|\psi\rangle}{\langle\psi|\psi\rangle} \geq E_0. \quad (16.1.3)$$

(We have not assumed the normalization condition  $\langle\psi|\psi\rangle = 1$ , as this is sometimes not convenient.) Equation (16.1.3) means that we can make a guess, or a class of guesses, for  $|\psi\rangle$ , and then evaluate the left side of eq. (16.1.3) to obtain an estimate for  $E_0$  that is guaranteed not to be lower than the true value. The guess is called a **trial state** or **trial wavefunction**.

In a typical application, we improve the quality of our guess by choosing a whole family of trial states  $|\psi(a)\rangle$ , where  $a$  denotes a continuously variable parameter (or, more generally, a set of parameters). Now we compute

$$E(a) = \frac{\langle\psi(a)|H|\psi(a)\rangle}{\langle\psi(a)|\psi(a)\rangle}, \quad (16.1.4)$$

and minimize it with respect to  $a$  to obtain  $a_{\min}$ , typically by solving

$$\frac{\partial E(a)}{\partial a} = 0. \quad (16.1.5)$$

Then  $E(a_{\min})$  is our corresponding best estimate and upper bound for  $E_0$ , and  $|\psi(a_{\min})\rangle$  is our best estimate for the ground state  $|E_0\rangle$ .

The variational method tends to forgive us if our guess is good, but not perfect. To see why, suppose the result of our best guess can be written as

$$|\psi(a_{\min})\rangle = |E_0\rangle + \sum_{n \neq 0} \delta_n |E_n\rangle, \quad (16.1.6)$$

where the  $\delta_n$  are some unknown, but hopefully small, numbers representing the fact that our guess was not perfect. (For simplicity, take the energy eigenvalues to be not degenerate for the illustration; this is not a crucial assumption.) Then we compute

$$\langle \psi | H | \psi \rangle = E_0 + \sum_{n \neq 0} E_n |\delta_n|^2, \quad (16.1.7)$$

$$\langle \psi | \psi \rangle = 1 + \sum_{n \neq 0} |\delta_n|^2, \quad (16.1.8)$$

which yields

$$\frac{\langle \psi | H | \psi \rangle}{\langle \psi | \psi \rangle} = E_0 + \frac{\sum_{n \neq 0} (E_n - E_0) |\delta_n|^2}{1 + \sum_{n \neq 0} |\delta_n|^2}. \quad (16.1.9)$$

The error in our estimate for  $E_0$  is seen to be quadratic, involving  $|\delta_n|^2$  rather than  $|\delta_n|$ . The lesson here is that the estimate for the energy is better than the estimate for the state, provided only that the best-estimate errors are not too horrible, obeying  $|\delta_n| < 1$ . A 10% error in the state ket typically leads to only a  $\sim 1\%$  error in the energy.

Still, we do want to make a good guess for our parameterized family of states  $|\psi(a)\rangle$ , so that the best of them has a chance to be as close as possible to the true ground state. We should choose trial wavefunctions  $\psi(\vec{r})$  such that  $|\psi(\vec{r})|$  is larger where the potential  $V(\vec{r})$  is smaller. Furthermore, the wavefunction should have as few wiggles and nodes as possible, consistent with the given potential  $V(\vec{r})$ , in order to minimize the kinetic contribution to the Hamiltonian. In three-dimensional problems, if the Hamiltonian has rotational symmetry, minimizing the kinetic contribution likewise implies that we should make guesses that have  $L^2$  eigenvalue  $l = 0$ .

## 16.2 Variational method for excited states

The variational method also can provide information about higher-energy states, by projecting out lower-energy states from the trial state guesses. For example, suppose that we want to obtain the energy eigenvalue for the first excited state(s)  $|E_1\rangle$ . If we somehow already knew the exact  $|E_0\rangle$ , then we could define the projector

$$P = I - |E_0\rangle\langle E_0|, \quad (16.2.1)$$



which satisfies  $P^2 = P$  and  $P|E_0\rangle = 0$ . Then, for all trial states  $P|\chi\rangle$ , one can prove the bound

$$\frac{\langle\chi|PHP|\chi\rangle}{\langle\chi|P|\chi\rangle} \geq E_1, \quad (16.2.2)$$

where  $E_1$  is the first excited energy eigenvalue.

In practice, we may only have a variational estimate  $|\psi_{\text{est}}\rangle$  for  $|E_0\rangle$ , for example  $|\psi_{\text{est}}\rangle = |\psi(a_{\text{min}})\rangle$  from a previous variational estimate of the ground state. In that case, it might be that the best we can do is to instead define

$$\tilde{P} = I - |\psi_{\text{est}}\rangle \langle\psi_{\text{est}}|, \quad (16.2.3)$$

(assuming that  $|\psi_{\text{est}}\rangle$  has been normalized to unity) and then minimize the expectation value of the Hamiltonian in a parameterized family of states  $\tilde{P}|\chi(b)\rangle$ , where  $|\chi(b)\rangle$  is chosen based on a prior understanding of the properties of the first excited state(s), and depends on a variational parameter  $b$ . Unfortunately, the accuracy of the resulting estimate for  $E_1$  will be limited by the goodness of the estimate  $|\psi_{\text{est}}\rangle \approx |E_0\rangle$ . To the extent that this is not a good approximation, there is some leakage from the ground state into the guess, and we do not have a strict bound on the result of the minimization of the Hamiltonian expectation value. For this reason, it is often computationally challenging to obtain good estimates for all but the lowest few energy eigenvalues.

Fortunately, strict bounds and variational estimates for some of the higher energy states can be obtained if the Hamiltonian is compatible with some other observable(s). Suppose that there is an observable  $\Omega$  that satisfies

$$[H, \Omega] = 0. \quad (16.2.4)$$

This implies that one can choose an orthobasis of common eigenstates of  $H$  and  $\Omega$ , so that the energy eigenstates can be sub-classified by their  $\Omega$  eigenvalues, denoted  $\omega$ . Common examples for  $\Omega$  are the total angular momentum operators  $J^2$ ,  $J_z$  and/or the parity operator  $\Pi$ . In such cases, we can define a family of trial kets  $|\psi_\omega(a)\rangle$  that are restricted to be eigenstates of  $\Omega$  with a fixed eigenvalue  $\omega$ , so

$$\Omega |\psi_\omega(a)\rangle = \omega |\psi_\omega(a)\rangle. \quad (16.2.5)$$

In most examples, it will be obvious how to do this, but if necessary we can define the projection operator for each  $\omega$ ,

$$P_\omega = \sum_{u_\omega} |\omega, u_\omega\rangle \langle\omega, u_\omega|, \quad (16.2.6)$$

where the  $u_\omega$  are degeneracy labels for  $\omega$ , and then take  $|\psi_\omega(a)\rangle = P_\omega |\psi(a)\rangle$ . In any case, given a family of trial states  $|\psi_\omega(a)\rangle$ , one finds

$$E(a, \omega) = \frac{\langle \psi_\omega(a) | H | \psi_\omega(a) \rangle}{\langle \psi_\omega(a) | \psi_\omega(a) \rangle} \geq E_{\min}(\omega), \quad (16.2.7)$$

where  $E_{\min}(\omega)$  is the true minimum energy eigenvalue within the subspace of states that have  $\Omega = \omega$ . Thus, we obtain a separate variational estimate and bound on the lowest energy eigenstate within each  $\omega$  subspace.

For example, if the Hamiltonian for a spinless particle has rotational symmetry, then we can take  $\Omega = L^2$ . Then, applying the variational principle in turn to the trial wavefunctions  $\psi(r, \theta, \phi) = R_A(r)$  and  $R_B(r)Y_1^0(\theta, \phi)$  and  $R_C(r)Y_2^0(\theta, \phi)$ , we will obtain estimates for the minimum energy eigenvalues and wavefunctions for the ground state with  $l = 0$ , and for the lowest energy state that has  $l = 1$ , and for the lowest energy state that has  $l = 2$ , respectively. (Recall that rotational invariance guarantees that the energy eigenvalues and the radial wavefunctions are actually independent of  $m$ , so we took  $m = 0$  in each case just to be specific. Any other value of  $m$  would work just as well.) It is not necessary in this case to go to the trouble of defining the projection operator in eq. (16.2.6), because the eigenstates of  $L^2$  are known in terms of the spherical harmonics.

### 16.3 Examples of the variational method

As an easy example, consider a system for which we already know the exact answer: the harmonic oscillator in one dimension, with

$$H = \frac{P^2}{2m} + \frac{1}{2}m\omega^2 X^2. \quad (16.3.1)$$

We now use a family of trial wavefunctions

$$\psi(a) = e^{-x^2/a^2}. \quad (16.3.2)$$

This is a very fortunate guess, as it just so happens to include the correct ground-state wavefunction as a special case. Now, calculate

$$\langle \psi(a) | \psi(a) \rangle = \int_{-\infty}^{\infty} dx e^{-2x^2/a^2} = \sqrt{\frac{\pi}{2}} a, \quad (16.3.3)$$

and

$$\langle \psi(a) | H | \psi(a) \rangle = \int_{-\infty}^{\infty} dx e^{-x^2/a^2} \left[ -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + \frac{1}{2}m\omega^2 x^2 \right] e^{-x^2/a^2} \quad (16.3.4)$$

$$= \int_{-\infty}^{\infty} dx \left[ \frac{\hbar^2}{2m} \left( \frac{2}{a^2} - \frac{4x^2}{a^4} \right) + \frac{1}{2}m\omega^2 x^2 \right] e^{-2x^2/a^2} \quad (16.3.5)$$

$$= \sqrt{\frac{\pi}{2}} \left( \frac{\hbar^2}{2ma} + \frac{1}{8}m\omega^2 a^3 \right). \quad (16.3.6)$$

Using these in eq. (16.1.4) gives

$$E(a) = \frac{\hbar^2}{2ma^2} + \frac{m\omega^2 a^2}{8}. \quad (16.3.7)$$

The first term is the kinetic contribution, which wants  $a$  to be large, and the second term is the potential contribution, which prefers  $a$  to be small. Minimizing  $E(a)$  with respect to  $a^2$  gives

$$a_{\min}^2 = \frac{2\hbar}{m\omega}, \quad (16.3.8)$$

and so the result of the variational method is

$$E_0 \leq E(a_{\min}) = \hbar\omega/2, \quad (16.3.9)$$

in agreement with the exact result. Of course,  $\psi(a_{\min})$  is also the exact ground state wavefunction (up to the normalization factor), as a result of our fortunate guess.

For a less trivial example, consider the Hamiltonian

$$H = \frac{P^2}{2m} + \lambda X^4, \quad (16.3.10)$$

which is a cousin of the harmonic oscillator but with a steeper, quartic, potential. Not feeling very imaginative at the moment, and remembering the forgiving nature of the variational principle, we again choose as the trial wavefunction

$$\psi(a) = e^{-x^2/a^2}, \quad (16.3.11)$$

which yields

$$E(a) = \frac{\hbar^2}{2ma^2} + \frac{3}{16}\lambda a^4. \quad (16.3.12)$$

(The computation of the first term is identical to that for the previous example, as the trial wavefunction and the  $P^2/2m$  part of the Hamiltonian have not changed.) Now minimizing  $E(a)$  with respect to  $a^2$  gives

$$a_{\min}^2 = \left( \frac{4\hbar^2}{3\lambda m} \right)^{1/3} \approx 1.101 \left( \frac{\hbar^2}{\lambda m} \right)^{1/3}, \quad (16.3.13)$$

and

$$E_{\min} = \frac{3}{8} \left( \frac{6\hbar^4\lambda}{m^2} \right)^{1/3} \approx 0.681 \left( \frac{\hbar^4\lambda}{m^2} \right)^{1/3} \quad (16.3.14)$$

as the estimate of the ground state energy.

To do better, we invent a three-parameter trial wavefunction

$$\psi(a, b, c) = (1 + bx^2 + cx^4)e^{-x^2/a^2}. \quad (16.3.15)$$

We do not bother including odd powers of  $x$ , because we expect that the parity of the ground state is  $+1$ . Using a computer to minimize  $E(a, b, c)$  with respect to all three of  $a, b, c$  gives

$$E(a_{\min}, b_{\min}, c_{\min}) \approx 0.668 \left( \frac{\hbar^4 \lambda}{m^2} \right)^{1/3} \quad (16.3.16)$$

as a better estimate. As always, the more general wavefunction gives an estimate that is lower, and closer to the true ground-state energy eigenvalue.

The first excited state for  $H$  in eq. (16.3.10) will have one node, at  $x = 0$ , and odd parity. To find it, we can therefore exploit eq. (16.2.7), by choosing a simple trial wavefunction

$$\psi(a) = xe^{-x^2/a^2}. \quad (16.3.17)$$

A straightforward calculation of the Hamiltonian's expectation value and the norm gives

$$E(a) = \frac{3\hbar^2}{2ma^2} + \frac{15}{16}\lambda a^4, \quad (16.3.18)$$

with minimum at

$$a_{\min}^2 = \left( \frac{4\hbar^2}{5\lambda m} \right)^{1/3}, \quad (16.3.19)$$

leading to the estimate and upper bound for  $E_1$  of

$$E(a_{\min}) = \frac{9}{4} \left( \frac{5\hbar^4 \lambda}{4m^2} \right)^{1/3} \approx 2.424 \left( \frac{\hbar^4 \lambda}{m^2} \right)^{1/3}. \quad (16.3.20)$$

A somewhat better result could follow from using a multi-parameter trial wavefunction of the form  $\psi(a, b, c, \dots) = (x + bx^3 + cx^5 + \dots)e^{-x^2/a^2}$ . Note that it is important to be careful to only include odd-parity terms in this trial wavefunction, to avoid contamination from the even-parity ground state. The actual first excited state energy eigenvalue  $E_1$  turns out to be

$$E_1 \approx 2.394 \left( \frac{\hbar^4 \lambda}{m^2} \right)^{1/3}, \quad (16.3.21)$$

as can be obtained from numerical minimization of, say, a 6-parameter trial wavefunction.

Next, consider as a three-dimensional example the hydrogen atom Hamiltonian, written in the spherical coordinate position representation as

$$H = T + V, \quad T = -\frac{\hbar^2 \nabla^2}{2m_e}, \quad V = -\frac{e^2}{r}. \quad (16.3.22)$$

Although we know the exact wavefunction, let us ignore the temptation to use it, and instead try a Gaussian trial wavefunction with a variational parameter  $k$ ,

$$\psi(k) = e^{-k^2 r^2}. \quad (16.3.23)$$

To find the expectation value of the Hamiltonian, we first compute the kinetic contribution,

$$\langle \psi | T | \psi \rangle = -\frac{\hbar^2}{2m_e} 4\pi \int_0^\infty dr r^2 e^{-k^2 r^2} \left( \frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} \right) e^{-k^2 r^2} = 4\pi \left( \frac{\hbar^2}{m_e k} \sqrt{\frac{\pi}{2}} \frac{3}{16} \right), \quad (16.3.24)$$

then the potential contribution,

$$\langle \psi | V | \psi \rangle = 4\pi \left( -\frac{e^2}{4k^2} \right), \quad (16.3.25)$$

and finally the normalization factor,

$$\langle \psi | \psi \rangle = 4\pi \left( \sqrt{\frac{\pi}{2}} \frac{1}{8k^3} \right). \quad (16.3.26)$$

Putting these together, we have

$$E(k) = \frac{3\hbar^2 k^2}{2m_e} - \sqrt{\frac{8}{\pi}} e^2 k. \quad (16.3.27)$$

The minimum is found from  $\partial E(k)/\partial k = 0$  to be at

$$k_{\min} = \sqrt{\frac{8}{\pi}} \frac{e^2 m_e}{3\hbar^2} = \sqrt{\frac{8}{\pi}} \frac{1}{3a_0}. \quad (16.3.28)$$

Plugging this in gives the estimate of the ground state energy

$$E(k_{\min}) = -\left( \frac{8}{3\pi} \right) \frac{e^2}{2a_0} \approx -0.849 \text{ Rydberg}. \quad (16.3.29)$$

Of course, the exact answer is  $-1$  Rydberg. Even though our choice of one-parameter trial wavefunctions was not a very good one, because the harmonic oscillator is a poor approximation to the Coulomb potential, the result was within about 15% of the correct answer.

## 16.4 Helium atom ground state from the variational method

We conclude this chapter with a more practical example in which the variational method shines: the helium atom consisting of two electrons with charge  $-e$  and a nucleus with charge  $+2e$ . The Hamiltonian for this system was already given in eqs. (15.3.1)–(15.3.3) of section 15.3, where we treated it using first-order perturbation theory.

To use the variational method in the most simple way, let us choose a factorized trial wavefunction of the form

$$\Psi(\tilde{Z}, \vec{r}_1, \vec{r}_2) = \psi(\tilde{Z}, \vec{r}_1) \psi(\tilde{Z}, \vec{r}_2), \quad (16.4.1)$$

where [compare eq. (11.1.42)]

$$\psi(\tilde{Z}, \vec{r}) = \sqrt{\frac{\tilde{Z}^3}{\pi a_0^3}} e^{-\tilde{Z}r/a_0} \quad (16.4.2)$$

is the normalized ground state wavefunction for a hydrogen-like atom with general atomic number  $\tilde{Z}$ . However, here we do not set  $\tilde{Z}$  equal to  $Z = 2$  in the trial wavefunction, but instead treat it as a continuous variational parameter, adjusting it to find the minimum expectation value of the total  $H$ . In contrast,  $Z$  in the Hamiltonian is fixed to be equal to the actual number of protons in the nucleus. For helium,  $Z = 2$ , but we will leave it general for the following discussion, so that our results apply to any ion with 2 electrons.

We therefore need to compute, and then minimize:

$$E(\tilde{Z}) = E_1(\tilde{Z}) + E_2(\tilde{Z}) + E_{12}(\tilde{Z}), \quad (16.4.3)$$

where  $E_1(\tilde{Z}) = E_2(\tilde{Z}) = \langle \Psi | H_1 | \Psi \rangle$ , and  $E_{12}(\tilde{Z}) = \langle \Psi | H_{12} | \Psi \rangle$ , since we have arranged for  $\langle \Psi | \Psi \rangle = 1$  in this example. Because each electron partly screens the charge  $+Ze$  nucleus from the other electron, we anticipate that the value  $\tilde{Z}_{\min}$  that minimizes  $E(\tilde{Z})$  will be in the range  $Z - 1 < \tilde{Z}_{\min} < Z$ , so that for helium,  $1 < \tilde{Z}_{\min} < 2$ .

To find the non-interaction contributions, we first compute

$$E_1(\tilde{Z}) = \int d^3\vec{r}_1 \psi(\vec{r}_1)^* \left( -\frac{\hbar^2}{2m_e} \nabla_1^2 - \frac{Ze^2}{r_1} \right) \psi(\vec{r}_1) \int d^3\vec{r}_2 |\psi(\vec{r}_2)|^2. \quad (16.4.4)$$

Note that the potential energy contribution is  $-Ze^2/r_1$ , not  $-\tilde{Z}e^2/r_1$ , because  $\tilde{Z}$  is our variational parameter, not the actual nuclear charge. In this expression,  $\int d^3\vec{r}_2 |\psi(\vec{r}_2)|^2 = 1$ , and the remaining integral factor evaluates simply, to give

$$E_1(\tilde{Z}) = \frac{e^2}{2a_0} (\tilde{Z}^2 - 2\tilde{Z}Z). \quad (16.4.5)$$

Of course,  $E_2(\tilde{Z}) = E_1(\tilde{Z})$  follows from an identical calculation with the labels 1 and 2 interchanged. The more difficult part is the interaction contribution

$$E_{12}(\tilde{Z}) = e^2 \int d^3\vec{r}_1 \int d^3\vec{r}_2 |\psi(\vec{r}_1)|^2 |\psi(\vec{r}_2)|^2 \frac{1}{|\vec{r}_1 - \vec{r}_2|}. \quad (16.4.6)$$

Fortunately, we have already evaluated this integral in eqs. (15.3.8)–(15.3.18), with the result

$$E_{12}(\tilde{Z}) = \frac{5\tilde{Z}e^2}{8a_0}. \quad (16.4.7)$$

Plugging this into eq. (16.4.3), along with  $E_1(\tilde{Z}) = E_2(\tilde{Z})$  from eq. (16.4.5), gives

$$E(\tilde{Z}) = \tilde{Z}(\tilde{Z} - 2Z + 5/8) \frac{e^2}{a_0}. \quad (16.4.8)$$

Minimizing this with respect to our variational parameter  $\tilde{Z}$  gives

$$\tilde{Z}_{\min} = Z - 5/16 = 27/16. \quad (16.4.9)$$

Note that indeed  $Z - 1 < \tilde{Z}_{\min} < Z$ , as we had anticipated.

Our variational method estimate for the ground state energy of the helium atom is therefore

$$E(\tilde{Z}_{\min}) = -\tilde{Z}_{\min}^2 \frac{e^2}{a_0} = -\left(\frac{27}{16}\right)^2 \frac{e^2}{a_0} = -5.695 \text{ Rydberg} = -77.49 \text{ eV}. \quad (16.4.10)$$

This can be compared to the experimental value (the minimum ionization energy needed to completely remove both electrons from the He atom in its ground state), which is

$$E_{\text{exp}} = -5.807 \text{ Rydberg} = -79.005 \text{ eV}. \quad (16.4.11)$$

As expected, our variational method estimate is higher than the true value, but by less than 2%. Recall that the first-order perturbation theory method in section 15.3 gave instead

$$E_{\text{pert.}} = -5.5 \text{ Rydberg} = -74.83 \text{ eV}, \quad (16.4.12)$$

which is about 5.3% higher than the experimental value. The variational method, even with the simplest trial wavefunction, performs better than first-order perturbation theory. If one uses a more sophisticated non-factorized trial wavefunction with more parameters, and asks a computer to do the integrations and minimization numerically, then the variational method will win even more convincingly for a given amount of human calculation effort expended. The variational method also scales nicely to more complex problems in atomic and molecular physics, where perturbation theory may have more difficulty.

## 16.5 Exercises

**Exercise 16.1.** Consider particle of mass  $m$  moving in 1 dimension in the very strongly attractive potential  $V(X) = \lambda X^6$ . Using a Gaussian trial wavefunction, estimate the ground state energy and wavefunction. What is the corresponding estimate for  $\langle X^2 \rangle$ ? Do the same for the first excited state, using a trial wavefunction of the form  $x$  multiplied by a Gaussian.

**Exercise 16.2.** A particle of mass  $m$  moves in a spherically symmetric potential  $V(R)$ . Denote by  $E_l$  the lowest bound state energy with orbital angular momentum  $l$  [that is, with  $L^2$  eigenvalue  $\hbar^2 l(l+1)$ ]. Use the variational principle to prove that  $E_l$  increases monotonically with  $l$ . (It obviously follows that the ground state must have  $l = 0$ .) Compare to Figures 10.3.2 and 11.1.1.

**Exercise 16.3.** Consider a particle of mass  $m$  moving in the 3-dimensional potential  $V(R) = \lambda R$ , corresponding to a constant restoring force. Use two different trial wavefunctions,  $\psi = e^{-r/a}$  and  $\psi = e^{-r^2/b^2}$ , to find the ground state energy. Which trial wavefunction does a better job?

**Exercise 16.4.** Consider a particle of mass  $m$  moving in a Yukawa (also known as screened Coulomb) potential

$$V(R) = -\frac{g}{R}e^{-R/a}, \quad (16.5.1)$$

which behaves at short distances like a Coulomb potential and vanishes exponentially at large distances. Consider a trial wavefunction  $\psi = e^{-\beta r/a}$ , where  $\beta$  is the variational parameter.

(a) Show that the energy expectation value for the trial state is

$$E(\beta) = \frac{\hbar^2}{ma^2} \left( \frac{1}{2\beta^2} - \frac{4G}{\beta(2+\beta)^2} \right), \quad (16.5.2)$$

where  $G = gma/\hbar^2$ .

(b) Find the condition (cubic in  $\beta$ , and linear in  $G$ ) that gives the minimum of  $E(\beta)$ .

(c) The condition for a bound state is  $E < 0$ . This requires  $G > G_c$ , where  $G_c$  is a certain critical value for the strength of the potential. Use the condition you found in the previous part to eliminate  $G$ , and  $E(\beta) = 0$ , to show that the estimate from this trial wavefunction is  $G_c = 1$ . (Note that this does not require solving a cubic equation.) To illustrate this, make a graph of  $E(\beta)$  for  $0 < \beta < 3$ , for the three cases  $G = 1/2$ ,  $G = 1$ , and  $G = 2$ . What is  $\beta_{\min}$  when  $G = G_c = 1$ ? (By using a more carefully chosen trial wavefunction, it is possible to show that the condition for a bound state to exist is more accurately given by  $gma/\hbar^2 > 0.8399$ .)

**Exercise 16.5.** Consider a particle of mass  $\mu$  moving in the 3-dimensional isotropic harmonic oscillator potential  $V(R) = \frac{1}{2}\mu\omega^2 R^2$ , where  $R$  is the radial coordinate operator. Instead of solving exactly as we did in section 10.5, let us see how well the variational principle works with a trial wavefunction  $\psi_{l,m}(n, k) = r^n e^{-kr} Y_l^m(\theta, \phi)$  in spherical coordinates. The quantities  $n$  and  $k$  are the variational parameters. We could take  $n$  to vary continuously, but we will consider it to only take on integer values in this problem.

(a) Compute the energy function for the trial wavefunction. Write your answer in the form  $E(n, k) = (\hbar^2 k^2 / 2m)A + (m\omega^2 / k^2)B$ , where  $A$  is a quadratic polynomial in  $l, n$  divided by a quadratic polynomial in  $n$ , and  $B$  is another quadratic polynomial in  $n$ . [Hint: to obtain the kinetic energy contribution, you will want to use eq. (8.6.12), and certainly not the explicit forms of the spherical harmonics.]

(b) Minimize  $E(n, k)$  with respect to  $k$ , to obtain  $E(n, k_{\min})$ . Work in terms of the symbols  $A$  and  $B$ , and only plug in what they are in the next step.

(c) Estimate the ground state energy for  $l = 0$ . Which integer  $n$  does the best job? (Plug in  $n = 0, 1, 2, 3, 4, 5, 6$  to find out.) How does it compare to the exact answer?

(d) Repeat part (c) for the lowest state with  $l = 1$ , and for the lowest state with  $l = 2$ .



# 17 Fine, hyperfine, and magnetic effects for the hydrogen atom

## 17.1 Relativistic kinetic, spin-orbit, and Darwin corrections

The hydrogen atom Hamiltonian was treated in Chapter 11 in a nonrelativistic approximation. There, it was argued that relativistic effects should affect the energy eigenvalues by a factor suppressed by the square of the fine-structure constant  $\alpha = 1/137.036 \dots$ . Now it is time to find those corrections, using stationary-state perturbation theory.

An expansion of the Dirac equation, carried out in section 27.4, shows that a more accurate approximation for the electron (with mass called  $m_e$  in this chapter) moving in a general spherically symmetric electrostatic potential<sup>†</sup>  $\Phi(R)$  is given by

$$H = H_0 + H_{\text{rel}} + H_{\text{SO}} + H_{\text{Darwin}}, \quad (17.1.1)$$

where

$$H_0 = \frac{P^2}{2m_e} - e\Phi(R) \quad (17.1.2)$$

is the nonrelativistic approximation we have already used, with  $P^2 = \vec{P} \cdot \vec{P}$ , and

$$H_{\text{rel}} = -\frac{(P^2)^2}{8m_e^3c^2}, \quad (17.1.3)$$

$$H_{\text{SO}} = -\frac{e}{2m_e^2c^2} \frac{1}{R} \frac{d\Phi}{dR} \vec{S} \cdot \vec{L}, \quad (17.1.4)$$

$$H_{\text{Darwin}} = -\frac{\hbar^2 e}{8m_e^2c^2} \nabla^2 \Phi \quad (17.1.5)$$

are called the relativistic kinematic, spin-orbit, and Darwin terms, respectively. (The last is named for Charles G. Darwin, grandson of the Charles R. Darwin famous for evolutionary biology.) These three terms give numerical contributions that are of the same order, and they are collectively called the **fine structure** corrections to the hydrogen atom energies. Each of them is a relativistic effect, as seen from the  $1/c^2$  factors, which would make them formally vanish if the speed of light were infinite. In this section, we will calculate their effects using perturbation theory. Before doing so, we briefly comment on the physical reasons behind the fine structure terms.

First,  $H_{\text{rel}}$  comes from the binomial expansion of the special relativistic expression for the energy of a massive particle with classical momentum  $p$ ,

$$E = \sqrt{m_e^2c^4 + p^2c^2} = m_e c^2 + \frac{p^2}{2m_e} - \frac{(p^2)^2}{8m_e^3c^2} + \dots \quad (17.1.6)$$

---

<sup>†</sup>For hydrogen-like atoms,  $\Phi(R) = Ze/R$ , but in eqs. (17.1.2)–(17.1.5) we choose to be more general.

The first term is just a constant, Einstein's equivalence of rest mass and energy. We can ignore it, since it causes all time-dependent states to be multiplied by the same phase. The second term is the kinetic term in  $H_0$ , and the third gives  $H_{\text{rel}}$  when  $p$  is promoted from classical momentum to the quantum momentum operator.

The  $H_{\text{SO}}$  term can be understood as the energy due to the interaction of the intrinsic magnetic moment of the electron with the magnetic field  $\vec{B}$  that it experiences because it is moving. Naively, this gives a contribution to the Hamiltonian  $-\vec{\mu}_e \cdot \vec{B}$ , where the magnetic field as seen in the electron's rest frame moving with velocity  $\vec{v}$  is, in gaussian cgs units,

$$\vec{B} = -\frac{\vec{v}}{c} \times \vec{E}. \quad (17.1.7)$$

Here, we can use  $\vec{v} = \vec{p}/m_e$  and, for a spherically symmetric potential,

$$\vec{E} = -\vec{\nabla}\Phi = -\frac{\vec{r}}{r} \frac{d\Phi}{dr}, \quad (17.1.8)$$

so that, promoting  $\vec{r}$  and  $\vec{p}$  to quantum operators,

$$\vec{B} = \frac{1}{m_e c} \frac{1}{R} \frac{d\Phi(R)}{dR} (\vec{P} \times \vec{R}) = -\frac{1}{m_e c} \frac{1}{R} \frac{d\Phi(R)}{dR} \vec{L}. \quad (17.1.9)$$

There is no operator ordering issue in writing the cross-product  $\vec{L} = \vec{R} \times \vec{P} = -\vec{P} \times \vec{R}$ , because different rectangular coordinate components of  $\vec{P}$  and  $\vec{R}$  commute. The magnetic moment of the electron is opposite the direction of the spin, because the charge  $-e$  is negative,

$$\vec{\mu}_e = -\frac{e}{m_e c} \vec{S}, \quad (17.1.10)$$

where we have used eqs. (4.3.18) and (4.3.19) with the (very good) approximation  $g_e \approx 2$ . So, naively we would have

$$H_{\text{SO, naive}} = -\vec{\mu}_e \cdot \vec{B} = -\frac{e}{m_e^2 c^2} \frac{1}{R} \frac{d\Phi}{dR} \vec{S} \cdot \vec{L}. \quad (17.1.11)$$

The extra factor of  $1/2$  in the true  $H_{\text{SO}}$  in eq. (17.1.4) is known as the Thomas precession effect. Llewellyn Thomas explained it in 1925 as a classical relativistic effect of the electron being at rest in a rotating (not inertial) reference frame. Alternatively, the derivation from the Dirac equation in section 27.4 gives the correct normalization more directly.

Finally, the physical origin of the Darwin term  $H_{\text{Darwin}}$  is related to the fact that, as we noted following eq. (11.1.71), the behavior of the electron is modified by relativistic effects on a small distance scale comparable to its reduced Compton wavelength,  $\lambda_e = \hbar/m_e c$ . This effectively makes the replacement  $\Phi(R) \rightarrow \Phi(R) - \frac{\lambda_e^2}{8} \nabla^2 \Phi$ , resulting in the correction of eq. (17.1.5). A proper derivation from the Dirac equation is again found in section 27.4.

Now we apply first-order perturbation theory to evaluate the  $H_{\text{rel}}$  contribution to the energies of the unperturbed states  $|n, l, m\rangle$ . These unperturbed states have degeneracies for  $n > 1$ , so we are really doing degenerate perturbation theory, but  $W = H_{\text{rel}}$  is already diagonal in this basis, so we can directly apply eq. (15.6.4). This gives

$$E_{n,l,m}^{(1),\text{rel}} = \langle H_{\text{rel}} \rangle_{n,l,m} = -\frac{1}{8m_e^3 c^2} \langle (P^2)^2 \rangle_{n,l,m}, \quad (17.1.12)$$

where here, and in the following,  $\langle \cdots \rangle_{n,l,m}$  denotes the expectation value  $\langle n, l, m | \cdots | n, l, m \rangle$ . To evaluate this efficiently, we can use a trick, by rewriting it using  $P^2 = 2m_e(H_0 + e^2/R)$ ,

$$E_{n,l,m}^{(1),\text{rel}} = -\frac{1}{2m_e c^2} \langle (H_0 + e^2/R)^2 \rangle_{n,l,m} \quad (17.1.13)$$

$$= -\frac{1}{2m_e c^2} (\mathcal{E}_n^2 + 2\mathcal{E}_n e^2 \langle 1/R \rangle_{n,l,m} + e^4 \langle 1/R^2 \rangle_{n,l,m}), \quad (17.1.14)$$

where  $\mathcal{E}_n = -e^2/2a_0 n^2$  is the eigenvalue of  $H_0$  for the state  $|n, l, m\rangle$ . The expectation values of  $1/R$  and  $1/R^2$  were given in eqs. (11.1.64) and (11.1.67), respectively. Plugging those results into eq. (17.1.14) gives

$$E_{n,l,m}^{(1),\text{rel}} = \alpha^2 \frac{e^2}{2a_0} \frac{1}{n^3} \left( \frac{3}{4n} - \frac{1}{l+1/2} \right), \quad (17.1.15)$$

where we have used  $a_0 = \hbar^2/m_e e^2$  to write the result in terms of the fine structure constant  $\alpha = e^2/\hbar c \approx 1/137.036$ , and  $e^2/2a_0 = 1$  Rydberg  $\approx 13.606$  eV.

By itself, eq. (17.1.15) is not very useful, because the spin-orbit and Darwin terms contribute at the same order. To evaluate the spin-orbit contribution, we note that for the special case of the hydrogen atom, the electrostatic potential is  $\Phi = e/R$ , so that eq. (17.1.4) becomes

$$H_{\text{SO}} = \frac{e^2}{2m_e^2 c^2} \frac{1}{R^3} \vec{S} \cdot \vec{L}. \quad (17.1.16)$$

This vanishes for all  $l = 0$  states, including the ground state  $n = 1$ , since they have no orbital angular momentum. Because the  $n > 1$  states have  $H_0$  eigenvalue degeneracies, we are really doing degenerate perturbation theory, which means that we must use a basis in which  $H_{\text{SO}}$  is diagonal. We use the standard trick for dot products of angular momentum operators,

$$\vec{S} \cdot \vec{L} = \frac{1}{2}(J^2 - L^2 - S^2) = \frac{\hbar^2}{2} [j(j+1) - l(l+1) - s(s+1)], \quad (17.1.17)$$

which is diagonal in the total angular momentum orthobasis  $|n, l, s, j, m_j\rangle$ . (In particular, we do *not* want to use the product orthobasis  $|n, l, s, m_l, m_s\rangle$  here, because then  $\vec{S} \cdot \vec{L}$  would not be diagonal, as required for degenerate perturbation theory.) Using  $s = 1/2$ , it follows that the first-order correction is

$$E_{n,l,j}^{(1),\text{SO}} = \frac{\hbar^2 e^2}{4m_e^2 c^2} [j(j+1) - l(l+1) - 3/4] \langle 1/R^3 \rangle_{n,l,m}, \quad (l \neq 0). \quad (17.1.18)$$

The expectation value of  $1/R^3$  was given in eq. (11.1.68), reproduced here for convenience,

$$\langle 1/R^3 \rangle_{n,l,m} = \frac{1}{a_0^3 n^3 l(l+1)(l+1/2)}. \quad (17.1.19)$$

It follows that

$$E_{n,l,j}^{(1),\text{SO}} = \frac{\hbar^2 e^2}{2m_e^2 c^2 a_0^3} \frac{j(j+1) - l(l+1) - 3/4}{n^3 l(l+1)(2l+1)}, \quad (l \neq 0). \quad (17.1.20)$$

Now, using  $j = l \pm 1/2$ , and writing the prefactor in terms of the fine structure constant and the Rydberg energy,

$$E_{n,l,j}^{(1),\text{SO}} = \alpha^2 \left( \frac{e^2}{2a_0} \right) \frac{-1 \pm (2l+1)}{2n^3 l(l+1)(2l+1)}, \quad (l \neq 0, j = l \pm 1/2), \quad (17.1.21)$$

to go along with  $E_{n,0,1/2}^{(1),\text{SO}} = 0$ .

Now we turn to the Darwin term of eq. (17.1.5), which for the hydrogen atom includes, in the position representation, a factor  $\nabla^2 \Phi = e \nabla^2 (1/r) = -4\pi e \delta^{(3)}(\vec{r})$ . It is left to Exercise 17.1 to compute the first-order energy correction

$$E^{(1),\text{Darwin}} = \delta_{l0} \alpha^2 \left( \frac{e^2}{2a_0} \right) \frac{1}{n^3}, \quad (17.1.22)$$

which only depends on  $n, l$  and is only non-zero for  $l = 0$ .

We are finally ready to combine the three fine-structure contributions,  $E^{\text{fine}} = E^{(1),\text{rel}} + E^{(1),\text{SO}} + E^{(1),\text{Darwin}}$ , from eqs. (17.1.15), (17.1.21), and (17.1.22). This can be done in three separate cases,

$$(l = 0, j = 1/2): \quad E^{\text{fine}} = \frac{\alpha^2}{n^3} \left( \frac{e^2}{2a_0} \right) \left( \frac{3}{4n} - 2 + 0 + 1 \right), \quad (17.1.23)$$

$$(l \neq 0, j = l + 1/2): \quad E^{\text{fine}} = \frac{\alpha^2}{n^3} \left( \frac{e^2}{2a_0} \right) \left( \frac{3}{4n} - \frac{2}{2l+1} + \frac{1}{(l+1)(2l+1)} + 0 \right), \quad (17.1.24)$$

$$(l \neq 0, j = l - 1/2): \quad E^{\text{fine}} = \frac{\alpha^2}{n^3} \left( \frac{e^2}{2a_0} \right) \left( \frac{3}{4n} - \frac{2}{2l+1} - \frac{1}{l(2l+1)} + 0 \right). \quad (17.1.25)$$

All three cases are seen to be encompassed by the simple formula

$$E_{n,j}^{\text{fine}} = \frac{\alpha^2}{n^3} \left( \frac{e^2}{2a_0} \right) \left( \frac{3}{4n} - \frac{1}{j+1/2} \right), \quad (17.1.26)$$

which notably depends only on  $n$  and  $j$ , and not separately on  $l$ , as one perhaps might have guessed. This result can also be obtained by expanding the exact solution of the Dirac equation; this will be done in section 27.5.

For the lowest few energy levels, in terms of the quantity

$$E_f = \alpha^2 e^2 / 2a_0 = 7.245 \times 10^{-4} \text{ eV}, \quad (17.1.27)$$

one has fine structure shifts for spectroscopic states for  $n = 1$ :

$$1S_{1/2} \quad (n = 1, l = 0, j = 1/2) \quad E_{\text{fine}} = -E_f/4, \quad (17.1.28)$$

and for  $n = 2$ :

$$\left. \begin{array}{l} 2S_{1/2} \quad (n = 2, l = 0, j = 1/2) \\ 2P_{1/2} \quad (n = 2, l = 1, j = 1/2) \end{array} \right\} \quad E_{\text{fine}} = -5E_f/64, \quad (17.1.29)$$

$$2P_{3/2} \quad (n = 2, l = 1, j = 3/2) \quad E_{\text{fine}} = -E_f/64, \quad (17.1.30)$$

where the  $2S_{1/2}$  and  $2P_{1/2}$  states stay degenerate, and for  $n = 3$ :

$$\left. \begin{array}{l} 3S_{1/2} \quad (n = 3, l = 0, j = 1/2) \\ 3P_{1/2} \quad (n = 3, l = 1, j = 1/2) \end{array} \right\} \quad E_{\text{fine}} = -E_f/36, \quad (17.1.31)$$

$$\left. \begin{array}{l} 3P_{3/2} \quad (n = 3, l = 1, j = 3/2) \\ 3D_{3/2} \quad (n = 3, l = 2, j = 3/2) \end{array} \right\} \quad E_{\text{fine}} = -E_f/108, \quad (17.1.32)$$

$$3D_{5/2} \quad (n = 3, l = 2, j = 5/2) \quad E_{\text{fine}} = -E_f/324, \quad (17.1.33)$$

where the  $3S_{1/2}$  and  $3P_{1/2}$  states stay degenerate, as do the  $3P_{3/2}$  and  $3D_{3/2}$  states. Including electron spin, each principal quantum number level  $n$  has  $(2s+1) \sum_{l=0}^{n-1} (2l+1) = 2n^2$  states. The fine structure splits each of those levels into  $n$  sub-levels, with  $j = 1/2, 3/2, \dots, n-1/2$ . The degeneracies are further enhanced by a factor of 2 for the proton spin, but then are further split by the hyperfine structure to be discussed next.

## 17.2 Hyperfine structure of hydrogen

The hyperfine structure takes into account that an atomic nucleus also has a magnetic dipole moment (along its spin direction) and an electric quadrupole moment (if the nuclear spin is at least 1; see Exercise 13.5). For the  $^1\text{H}$  isotope of hydrogen with a nucleus consisting of one proton and no neutrons, the magnetic moment of the proton is related to its spin-1/2 operator, which we denote  $\vec{S}_p$ , by

$$\vec{\mu}_p = \frac{g_p e}{2m_p c} \vec{S}_p, \quad (17.2.1)$$

(see section 4.3), where  $m_p$  is the proton's mass, and  $g_p = 5.5856946893(16)$  is the proton's  $g$ -factor. The hyperfine contributions to the energy eigenvalues are numerically suppressed compared to the fine structure contributions because of the fact that  $m_p \gg m_e$ .

The interaction between the magnetic field created by the proton's magnetic moment and the moving electron introduces a proton spin-orbit coupling term in the Hamiltonian,

$$H_{\text{SO}}^{\text{proton}} = \frac{e}{m_e c R^3} \vec{\mu}_p \cdot \vec{L} = \frac{g_p e^2}{2m_e m_p c^2} \frac{1}{R^3} \vec{S}_p \cdot \vec{L}. \quad (17.2.2)$$

In addition, there is the energy of the magnetostatic interaction of the dipole moments of the proton and the electron,

$$H_{\text{dipole-dipole}} = \frac{1}{R^3} \left[ \vec{\mu}_e \cdot \vec{\mu}_p - \frac{3}{R^2} (\vec{\mu}_e \cdot \vec{R}) (\vec{\mu}_p \cdot \vec{R}) \right] - \frac{8\pi}{3} \vec{\mu}_e \cdot \vec{\mu}_p \delta^{(3)}(\vec{R}), \quad (17.2.3)$$

in which the final term is called a contact term. These two contributions are of the same numerical order, and so, using

$$\vec{\mu}_e = -\frac{g_e e}{2m_e c} \vec{S} \quad (17.2.4)$$

with  $g_e \approx 2$  and  $\vec{S}$  the electron's spin, the full hyperfine contribution to the Hamiltonian for hydrogen is

$$H_{\text{hf}} = H_{\text{SO}}^{\text{proton}} + H_{\text{dipole-dipole}} \quad (17.2.5)$$

$$= \frac{g_p e^2}{2m_e m_p c^2} \left[ \frac{1}{R^3} (\vec{L} - \vec{S}) \cdot \vec{S}_p + \frac{3}{R^5} (\vec{R} \cdot \vec{S}) (\vec{R} \cdot \vec{S}_p) + \frac{8\pi}{3} \vec{S} \cdot \vec{S}_p \delta^{(3)}(\vec{R}) \right]. \quad (17.2.6)$$

We now treat this as a perturbation of the fine-structure results found in the previous section.

Even after taking into account the fine structure, there were still energy degeneracies. Therefore, we again need to do degenerate perturbation theory by choosing a good set of unperturbed orthobasis states, so that  $H_{\text{hf}}$  is diagonal in that basis. The appropriate basis is again a total angular momentum basis, but now taking into account the proton's spin in addition to the electron's spin and orbital angular momentum. We define

$$\vec{J} = \vec{L} + \vec{S}, \quad (17.2.7)$$

$$\vec{F} = \vec{J} + \vec{S}_p = \vec{L} + \vec{S} + \vec{S}_p, \quad (17.2.8)$$

so that  $\vec{F}$  ( $\vec{J}$ ) is the total angular momentum including (excluding) the proton's spin. The eigenvalues for  $J^2$  and  $J_z$  are denoted  $\hbar^2 j(j+1)$  and  $\hbar m_j$ , where the allowed values for  $j$  are  $l-1/2$  and  $l+1/2$ , and in each case  $m_j$  takes on the  $2j+1$  values  $-j, -j+1, \dots, j-1, j$ . Because  $\vec{F}$  arises from adding the orbital angular momentum to two spin-1/2 angular momenta, the eigenvalues for  $F^2$  and  $F_z$  are  $\hbar^2 f(f+1)$  and  $\hbar m_f$ , with integer values  $f = j-1/2$  and  $j+1/2$ , and in each case  $m_f = -f, -f+1, \dots, f-1, f$ . The total angular momentum orthobasis states are constructed by first combining  $\vec{L}$  and  $\vec{S}$  eigenstates to obtain an orthobasis  $|l, s, j, m_j\rangle$ , and then taking the tensor product with the proton spin eigenstates  $|s_p, m_{s_p}\rangle$  and going to the total angular momentum basis to obtain states labeled  $|l, s, j, s_p, f, m_f\rangle$ . Here the  $s$  and  $s_p$  spin labels are trivial and can be omitted, as they are both fixed to be 1/2. So, including the principal quantum number  $n$ , the CSCO =  $\{H_0, L^2, J^2, F^2, F_z\}$  eigenstate orthobasis states can be labeled

$$|n, l, j, f, m_f\rangle. \quad (17.2.9)$$

We now want to show that  $H_{\text{hf}}$  is diagonal in this basis, and obtain its expectation values. It is convenient to treat separately the case  $l = 0$  (including the ground state) and the case  $l \neq 0$ .

For  $l = 0$ , the first two terms in eq. (17.2.6) do not contribute. To see this, note that the spherical symmetry of  $l = 0$  states guarantees that the expectation value of these two terms is proportional to the angular integral of  $3(\hat{r} \cdot \vec{S})(\hat{r} \cdot \vec{S}_p) - \vec{S} \cdot \vec{S}_p$ , where we have used  $\vec{L} = 0$  for  $l = 0$ . Such an integral is a scalar, and is linear in each of the two spins, and depends on no other vectors, so it can only be proportional to their dot product. That is, we must have

$$\int d\Omega \left[ 3(\hat{r} \cdot \vec{S})(\hat{r} \cdot \vec{S}_p) - \vec{S} \cdot \vec{S}_p \right] = a \vec{S} \cdot \vec{S}_p \quad (17.2.10)$$

for some constant number  $a$ . The value of  $a$  can now be discovered to be 0 by evaluating the left side for  $\vec{S} = \vec{S}_p = \hat{z}$ , using  $\hat{r} \cdot \hat{z} = \cos \theta$ .

The last term in eq. (17.2.6), proportional to  $\delta^{(3)}(\vec{R})$ , does contribute for  $l = 0$ , and only in that case. To see this, note that

$$\langle n, l, m_l | \delta^{(3)}(\vec{R}) | n, l, m_l \rangle = |\psi_{n,l,m_l}(0,0,0)|^2, \quad (17.2.11)$$

the square of the wavefunction at the origin. Using the radial wavefunction at  $r = 0$  from eq. (11.1.41), this is non-zero only if  $l = 0$ , which of course implies  $m_l = 0$  also, and

$$\langle n, l, m_l | \delta^{(3)}(\vec{R}) | n, l, m_l \rangle = \delta_{l0} \delta_{m_l 0} \frac{1}{\pi a_0^3 n^3}. \quad (17.2.12)$$

Now, with  $l = 0$ ,  $\vec{F} = \vec{S} + \vec{S}_p$ , so

$$\vec{S} \cdot \vec{S}_p = \frac{1}{2}(F^2 - S^2 - S_p^2). \quad (17.2.13)$$

This is indeed diagonal in the total angular momentum basis  $|n, l, j, f, m_f\rangle$ , with  $F^2 - S^2 - S_p^2 = \hbar^2[f(f+1) - 3/2]$ , where the possible values of  $f$  are 0 and 1. Putting together the factors,

$$\langle H_{\text{hf}} \rangle = \frac{g_p e^2}{2m_e m_p c^2} \frac{8\pi}{3} \frac{\hbar^2}{2} [f(f+1) - 3/2] \frac{1}{\pi a_0^3 n^3} \quad (l=0), \quad (17.2.14)$$

so we arrive at the hyperfine energy correction for  $l = 0$ ,  $j = 1/2$  states,

$$E_{\text{hf}}^{(l=0)} = \langle H_{\text{hf}} \rangle = \frac{g_p m_e}{m_p} \frac{\alpha^2}{n^3} \left( \frac{e^2}{2a_0} \right) \times \begin{cases} 2/3 & \text{for } f = 1, \\ -2 & \text{for } f = 0. \end{cases} \quad (17.2.15)$$

In particular, the hyperfine interaction splits the  $n = 1$  ground state by an amount

$$E_\gamma = E_{\text{hf}}^{(f=1)} - E_{\text{hf}}^{(f=0)} = \frac{8g_p m_e}{3m_p} \alpha^2 \left( \frac{e^2}{2a_0} \right) = 5.878 \times 10^{-6} \text{ eV}. \quad (17.2.16)$$

We had already discussed this in eqs. (12.2.27)–(12.2.29); here we have shown how the numerical prefactor arises. The actual experimental value  $5.87433 \times 10^{-6} \text{ eV}$  is very slightly less than

eq. (17.2.16), due to higher order effects. The  $f = 1$  state is higher in energy than the  $f = 0$  state, and when the former decays to the latter it emits a photon with this energy splitting, whose frequency is  $\nu = \omega/2\pi = E_\gamma/2\pi\hbar = 1.42041 \times 10^9$  Hz, corresponding to a wavelength

$$\lambda = c/\nu = 21.106 \text{ cm.} \quad (17.2.17)$$

This is the “21 centimeter line” of radio astronomy, which comes from a photon emission or absorption between states with different orientations of the electron and proton spins in the hydrogen atom ground state.

Now take the case  $l \neq 0$ . We will use the Landé projection formula eq. (13.2.10) derived in section 13.2 for a general vector operator, given in our present problem by the vector that appears dotted into  $\vec{S}_p$  in eq. (17.2.6),

$$\vec{V} = \frac{1}{R^3}(\vec{L} - \vec{S}) + \frac{3}{R^5}(\vec{R} \cdot \vec{S}) \vec{R}, \quad (17.2.18)$$

so that

$$H_{\text{hf}}^{l \neq 0} = \frac{g_p e^2}{2m_e m_p c^2} \vec{V} \cdot \vec{S}_p. \quad (17.2.19)$$

Here we have used the fact that the  $\delta^{(3)}(\vec{R})$  term does not contribute for  $l \neq 0$ , as noted in the calculation of eq. (17.2.12). Because  $\vec{V}$  itself does not involve  $\vec{S}_p$  at all, for the purposes of evaluating its matrix elements we can temporarily ignore the  $f$  and  $m_f$  quantum numbers when using eqs. (13.2.10) and (13.2.11), and so take the total angular momentum operator to be just  $\vec{J} = \vec{L} + \vec{S}$ , the generator of rotations for the orbital and electron spin degrees of freedom.<sup>†</sup> Also, in our present problem, we are calculating an energy correction as an expectation value, so the degeneracy labels  $\alpha$  and  $\beta$  appearing in eq. (13.2.10) are the same,  $\alpha = (n, l)$ , and the Landé projection formula eq. (13.2.10) gives

$$\langle \alpha, j, m'_j | \vec{V} | \alpha, j, m_j \rangle = \langle j, m'_j | \vec{J} | j, m_j \rangle \frac{\langle \alpha, j | \vec{J} \cdot \vec{V} | \alpha, j \rangle}{\hbar^2 j(j+1)}, \quad (17.2.20)$$

where the reduced matrix element is the scalar expectation value

$$\langle \alpha, j | \vec{J} \cdot \vec{V} | \alpha, j \rangle = \langle \alpha, j, m_j | \vec{J} \cdot \vec{V} | \alpha, j, m_j \rangle. \quad (17.2.21)$$

Recall that, because of Theorem 13.2.1, since  $\vec{J} \cdot \vec{V}$  is a scalar operator,  $\langle \alpha, j | \vec{J} \cdot \vec{V} | \alpha, j \rangle$  does not actually depend on  $m_j$  at all, despite its appearance on the right side of eq. (17.2.21).

---

<sup>†</sup>If  $\vec{V}$  had contained  $\vec{S}_p$ , then the Landé projection formula eq. (13.2.10) would apply only if  $\vec{J}$  were replaced by  $\vec{F}$ , the total angular momentum operator including the proton spin.



To evaluate  $\langle \alpha, j | \vec{J} \cdot \vec{V} | \alpha, j \rangle$  we therefore need the expectation values  $\langle \vec{J} \cdot (\vec{L} - \vec{S}) / R^3 \rangle$  and  $\langle (\vec{R} \cdot \vec{J})(\vec{R} \cdot \vec{S}) / R^5 \rangle$  in a state  $|\alpha, j, m_j\rangle$ . For the first of these, we calculate:

$$\vec{J} \cdot (\vec{L} - \vec{S}) = (\vec{L} + \vec{S}) \cdot (\vec{L} - \vec{S}) = L^2 - S^2 = \hbar^2[l(l+1) - 3/4]. \quad (17.2.22)$$

For the second, we use  $\vec{R} \cdot \vec{L} = \vec{R} \cdot (\vec{R} \times \vec{P}) = 0$  to obtain  $\vec{R} \cdot \vec{J} = \vec{R} \cdot \vec{S}$ , and therefore

$$(\vec{R} \cdot \vec{J})(\vec{R} \cdot \vec{S}) = (\vec{R} \cdot \vec{S})(\vec{R} \cdot \vec{S}) = \frac{\hbar^2}{4}(\vec{R} \cdot \vec{\sigma})(\vec{R} \cdot \vec{\sigma}) = \hbar^2 R^2 / 4, \quad (17.2.23)$$

where we have used the Pauli matrix representation of spin, and then the identity eq. (8.2.18). Putting eqs. (17.2.22) and (17.2.23) into eq. (17.2.21) gives

$$\langle \alpha, j | \vec{J} \cdot \vec{V} | \alpha, j \rangle = \hbar^2 l(l+1) \langle n, l | \frac{1}{R^3} | n, l \rangle \quad (17.2.24)$$

for the operator  $\vec{V}$  defined in eq. (17.2.18). So, we can write for eq. (17.2.19), within matrix elements with common  $n, l, j$ ,

$$H_{\text{hf}}^{l \neq 0} = \frac{g_p e^2}{2m_e m_p c^2} \frac{l(l+1)}{j(j+1)} \vec{J} \cdot \vec{S}_p \frac{1}{R^3}. \quad (17.2.25)$$

In the orthobasis  $|n, l, j, f, m_f\rangle$ , the angular momentum operator factor in eq. (17.2.25) is diagonal, and evaluates to

$$\vec{J} \cdot \vec{S}_p = \frac{1}{2}(F^2 - J^2 - S_p^2) = \frac{\hbar^2}{2}(f(f+1) - j(j+1) - 3/4). \quad (17.2.26)$$

The radial part of the expectation value was found in eq. (11.1.68),

$$\langle 1/R^3 \rangle = \frac{1}{a_0^3 n^3 l(l+1)(l+1/2)}. \quad (17.2.27)$$

Using eqs. (17.2.26) and (17.2.27) to find the expectation value of eq. (17.2.25), and eliminating  $j$  in favor of  $f$ , gives

$$E_{\text{hf}} = \langle H_{\text{hf}} \rangle = \frac{g_p m_e}{m_p} \frac{\alpha^2}{n^3} \left( \frac{e^2}{2a_0} \right) \frac{\pm 1}{(l+1/2)(2f+1)}, \quad (17.2.28)$$

where the  $+$  sign applies if  $f = j + 1/2$  and the  $-$  sign if  $f = j - 1/2$ . Actually, this agrees with the  $l = 0, j = 1/2$  formula in eq. (17.2.15), so that eq. (17.2.28) is an all-purpose result, correct for all states  $|n, l, j, f, m_f\rangle$  in the total angular momentum basis.

Note that the hyperfine energy contribution is independent of the quantum number  $m_f$ . This had to be true, because the energy eigenvalues for the hydrogen atom cannot possibly depend on the orientation of the total angular momentum with respect to an arbitrarily chosen  $\hat{z}$  direction. Therefore, each state with total angular momentum quantum number  $f$  has a remaining exact

degeneracy of  $2f + 1$ . The only way to break this degeneracy is to introduce some external effect that picks out a special direction in space, for example an external electric or magnetic field.

The average hyperfine energy contribution for any given  $n, l, j$  level is 0. Of the  $2(2j + 1)$  states before the hyperfine splitting is taken into account, the  $2j + 2$  states with  $f = j + 1/2$  are raised by an amount multiplied by  $1/(2j + 2)$ , while the  $2j$  states with  $f = j - 1/2$  are lowered by the same amount multiplied by  $1/2j$ .

In terms of the total  $n = 1$  hyperfine splitting  $E_\gamma$  defined in eq. (17.2.16), the hyperfine splittings of the  $n = 2$  states are

$$2S_{1/2}(l = 0, j = 1/2) : \quad E_{\text{hf}} = \begin{cases} E_\gamma/32 & (f = 1) \\ -3E_\gamma/32 & (f = 0) \end{cases} \rightarrow \Delta E_{\text{hf}}^{2S_{1/2}} = E_\gamma/8, \quad (17.2.29)$$

$$2P_{1/2}(l = 1, j = 1/2) : \quad E_{\text{hf}} = \begin{cases} E_\gamma/96 & (f = 1) \\ -E_\gamma/32 & (f = 0) \end{cases} \rightarrow \Delta E_{\text{hf}}^{2P_{1/2}} = E_\gamma/24, \quad (17.2.30)$$

$$2P_{3/2}(l = 1, j = 3/2) : \quad E_{\text{hf}} = \begin{cases} E_\gamma/160 & (f = 2) \\ -E_\gamma/96 & (f = 1) \end{cases} \rightarrow \Delta E_{\text{hf}}^{2P_{3/2}} = E_\gamma/60. \quad (17.2.31)$$

So far, we have evaluated the fine and hyperfine contributions, by working at first order in degenerate perturbation theory. However, there is a second-order effect, called the **Lamb shift** after Willis E. Lamb Jr., which for  $l = 0$  states is numerically larger than the hyperfine effect. The Lamb shift arises due to the electron emitting and reabsorbing virtual photons, quanta of the electromagnetic field. To calculate it in a systematic way requires the toolbox of quantum electrodynamics, including renormalization of ultraviolet divergent contributions and a careful treatment of infrared effects. Understanding these issues played a critical role in the development of relativistic quantum field theory, but here we can only summarize the results numerically. The Lamb shift for atomic hydrogen states  $|n, l, j\rangle$  has been calculated to be, approximately,

$$\Delta E_{\text{Lamb}} = \frac{\alpha^3}{n^3} \left( \frac{e^2}{2a_0} \right) \times \begin{cases} 6.50 & (\text{for } l = 0, j = 1/2), \\ 0.026 \pm \frac{1}{\pi(j + 1/2)(2l + 1)} & (\text{for } l \neq 0, j = l \pm 1/2), \end{cases} \quad (17.2.32)$$

where the decimal coefficients actually have a slight dependence on  $n$ . Note that the Lamb shift is much larger for  $s$ -wave ( $l = 0$ ) states than for  $l \neq 0$  states. The extra factor of  $\alpha$  here (compared to the  $\alpha^2$  factor for fine and hyperfine energy corrections) is indicative of an effect of second order in perturbation theory. Effects that are even higher order in  $\alpha = 1/137.036$  slightly modify each of the fine, hyperfine, and Lamb shift contributions, so each of the energy splittings that we have calculated here cannot be trusted beyond about the per cent level. More heroic calculations, not reviewed here, have been done to improve beyond that accuracy.

The fine, hyperfine, and Lamb shift splitting effects on the  $n = 2$  states of atomic hydrogen are summarized and illustrated in Figure 17.2.1.




Figure 17.2.1: The fine, hyperfine, and Lamb shift energy splittings in  $\mu\text{eV}$  for the  $n = 2$  energy levels of the hydrogen atom, with  $nl_j$  spectroscopic notation where  $l$  is the orbital angular momentum replaced by its (capitalized) letter code as in eq. (11.1.20),  $j$  is the total angular momentum quantum number excluding the proton spin, and  $f$  is the total angular momentum quantum number including the proton spin. Relative energy spacings are indicated qualitatively, but not to scale. The Lamb-shift splitting between the  $2S_{1/2}$  and  $2P_{1/2}$  levels is the experimental one, which differs slightly from the calculation quoted in eq. (17.2.32) due to higher order effects. Each energy level shown has a degeneracy  $2f + 1$ .

### 17.3 Hydrogen atom in external magnetic field (Zeeman and Paschen–Back effects)

Atomic and molecular state properties can be probed using an external magnetic field. In this section, we consider the energy levels of the hydrogen atom in the presence of a uniform constant magnetic field, which is taken without loss of generality to point in the  $\hat{z}$  direction,

$$\vec{B} = B\hat{z}. \quad (17.3.1)$$

This is associated with the Coulomb-gauge vector potential in spherical coordinates,

$$\vec{A} = \frac{1}{2}Br \sin \theta \hat{\phi}. \quad (17.3.2)$$

We wish to apply this in the Hamiltonian for a charged particle moving in an electromagnetic field, eq. (4.3.35), which includes  $\vec{A} \cdot \vec{P}$ . Working in the position representation with  $\vec{P} = -i\hbar\vec{\nabla}$ ,

$$\vec{A} \cdot \vec{P} = \frac{1}{2}B \left( -i\hbar \frac{\partial}{\partial \phi} \right) = \frac{1}{2}BL_z, \quad (17.3.3)$$

where the last equality uses the representation of  $L_z$  of eq. (8.6.7). So, from eq. (4.3.35), the external magnetic field contribution to the electron's Hamiltonian in the hydrogen atom is

$$\Delta H = \frac{eB}{2m_e c} (L_z + g_e S_z) + \frac{e^2 B^2}{8m_e c^2} r^2 \sin^2 \theta. \quad (17.3.4)$$

Let us first get a rough idea of the order of magnitude of these contributions, for typical laboratory magnetic fields. Putting in the numerical values of the constants gives

$$\Delta H = (5.8 \times 10^{-5} \text{ eV}) \frac{1}{\hbar} (L_z + g_e S_z) \frac{B}{\text{Tesla}} + (6.2 \times 10^{-11} \text{ eV}) \frac{r^2}{a_0^2} \sin^2 \theta \left( \frac{B}{\text{Tesla}} \right)^2. \quad (17.3.5)$$

From this, we estimate that the term linear in  $B$  will contribute comparably to the fine-structure effects in atomic hydrogen if  $B$  is of order 1 Tesla. For fields typically accessible in laboratories (of order 10 Tesla or less), the term quadratic in  $B$  is much smaller, so we will neglect it. Taking  $g_e = 2$ , the new Hamiltonian contribution to be considered in this section is therefore

$$H_B = \mu_B B (L_z + 2S_z) / \hbar, \quad (17.3.6)$$

where the Bohr magneton  $\mu_B$  was defined in eq. (4.3.26). Our goal is to understand the energy splitting of hydrogen atomic states due to this Hamiltonian perturbation, as a function of the magnetic field strength.

Let us first consider the weak-field limit, in which  $\mu_B B$  can be treated as a perturbation that is small compared to the fine-structure contributions, but still large compared to the hyperfine energy splittings, which we will neglect. This is called the **Zeeman effect**, after Pieter Zeeman. We apply degenerate first-order perturbation theory to the fine-structure eigenstates  $|n, l, j, m\rangle$  with unperturbed energies

$$E = \frac{e^2}{2a_0} \left[ -\frac{1}{n^2} + \frac{\alpha^2}{n^3} \left( \frac{3}{4n} - \frac{1}{j+1/2} \right) \right]. \quad (17.3.7)$$

Following the general discussion of degenerate perturbation theory in section 15.6, the energy shifts due to  $B$  will be the eigenvalues of the  $(2j+1) \times (2j+1)$  perturbation matrix

$$(H_B)_{m',m} = \frac{\mu_B B}{\hbar} \langle n, l, j, m' | (L_z + 2S_z) | n, l, j, m \rangle, \quad (17.3.8)$$

for each energy level with fixed  $n, l, j$ . (There is no need to consider matrix elements between different values of  $l$ , because the perturbation  $L_z + 2S_z$  cannot connect states with different  $l$ . We will also suppress the common label  $n$  in the following.) Now we write

$$L_z + 2S_z = J_z + S_z, \quad (17.3.9)$$

and replace  $J_z$  by its eigenvalue, using  $\langle l, j, m' | J_z | l, j, m \rangle = \hbar m \delta_{mm'}$ . To finish the calculation, we need the matrix elements  $\langle l, j, m' | S_z | l, j, m \rangle$ . Here, the Landé projection formula of eqs. (13.2.10) and (13.2.11) comes to our aid, with  $\vec{S}$  in the role of  $\vec{V}$ . It tells us that the matrix elements of  $S_z$  are proportional to those of  $J_z$ , with

$$\langle l, j, m' | S_z | l, j, m \rangle = \langle l, j, m' | J_z | l, j, m \rangle \frac{\langle l, j, m | \vec{J} \cdot \vec{S} | l, j, m \rangle}{\hbar^2 j(j+1)}. \quad (17.3.10)$$

Now, we again evaluate  $\langle l, j, m' | J_z | l, j, m \rangle = \hbar m \delta_{mm'}$ , and are delighted to see that the whole  $H_B$  is proportional to  $\delta_{mm'}$ , and so is actually a diagonal matrix on the subspace of degenerate total angular momentum basis states. It remains to calculate the matrix element  $\langle l, j, m | \vec{J} \cdot \vec{S} | l, j, m \rangle$ . This can be done using the standard trick for dot products of angular momenta,

$$\vec{J} \cdot \vec{S} = \frac{1}{2} [J^2 + S^2 - (\vec{J} - \vec{S})^2] = \frac{1}{2} [J^2 + S^2 - L^2], \quad (17.3.11)$$

which is replaced by its eigenvalue,  $\hbar^2[j(j+1) + s(s+1) - l(l+1)]/2$ , when acting on  $|l, j, m\rangle$ .

Putting the ingredients of eq. (17.3.8) together, the energy shifts are

$$\Delta E_B = g \mu_B B m, \quad (17.3.12)$$

where

$$g = \frac{3}{2} + \frac{s(s+1) - l(l+1)}{2j(j+1)} \quad (17.3.13)$$

is called the **Landé  $g$ -factor**. These energy splittings are directly proportional to the magnetic quantum number  $m$  for the total angular momentum. Using the facts that  $s = 1/2$  and  $j = l \pm 1/2$ , we can rewrite the Landé  $g$ -factor for the hydrogen atom as

$$g = \begin{cases} 1 + \frac{1}{2j} & (\text{for } j = l + 1/2), \\ 1 - \frac{1}{2j+2} & (\text{for } j = l - 1/2). \end{cases} \quad (17.3.14)$$

The weak-field Zeeman energy shifts for the lowest few values of  $j$  are therefore

$$\Delta E_{S_{1/2}} = 2\mu_B B m, \quad (m = \pm 1/2), \quad (17.3.15)$$

$$\Delta E_{P_{1/2}} = \frac{2}{3}\mu_B B m, \quad (m = \pm 1/2), \quad (17.3.16)$$

$$\Delta E_{P_{3/2}} = \frac{4}{3}\mu_B B m, \quad (m = \pm 3/2, \pm 1/2), \quad (17.3.17)$$

$$\Delta E_{D_{3/2}} = \frac{4}{5}\mu_B B m, \quad (m = \pm 3/2, \pm 1/2), \quad (17.3.18)$$

$$\Delta E_{D_{5/2}} = \frac{6}{5}\mu_B B m, \quad (m = \pm 5/2, \pm 3/2, \pm 1/2), \quad (17.3.19)$$

etc. By applying a weak magnetic field, the energy degeneracies for each principal quantum number  $n$  are eliminated, and the observed splittings can be used to count the (previously

degenerate) states and deduce their angular momentum quantum numbers. This phenomenon is the historical reason for calling  $m$  the “magnetic quantum number”.

Next, consider the opposite limit, called the **Paschen–Back** regime after Friedrich Paschen and Ernst Back, in which the effect of the external magnetic field is taken to be much stronger than the fine-structure effect, which will be treated as a perturbation. The Hamiltonian  $H_0 = P^2/2m_e - e^2/R$  commutes with the magnetic field Hamiltonian  $H_B$  in eq. (17.3.6), so the exact energy eigenstates of  $H_0 + H_B$  are the product angular momentum states  $|n, l, m_l, m_s\rangle$ , with energy eigenvalues  $E = -e^2/2a_0n^2 + \Delta E_B$ , where

$$\Delta E_B = \mu_B B(m_l + 2m_s). \quad (17.3.20)$$

Now we must apply the fine-structure Hamiltonian to these states as a perturbation. A key point is that when doing so, we cannot use the states  $|n, l, j, m\rangle$  as the unperturbed states, as we did in the previous treatment of the fine structure with a vanishing or weak external magnetic field. This is because the strong magnetic field has eliminated the spherical symmetry and the degeneracy associated with it. Instead, degenerate perturbation theory tells us that the fine-structure energy shifts are obtained as the expectation values of the fine-structure Hamiltonian in the product basis states  $|n, l, m_l, m_s\rangle$ , with their degeneracies for fixed  $m_l + 2m_s$ .

First, consider the easier case that  $l = 0$ . Then the spin-orbit Hamiltonian vanishes, and the fine-structure contribution to the energies are the same as found in eq. (17.1.23) from the  $H_{\text{rel}}$  and  $H_{\text{Darwin}}$  terms. Since  $m_l = 0$  in this case, the combined energy shift due to the strong external  $B$  field and the fine structure is

$$\Delta E_{B, \text{fine}}^{l=0} = 2\mu_B B m_s + \frac{\alpha^2}{n^3} \left( \frac{e^2}{2a_0} \right) \left( \frac{3}{4n} - 1 \right). \quad (17.3.21)$$

Now consider the case  $l \neq 0$ . From the spin-orbit Hamiltonian in eq. (17.1.16), the energy correction from first-order perturbation theory is

$$\langle n, l, m_l, m_s | H_{\text{SO}} | n, l, m_l, m_s \rangle = \frac{e^2}{2m_e^2 c^2} \langle 1/R^3 \rangle \langle l, m_l, m_s | \vec{S} \cdot \vec{L} | l, m_l, m_s \rangle. \quad (17.3.22)$$

The expectation value  $\langle 1/R^3 \rangle$  was already given in eq. (17.1.19). To evaluate the expectation value of  $\vec{S} \cdot \vec{L}$ , it is convenient to use

$$\vec{S} \cdot \vec{L} = \frac{1}{2}(S_+ L_- + S_- L_+) + S_z L_z, \quad (17.3.23)$$

because in the product basis states the expectation value of  $S_z L_z$  evaluates to  $\hbar^2 m_s m_l$ , and the expectation values of  $S_{\pm}$  and  $L_{\pm}$  vanish. (Note the difference from the previously treated spin-orbit case with a weak magnetic field, where one must instead evaluate the expectation value of  $\vec{L} \cdot \vec{S}$  in the total angular momentum basis states.) Combining the spin-orbit result

with the  $H_{\text{rel}}$  contribution from eq. (17.1.15), and recalling from eq. (17.1.22) that the Darwin contribution vanishes for  $l \neq 0$ , we obtain the total first-order perturbation contribution

$$\Delta E_{B,\text{fine}}^{l \neq 0} = \mu_B B(m_l + 2m_s) + \alpha^2 \frac{e^2}{2a_0} \frac{1}{n^3} \left( \frac{m_l m_s}{l(l+1/2)(l+1)} + \frac{3}{4n} - \frac{1}{l+1/2} \right) \quad (17.3.24)$$

from the strong external  $B$  field and the fine-structure effects.

Finally, let us consider the intermediate case in which the effects of the fine structure  $H_{\text{fine}} = H_{\text{rel}} + H_{\text{SO}} + H_{\text{Darwin}}$  and external magnetic field  $H_B$  Hamiltonians are comparable to each other, so that  $H_{\text{fine}} + H_B$  must be treated together as a single perturbation to  $H_0$ . The preceding results for weak  $B$  and for strong  $B$  can be obtained as special-case limits of the following analysis. As before, we will ignore the smaller hyperfine effects. For the unperturbed states, it is not a bad idea to choose<sup>†</sup> the total angular momentum basis elements  $|n, l, j, m\rangle$ , so that the matrix for the fine-structure Hamiltonian  $H_{\text{fine}}$  is diagonal and we can make use of results already found in section 17.1.

For  $l = 0$  states, there is no distinction between  $S_z$  and  $J_z$ , so  $H_B = 2\mu_B B m_s$  is also diagonal in the total angular momentum basis. This means that the energy shift for the  $l = 0$  states is the same as we already found in eq. (17.3.21),

For  $l \neq 0$  states, things are more complicated. Let us consider only the  $n = 2$  level with  $l = 1$ , where the six unperturbed states are

$$|j, m\rangle = \left| \frac{3}{2}, \frac{3}{2} \right\rangle, \left| \frac{3}{2}, \frac{1}{2} \right\rangle, \left| \frac{3}{2}, -\frac{1}{2} \right\rangle, \left| \frac{3}{2}, -\frac{3}{2} \right\rangle, \quad (2P_{3/2}), \quad (17.3.25)$$

$$\left| \frac{1}{2}, \frac{1}{2} \right\rangle, \left| \frac{1}{2}, -\frac{1}{2} \right\rangle, \quad (2P_{1/2}). \quad (17.3.26)$$

The matrix elements of  $L_z + 2S_z$  between any pair of these states can be quickly evaluated using eqs. (12.3.16)–(12.3.21), which give these  $|j, m\rangle$  states in terms of the product orthobasis states  $|m_l, m_s\rangle$ , allowing the operator  $L_z + 2S_z$  to be replaced by  $\hbar(m_l + 2m_s)$ . It is useful to note that the only non-zero matrix elements in the  $|j, m\rangle$  basis are between states with the same  $m$ , because  $L_z + 2S_z$  commutes with  $J_z$ . Therefore, most of the elements of the  $6 \times 6$  matrix of perturbations for the  $2P_{3/2}$  and  $2P_{1/2}$  states vanish. For  $H_B$  in eq. (17.3.6), the only relevant non-zero matrix elements in the  $|j, m\rangle$  basis with positive  $m$  are

$$\left\langle \frac{3}{2}, \frac{3}{2} \right| (L_z + 2S_z) / \hbar \left| \frac{3}{2}, \frac{3}{2} \right\rangle = 2, \quad (17.3.27)$$

$$\left\langle \frac{3}{2}, \frac{1}{2} \right| (L_z + 2S_z) / \hbar \left| \frac{3}{2}, \frac{1}{2} \right\rangle = 2/3, \quad (17.3.28)$$

$$\left\langle \frac{3}{2}, \frac{1}{2} \right| (L_z + 2S_z) / \hbar \left| \frac{1}{2}, \frac{1}{2} \right\rangle = \left\langle \frac{1}{2}, \frac{1}{2} \right| (L_z + 2S_z) / \hbar \left| \frac{3}{2}, \frac{1}{2} \right\rangle = -\sqrt{2}/3, \quad (17.3.29)$$

$$\left\langle \frac{1}{2}, \frac{1}{2} \right| (L_z + 2S_z) / \hbar \left| \frac{1}{2}, \frac{1}{2} \right\rangle = 1/3, \quad (17.3.30)$$

---

<sup>†</sup>It is also possible to choose the product basis eigenstates of  $H_B$  as the unperturbed states. Then the  $H_{\text{fine}}$  matrix elements will be off-diagonal, but the final results for the energy corrections will be the same.

while the corresponding matrix elements with negative  $m$  each have the opposite sign. From eq. (17.1.26),  $H_{\text{fine}}$  is diagonal, with non-zero matrix elements

$$\langle \frac{3}{2}, m | H_{\text{fine}} | \frac{3}{2}, m \rangle = -\frac{1}{64} \alpha^2 \frac{e^2}{2a_0}, \quad (m = \pm 3/2, \pm 1/2), \quad (17.3.31)$$

$$\langle \frac{1}{2}, m | H_{\text{fine}} | \frac{1}{2}, m \rangle = -\frac{5}{64} \alpha^2 \frac{e^2}{2a_0}. \quad (m = \pm 1/2). \quad (17.3.32)$$

We can now put together these results to find the matrix elements of  $W = H_{\text{fine}} + H_B$ . Using the notations

$$a = \frac{\alpha^2 e^2}{64 \cdot 2a_0}, \quad b = \mu_B B \quad (17.3.33)$$

for simplicity, and ordering the basis elements as  $|\frac{3}{2}, \frac{3}{2}\rangle, |\frac{3}{2}, -\frac{3}{2}\rangle, |\frac{3}{2}, \frac{1}{2}\rangle, |\frac{1}{2}, \frac{1}{2}\rangle, |\frac{3}{2}, -\frac{1}{2}\rangle, |\frac{1}{2}, -\frac{1}{2}\rangle$ , the matrix representation for the perturbation is

$$W = \begin{pmatrix} -a+2b & 0 & 0 & 0 & 0 & 0 \\ 0 & -a-2b & 0 & 0 & 0 & 0 \\ 0 & 0 & -a+\frac{2}{3}b & -\frac{\sqrt{2}}{3}b & 0 & 0 \\ 0 & 0 & -\frac{\sqrt{2}}{3}b & -5a+\frac{1}{3}b & 0 & 0 \\ 0 & 0 & 0 & 0 & -a-\frac{2}{3}b & \frac{\sqrt{2}}{3}b \\ 0 & 0 & 0 & 0 & \frac{\sqrt{2}}{3}b & -5a-\frac{1}{3}b \end{pmatrix}. \quad (17.3.34)$$

This contains only  $1 \times 1$  and  $2 \times 2$  non-zero blocks, with eigenvalues

$$\Delta E_{B, \text{fine}}^{n=2, l=1} = -a \pm 2b, \quad (17.3.35)$$

$$-3a + b/2 \pm \sqrt{4a^2 + 2ab/3 + b^2/4}, \quad (17.3.36)$$

$$-3a - b/2 \pm \sqrt{4a^2 - 2ab/3 + b^2/4}. \quad (17.3.37)$$

In the same notation, the  $l = 0$  energy corrections from eq. (17.3.21) are

$$\Delta E_{B, \text{fine}}^{n=2, l=0} = -5a \pm b. \quad (17.3.38)$$

These eight energy corrections to the  $n = 2$  states are graphed in Figure 17.3.1 as a function of the external magnetic field  $B$ , also including the small positive Lamb shift of  $4.375 \mu\text{eV}$  for the  $l = 0$  eigenstates. It is left as an exercise to confirm that expanding eqs. (17.3.35)–(17.3.38) in small  $b/a$  recovers the weak-field limit of eqs. (17.1.26) and (17.3.15)–(17.3.17), and expanding in small  $a/b$  recovers the strong-field limit found in eqs. (17.3.21) and (17.3.24).






Figure 17.3.1: Energy corrections (in  $\mu\text{eV}$ ) to the  $n = 2$  states of atomic hydrogen, due to the fine structure Hamiltonian and an external magnetic field  $B$  (in Tesla), as found in eqs. (17.3.35)–(17.3.38). At small  $B$ , the weak-field Zeeman effect energy eigenstates approach total angular momentum eigenstates  $|l, s, j, m\rangle$ ; the group of upper four states are  $2P_{3/2}$ , and the lower four states are  $2P_{1/2}$  and  $2S_{1/2}$ . The small Lamb shift of  $4.375 \mu\text{eV}$  has also been included for the  $2S_{1/2}$  states, and is barely visible. At large  $B$ , the strong-field (Paschen–Back regime) energy eigenstates approach product angular momentum eigenstates  $|l, s, m_l, m_s\rangle$ , with  $m_l + 2m_s = 2, 1, 0, -1$ , and  $-2$ , from top to bottom.

## 17.4 Exercises

**Exercise 17.1.** Consider the normalized Hydrogen atom stationary state wavefunctions, which are given by  $\psi_{n,l,m}(\vec{r}) = R_{n,l}(r)Y_l^m(\theta, \phi)$ . For general  $n, l, m$ , compute the probability density for the electron to be at the origin,  $|\psi_{n,l,m}(0)|^2$ . [Hint: use eq. (11.1.41).] Use the result to obtain the Darwin term energy correction in eq. (17.1.22).

**Exercise 17.2.** Make a diagram of the relative energy for the  $n = 3$  level of atomic hydrogen, similar to Figure 17.2.1 for the  $n = 2$  level. Don't try to draw the energy splittings to scale, but include numerical values in  $\mu\text{eV}$  for the fine structure, hyperfine structure, and Lamb shift splittings. Use spectroscopic notation to label the states before the hyperfine effect, and label the hyperfine-split states by their  $f$  (grand total angular momentum) quantum numbers.

**Exercise 17.3.** Deuterium, also known as heavy hydrogen, is an atom with one electron and a nucleus consisting of a deuteron (a proton-neutron bound state) with spin 1. The magnetic moment operator of the deuteron is

$$\vec{\mu}_d = \frac{g_d e}{2m_d c} \vec{S}_d, \quad (17.4.1)$$

where  $\vec{S}_d$  is the spin-1 operator,  $m_d = 1875.6 \text{ MeV}/c^2$  is the mass, and  $g_d = 1.713$ . Work out the hyperfine structure of the  $n = 1$  state of deuterium. What are the energy, frequency, and wavelength of the photon corresponding to the hyperfine transition of the ground state?

**Exercise 17.4.** An electron is moving in a spherically symmetric but otherwise generic (no other special properties) potential  $V(r)$ , with a perturbation Hamiltonian of the spin-orbit type,  $W = a\vec{L} \cdot \vec{S}$ , where  $a$  is a positive constant and  $\vec{L}$  is the orbital angular momentum and  $\vec{S}$  is the spin operator. Consider a set of unperturbed states that are degenerate in energy (before including  $W$ ), and have  $L^2$  eigenvalue  $12\hbar^2$ .

(a) Work in the total angular momentum basis  $|j, m_j\rangle$  to answer the following questions. What are the energy corrections due to  $W$ ? What are the remaining degeneracies of the corresponding energy eigenstates?

(b) Repeat part (a), but this time do your work in terms of the matrix elements of  $W$  in the product angular momentum basis  $|m_l, m_s\rangle$ . [Hint: you may want to use eq. (17.3.23).]

(c) Repeat part (a), but this time pretend that you live in a universe where the electron has spin  $3/2$  (instead of spin  $1/2$ , as in our universe).

**Exercise 17.5.** Consider a hydrogen atom in a constant uniform magnetic field. Repeat the analysis leading to eqs. (17.3.35)–(17.3.35) to obtain the energy shifts, but this time for the  $n = 3$  states.




Figure 18.1.1: In elastic scattering of indistinguishable particles (for example electrons)  $1, 2 \rightarrow 3, 4$ , each of the final-state particles labeled 3 and 4 have an equal claim to be the same as each of the initial-state particles 1 and 2 of the same type. It is allowed and useful to label the identical particles, but physical results cannot depend on our labeling choice.

## 18 Identical particles and multi-electron atoms

### 18.1 Intrinsic indistinguishability of identical particles

Particles are said to be **identical** if there is no way of distinguishing them, even in principle. In particular, they must have the same mass, spin magnitude, charge, and any other intrinsic properties. Classically, one can keep track of individual particles, even if they are identical, by following their trajectories, which are determined by the equations of motion. However, in quantum mechanics, this is impossible.

For example, if we scatter two electrons off of each other, as illustrated in Figure 18.1.1, we may choose to label the electrons in the initial state by 1, 2, and the electrons in the final state by 3, 4. However, it has no meaning to say that final-state electron 3 is uniquely the same as the initial-state electron 1. Both of the outgoing electrons 3 and 4 have an equal claim to being the same as the incoming electron 1 (or 2). Although we may choose a labeling scheme, and it is generally quite useful to do so, the physics results cannot depend at all on that choice.

One way to describe a state with  $N$  identical particles is to take a tensor product of  $N$  identical individual state spaces. Suppose that we have an orthobasis of single-particle kets  $|\alpha_n\rangle$ , where each  $\alpha_n$  represents a whole set of CSCO labels for one particle. An orthobasis of kets for the  $N$ -particle Hilbert space is, mathematically,<sup>†</sup>

$$|\alpha_1\rangle \otimes |\alpha_2\rangle \otimes \cdots \otimes |\alpha_N\rangle. \quad (18.1.1)$$

For example, we could take  $\alpha_n = (\vec{r}_n, m_n)$ , and the orthobasis elements would be

$$|\vec{r}_1, m_1\rangle \otimes |\vec{r}_2, m_2\rangle \otimes \cdots \otimes |\vec{r}_N, m_N\rangle, \quad (18.1.2)$$

where the interpretation of  $|\vec{r}_n, m_n\rangle$  is that we are certain to find the particle at position  $\vec{r}_n$  with spin component  $S_z = \hbar m_n$ . Alternatively,  $\alpha_n$  could instead include the momentum  $\vec{p}_n$

---

<sup>†</sup>We say “mathematically” here because the physical  $N$ -identical-particle Hilbert space is a proper subset of the mathematical  $N$ -identical-particle Hilbert space, as we will soon see.

rather than the position, or it could include the energy,  $L^2$ , and  $L_z$  quantum numbers of the  $n$ th particle, or some other choice of CSCO eigenvalues. In any case, we can dispense with the tensor product notation and write eq. (18.1.1) as a single ket for a given orthobasis element,

$$|\alpha_1, \alpha_2, \dots, \alpha_N\rangle. \quad (18.1.3)$$

The fact that the particles are identical means that although we may choose to label them by integers  $n = 1, 2, \dots, N$ , we cannot distinguish them, and so physical results cannot depend on our arbitrary choice of labels. In particular, this means that any observable operator must be unchanged if we exchange any pair of labels.

For example, the Hamiltonian for two identical particles of mass  $m$  can be of the form

$$H = \frac{P_1^2}{2m} + \frac{P_2^2}{2m} + V(\vec{R}_1) + V(\vec{R}_2) + V_{\text{int}}(|\vec{R}_1 - \vec{R}_2|), \quad (18.1.4)$$

since this is invariant under the exchange of labels  $1 \leftrightarrow 2$ . It is important here that the potentials  $V(\vec{r})$  for the individual identical particles 1 and 2 are the same function. The total momentum operator for  $N$  identical particles,

$$\vec{P} = \vec{P}_1 + \vec{P}_2 + \dots + \vec{P}_N, \quad (18.1.5)$$

is likewise an observable, but the individual operators  $\vec{P}_n$  appearing in it are not observables. Similarly, we can define the observable operator for the density at a fixed position  $\vec{r}$  as

$$\rho(\vec{r}) = \sum_{n=1}^N \delta^{(3)}(\vec{r} - \vec{R}_n). \quad (18.1.6)$$

Here  $\vec{r}$  is an ordinary 3-vector that labels which operator we are talking about, and  $\vec{R}_n$  are the individual position operators for the identical particles.

**Pair-exchange operators** swap the labels of two identical particles, and are defined by

$$P_{ij}|\dots, \alpha_i, \dots, \alpha_j, \dots\rangle = |\dots, \alpha_j, \dots, \alpha_i, \dots\rangle. \quad (18.1.7)$$

So, for example,  $P_{13}|\alpha_1, \alpha_2, \alpha_3\rangle = |\alpha_3, \alpha_2, \alpha_1\rangle$  in the special case of a system of three identical particles. Note that  $P_{ij}$  is both unitary and Hermitian (like the parity operator), and  $P_{ij} = P_{ji}$ , and  $(P_{ij})^2 = I$ . It follows from the last property that if  $P_{ij}|\psi\rangle = \lambda|\psi\rangle$ , then  $\lambda^2 = 1$  unless  $|\psi\rangle$  is the null ket, so the only possible eigenvalues of  $P_{ij}$  are  $\lambda = \pm 1$ . Despite being Hermitian,  $P_{ij}$  is not itself an observable, since, for example,  $P_{12}$  is not left unchanged if we exchange identical particles 1 and 3.

Now, since observables must be symmetric under the pair exchange of labels, we have

**Theorem 18.1.1.** *For any observable  $A$ , and any pair-exchange operator  $P_{ij}$  for identical particles,*

$$[P_{ij}, A] = 0. \quad (18.1.8)$$

Proof: write  $A = A(a_1, a_2, \dots, a_N)$ , where the  $a_n$  are operators that act non-trivially only on orthobasis kets  $|\alpha_n\rangle$  for the  $n$ th particle. Since  $A$  must be symmetric under exchange of labels,

$$A(\dots, a_i, \dots, a_j, \dots) = A(\dots, a_j, \dots, a_i, \dots). \quad (18.1.9)$$

It follows that

$$P_{ij}A|\dots, \alpha_i, \dots, \alpha_j, \dots\rangle = A|\dots, \alpha_j, \dots, \alpha_i, \dots\rangle = AP_{ij}|\dots, \alpha_i, \dots, \alpha_j, \dots\rangle, \quad (18.1.10)$$

where the first equality made used of eq. (18.1.9). Since the states  $|\dots, \alpha_i, \dots, \alpha_j, \dots\rangle$  are an orthobasis, the theorem follows by linearity since  $P_{ij}A$  must equal  $AP_{ij}$  acting on every state.  $\square$

In particular, the Hamiltonian must commute with every  $P_{ij}$ ,

$$[H, P_{ij}] = 0. \quad (18.1.11)$$

Now suppose that  $|E\rangle$  is an eigenket of  $H$ , with energy eigenvalue  $E$ . Then,

$$H(P_{ij}|E\rangle) = P_{ij}H|E\rangle = EP_{ij}|E\rangle, \quad (18.1.12)$$

so  $P_{ij}|E\rangle$  is also an eigenket of  $H$  with the same energy eigenvalue  $E$ . This fact sometimes goes by the name of **exchange degeneracy**.

However, while mathematically useful, the exchange degeneracy is really a fake physically, because it only applies to mathematical kets, and not to physical states. The reason is the **spin-statistics principle**: physical states are required to be eigenstates of all exchange operators  $P_{ij}$  for pairs of identical particles, with eigenvalue  $+1$  if the identical particles being exchanged are bosons (have integer spin), and eigenvalue  $-1$  if they are fermions (have spin  $1/2, 3/2, \dots$ ). These are the only two possible eigenvalues, as noted after eq. (18.1.7). Fermions are said to obey **Fermi–Dirac statistics**, and bosons obey **Bose–Einstein statistics**.

The spin-statistics principle is sometimes taken as a postulate, but it can be proved as a theorem in 4-d relativistic quantum field theory, the fundamental quantum mechanical framework that underlies our best understanding of the universe at small distance scales. Since it can be proved in that context, rather than assumed, it was not listed among the postulates of quantum mechanics in section 3.1. On the other hand, if a quantum system is not assumed to obey special relativity, then the spin-statistics principle cannot be proved, but can be adopted

(or not) as an assumption. Alternatively, one can view the spin-statistics principle simply as an experimentally observed fact, subject at any time to potential falsification.

The spin-statistics principle applies not just for the particles that we currently believe to be fundamental, including bosons (photons, gluons,  $W$  and  $Z$  bosons, and Higgs bosons) and fermions (electrons, muons, tau leptons, neutrinos, quarks), but even for composite bosons (for example, H atoms, deuterons,  $^4\text{He}$  nuclei, pions) and composite fermions (protons, neutrons,  $^3\text{He}$  nuclei). Whether a composite particle is a boson or a fermion is determined simply by whether its constituents include an even or odd number of fermions, respectively. For example, a  $^4\text{He}$  nucleus consists of two protons and two neutrons. Doing a pair exchange of two  $^4\text{He}$  nuclei therefore involves the simultaneous exchange of four fermions, resulting in a factor of  $(-1)^4$ . The proton can be viewed as consisting of  $3 + n$  quarks and  $n$  antiquarks, where naively  $n = 0$ , but the presence of virtual particles inside the proton means that  $n$  is actually indeterminate. Nevertheless,  $n$  takes on integer values, so the pair exchange of two protons always results in factor of  $(-1)^{3+n}(-1)^n = -1$ , making the proton a fermion.

The exchange degeneracy described by eq. (18.1.12) is completely eliminated from physical states by the spin-statistics principle. For example, if there are two identical particles, then  $|\alpha_1, \alpha_2\rangle$  and  $|\alpha_2, \alpha_1\rangle$  are not eigenstates of  $P_{12}$  unless  $\alpha_1 = \alpha_2$ . If  $\alpha_1 \neq \alpha_2$ , then using

$$P_{12}|\alpha_1, \alpha_2\rangle = |\alpha_2, \alpha_1\rangle, \quad P_{12}|\alpha_2, \alpha_1\rangle = |\alpha_1, \alpha_2\rangle, \quad (18.1.13)$$

one finds that the linear combinations

$$|\alpha_1, \alpha_2\rangle_S = \frac{1}{\sqrt{2}}(|\alpha_1, \alpha_2\rangle + |\alpha_2, \alpha_1\rangle), \quad (18.1.14)$$

$$|\alpha_1, \alpha_2\rangle_A = \frac{1}{\sqrt{2}}(|\alpha_1, \alpha_2\rangle - |\alpha_2, \alpha_1\rangle), \quad (18.1.15)$$

are normalized eigenkets of  $P_{12}$ , with eigenvalues  $+1$  and  $-1$ , respectively. Therefore, according to the spin-statistics principle, only the linear combination  $|\alpha_1, \alpha_2\rangle_S$  is a physical state if the two particles are identical bosons, and only  $|\alpha_1, \alpha_2\rangle_A$  is a physical state if the two particles are identical fermions. In either case, if  $|\alpha_1, \alpha_2\rangle$  is an energy eigenstate, then exactly one linear combination is physical, and there is no true energy degeneracy associated with the exchange of identical particles. (However, there may be energy degeneracies for other reasons.)

If instead  $\alpha_1 = \alpha_2$ , then  $|\alpha_1, \alpha_1\rangle_S = |\alpha_1, \alpha_1\rangle$  is a physical state if the particles are bosons; note that only the normalization has changed compared to eq. (18.1.14). But if the particles are fermions, then  $|\alpha_1, \alpha_1\rangle_A$  is the null ket, so there is no such physical state. This is the **Pauli exclusion principle**; two identical fermions are not allowed to be in the same state. In particular, they are not allowed to have the same eigenvalues for a CSCO.

To construct physical states that satisfy the spin-statistics principle for  $N > 2$  identical particles, we first construct general **permutation operators**, which are arbitrary products of pair-exchange operators, for example  $P = P_{12}P_{13}P_{45}$ . (Note that the order of the  $P_{ij}$  matters, because they do not always commute.) There are many different ways of writing a given permutation operator as a product in this way, because of identities like  $P_{12}P_{23}P_{12} = P_{13}$ , but the total number of distinct permutation operators is exactly  $N!$ . They can all be classified as odd or even, according to whether the number of pair-exchange operators in the product is even or odd. If  $P$  is an even permutation, we define  $(-1)^P = +1$ , and if  $P$  is an odd permutation, we write  $(-1)^P = -1$ .

We can then define the **symmetrization and antisymmetrization operators** as sums over all  $N!$  distinct permutation operators for  $N$  identical particles,

$$\mathcal{S} = \frac{1}{N!} \sum_P P, \quad (18.1.16)$$

$$\mathcal{A} = \frac{1}{N!} \sum_P (-1)^P P. \quad (18.1.17)$$

These are projection operators onto the subspaces of completely symmetric and antisymmetric kets, respectively, and satisfy the properties  $\mathcal{S}^2 = \mathcal{S}$  and  $\mathcal{A}^2 = \mathcal{A}$  and  $\mathcal{S}\mathcal{A} = \mathcal{A}\mathcal{S} = 0$ . Furthermore, we have  $P_{ij}\mathcal{S} = \mathcal{S}$  and  $P_{ij}\mathcal{A} = -\mathcal{A}$  for all  $i, j$ , so that any ket acted on by  $\mathcal{S}$  will be an eigenstate of every  $P_{ij}$  with eigenvalue 1, and any ket acted on by  $\mathcal{A}$  will be an eigenstate of every  $P_{ij}$  with eigenvalue  $-1$ .

It follows that the physical orthobasis states for  $N$  identical fermions are proportional to  $\mathcal{A}|\alpha_1, \alpha_2, \dots, \alpha_N\rangle$ ,

$$|\alpha_1, \alpha_2, \dots, \alpha_N\rangle_A = \frac{1}{\sqrt{N!}} \sum_P (-1)^P P|\alpha_1, \alpha_2, \dots, \alpha_N\rangle. \quad (18.1.18)$$

These are the simultaneous eigenstates for every pair-exchange operator  $P_{ij}$  with eigenvalue  $-1$ , unique up to normalization, that one can build out of the 1-particle orthobasis states  $|\alpha_i\rangle$ . The prefactor has been chosen to make the ket have unit norm if the  $|\alpha_i\rangle$  are orthonormal and all distinct from each other. An equivalent way to write this is the **Slater determinant**, named after John C. Slater,

$$|\alpha_1, \alpha_2, \dots, \alpha_N\rangle_A = \frac{1}{\sqrt{N!}} \begin{vmatrix} |\alpha_1\rangle_1 & |\alpha_1\rangle_2 & \dots & |\alpha_1\rangle_N \\ |\alpha_2\rangle_1 & |\alpha_2\rangle_2 & \dots & |\alpha_2\rangle_N \\ \vdots & \vdots & \ddots & \vdots \\ |\alpha_N\rangle_1 & |\alpha_N\rangle_2 & \dots & |\alpha_N\rangle_N \end{vmatrix}, \quad (18.1.19)$$

where the products obtained from the determinant are to be understood as

$$|\alpha_i\rangle_1 |\alpha_j\rangle_2 \dots |\alpha_k\rangle_N = |\alpha_i, \alpha_j, \dots, \alpha_k\rangle. \quad (18.1.20)$$

If any two of the  $\alpha_i$  are the same, then the determinant vanishes, and there is no corresponding physical state at all. This is the more general version of the Pauli exclusion principle.

Similarly, the physical orthobasis states for  $N$  identical bosons are proportional to completely symmetrized kets  $\mathcal{S}|\alpha_1, \alpha_2, \dots, \alpha_N\rangle$ , because these are the simultaneous eigenstates for all of the pair-exchange operators  $P_{ij}$  with eigenvalue  $+1$ , unique up to normalization, that one can build out of the 1-particle orthobasis states. Here, the normalization factor is complicated slightly by the fact that for bosons the  $\alpha_i$  can be the same. If we write  $n_i$  for the number of times each distinct  $\alpha_i$  is repeated in the list, and take the  $|\alpha_i\rangle$  to be orthonormal, then the normalized physical orthobasis states are

$$|\alpha_1, \alpha_2, \dots, \alpha_N\rangle_S = \sqrt{\frac{N!}{n_1!n_2!\dots n_N!}} \mathcal{S}|\alpha_1, \alpha_2, \dots, \alpha_N\rangle. \quad (18.1.21)$$

This can also be written as

$$|\alpha_1, \alpha_2, \dots, \alpha_N\rangle_S = \sqrt{\frac{n_1!n_2!\dots n_N!}{N!}} \sum'_P P|\alpha_1, \alpha_2, \dots, \alpha_N\rangle, \quad (18.1.22)$$

where  $\sum'_P$  means that the  $n_1!n_2!\dots n_N!$  redundancies in the sum are eliminated, by including each distinct ket  $P|\alpha_1, \alpha_2, \dots, \alpha_N\rangle$  only once.

For example, consider  $N = 3$  identical particles. Let us construct the physical states that can be made using a finite number  $n$  of orthobasis states for the individual particles,  $|\alpha_i\rangle$  with  $i = 1, 2, \dots, n$ . If the three identical particles are bosons, then we have  $n(n+1)(n+2)/6$  physical orthobasis states, consisting of  $n(n-1)(n-2)/6$  states with distinct  $i, j, k$ ,

$$\begin{aligned} |\alpha_i, \alpha_j, \alpha_k\rangle_S &= \frac{1}{\sqrt{6}} \left( |\alpha_i, \alpha_j, \alpha_k\rangle + |\alpha_i, \alpha_k, \alpha_j\rangle + |\alpha_j, \alpha_k, \alpha_i\rangle + |\alpha_j, \alpha_i, \alpha_k\rangle \right. \\ &\quad \left. + |\alpha_k, \alpha_i, \alpha_j\rangle + |\alpha_k, \alpha_j, \alpha_i\rangle \right), \end{aligned} \quad (18.1.23)$$

and  $n(n-1)$  orthobasis states with distinct  $i$  and  $j$ ,

$$|\alpha_i, \alpha_i, \alpha_j\rangle_S = \frac{1}{\sqrt{3}} \left( |\alpha_i, \alpha_i, \alpha_j\rangle + |\alpha_i, \alpha_j, \alpha_i\rangle + |\alpha_j, \alpha_i, \alpha_i\rangle \right), \quad (18.1.24)$$

and  $n$  orthobasis states,

$$|\alpha_i, \alpha_i, \alpha_i\rangle_S = |\alpha_i, \alpha_i, \alpha_i\rangle. \quad (18.1.25)$$

Each of these is an eigenstate of each of  $P_{12}$ ,  $P_{13}$ , and  $P_{23}$ , with eigenvalue  $+1$ . For fermions, there are only the  $n(n-1)(n-2)/6$  orthobasis states,

$$\begin{aligned} |\alpha_i, \alpha_j, \alpha_k\rangle_A &= \frac{1}{\sqrt{6}} \left( |\alpha_i, \alpha_j, \alpha_k\rangle - |\alpha_i, \alpha_k, \alpha_j\rangle + |\alpha_j, \alpha_k, \alpha_i\rangle - |\alpha_j, \alpha_i, \alpha_k\rangle \right. \\ &\quad \left. + |\alpha_k, \alpha_i, \alpha_j\rangle - |\alpha_k, \alpha_j, \alpha_i\rangle \right) \end{aligned} \quad (18.1.26)$$



for distinct  $i, j, k$ .

An alternative way of specifying the orthobasis states for  $N$  identical particles is by giving the **occupation number** for each of the 1-particle orthobasis states  $|\alpha_i\rangle$ . The occupation number is simply the number of times  $n_i$  that each  $\alpha_i$  appears in the symmetrized or antisymmetrized state ket. For bosons, the occupation numbers are just the same as the numbers  $n_1, n_2, \dots$  appearing in eq. (18.1.22). For fermions, each occupation number is always 0 or 1, due to the Pauli exclusion principle. Then, in both cases, the orthobasis states can be written using the occupation-number notation  $|n_1, n_2, n_3, \dots\rangle$ , where some ordering has been chosen for the  $\alpha_i$ . The individual occupation numbers can change with time, due to interactions. However, we will usually treat the total of the occupation numbers as fixed at  $\sum_i n_i = N$ . (Quantum field theory is what happens to you when you allow  $N$  to be variable.)

So far, we have considered orthobasis states for  $N$  identical particles, built out of the orthobasis states  $|\alpha_i\rangle$  for the individual particles. It is crucial that the resulting states are indeed an orthobasis for the full Hilbert state space of  $N$  identical particles, whether they are bosons or fermions. This means that if we define any  $N$  linear combinations of 1-particle states

$$|\phi_j\rangle = \sum_i c_{ji} |\alpha_i\rangle, \quad (j = 1, \dots, N), \quad (18.1.27)$$

which may be taken to have unit norm but are not necessarily the elements of an orthobasis, then the allowed physical states for identical fermions can be written as

$$|\phi_1, \phi_2, \dots, \phi_N\rangle_A = \mathcal{A}|\phi_1, \phi_2, \dots, \phi_N\rangle, \quad (18.1.28)$$

and the allowed physical states for identical bosons are

$$|\phi_1, \phi_2, \dots, \phi_N\rangle_S = \mathcal{S}|\phi_1, \phi_2, \dots, \phi_N\rangle, \quad (18.1.29)$$

up to normalization. For fermions, a general Slater determinant ket

$$|\phi_1, \phi_2, \dots, \phi_N\rangle_A = \begin{vmatrix} |\phi_1\rangle_1 & |\phi_1\rangle_2 & \cdots & |\phi_1\rangle_N \\ |\phi_2\rangle_1 & |\phi_2\rangle_2 & \cdots & |\phi_2\rangle_N \\ \vdots & \vdots & \ddots & \vdots \\ |\phi_N\rangle_1 & |\phi_N\rangle_2 & \cdots & |\phi_N\rangle_N \end{vmatrix} \quad (18.1.30)$$

is physical (provided that it is not null), and can be shown to be a unique linear combination of the orthobasis states  $|\alpha_1, \alpha_2, \dots, \alpha_N\rangle_A$  defined by eq. (18.1.19). Here, the states  $|\phi_i\rangle$  must be linearly independent, because otherwise the Slater determinant will vanish. Similarly, if the identical particles are bosons, then the state

$$|\phi_1, \phi_2, \dots, \phi_N\rangle_S = \sum_P P|\phi_1, \phi_2, \dots, \phi_N\rangle, \quad (18.1.31)$$

is an allowed physical state, even if the  $|\phi_i\rangle$  are not orthobasis states or even linearly independent, and any state of this form can be written in a unique way as a linear combination of the orthobasis states  $|\alpha_1, \alpha_2, \dots, \alpha_N\rangle_S$  defined by eq. (18.1.22).

Consider  $N$  identical particles, each of which has a single-particle Hamiltonian of the form

$$H_i = \frac{P_i^2}{2m} + V(\vec{R}_i). \quad (18.1.32)$$

For now, we are neglecting any kind of Hamiltonian interaction between the particles [like  $V_{\text{int}}$  in eq. (18.1.4)]. Suppose that the single-particle energy eigenstate position wavefunctions (neglecting spin) are

$$\phi_0(\vec{r}), \phi_1(\vec{r}), \phi_2(\vec{r}), \phi_3(\vec{r}), \dots, \quad (18.1.33)$$

with individual  $H_i$  energy eigenvalues

$$\mathcal{E}_0 < \mathcal{E}_1 \leq \mathcal{E}_2 \leq \mathcal{E}_3 \leq \dots. \quad (18.1.34)$$

We would like to now consider the eigenstates of the total unperturbed Hamiltonian  $H_0 = \sum_i H_i$ . Interactions between the particles can be treated later as a perturbation.

If  $N = 2$ , then the ground state has  $E = 2\mathcal{E}_0$ , with wavefunction

$$\psi(\vec{r}_1, \vec{r}_2) = \phi_0(\vec{r}_1)\phi_0(\vec{r}_2). \quad (18.1.35)$$

Since this position-wavefunction part of the state is symmetric under exchange of the labels 1, 2, the spin state must also be symmetric if the particles are bosons, and the spin state must be antisymmetric if the particles are fermions. Call the spins  $\vec{S}_1$  and  $\vec{S}_2$ , so that the total spin operator is  $\vec{S} = \vec{S}_1 + \vec{S}_2$ , and denote the eigenvalue of the operator  $S^2$  as  $\hbar^2 s(s+1)$  where  $s$  is the total-spin quantum number. For spin-0 bosons, the total-spin quantum number is of course  $s = 0$ , and for spin-1 bosons it must be either  $s = 0$  or  $s = 2$ , because of the addition of angular momentum rule

$$1 \otimes 1 = 0_S \oplus 1_A \oplus 2_S. \quad (18.1.36)$$

For spin-1/2 fermions, the spin state must be  $s = 0$ , the antisymmetric combination in

$$\frac{1}{2} \otimes \frac{1}{2} = 0_A \oplus 1_S. \quad (18.1.37)$$

We will explore this in more detail in the next two sections.

For any number  $N \geq 2$  bosons, one can have a completely symmetric position wavefunction

$$\psi(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N) = \phi_0(\vec{r}_1)\phi_0(\vec{r}_2)\phi_0(\vec{r}_3) \cdots \phi_0(\vec{r}_N), \quad (18.1.38)$$

together with a symmetric spin state, with a total spin quantum number that is an even integer, due to eq. (12.4.44). To minimize the energy, it is therefore favorable to simply maximize the occupation number of the single-particle ground state, forming a **Bose–Einstein condensate** with ground-state energy eigenvalue

$$E = N\mathcal{E}_0. \quad (18.1.39)$$

The ground state in the occupation number notation is  $|N, 0, 0, \dots\rangle$ , where the ordering of 1-particle state labels is the same as for the energy eigenvalues.

However, for  $N \geq 2$  spin-1/2 fermions, at most two can be put into each spatial wavefunction, due to the Pauli exclusion principle. So, one can assign two fermions to  $\phi_0$  (with  $s = 0$  for their antisymmetric combined spin state), two more to  $\phi_1$  (again with  $s = 0$  for their spin state) and so on. If  $N$  is even, then the last two fermions in an  $s = 0$  state will occupy the  $\phi_{\frac{N}{2}-1}$  wavefunction, while if  $N$  is odd then the last electron will have wavefunction  $\phi_{\frac{N-1}{2}}$ . The total unperturbed energy for  $N$  fermions is therefore

$$E = \begin{cases} 2\mathcal{E}_0 + 2\mathcal{E}_1 + \dots + 2\mathcal{E}_{\frac{N}{2}-1} & (\text{even } N), \\ 2\mathcal{E}_0 + 2\mathcal{E}_1 + \dots + 2\mathcal{E}_{\frac{N-3}{2}} + \mathcal{E}_{\frac{N-1}{2}} & (\text{odd } N). \end{cases} \quad (18.1.40)$$

One application of this counting is to the shell model for electrons in atoms. If one ignores the electron-electron Coulomb repulsion, fine, and hyperfine interactions, and just considers electrons as moving in a central potential, then each single particle state with eigenvalue  $E_{n,l,m}$  for  $m = -l, \dots, l$  can be occupied by 2 electrons in an  $s = 0$  state. So, in the ground state of the multi-electron atom, the unperturbed energy level  $E_{n,l}$  can contain up to  $2(2l+1)$  electrons. We will explore this in more detail in section 18.4.

## 18.2 Wavefunctions and spin for two identical particles

In practice, the spin degrees of freedom often play a special role in specifying the orthobasis of states for identical particles. As an example, consider a system of two electrons. We can write the orthobasis kets as

$$|\alpha_1, \alpha_2\rangle = |\vec{r}_1, m_1, \vec{r}_2, m_2\rangle, \quad (18.2.1)$$

with possible values  $\pm 1/2$  for each of the spin magnetic quantum numbers  $m_1$  and  $m_2$ . The total spin is  $\vec{S} = \vec{S}_1 + \vec{S}_2$ , and the operator  $S^2$  has eigenvalues  $\hbar^2 s(s+1)$  where  $s = 0, 1$ , while the  $S_z$  eigenvalue  $\hbar m_S$  can have  $m_S = 0$  (if  $s = 0$ ) and  $m_S = 1, 0, -1$  (if  $s = 1$ ). The state can be specified in a hybrid form, with wavefunctions for the spatial degrees of freedom and kets for the spin degrees of freedom, as

$$\psi_{0,0}(\vec{r}_1, \vec{r}_2) \chi_{0,0} + \sum_{m_S=-1}^1 \psi_{1,m_S}(\vec{r}_1, \vec{r}_2) \chi_{1,m_S}. \quad (18.2.2)$$

Here  $\chi_{s,m_S}$  is the orthobasis ket for a state with total spin quantum numbers  $(s, m_S)$ , so that in terms of the product orthobasis kets,

$$\chi_{0,0} = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle), \quad (18.2.3)$$

$$\chi_{1,1} = |\uparrow\uparrow\rangle, \quad (18.2.4)$$

$$\chi_{1,0} = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle), \quad (18.2.5)$$

$$\chi_{1,-1} = |\downarrow\downarrow\rangle. \quad (18.2.6)$$

The pair-exchange operator separates into spatial and spin parts, as

$$P_{12} = P_{12}^{\text{spatial}} P_{12}^{\text{spin}}. \quad (18.2.7)$$

The total spin singlet ( $s = 0$ ) state is antisymmetric (eigenvalue  $-1$  for  $P_{12}^{\text{spin}}$ ), while the total spin triplet ( $s = 1$ ) states are each symmetric (eigenvalue  $+1$  for  $P_{12}^{\text{spin}}$ ). Since the total state must be antisymmetric to describe identical fermions, we must have

$$\psi_{0,0}(\vec{r}_1, \vec{r}_2) = \psi_{0,0}(\vec{r}_2, \vec{r}_1), \quad (18.2.8)$$

$$\psi_{1,m_S}(\vec{r}_1, \vec{r}_2) = -\psi_{1,m_S}(\vec{r}_2, \vec{r}_1). \quad (18.2.9)$$

Thus  $\psi_{0,0}$  is a symmetric spatial wavefunction for the total spin singlet, and  $\psi_{1,m_S}$  are three distinct antisymmetric spatial wavefunctions for the total spin triplets. In the context of the spatial wavefunctions, “symmetric” (or “antisymmetric”) means eigenvalue  $+1$  (or  $-1$ ) for  $P_{12}^{\text{spatial}}$ . If the operator  $S^2$  commutes with the Hamiltonian  $H$ , then they have common eigenstates, so one can label the energy levels as either  $s = 1$  or  $s = 0$ .

Suppose the two electrons are in a potential  $V(\vec{r})$  that is large in magnitude compared to their Coulomb interaction with each other and any spin interactions, which we therefore neglect. Then, in that approximation,

$$H = \frac{1}{2m_e}(P_1^2 + P_2^2) + V(\vec{R}_1) + V(\vec{R}_2), \quad (18.2.10)$$

which could be the unperturbed Hamiltonian in a perturbative approach, and the Schrödinger equation for the wavefunction  $\psi(\vec{r}_1, \vec{r}_2)$  is

$$\left[ -\frac{\hbar^2}{2m_e}(\nabla_1^2 + \nabla_2^2) + V(\vec{r}_1) + V(\vec{r}_2) - E \right] \psi = 0. \quad (18.2.11)$$

Let  $\phi_a(\vec{r})$  be the wavefunction solutions to the single-particle eigenvalue differential equation

$$\left[ -\frac{\hbar^2}{2m_e}\nabla^2 + V(\vec{r}) - \mathcal{E}_a \right] \phi_a(\vec{r}) = 0. \quad (18.2.12)$$

Then, there are spin singlets (but not spin triplets) with energies  $E = 2\mathcal{E}_a$ , and total spatial wavefunctions

$$\psi_{0,0}(\vec{r}_1, \vec{r}_2) = \phi_a(\vec{r}_1)\phi_a(\vec{r}_2). \quad (18.2.13)$$

This will apply, for example, if  $\mathcal{E}_a$  is the non-degenerate lowest eigenvalue for eq. (18.2.12), so that, as already noted in the previous section, the ground state of the two-electron system is a spin singlet state. Also, for each pair of distinct eigenfunctions  $\phi_a(\vec{r})$  and  $\phi_b(\vec{r})$  with eigenvalues  $\mathcal{E}_a$  and  $\mathcal{E}_b$ , we have four energy eigenstates with  $E = \mathcal{E}_a + \mathcal{E}_b$ , with spatial wavefunctions

$$\psi_{0,0}(\vec{r}_1, \vec{r}_2) = \frac{1}{\sqrt{2}} [\phi_a(\vec{r}_1)\phi_b(\vec{r}_2) + \phi_b(\vec{r}_1)\phi_a(\vec{r}_2)], \quad (s = 0), \quad (18.2.14)$$

$$\psi_{1,m_S}(\vec{r}_1, \vec{r}_2) = \frac{1}{\sqrt{2}} [\phi_a(\vec{r}_1)\phi_b(\vec{r}_2) - \phi_b(\vec{r}_1)\phi_a(\vec{r}_2)], \quad (s = 1, m_S = 1, 0, -1), \quad (18.2.15)$$

which are total spin singlets and triplets, respectively. The eigenvalues  $\mathcal{E}_a$  may have degeneracies, so that in the notation we are using here,  $\mathcal{E}_a$  and  $\mathcal{E}_b$  could be equal even if  $\phi_a(\vec{r})$  and  $\phi_b(\vec{r})$  are distinct eigenfunctions.

For each of these possibilities, consider the probability to find one electron within a volume  $d^3\vec{r}_1$  near  $\vec{r}_1$  and the other within a volume  $d^3\vec{r}_2$  near  $\vec{r}_2$ . From Postulate 4 in section 3.1,

$$d\mathcal{P}(\vec{r}_1, \vec{r}_2) = d^3\vec{r}_1 d^3\vec{r}_2 |\psi(\vec{r}_1, \vec{r}_2)|^2, \quad (18.2.16)$$

where

$$|\psi(\vec{r}_1, \vec{r}_2)|^2 = \frac{1}{2} (|\phi_a(\vec{r}_1)|^2|\phi_b(\vec{r}_2)|^2 + |\phi_b(\vec{r}_1)|^2|\phi_a(\vec{r}_2)|^2 \pm 2\text{Re}[\phi_a(\vec{r}_1)\phi_b^*(\vec{r}_1)\phi_b(\vec{r}_2)\phi_a^*(\vec{r}_2)]), \quad (18.2.17)$$

with the  $+$  sign for spin-singlet states and the  $-$  sign for spin-triplet states. Note that for either sign, eq. (18.2.17) is invariant under the exchange  $\vec{r}_1 \leftrightarrow \vec{r}_2$ .

The  $\pm$  term in eq. (18.2.17) is known as the **exchange density**. As a consequence of it, when the electrons are in the spin-triplet state, they avoid each other. To see this, note that the probability density vanishes when  $\vec{r}_1 = \vec{r}_2$ , and therefore is small (by continuity) when  $\vec{r}_1 \approx \vec{r}_2$ . Conversely, in the spin-singlet state, the two electrons “attract”; there is an enhanced probability density for  $\vec{r}_1 \approx \vec{r}_2$  due to constructive interference. This attraction and repulsion is not due to any electromagnetic interaction or other Hamiltonian interaction between the identical fermions, but rather due to the Fermi–Dirac statistics.

Electrons in Argonne are identical to those in Batavia. So, it is natural to wonder why do we not need to worry about significant effects due to correlation effects from antisymmetrization of wavefunctions belonging to very distant electrons. Suppose that we define normalized single-particle stationary-state wavefunctions that are peaked in Argonne and Batavia, with very little

overlap, so that  $\phi_A(\vec{r})$  is insignificant outside of a small volume somewhere in Argonne, and  $\phi_B(\vec{r})$  is insignificant outside of a small volume in Batavia, and therefore

$$\phi_A(\vec{r})\phi_B(\vec{r}) \approx 0 \quad (18.2.18)$$

everywhere. The total wavefunction is

$$\psi(\vec{r}_1, \vec{r}_2) = \frac{1}{\sqrt{2}} [\phi_A(\vec{r}_1)\phi_B(\vec{r}_2) \pm \phi_B(\vec{r}_1)\phi_A(\vec{r}_2)], \quad (18.2.19)$$

where the  $+$  sign applies for spin singlets and the  $-$  sign for spin triplet states. Note that there is no sense in which the electron labeled 1 is nearer to Argonne, or nearer to Batavia; it is always equally likely to be found in either place.

Now we can ask: what is the probability of seeing an electron within an infinitesimal volume  $d^3\vec{r}$  near  $\vec{r}$ ? The answer is

$$d\mathcal{P}(\vec{r}) = d^3\vec{r} \int d^3\vec{r}_2 |\psi(\vec{r}, \vec{r}_2)|^2 + d^3\vec{r} \int d^3\vec{r}_1 |\psi(\vec{r}_1, \vec{r})|^2. \quad (18.2.20)$$

The first term is the probability that the electron labeled 1 is near  $\vec{r}$ , summed over all possible electron 2 positions, and the second term is the same with  $1 \leftrightarrow 2$ . This can be rewritten as

$$\begin{aligned} d\mathcal{P}(\vec{r}) = & d^3\vec{r} \left( |\phi_A(\vec{r})|^2 \int d^3\vec{x} |\phi_B(\vec{x})|^2 + |\phi_B(\vec{r})|^2 \int d^3\vec{x} |\phi_A(\vec{x})|^2 \right. \\ & \left. \pm 2\text{Re} \left[ \phi_A(\vec{r})\phi_B(\vec{r})^* \int d^3\vec{x} \phi_A(\vec{x})^*\phi_B(\vec{x}) \right] \right). \end{aligned} \quad (18.2.21)$$

The important point is that the last term with  $\pm$  is (doubly!) negligible, because of eq. (18.2.18). In each of the first two terms, the integral is 1 by the assumed normalization, so

$$d\mathcal{P}(\vec{r}) = d^3\vec{r} (|\phi_A(\vec{r})|^2 + |\phi_B(\vec{r})|^2) \quad (18.2.22)$$

to a very good approximation. This shows that, despite the form of the wavefunction in eq. (18.2.19), and despite the fact that our labeling 1, 2 of the electrons did not distinguish either one as being “the electron in Batavia”, the probability density decouples into two separate contributions with negligible interference, for sufficiently separated identical particle wavefunctions. It is the lack of wavefunction overlap for single-particle (approximate) stationary states that enforces the decoupling, not the electron labels. The same type of argument applies to the decoupling of expectation values of observables, including the Hamiltonian. When in Batavia, we can usually ignore electrons in Argonne, and vice versa.

Suppose that two identical particles are bosons; let us consider the cases of spin 0 and spin 1. For spin 0, the spin degrees of freedom are trivial, and in place of eq. (18.2.2) we have simply a symmetric wavefunction

$$\psi_{0,0}(\vec{r}_1, \vec{r}_2) = \psi_{0,0}(\vec{r}_2, \vec{r}_1). \quad (18.2.23)$$

The case of spin 1 bosons is more interesting, because the two individual spins can combine to form nine different total spin states  $\chi_{s,m_S}$ , with  $s = 0, m_S = 0$  and  $s = 1, m_S = 1, 0, -1$  and  $s = 2, m_S = 2, 1, 0, -1, -2$ . By the addition of angular momentum method discussed in section 12.4, one can write these in terms of the product orthobasis kets  $|m_{s_1} : m_{s_2}\rangle$ , where the eigenvalues of the individual spin operators  $S_{1z}$  and  $S_{2z}$  are  $\hbar m_{s_1}$  and  $\hbar m_{s_2}$ , with  $m_{s_1}, m_{s_2} = -1, 0, 1$ . The total spin  $S^2, S_z$  eigenstates are, from eqs. (12.4.34)–(12.4.42),

$$\chi_{2,\pm 2} = |\pm 1 : \pm 1\rangle, \quad (18.2.24)$$

$$\chi_{2,\pm 1} = \frac{1}{\sqrt{2}}(|\pm 1 : 0\rangle + |0 : \pm 1\rangle), \quad (18.2.25)$$

$$\chi_{2,0} = \frac{1}{\sqrt{6}}(|1 : -1\rangle + 2|0 : 0\rangle + |-1 : 1\rangle), \quad (18.2.26)$$

$$\chi_{1,\pm 1} = \frac{1}{\sqrt{2}}(|\pm 1 : 0\rangle - |0 : \pm 1\rangle), \quad (18.2.27)$$

$$\chi_{1,0} = \frac{1}{\sqrt{2}}(|1 : -1\rangle - |-1 : 1\rangle), \quad (18.2.28)$$

$$\chi_{0,0} = \frac{1}{\sqrt{3}}(|1 : -1\rangle - |0 : 0\rangle + |-1 : 1\rangle). \quad (18.2.29)$$

These expressions show that  $\chi_{s,m_S}$  has  $P_{12}^{\text{spin}}$  eigenvalue equal to  $(-1)^s$ . This same fact can be expressed in the shorthand form for addition of angular momentum

$$1 \otimes 1 = 2_S \oplus 1_A \oplus 0_S. \quad (18.2.30)$$

where the subscripts  $S, A$  indicate symmetry or antisymmetry under exchange of the two spins. Therefore, the total state for two identical spin-1 bosons is, again in a hybrid notation with wavefunctions for spatial degrees of freedom and kets for spin degrees of freedom,

$$\Psi = \psi_{0,0}(\vec{r}_1, \vec{r}_2)\chi_{0,0} + \sum_{m_S=-1}^1 \psi_{1,m_S}(\vec{r}_1, \vec{r}_2)\chi_{1,m_S} + \sum_{m_S=-2}^2 \psi_{2,m_S}(\vec{r}_1, \vec{r}_2)\chi_{2,m_S}. \quad (18.2.31)$$

This contains nine component wavefunctions satisfying

$$\psi_{s,m_S}(\vec{r}_1, \vec{r}_2) = (-1)^s \psi_{s,m_S}(\vec{r}_2, \vec{r}_1). \quad (18.2.32)$$

The position wavefunction for the total spin 0 and 2 states are symmetric, and the position wavefunctions for total spin 1 are antisymmetric, under exchange of the boson labels.

### 18.3 Excited states of the helium atom

The binding energy of the ground state of the two-electron helium atom was studied in sections 15.3 and 16.4. In doing so, we ignored the spin degrees of freedom, as well as the fact that the

electrons are identical particles. This is now seen to be justified by the fact that the electrons can arrange themselves into a total spin  $s = 0$  state. Then the total spatial wavefunction is the symmetric  $\psi_{0,0}(\vec{r}_1, \vec{r}_2) = \psi_{0,0}(\vec{r}_2, \vec{r}_1)$  appearing in eqs. (18.2.2), (18.2.8), and (18.2.13), just as we had assumed (without bothering to justify it in detail) in sections 15.3 and 16.4. As we found in section 16.4, the helium ground state spatial wavefunction is approximately just that of the product of two identical  $1s$  hydrogen-like states with  $Z_{\text{eff}} \approx 1.7$ . The ground state of helium does not have a counterpart with total spin  $s = 1$ , because that would force the spatial wavefunction to be antisymmetric under exchange of the two electron levels, which is inconsistent with it being the product of two identical lowest-energy single particle states.

However, for the excited states of helium, we must explicitly take into account both spin and the Fermi–Dirac statistics of the electrons. The Hamiltonian is once again given by eqs. (15.3.1)–(15.3.3). We can start by neglecting the  $H_{12} = e^2/|\vec{r}_1 - \vec{r}_2|$  interaction Hamiltonian, and later include it as a perturbation. Consider the spatial wavefunctions that can be formed from the individual  $Z = 2$  hydrogen-like states, with one electron in the ground state ( $1s$ ) and the other in a higher level,  $2s$  or  $2p$  or  $3s$  or  $3p$  or  $3d$  or  $4s$ , etc.:

$$\psi(\vec{r}_1, \vec{r}_2) = \frac{1}{\sqrt{2}} [\phi_{1,0,0}(\vec{r}_1)\phi_{n,l,m}(\vec{r}_2) \pm \phi_{1,0,0}(\vec{r}_2)\phi_{n,l,m}(\vec{r}_1)]. \quad (18.3.1)$$

Here, if the  $+$  sign applies, then the spatial wavefunction is symmetric, and so the total-spin state must have  $s = 0$ ; these are called para-helium states. (The ground state is also a para-helium state.) Conversely, if the  $-$  sign applies, then the spatial wavefunction is antisymmetric, and so the total-spin state must be symmetric with  $s = 1$ ; these are called ortho-helium states. For both of these, the unperturbed energy eigenvalues of  $H_0 = H_1 + H_2$  are

$$\mathcal{E}_{n,l,m} = -\frac{Z^2 e^2}{2a_0} (1 + 1/n^2). \quad (18.3.2)$$

Now, treating  $H_{12}$  as a perturbation, we obtain the first-order energy correction

$$E_{n,l,m}^{(1)} = \langle \psi | H_{12} | \psi \rangle = e^2 (I_{n,l,m} \pm J_{n,l,m}), \quad (18.3.3)$$

where the  $+$  sign applies to para-helium ( $s = 0$ ) and the  $-$  sign to ortho-helium ( $s = 1$ ), and

$$I_{n,l,m} = \int d^3\vec{r}_1 \int d^3\vec{r}_2 |\phi_{1,0,0}(\vec{r}_1)|^2 |\phi_{n,l,m}(\vec{r}_2)|^2 \frac{1}{|\vec{r}_1 - \vec{r}_2|}, \quad (18.3.4)$$

$$J_{n,l,m} = \int d^3\vec{r}_1 \int d^3\vec{r}_2 \phi_{1,0,0}(\vec{r}_1)\phi_{1,0,0}(\vec{r}_2) \phi_{n,l,m}(\vec{r}_1)\phi_{n,l,m}(\vec{r}_2)^* \frac{1}{|\vec{r}_1 - \vec{r}_2|}. \quad (18.3.5)$$

Here  $I_{n,l,m}$ , which is manifestly real and positive, is called the “direct integral”, while  $J_{n,l,m}$  is called the “exchange integral”. We will not evaluate them explicitly, but it is worth knowing that  $J_{n,l,m}$  is also real and positive, but smaller in magnitude than  $I_{n,l,m}$ . (The example of



$n, l, m = 2, 0, 0$  is left to Exercise 18.4.) This shows that each ortho-helium ( $s = 1$ ) state is always lower in energy than the corresponding para-helium ( $s = 0$ ) state. A way of understanding this qualitatively is to note that in the  $s = 0$  combination with antisymmetric spins, the symmetric spatial wavefunction results in a higher electrostatic repulsion between the two electrons, due to their greater wavefunction overlap, than for the  $s = 1$  combination with antisymmetric spatial wavefunction.

For  $s = 0$ , the total angular momentum quantum number  $j$  is the same as the orbital angular momentum of the excited electron state,  $l$ . For  $l = 0$ ,  $j$  is the same as  $s$ . For  $s = 1$  and  $l \geq 1$ , one can instead have  $j$  values  $l - 1$ ,  $l$ , or  $l + 1$ . A sketch of the observed energy levels for the lowest few states of the neutral helium atom is shown in Figure 18.3.1, classified by their unperturbed quantum numbers on the left, and by their  $^{2s+1}l_j$  angular momentum labels on the right. The excited states are each split into  $s = 0$ , para-helium (upper) and  $s = 1$ , ortho-helium (lower) levels. Transitions between para-helium states and ortho-helium states are highly suppressed, because they require spin flips, so to a good approximation they form two distinct sets of spectral lines. For this reason it was thought by early investigators that they might actually be two separate kinds of helium.

Each of the ortho-helium levels with  $l > 1$  is split by small fine-structure effects into its separate  $j = l - 1$ ,  $l$  and  $l + 1$  states. These splittings are not shown in Figure 18.3.1, but move the states with larger  $j$  slightly lower in energy. We will discuss this in the next section in the more general context of multi-electron atoms.

Regarding hyperfine effects, the helium nucleus exists in two stable isotopes. Almost all naturally occurring helium nuclei are  $^4\text{He}$ , a boson with spin 0. A spin-0 particle has no special direction in its rest frame, and therefore cannot have a magnetic moment or electric quadrupole moment, so  $^4\text{He}$  atoms have no hyperfine splitting. A small fraction of naturally occurring helium nuclei are  $^3\text{He}$ , a fermion with spin  $1/2$ . The  $^3\text{He}$  atomic states with a given  $j$  are therefore further split by the hyperfine contribution.

He<sup>+</sup> ion, 1s, -54.42 eV

---




Figure 18.3.1: The lowest few energy levels of the neutral helium atom, with electron configurations,  $^{2s+1}l_j$  spectroscopic notations, and energies in eV. Relative energies spacings are indicated qualitatively, but not to scale. The ground state has both electrons in the lowest energy level  $n = 1$  and opposite spins ( $s = 0$ , para-helium). The remaining states have one electron in the lowest level 1s and one in an excited energy level  $n > 1$ , and are split into para-helium (upper, blue lines,  $s = 0$ ) and ortho-helium (lower, red lines,  $s = 1$ ) states. Would-be states with both electrons in an excited level are above the ground-state energy of the He<sup>+</sup> ion, shown as the dashed line, for which one electron is in the ground state and the other has been completely ionized away. Not shown are other states with (1s)(4s) and (1s)(4p) etc., which fall below the dashed line and asymptotically approach it for large  $n$  for one electron. Small fine structure effects, also not shown, split the ortho-helium levels  $^3P_{0,1,2}$ ,  $^3D_{1,2,3}$ , etc., into their different  $j$  components, with larger  $j$  slightly lower in energy.

## 18.4 Multi-electron atoms

Consider the problem of finding the energy eigenstates for multi-electron atoms. To a good approximation, the Hamiltonian depends on the nonrelativistic kinetic energy of the electrons, their electrostatic attraction to the nucleus, their pairwise electrostatic repulsion, and the couplings between the electron spins and their angular momenta. These effects alone are enough to provide a challenge that we cannot hope to solve exactly, and approximation methods must be used. In the following, we will not consider other, usually smaller, effects, including the contributions due to the relativistic correction to the electron kinetic energies, Darwin terms,

and hyperfine contributions due to the magnetic dipole and electric quadrupole moments of the nucleus. These can be added as further perturbations.

One way to make progress is the **central-field method**. We invent a central potential  $U(r)$ , a rotationally invariant function (of the radial coordinate only) that is chosen to approximate the electrostatic potential seen by an electron at a given point due to the nucleus and the suitably averaged effects of the other electrons. Although it is not easy to make a rigorous definition of “suitably averaged”, there are motivated techniques for estimating  $U(r)$  by iteration (notably the Hartree and Hartree–Fock approximations, due to Douglas Hartree and improved by Vladimir Fock) that will not be discussed here. In any case, the choice of  $U(r)$  is arbitrary, in principle, because we can add it to one part of the Hamiltonian (the unperturbed part, or the part used to define trial states for use with the variational principle) and subtract it from another (the perturbation part), so that the total Hamiltonian does not depend on it.

More specifically, we can write the approximate Hamiltonian as

$$H = H_0 + W + H_{\text{SO}}, \quad (18.4.1)$$

$$H_0 = \sum_{i=1}^{N_e} \left( \frac{P_i^2}{2m_e} + U(R_i) \right), \quad (18.4.2)$$

$$W = \sum_{i=1}^{N_e} \left( -U(R_i) - \frac{Ze^2}{R_i} + \sum_{j=1}^{i-1} \frac{e^2}{|\vec{R}_i - \vec{R}_j|} \right), \quad (18.4.3)$$

$$H_{\text{SO}} = \sum_{i=1}^{N_e} \vec{S}_i \cdot \vec{\Lambda}_i. \quad (18.4.4)$$

This Hamiltonian is invariant under the exchange of any two electrons, as it must be. In the spin-orbit term eq. (18.4.4),

$$\vec{\Lambda}_i = -\frac{e}{2m_e^2 c^2} (\vec{\nabla} \Phi_i) \times \vec{P}_i \quad (18.4.5)$$

where the electric potential seen by the  $i$ th electron is

$$\Phi_i(\vec{R}_i) = \frac{Ze}{R_i} - \sum_{j \neq i} \frac{e}{|\vec{R}_i - \vec{R}_j|}. \quad (18.4.6)$$

The contribution to the spin-orbit interaction  $H_{\text{SO}}$  from each electron generalizes the case of a spherically symmetric potential in eq. (17.1.4), and for hydrogen in eq. (17.1.16), which would be recovered for  $\Phi = e/R$ . Note that there is no operator ordering problem associated with the placement of  $\vec{P}_i$  in eq. (18.4.5), because of the vector calculus identity  $\vec{\nabla} \times (\vec{\nabla} \Phi) = 0$ . We have distinguished the number of protons in the nucleus  $Z$  from the number of electrons  $N_e$ , so that the formulas apply to charged ions as well as neutral atoms.




Figure 18.4.1: The solid line shows the qualitative shape of a choice for the central-field potential energy function  $U(r)$  for the electrons in a neutral atom with  $N_e = Z = 10$ . The upper dashed and lower dashed lines show the potentials  $-e^2/r$  (maximally screened by 9 electrons closer to the nucleus) and  $-10e^2/r$  (no screening, all electrons farther from the nucleus), respectively.

One should aim to choose  $U(R)$  in such a way that the eigenstates of  $H_0$  are as close as possible to those of the full Hamiltonian. A reasonable choice might interpolate between

$$U(r) \approx \begin{cases} -Ze^2/r & (\text{small } r), \\ -(Z - N_e + 1)e^2/r & (\text{large } r). \end{cases} \quad (18.4.7)$$

The idea behind this choice is the principle from classical electrostatics, following from Gauss' Law, that the electric field and electric potential at a point  $\vec{r}$  in a spherically symmetric charge distribution are both determined only by the total charge contained within  $r$ . Thus, at small  $r$  there is no screening of the charge  $+Ze$  nuclear potential due to other electrons, because they are likely to be found farther from the nucleus, while at large  $r$  the other  $N_e - 1$  electrons are all likely to be closer to the nucleus and so the effective number of charges seen by an outermost very distant electron is only  $Z - N_e + 1$ . A sketch of a plausible choice of  $U(r)$  is shown in Figure 18.4.1, compared to the asymptotic limits of no screening at small  $r$  and maximum screening at large  $r$ . A choice of  $U(r)$  should be justified *a posteriori* by the successful convergence of perturbation theory or the variational method.

The single-electron eigenstates of the central-field Hamiltonian  $H_0$ , called **orbital states** or **orbitals**, will have wavefunctions of the form

$$\phi_{n,l,m_l,m_s}(r, \theta, \phi) = R_{n,l}^U(r) Y_l^{m_l}(\theta, \phi) \chi_{m_s}, \quad (18.4.8)$$

where  $\chi_{m_s} = |\uparrow\rangle$  or  $|\downarrow\rangle$  for the spin states. The radial wavefunctions  $R_{n,l}^U(r)$  will not be the same as for a hydrogen-like atom, because  $U(r)$  is not proportional to  $1/r$ . However, they will be qualitatively similar. In particular, the label  $n$  is a positive integer, and then  $0 \leq l \leq n-1$ , with  $n-l-1$  equal to the number of radial zeros for  $r > 0$ . In general, the orbital energy eigenvalues of  $H_0$  will increase with  $n$ , but will also depend on  $l$ , unlike hydrogen-like atom energy eigenstates. The unperturbed eigenstates of  $H_0$  will be Slater determinants formed out of  $N_e$  states of the form of eq. (18.4.8). Then the effects of  $W$  and  $H_{\text{SO}}$  can be added as perturbations.

The set of orbitals with the same values of  $n$  are called a **shell**, and those with the same values of both  $n$  and  $l$  are called a **subshell**. Each subshell consists of

$$g_{n,l}^{\text{subshell}} = 2(2l + 1) \quad (18.4.9)$$

states, due to the  $m_s$  and  $m_l$  quantum numbers. (This neglects any further degeneracy due to nuclear spins, which are different for distinct isotopes of the same element.) The orbital states and subshells are labeled  $nl$ , but with  $l = 0, 1, 2, 3, \dots$  replaced by the spectroscopic code letter  $s, p, d, f, \dots$ , just as for the hydrogen atom states. The state of a given atom can be specified by giving the occupation numbers of each subshell, called the **electron configuration**, subject to the Pauli exclusion principle, enforced by the Slater determinant, that we cannot put two electrons in the same state due to their Fermi–Dirac statistics. Using eq. (18.4.9), the maximum number of electrons in an  $s, p, d, f$  subshell is 2, 6, 10, 14, respectively. For a given  $l$ , smaller  $n$  tends to have lower energy, but this does not always mean that the subshells fill up from smaller to larger  $n$ , as we will soon see. For a given  $n$ , the states with smaller  $l$  tend to have lower energies. The reason is that wavefunctions with smaller  $l$  have support closer to the nucleus, where the attractive force of the nucleus is less screened by the other electrons in the complete subshells.

The  $1s$  shell can fit up to 2 electrons, and so is the only shell necessary for H and He, but is insufficient to accommodate larger  $Z$  atoms. The next to fill is the  $2s$  subshell, which again fits up to 2 electrons, and so is the last subshell needed for Li and Be. The  $2p$  subshell fills next, and can fit up to 6 electrons, so it is enough to accommodate the neutral atoms with  $Z$  up to 10, namely B, C, N, O, Fe, and Ne. Likewise, the  $3s$  subshell accommodates the last electron for Na and Mg, while the  $3p$  subshell is used for Al, Si, P, S, Cl, Ar. However, the  $4s$  subshell tends to fill before the  $3d$  subshell, in K, Ca, Sc, Ti, V, Mn, Fe, Co, Ni, and Zn. This tendency has two exceptions; within this sequence from  $Z = 19$  to  $Z = 30$ , Cr and Cu break the trend, by having only one electron in the  $4s$  subshell, with the  $3d$  subshell containing the other five and ten electrons, respectively.

The subshell filling order for multi-electron atoms is, empirically,

$$1s, 2s, 2p, 3s, 3p, \begin{pmatrix} 4s \\ 3d \end{pmatrix}, 4p, \begin{pmatrix} 5s \\ 4d \end{pmatrix}, 5p, \begin{pmatrix} 6s \\ 4f \\ 5d \end{pmatrix}, 6p, \begin{pmatrix} 7s \\ 5f \\ 6d \end{pmatrix}, \quad (18.4.10)$$

where the cases in parentheses correspond to ambiguities due to close energies, with the most common (but not universally followed) filling order from top to bottom. The fact noted earlier that smaller  $l$  tends to give lower energy for a given  $n$  explains why  $2s$  fills before  $2p$ , and why the  $4s$  can compete with the  $3d$ , and the  $5s$  can compete with the  $4d$ , etc.

Some example electron configurations are

$$\begin{array}{llll} \text{H: } 1s^1, & \text{He: } 1s^2, & \text{Li: } 1s^2 2s^1, & \text{C: } 1s^2 2s^2 2p^2, \\ \text{N: } 1s^2 2s^2 2p^3, & \text{Na: } 1s^2 2s^2 2p^6 3s^1, & & \\ \text{Hg: } 1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^{10} 4p^6 5s^2 4d^{10} 5p^6 6s^2 4f^{14} 5d^{10}. & & & \end{array}$$

The chemical properties of a given element are mostly dependent on the electron configuration, and the similarities between elements in a given column of the periodic table are due to having similar configurations of electrons in the outermost (partially filled) subshells.

The inert (noble) gases He, Ne, Ar, Kr, Xe, Rn are those that have a full  $p$  subshell, and all previous subshells filled, leading to a very low tendency to form chemical bonds. (The  $p$  subshells are special, because the next higher energy subshell is always  $s$ . This means a big energy gap to excite an electron out of a  $p$  subshell.) It is common to give the electron configuration for any element by just listing the filled orbitals in excess of that for the previous inert gas. For example, for the inert gases themselves, one can write the electron configurations as

$$\begin{array}{llll} \text{He: } 1s^2, & \text{Ne: } [\text{He}]2s^2 2p^6, & \text{Ar: } [\text{Ne}]3s^2 3p^6, & \text{Kr: } [\text{Ar}]4s^2 3d^{10} 4p^6, \\ \text{Xe: } [\text{Kr}]5s^2 4d^{10} 5p^6, & \text{Rn: } [\text{Xe}]6s^2 4f^{14} 5d^{10} 6p^6. & & \end{array}$$

For the alkali metals, there is an  $s$  subshell with only one electron,

$$\begin{array}{lll} \text{Li: } [\text{He}]2s^1, & \text{Na: } [\text{Ne}]3s^1, & \text{K: } [\text{Ar}]4s^1, \\ \text{Rb: } [\text{Kr}]5s^1, & \text{Cs: } [\text{Xe}]6s^1, & \text{Fr: } [\text{Rn}]7s^1, \end{array}$$

while the halogens are all missing 1 electron in a  $p$  subshell,

$$\begin{array}{lll} \text{F: } [\text{He}]2s^2 2p^5, & \text{Cl: } [\text{Ne}]3s^2 3p^5, & \text{Br: } [\text{Ar}]3d^{10} 4s^2 4p^5, \\ \text{I: } [\text{Kr}]4d^{10} 5s^2 5p^5, & \text{At: } [\text{Xe}]4f^{14} 5d^{10} 6s^2 6p^5. & \end{array}$$

A good periodic table of the elements will give the electron configuration for the atomic ground state of each element.

The energy needed to remove a single electron from the ground state of a neutral atom is called the **first-ionization energy**. Experimental results for the first-ionization energy as a function of the atomic number  $Z$  are shown in Figure 18.4.2, including as the first two data points 13.6 eV for H and 24.6 eV for He. (Note that the result for He can be obtained from numbers given in Figure 18.3.1.) Because of the shielding of the nuclear charge by the innermost electrons, the results do not depend very strongly on  $Z$ ; the outermost electrons are attracted to the nucleus as if they “see” a reduced net charge of order  $e$  rather than the full nuclear charge




Figure 18.4.2: The energy (in eV) needed to remove a single electron from the ground state of a neutral atom, as a function of the atomic number  $Z$ . Atoms with the same highest-energy subshells ( $1s$ ,  $2s$ ,  $2p$ ,  $3s$ , ...) are connected by lines to guide the eye, as labeled. The inert gas elements (He, Ne, Ar, Kr, Xe, Rn) each have completely filled  $s$ - and  $p$ -subshells and thus a large first-ionization energy, while the alkali metal atoms (Li, Na, K, Rb, Cs, Fr) each have a lone  $s$ -subshell electron and a small first ionization energy.

$Ze$ . The first-ionization energies are largest for the inert gases, and are smallest for alkali metals, which each have a single, relatively weakly bound,  $s$ -subshell electron available for ionization.

A more complete specification of the structure of the atoms will include information about the angular momentum quantum numbers. Suppose that we can treat the spin-orbit contribution to the Hamiltonian  $H_{SO}$  as small compared to  $H_0 + W$  in eqs. (18.4.1)–(18.4.4). In practice, this is especially likely to be a good approximation if  $Z$  is not too large. Now, all three of the summed orbital, summed spin, and total angular momentum operators

$$\vec{L} = \sum_{i=1}^{N_e} \vec{L}_i, \quad \vec{S} = \sum_{i=1}^{N_e} \vec{S}_i, \quad \vec{J} = \vec{L} + \vec{S} \quad (18.4.11)$$

commute with  $H_0$  (as well as  $W$ ). One can therefore choose unperturbed orthobasis states that are not only eigenstates of  $H_0$ , but also eigenstates of the operators  $L^2$ ,  $S^2$ ,  $L_z$  and  $S_z$  with eigenvalues  $\hbar^2 L(L+1)$ ,  $\hbar^2 S(S+1)$ ,  $\hbar m_L$ , and  $\hbar m_S$  respectively. These states also carry other quantum number labels (including the unperturbed  $H_0$  energy eigenvalue) which we will symbolically denote as  $N$ , so that the states are labeled

$$|NLSm_Lm_S\rangle. \quad (18.4.12)$$

The  $H_0$  energy eigenvalues have energy degeneracy  $g_N = (2S + 1)(2L + 1)$ , due to the magnetic quantum numbers  $m_L, m_S$ . Alternatively, one can choose as an orthobasis the linear combinations of the same unperturbed states that are eigenstates of  $H_0$ ,  $S^2$ ,  $L^2$ , and  $J^2$  [with eigenvalue  $\hbar^2 J(J + 1)$ ], and  $J_z$  (with eigenvalue  $\hbar m_J$ ),

$$|NLSJm_J\rangle = \sum_{m_L, m_S} C_{m_L m_S m_J}^{LSJ} |NLSm_L m_S\rangle, \quad (18.4.13)$$

where  $C_{m_L m_S m_J}^{LSJ}$  are the Clebsch–Gordan coefficients. The total number of such states for fixed  $N, L, S$  is again  $(2S + 1)(2L + 1)$ . The approximation in which the states constructed in this way from eigenstates of  $H_0$  (or perhaps  $H_0 + W$ ) are close to being eigenstates of the full Hamiltonian is called the **LS coupling** (or **Russell–Saunders coupling**, after Henry N. Russell and Frederick Saunders) approximation. Note that this approximation is only good to the extent that  $H_{SO}$  can be treated as small, since it does not commute with  $S^2$  or  $L^2$  if there is more than one electron.

We now investigate the spin-orbit fine-structure energy corrections, using degenerate perturbation theory. To do so, we must find the matrix elements of  $H_{SO}$  on the degenerate subspaces of  $H_0$ , and diagonalize it. We start by working with the product orthobasis kets, which decompose into linear combinations of tensor products of orbital and spin angular momentum eigenkets,

$$|NLSm_L m_S\rangle = \sum_{\alpha, \beta} c_{\alpha, \beta} |\alpha N L m_L\rangle \otimes |\beta S m_S\rangle. \quad (18.4.14)$$

Here  $\alpha$  is a degeneracy label whose presence reflects the fact that there can be more than one state with eigenvalues  $L, m_L$ , and similarly  $\beta$  labels the different spin states with eigenvalues  $S, m_S$ . The whole state  $|NLSm_L m_S\rangle$  is a Slater determinant incorporating the antisymmetry under interchange of any two electrons. This puts constraints, which we will not explicitly identify or make use of, on the coefficients  $c_{\alpha, \beta}$ .

The matrix elements of  $H_{SO}$  from eq. (18.4.4) on the degenerate subspace of states are

$$\begin{aligned} \langle NLSm'_L m'_S | H_{SO} | NLSm_L m_S \rangle &= \\ \sum_{\alpha, \beta} \sum_{\alpha', \beta'} c_{\alpha, \beta} c_{\alpha', \beta'}^* \sum_{i=1}^{N_e} \langle \alpha' N L m'_L | \vec{\Lambda}_i | \alpha N L m_L \rangle \cdot \langle \beta' S m'_S | \vec{S}_i | \beta S m_S \rangle. \end{aligned} \quad (18.4.15)$$

Electrons in complete subshells can be omitted, because pairs of opposite spin expectation values with the same orbital quantum numbers will always cancel in the sum over  $i$ . Now we will invoke the Landé projection formula, eq. (13.2.10), twice. First, note that each  $\vec{\Lambda}_i$  is a vector operator with respect to the total orbital angular momentum operator  $\vec{L}$ , without involving spin. Therefore, for  $L \neq 0$ , we have

$$\langle \alpha' N L m'_L | \vec{\Lambda}_i | \alpha N L m_L \rangle = \langle L m'_L | \vec{L} | L m_L \rangle \frac{\langle \alpha' N L || \vec{L} \cdot \vec{\Lambda}_i || \alpha N L \rangle}{\hbar^2 L(L + 1)}. \quad (18.4.16)$$



As noted in Theorem 13.2.2, for the case  $L = 0$ , the matrix element simply vanishes, and there is no spin-orbit splitting. Similarly, each of the individual electron spin operators  $\vec{S}_i$  is a vector with respect to rotations generated by the total spin angular momentum operator  $\vec{S}$ , without involving orbital angular momentum, so we can write, for  $S \neq 0$ ,

$$\langle \beta' S m'_S | \vec{S}_i | \beta S m_S \rangle = \langle S m'_S | \vec{S} | S m_S \rangle \frac{\langle \beta' S | \vec{S} \cdot \vec{S}_i | \beta S \rangle}{\hbar^2 S(S+1)}. \quad (18.4.17)$$

while for  $S = 0$  the matrix element again vanishes. Therefore, putting things together,

$$\langle N L S m'_L m'_S | H_{\text{SO}} | N L S m_L m_S \rangle = \zeta(N, L, S) \langle L S m'_L m'_S | \vec{S} \cdot \vec{L} | L S m_L m_S \rangle, \quad (18.4.18)$$

where the factor

$$\zeta(N, L, S) = \sum_{\alpha, \beta} \sum_{\alpha', \beta'} c_{\alpha, \beta} c_{\alpha', \beta'}^* \sum_{i=1}^{N_e} \frac{\langle \alpha' N L | \vec{L} \cdot \vec{\Lambda}_i | \alpha N L \rangle}{\hbar^2 L(L+1)} \frac{\langle \beta' S | \vec{S} \cdot \vec{S}_i | \beta S \rangle}{\hbar^2 S(S+1)} \quad (18.4.19)$$

may be quite difficult to evaluate exactly, but has the important property that it does not depend on the magnetic quantum numbers  $m_L, m_S, m'_L, m'_S$  at all, because the reduced matrix elements  $\langle \alpha' N L | \vec{L} \cdot \vec{\Lambda}_i | \alpha N L \rangle$  and  $\langle \beta' S | \vec{S} \cdot \vec{S}_i | \beta S \rangle$  do not. For a given atom,  $\zeta(N, L, S)$  can, in principle, be evaluated, perhaps in approximations such as replacing the electric potential  $\Phi_i(\vec{R}_i)$  in eq. (18.4.5) by a spherically symmetric approximation. This is beyond our scope here; we will merely consider it to be an empirical function of  $N, L, S$ .

Equation (18.4.18) shows that the matrix elements of  $H_{\text{SO}}$  are proportional to those of  $\vec{S} \cdot \vec{L} = \hbar^2(J^2 - L^2 - S^2)/2$ , where  $J^2, L^2$  and  $S^2$  are understood to be operators. To properly conduct degenerate perturbation theory, we should switch to the orthobasis  $|N L S J m_J\rangle$ , in which this operator is diagonal. We therefore arrive at a formula for the spin-orbit correction to the energies, at first order in degenerate perturbation theory,

$$\Delta E(N, L, S, J) = \zeta(N, L, S) \hbar^2 [J(J+1) - L(L+1) - S(S+1)]/2, \quad (18.4.20)$$

where  $J, L$ , and  $S$  are now numbers. This vanishes in the special cases  $L = 0$  (which then implies  $J = S$ ) and  $S = 0$  (which then implies  $J = L$ ), consistent with observations already made in the previous paragraph. Equation (18.4.20) shows that the degeneracy  $(2S+1)(2L+1)$  is partially broken by  $H_{\text{SO}}$ , but states with a given  $J$  are not split, with remaining degeneracies  $g_J = 2J+1$ , since  $\Delta E(N, L, S, J)$  does not depend on  $m_J$ .

We can now use eq. (18.4.20) to obtain a simple formula for the energy differences between successive eigenvalues  $J-1$  and  $J$ , for fixed  $N, L$ , and  $S$ ,

$$\Delta E(N, L, S, J) - \Delta E(N, L, S, J-1) = \hbar^2 \zeta(N, L, S) J. \quad (18.4.21)$$

This result, that the spin-orbit fine-structure energy splitting between states  $J$  and  $J-1$  with fixed  $N, L, S$  is proportional to  $J$ , is called the **Landé interval rule**, as observed in spectral lines from transitions between atomic states. It can be used as a check of the validity of the  $LS$  coupling scheme approximation.

The preceding discussion shows how, within the  $LS$  coupling approximation, energy eigenstates can be built by first adding all of the spins, then adding all of the orbital angular momenta, and then adding the two to get the total angular momentum. The angular momentum quantum numbers of these states are then traditionally specified with the **term symbol**

$$^{2S+1}L_J = \text{Russell-Saunders } (LS) \text{ spectroscopic notation,} \quad (18.4.22)$$

where the eigenvalues of operators  $J^2$ ,  $S^2$ , and  $L^2$  are numbers  $\hbar^2 J(J+1)$ ,  $\hbar^2 S(S+1)$ , and  $\hbar^2 L(L+1)$ , respectively, and then the value of  $L$  is replaced by its spectroscopic code  $S, P, D, F, G, H, \dots$  for  $L = 0, 1, 2, 3, 4, 5, \dots$ . Thus, for example, the hydrogen and helium atom ground states are given as  $^2S_{1/2}$  and  $^1S_0$ , respectively. Many editions of the periodic table of elements list the Russell-Saunders notation for the atomic ground states along with the electron configuration.

The list of possible term symbols  $^{2S+1}L_J$  of an atom can be obtained from looking only at the **valence electrons**, the ones in the incompletely filled subshells. This is because each electron in a filled subshell with a given  $m_s$ ,  $m_l$ , and  $m_j$  always has a partner with the opposite values of those quantum numbers, giving no net contribution to  $S$ ,  $L$ , and  $J$ . The parity of the state is useful for selection rules for transitions. From eq. (8.7.7), it is  $\pi = (-1)^{\sum_i l_i}$ , where  $l_i$  are the individual orbital angular momenta, specified by the electronic configuration. [Note that for orbitals the parity is defined with respect to the fixed nucleus as the origin, not the center of mass of the valence electrons, so even for two electrons  $\pi$  is not necessarily equal to  $(-1)^L$ .]

For an atom with a partly filled subshell, there may be several candidates for the ground state term symbol. Three empirical rules, known as **Hund's Rules** after Friedrich Hund, can be used to predict which the ground state will be, given the electron configuration. They are:

Hund's Rule 1: The  $L, S, J$  multiplet with the lowest energy will have the largest  $S$  from among the candidates.

(Qualitative justification: for larger  $S$ , the spin wavefunction is more symmetric, which means the spatial wavefunction is more antisymmetric, which means the electrons are less often near each other, which gives less repulsive forces between them, and so lower energy. We have already seen this effect in action; it is why each ortho-helium  $S = 1$  state is lower in energy than its para-helium  $S = 0$  counterpart.)

Hund's Rule 2: If there are two or more candidates selected by the first rule that have the same largest  $S$ , then the lowest energy will have the largest  $L$  among them.

(Qualitative justification: larger  $L$  means that the orbital wavefunctions can be arranged to have less overlap, again leading to smaller repulsive forces and thus lower energy.)

Hund's Rule 3: If there are two or more candidates selected by the first two rules with the same  $S$  and  $L$ , then the spin-orbit coupling splits apart the different levels of  $J = |L - S|, \dots, L + S$ . If an incomplete subshell is not more than half-filled, then the ground state has  $J = |L - S|$ , otherwise the ground state has  $J = L + S$ .

(Qualitative justification: the form of eq. (18.4.20) immediately tells us that for fixed  $L, S$ , the minimum energy will occur for one of the two extremes for  $J$ , either  $|L - S|$  or  $L + S$ . Hund's Rule 3 is equivalent to the empirical fact that  $\zeta(N, L, S)$  in eq. (18.4.20) turns out to be positive if the subshell is not more than half-filled, and negative otherwise.)

We now illustrate the prediction of term symbols and ground states, using Hund's rules where necessary, with examples from the beginning of the periodic table. For the first four elements, Hund's rules are not needed.

Hydrogen (H,  $Z = 1$ ) has electron configuration  $1s^1$ . Because there is one electron with  $S = 1/2$  and  $L = 0$ , the only possibility is  $J = 1/2$ , and the term symbol is  $^2S_{1/2}$ .

Helium (He,  $Z = 2$ ) has electron configuration  $1s^2$ . As always when there are no partially filled subshells, the ground state is  $^1S_0$ . The same result is obtained for all inert (noble) gases.

Lithium (Li,  $Z = 3$ ) has electron configuration  $1s^2 2s^1$ . Just as for hydrogen, there is one electron that is not in a filled subshell, with  $S = 1/2$  and  $L = 0$ , so the term symbol is  $^2S_{1/2}$ . The same applies for all alkali metals, the elements in the same periodic table column as Li, which have a single  $s$  subshell electron.

Beryllium (Be,  $Z = 4$ ) has electron configuration  $1s^2 2s^2$ , with only completely filled subshells, so the ground state is  $^1S_0$ . The same applies for all elements in the Be periodic table column.

Boron (B,  $Z = 5$ ) has electron configuration  $1s^2 2s^2 2p^1$ . Here, the  $1s$  and  $2s$  subshells are irrelevant, since they are completely filled. In the incomplete  $2p$  subshell, we have one electron with  $L = 1$  and  $S = 1/2$ . Since this is the only choice, we do not need Hund's rules 1 or 2. The possible candidates have  $J = 1/2$  and  $J = 3/2$ , with term symbols  $^2P_{1/2}$  and  $^2P_{3/2}$ . Because the last incomplete subshell is less than half filled (1 out of 6), Hund's rule 3 tells us that the ground state has  $J = |L - S| = 1/2$  and is  $^2P_{1/2}$ . The same result is obtained for all elements in the B column of the periodic table, which have a single  $p$  electron.

Carbon (C,  $Z = 6$ ) has electron configuration  $1s^2 2s^2 2p^2$ . We start by computing the total number of allowed states associated with the incomplete  $2p$  subshell. The 2 electrons have 6 orbital states from which to choose. In general, the number of ways of choosing  $k$  things from

$n$  possibilities, without regard to order, is equal to the binomial coefficient

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}. \quad (18.4.23)$$

So in our present case there are  $(6 \cdot 5)/(2 \cdot 1) = 15$  completely antisymmetric states. Now, let us see how these 15 states are divided into  $^{2S+1}L_J$  multiplets. Combining the two spin- $1/2$ 's gives  $S = 0$  (the antisymmetric combination) and  $S = 1$  (the symmetric combination). Combining the two orbital angular momenta, each with  $l = 1$ , gives  $L = 0$  (symmetric), and  $L = 1$  (antisymmetric), and  $L = 2$  (symmetric). Therefore, requiring that the full state be antisymmetric under exchange of the two electrons, we have the following allowed combinations. First, there is  $L = 2, S = 0$ , which implies  $J = 2$ . This possibility  $^1D_2$  consists of  $2J + 1 = 5$  states. Second, there is  $L = 1, S = 1$ , which allows  $J = 0, 1, 2$ , so the terms are  $^3P_0$ ,  $^3P_1$ , and  $^3P_2$ , with 1, 3, and 5 states respectively. Third, there is  $L = 0, S = 0$ , which of course allows only one state,  $^1S_0$  with  $J = 0$ . As a check, the total of these possibilities is indeed 15. Now we are ready to use Hund's rules. Hund's rule 1 selects the largest possible value of  $S$ , which is  $S = 1$ . Hund's rule 2 tells us nothing, because the remaining competitors all have  $L = 1$ . Finally, for Hund's rule 3, we note that the  $2p$  subshell is not more than half-filled (2 out of 6), so that  $J = |L - S| = 0$  in the ground state. We therefore have  $S = 1$ ,  $L = 1$ ,  $J = 0$ , and the ground state of carbon is  $^3P_0$ . The same logic and result applies to all elements in the C column of the periodic table, which have two  $p$  electrons.

Nitrogen (N,  $Z = 7$ ) has electron configuration  $1s^2 2s^2 2p^3$ . The total number of possible electron states associated with the three electrons in the  $2p$  subshell is  $(6 \cdot 5 \cdot 4)/(3 \cdot 2 \cdot 1) = 20$  antisymmetric states. To see which  $L, S, J$  combinations these correspond to, first consider the combination of the orbital angular momenta, which gives  $1 \otimes 1 \otimes 1 = 0 \oplus 1 \oplus 1 \oplus 2 \oplus 3$  for the possible values of  $L$ . If  $L = 3$ , one can check that the spatial wavefunction constructed by addition of angular momentum would be totally symmetric under exchange of the electron labels, but there is no totally antisymmetric spin state for  $\geq 3$  electrons; we conclude from this that  $L$  cannot be 3. If  $L = 0$ , then the spatial wavefunction is instead totally antisymmetric, from which we conclude that the spin state must be totally symmetric, so  $S = 3/2$ . Conversely, if  $S = 3/2$ , then only the totally antisymmetric  $L = 0$  combination is allowed. We conclude that the only  $(L, S)$  combinations that are allowed are  $L = 0, S = 3/2$  with  $J = 3/2$  (term  $^4S_{3/2}$ , 4 states), and  $L = 1, S = 1/2$  with  $J = 1/2$  and  $3/2$  (terms  $^2P_{1/2}$  and  $^2P_{3/2}$ , 6 states), and  $L = 2, S = 1/2$  with  $J = 3/2$  and  $J = 5/2$  (terms  $^2D_{3/2}$  and  $^2D_{5/2}$ , 10 states). As a check, these indeed add up to 20 states. Hund's rule 1 tells us that the ground state should have maximal  $S$ , which selects the  $L = 0, S = 3/2, J = 3/2$  combination. This is the only one that realizes that maximal value of  $S$ , so the other two Hund's rules are not needed, and the ground state

term is  ${}^4S_{3/2}$ . The same logic and result applies to all elements in the N column of the periodic table, which have three  $p$  electrons.

Oxygen (O,  $Z = 8$ ) has electron configuration  $1s^2 2s^2 2p^4$ . As in the previous examples, we start by doing the counting of the total number of physical antisymmetric electron states for the incomplete subshell. Since there are 4 electrons and 6 available states, that number is  $(6 \cdot 5 \cdot 4 \cdot 3)/(4 \cdot 3 \cdot 2 \cdot 1) = 15$ . However, as a useful trick it is better to consider the 2 “holes”, by which we mean the electrons omitted from the completely filled  $2p$  subshell. Treating the holes as equivalent spin-1/2 particles, we again obtain  $(6 \cdot 5)/(2 \cdot 1) = 15$  states. A complete  $2p$  subshell would have  $L = S = 0$ , so we treat the 2 holes as carrying the same  $S$  and  $L$  as the 4 electrons. So, the allowed candidate multiplets are the same as we found for carbon with  $2p^2$ , namely  $(L, S, J) = (2, 0, 2), (1, 1, 0), (1, 1, 1), (1, 1, 2),$  and  $(0, 0, 0)$ , with term symbols  ${}^1D_2, {}^3P_0, {}^3P_1, {}^3P_2,$  and  ${}^1S_0$ . Hund’s rule 1 says the ground state has  $S = 1$ , and Hund’s rule 2 does not differentiate between the remaining candidates which all have  $L = 1$ . To apply Hund’s rule 3, we note that the  $2p$  subshell is now more than half-filled with electrons (4 out of 6), so  $J = L + S = 2$  for the ground state. (Note that this is the difference between carbon and oxygen; although the enumeration of the candidate terms is easier in terms of the holes, Hund’s rule 3 for the selection of the ground state still applies to the actual electrons, not the holes.) So, the ground state term for oxygen is  ${}^3P_2$ . The same logic and result applies to all elements in the O column of the periodic table, with four  $p$  electrons.

Fluorine (F,  $Z = 9$ ) has electron configuration  $1s^2 2s^2 2p^5$ . Using the trick introduced in the oxygen example, we get the allowed  $(L, S, J)$  multiplets by considering the  $2p$  subshell to consist of just one hole, rather than 5 electrons. This means that, just as for boron, the only combinations are  $L = 1, S = 1/2$  with  $J = 1/2$  or  $3/2$ . Since these have the same  $L$  and  $S$ , Hund’s rules 1 and 2 provide no information. To apply Hund’s rule 3, we note that the  $2p$  subshell is more than half full (5 out of 6), and so the ground state has  $J = L + S = 3/2$ , with term symbol  ${}^2P_{3/2}$ . The same applies to all halogens, the elements in the F column of the periodic table, with five  $p$  electrons.

Hund’s rules correctly predict the ground state of all neutral atoms for which the ground state term symbol is unambiguously measured. They also often give the correct result for charged ions, and for neutral atoms with an electron configuration different from the ground state. For example, the singly ionized carbon ion  $C^+$  has electron configuration  $1s^2 2s^2 2p$ , which is the same as boron, and has the same term symbol  ${}^2P_{1/2}$ , as correctly predicted by Hund’s rules. Also, the neutral carbon atom has excited states with electron configuration  $1s^2 2s 2p^3$ . Among those excited states, Hund’s rules correctly predict that the lowest energy term is  ${}^5S_2$ .

An important experimental tool for exposing the properties of atomic states is to observe

the behavior of energy levels and transition spectra as a function of an applied magnetic field, called the Zeeman effect. In section 17.3, we studied the Zeeman effect for atomic hydrogen, and much of that discussion can be adapted to the multi-electron case. Adding the contributions to the Hamiltonian linear in a constant, uniform magnetic field  $\vec{B} = B\hat{z}$ , one obtains a result of the same form as eq. (17.3.6),

$$H_B = \mu_B B (L_z + 2S_z)/\hbar, \quad (18.4.24)$$

but now with  $L_z$  and  $S_z$  respectively equal to the total orbital and spin angular momentum operators, obtained as the sums of the operators for the individual electrons. Within the  $LS$  coupling scheme approximation, the analysis for the weak-field (small- $B$ ) Zeeman effect in multi-electron atoms just mimics what we already did for hydrogen from eqs. (17.3.8)–(17.3.13). So, one finds that atomic states characterized by  $N, L, S, J$  obtain energy splittings proportional to the total angular momentum quantum number  $m_J$ ,

$$\Delta E_B = g \mu_B B m_J, \quad (m_J = -J, -J+1, \dots, J-1, J), \quad (18.4.25)$$

where the Landé  $g$ -factor is

$$g = \frac{3}{2} + \frac{S(S+1) - L(L+1)}{2J(J+1)}. \quad (18.4.26)$$

(Of course, unlike the atomic hydrogen case, in general  $S$  need not be equal to  $1/2$ , and  $J$  need not be equal to  $L \pm 1/2$ .) For weak external magnetic fields, the magnetic moment of the atom is therefore related to its total angular momentum by  $-g\mu_B \vec{J}/\hbar$ , and the splitting exposes the degeneracies and angular momentum quantum numbers for the multi-electron atomic states.

The strong magnetic field (Paschen–Back) limit for multi-electron atoms is also quite similar to the example of the hydrogen atom, as given in eqs. (17.3.20)–(17.3.24). This includes the use of  $|NLSm_Lm_S\rangle$ , in which  $H_B$  is diagonal, as the degenerate unperturbed states to which  $H_{SO}$  is applied as a perturbation. For multi-electron atoms, the counterpart of eq. (17.3.22) is

$$\langle NLSm_Lm_S | H_{SO} | NLSm_Lm_S \rangle = \zeta(N, L, S) \langle NLSm_Lm_S | \vec{S} \cdot \vec{L} | NLSm_Lm_S \rangle, \quad (18.4.27)$$

where  $\zeta(N, L, S)$  is the quantity appearing in eqs. (18.4.19)–(18.4.21) and  $\vec{S} \cdot \vec{L} = \frac{1}{2}(S_+L_- + S_-L_+) + S_zL_z$  evaluates to  $\hbar^2 m_S m_L$ . Therefore, we have in the strong-field limit

$$\Delta E_{B, \text{fine}} = \mu_B B (m_S + 2m_L) + \hbar^2 \zeta(N, L, S) m_S m_L, \quad (18.4.28)$$

where we have kept only the spin-orbit part of the fine-structure effect, which contains the dependence on  $m_S$  and  $m_L$ . For the intermediate case that the fine-structure and external

magnetic fields are comparable, things are more complicated and one should find the energy shift by treating  $H_B + H_{\text{SO}}$  together as a single perturbation, to be diagonalized using either the basis of states  $|NLSJm_J\rangle$  or the basis  $|NLSm_Lm_S\rangle$ .

It is important to keep in mind that the  $LS$  coupling scheme is just an approximation. The assumption that  $H_{\text{SO}}$  can be treated as a small effect becomes less appropriate for atoms with larger  $Z$ , and so the  $LS$  coupling scheme approximation is not as good for heavier atoms. To understand this, consider an approximation in which the potential in the spin-orbit interaction is spherically symmetric, so that we have a sum of terms like eq. (17.1.4) for each electron,

$$H_{\text{SO}} = \sum_{i=1}^{N_e} \xi(R_i) \vec{S}_i \cdot \vec{L}_i, \quad \xi(r) = -\frac{e}{2m_e^2 c^2} \frac{1}{r} \frac{\partial \Phi}{\partial r}. \quad (18.4.29)$$

Now, as a qualitative approximation, suppose that the electrons are in hydrogen-like orbitals arising from a potential  $\Phi(r) = -Z_{\text{eff}}e^2/r$ , where  $Z_{\text{eff}} < Z$  roughly takes into account the effects of partial screening of the nuclear charge by the other electrons. Then  $H_{\text{SO}}$  scales like  $Z_{\text{eff}}/R^3$ , and the characteristic size of orbitals scales like  $1/Z_{\text{eff}}$ , as can be seen from eq. (11.1.65), for example. This implies that the effects of  $H_{\text{SO}}$  scale like  $Z_{\text{eff}}^4$ .

In the limit that the effects of  $H_{\text{SO}}$  are too large to be treated as a perturbation, one can instead use the **jj coupling approximation**. Since eq. (18.4.29) involves the individual  $\vec{S}_i \cdot \vec{L}_i$ , it is useful to first perform addition of angular momentum on each electron,  $\vec{J}_i = \vec{L}_i + \vec{S}_i$ , and then construct  $\vec{J} = \sum_{i=1}^{N_e} \vec{J}_i$ . Then  $H_0 + H_{\text{SO}}$  is diagonal in an orthobasis labeled by the quantum numbers  $l_i$ ,  $s_i$ , and  $j_i$  associated with the operators  $L_i^2$ ,  $S_i^2$ , and  $J_i^2$ . In particular, the dominant spin-orbit coupling term is diagonal in that basis and can be evaluated according to

$$H_{\text{SO}} = \frac{\hbar^2}{2} \sum_{i=1}^{N_e} \xi(R_i) [j_i(j_i + 1) - l_i(l_i + 1) - s_i(s_i + 1)], \quad (18.4.30)$$

where  $s_i = 1/2$ , and  $l_i$  is specified by the electron configuration. The  $W$  part of the Hamiltonian in eq. (18.4.3) does not commute with the operators  $L_i^2$  and  $J_i^2$ , so it is treated as a perturbation, which then splits the states into different values of the total angular momentum quantum number  $J$ , with a remaining degeneracy  $2J + 1$  (neglecting small hyperfine effects associated with electronic interactions with the nucleus) due to  $m_J$ . In the  $jj$  coupling scheme, the term symbols specifying the angular momentum are thus written as

$$(j_1, j_2, \dots, j_n)_J, \quad (18.4.31)$$

where  $n$  is the number of electrons in the incomplete subshell.

The standard example of the  $jj$  coupling scheme is provided by the lead atom (Pb,  $Z = 82$ ). The electron configuration is  $[\text{Xe}]4f^{14}5d^{10}6s^26p^2$ , so what matters is the two  $l = 1$  electrons in



the unfilled  $6p$  subshell. Just as for carbon, the number of antisymmetric states that we can form from two  $p$  electrons in the same subshell is  $6 \cdot 5/2 = 15$ . Each electron has total angular momentum  $j = 1/2$  or  $3/2$ . Now we can use

$$\frac{1}{2} \otimes \frac{1}{2} = 0_A \oplus 1_S \quad (18.4.32)$$

$$\frac{3}{2} \otimes \frac{3}{2} = 0_A \oplus 1_S \oplus 2_A \oplus 3_S, \quad (18.4.33)$$

$$\frac{1}{2} \otimes \frac{3}{2} = 1 \oplus 2 \quad (18.4.34)$$

to find that the possible  $jj$  coupling terms for two  $p$  electrons are

$$(j_1, j_2)_J = \left(\frac{1}{2}, \frac{1}{2}\right)_0, \quad \left(\frac{1}{2}, \frac{3}{2}\right)_1, \quad \left(\frac{1}{2}, \frac{3}{2}\right)_2, \quad \left(\frac{3}{2}, \frac{3}{2}\right)_2, \quad \left(\frac{3}{2}, \frac{3}{2}\right)_0, \quad (18.4.35)$$

with degeneracies 1, 3, 5, 5, 1, respectively (which indeed add up to 15 as a check). Experimentally, it turns out that the order given in eq. (18.4.35) is in increasing energy, so the ground state has  $J = 0$ . By way of comparison, the Hund's rules prediction for the  $LS$ -coupling term for Pb is  $^3P_0$  (by an argument identical to the one for carbon), which also has  $J = 0$ , so the two schemes agree in that respect. However, the  $jj$  coupling scheme provides a somewhat better numerical approximation to the energy levels and other properties of the lowest few eigenstates of Pb than the  $LS$  coupling scheme does. For atoms with high  $Z$ , neither the  $LS$  scheme nor the  $jj$  scheme is very accurate, and more complicated descriptions are appropriate.

## 18.5 Exercises

**Exercise 18.1.** Two identical particles are moving in a 3-dimensional isotropic harmonic oscillator potential with natural frequency  $\omega$ . The only Hamiltonian interaction between the two particles is the spin-spin  $W = a\vec{S}_1 \cdot \vec{S}_2$ , where  $a$  is a constant.

- (a) If the particles are bosons with spin 1, find the energy, the spectroscopic term notation, and the degeneracy of the ground state. Give separate answers for positive and negative  $a$ .
- (b) Repeat part (a) if the particles have spin 0.
- (c) Repeat part (a) if the particles have spin 2.
- (d) Repeat part (a) if the particles are fermions with spin  $1/2$ .

**Exercise 18.2.** Two identical spin- $1/2$  fermions are free to move inside a cubic box of side  $a$  with one corner at the origin. The potential is infinite outside of the box. The fermions do not have any Hamiltonian interaction with each other.

- (a) The *one-particle* energy eigenstate wavefunctions are

$$\psi_{n_x, n_y, n_z}(x, y, z) = \left(\frac{2}{a}\right)^{3/2} \sin\left(\frac{n_x \pi x}{a}\right) \sin\left(\frac{n_y \pi y}{a}\right) \sin\left(\frac{n_z \pi z}{a}\right). \quad (18.5.1)$$



What are the corresponding one-particle energies?

(b) In the remainder of this problem, we will consider two-particle energy eigenstates, organized in terms of the total spin,  $S = 0$  or  $S = 1$ . But notice that in this problem we do not have spherical symmetry, and the Hamiltonian does not commute with  $L^2$  or  $L_z$ . Therefore, we cannot organize energy eigenstates in terms of orbital angular momentum. Instead, consider states that are either of the form

$$\chi_{S,m_S} \psi_N(x_1, y_1, z_1) \psi_N(x_2, y_2, z_2), \quad (18.5.2)$$

when the two particles have the same spatial wavefunction  $\psi_N$ , and

$$\chi_{S,m_S} \frac{1}{\sqrt{2}} [\psi_N(x_1, y_1, z_1) \psi_M(x_2, y_2, z_2) \pm \psi_M(x_1, y_1, z_1) \psi_N(x_2, y_2, z_2)], \quad (18.5.3)$$

if they have different spatial wavefunctions, where  $\chi_{S,m_S}$  is the spin ket for  $S = 0$  or  $S = 1$ , and  $N$  or  $M$  represents  $(n_x, n_y, n_z)$  for a given 1-particle wavefunction. Find the energy eigenvalues and the corresponding degeneracies for the lowest three energy levels of the two-fermion system. [Hints: the one-particle energies simply add to give the two-particle energies, since we are not including any interactions between the particles. Also, recall that for  $S = 0$ , the spatial wavefunction is symmetric under interchange of the labels 1 and 2, while for  $S = 1$ , the spatial wavefunction must be anti-symmetric under interchange of the labels 1 and 2. You should find that the total number of states in the lowest three energy levels is a number greater than 35.]

**Exercise 18.3.** Consider three identical spin-1/2 fermions bound in a 3-d isotropic harmonic oscillator potential characterized by frequency  $\omega$ . The fermions do not interact with each other.

(a) Use the Pauli exclusion principle to find the energies and the degeneracies of the ground state energy level and the first excited energy level. Give the spectroscopic term notation  $^{2S+1}L_J$  for the allowed states in each case. [Hint: combine two of the fermions, then add the third. For the ground state energy level, you should find two distinct spectroscopic terms. For the first excited state energy level, there are 11 spectroscopic terms, some of which are the same.]

(b) A spin-orbit perturbation Hamiltonian is added,  $W = a\vec{L} \cdot \vec{S}$ , where  $\vec{L}$  and  $\vec{S}$  are the total orbital and spin angular momenta of the fermions. What are the energies and spectroscopic term notations of the lowest three energy eigenstates states if  $a$  is positive? What if  $a$  is negative?

(c) Now suppose that in addition to the perturbation  $W$ , there is a much weaker perturbation  $W' = bL_z$ . Use degenerate perturbation theory to find the energy splitting of the lowest level you found in part (b), by finding the the matrix representation for  $W'$  and finding its eigenvalues. The cases of positive  $a$  and negative  $a$  are different, so treat them separately. [Hint: you may find eqs. (12.3.16)-(12.3.21) useful.]

**Exercise 18.4.** Compute the direct and exchange integrals  $I_{2,0,0}$  and  $J_{2,0,0}$ , as defined in eqs. (18.3.4) and (18.3.5), for the  $(1s)(2s)$  excited states of the helium atom. Note that from eqs. (11.1.42) and (11.1.43), the one-electron wavefunctions with  $Z = 2$  are  $\phi_{1,0,0}(r) = e^{-2r/a_0} \sqrt{8/\pi a_0^3}$  and  $\phi_{2,0,0}(r) = e^{-r/a_0}(1 - r/a_0)/\sqrt{\pi a_0^3}$ . Check that  $0 < J_{2,0,0} < I_{2,0,0}$ . Use these results to estimate the energies of the  $^1S_0$  and  $^3S_1$  states, and their energy splitting  $\Delta E = 2e^2 J_{2,0,0}$ , at first order in perturbation theory. Compare to the results quoted in Figure 18.3.1. [Hints: to set up the integrals, follow the example of the method used in eqs. (15.3.8)-(15.3.17). Before integrating, you are encouraged to change to dimensionless variables  $u = r_1/a_0$  and  $v = r_2/a_0$ . The perturbation theory result for  $\Delta E$  is far from a triumph, as you should find that it is numerically about a factor of 3 larger than the experimental value. However, it does have two important qualitative features: the correct sign, and a magnitude much smaller than the energy shift from the direct integral.]

**Exercise 18.5.** For each of calcium ( $Z = 20$ ), scandium ( $Z = 21$ ), titanium ( $Z = 22$ ), nickel ( $Z = 28$ ), germanium ( $Z = 32$ ) and selenium ( $Z = 34$ ), use the shell model to find the electronic configuration. Assume that the  $4s$  shell fills before the  $3d$  shell; although this is not always the case, it is true for these examples. In Russell-Saunders approximation, find the spectroscopic term notation for the electronic configurations you found, and apply Hund's rules to select the ground state in each case.

**Exercise 18.6.** In the Bizarro Universe, long ago and very far away, everything is just like in our universe except that electrons are fermions with spin  $3/2$  instead of spin  $1/2$ . Consider atomic states in the Bizarro Universe.

- How many Bizarro electrons fit into each  $s$ ,  $p$ , and  $d$  orbital ( $l = 0, 1, 2$ , respectively)?
- What are the atomic numbers of the lightest two Bizarro noble gases (the lightest has a filled  $1s$  orbital, the next has a filled  $2p$  orbital)? What about the lightest two Bizarro alkali metals (one  $s$ -subshell electron) and Bizarro halogens (one missing  $p$ -subshell electron)?
- Consider Bizarro neon, a neutral atom with  $Z = 10$ . What is the electron configuration, instead of  $(1s)^2(2s)^2(2p)^6$  as in our universe? What are all of the  $^{2S+1}L_J$  spectroscopic terms for the electron configuration you found, and which of them is the ground state?

## 19 Simple molecules

### 19.1 Hierarchies of scales and the Born-Oppenheimer approximation

Molecules are bound states consisting of multiple atoms, or more fundamentally electrons and nuclei. Understanding the quantum energy levels and structure of molecules is a complicated problem requiring strategic thinking, as there are many degrees of freedom.

One way of simplifying the problem is to make use of the hierarchies of energy and distance scales caused by the large ratio (more than 3 orders of magnitude) of the nuclear masses to the electron masses. In a first approximation, the slowly moving nuclei can be treated as essentially at rest on the time scales relevant for the motion of electrons. The nuclear displacements from their equilibrium positions are small compared to the size of the molecule. One can therefore solve first for the electronic states while treating the nuclei as fixed. The dynamics of the nuclear states can then be treated separately, by recognizing that the electrons will adjust themselves relatively quickly to changes in the nuclear positions. To determine the state describing the nuclear relative position degrees of freedom, one uses a potential that can be thought of as the result of averaging over the electronic positions, using the electronic wavefunction. This way of organizing the calculation in terms of a separation between light, fast degrees of freedom (the electrons) and heavy, slow degrees of freedom (the nuclei) is called the **Born-Oppenheimer approximation**, after the work of Max Born and J. Robert Oppenheimer in 1927. It is an example of a more general strategy called the **adiabatic approximation**, in which slow degrees of freedom are treated as nearly constant.

Let us start by making parametric estimates of the distance and energy scales involved. For fixed nuclear positions, dimensional analysis tells us that the characteristic size  $\Delta x_e$  of electronic states is comparable to the Bohr radius, since it is the only length scale that we can construct out of  $\hbar$ , the electron mass  $m$ , and the electronic charge  $e$ . The corresponding electronic energies  $E_e$  should be of order a Rydberg. So:

$$\Delta x_e \sim a_0 = \frac{\hbar^2}{e^2 m}, \quad E_e \sim \frac{e^2}{2a_0} = \frac{e^4 m}{2\hbar^2} = \frac{\hbar^2}{2ma_0^2}. \quad (19.1.1)$$

The vibrations of the nuclei about their equilibrium positions can be considered as determined by an effective harmonic oscillator with natural frequency that we can call  $\omega_N$ . To estimate it crudely, we note that if we were to displace the position of a nucleus by a distance of order  $a_0$ , the change in energy should be of order a Rydberg. Using  $M$  to denote the nuclear mass, we therefore estimate the energy of the oscillator as  $e^4 m / 2\hbar^2 \sim M\omega_N^2 a_0^2 / 2$ , which implies

$$\omega_N \sim \left(\frac{m}{M}\right)^{1/2} \frac{e^2}{\hbar a_0}. \quad (19.1.2)$$

The corresponding length and minimum energy scales associated with nuclear vibrations can therefore be estimated, using the harmonic oscillator results of eqs. (7.2.4) and (7.2.17), as

$$\Delta x_N \sim \sqrt{\frac{\hbar}{M\omega_N}} \sim \left(\frac{m}{M}\right)^{1/4} a_0, \quad E_{\text{vib}} \sim \hbar\omega_N \sim \left(\frac{m}{M}\right)^{1/2} \frac{e^2}{a_0}. \quad (19.1.3)$$

Molecules also have rotational degrees of freedom, whose minimum energy scale can be estimated by noting that the largest moment of inertia should be of order  $I \sim Ma_0^2$ , and the smallest angular momentum difference is of order  $L \sim \hbar$ . Therefore, we estimate

$$E_{\text{rot}} \sim \frac{L^2}{2I} \sim \frac{\hbar^2}{Ma_0^2} \sim \frac{m}{M} \frac{e^2}{a_0}. \quad (19.1.4)$$

The hierarchy of nuclear and electron masses therefore leads to a hierarchy of electronic, vibrational, and rotational energies, with very roughly

$$E_e : E_{\text{vib}} : E_{\text{rot}} = 1 : \left(\frac{m}{M}\right)^{1/2} : \frac{m}{M}. \quad (19.1.5)$$

Putting in the numbers, transitions between electronic states will result in emission of light that is typically in the UV or visible ranges, while transitions between nuclear vibrational modes are associated with the IR, and rotational modes with microwave radiation.

Consider a molecule consisting of  $N_e$  electrons and  $N_n$  nuclei. Let us work in position space, neglecting spin degrees of freedom for simplicity. The electrons have coordinates  $\vec{r}_i$  with  $i = 1, \dots, N_e$ , and the nuclei have coordinates  $\vec{R}_I$  where  $I = 1, \dots, N_n$ , with masses  $M_I$  and integer charges  $Z_I$ . To realize the Born-Oppenheimer strategy, the Hamiltonian can be divided into parts that do and do not involve the nuclear masses:

$$H = H_e + H_N, \quad (19.1.6)$$

$$H_e = -\sum_i \frac{\hbar^2 \nabla_i^2}{2m} + \frac{e^2}{2} \left( \sum_{i \neq j} \frac{1}{|\vec{r}_i - \vec{r}_j|} - 2 \sum_i \sum_I \frac{Z_I}{|\vec{r}_i - \vec{R}_I|} + \sum_{I \neq J} \frac{Z_I Z_J}{|\vec{R}_I - \vec{R}_J|} \right), \quad (19.1.7)$$

$$H_N = -\sum_I \frac{\hbar^2 \nabla_I^2}{2M_I}. \quad (19.1.8)$$

The first step in carrying out the Born-Oppenheimer approach is to find solutions for the electronic eigenvalue equation

$$H_e \phi_{n,\vec{R}}(\vec{r}_i) = \mathcal{E}_{n,\vec{R}} \phi_{n,\vec{R}}(\vec{r}_i), \quad (19.1.9)$$

in which the nuclear coordinates  $\vec{R}_I$  are treated as fixed, and the labels  $n$  distinguish different bound eigenstates. The presence of the subscript  $\vec{R}$  on the energy eigenvalues  $\mathcal{E}_{n,\vec{R}}$  and the multi-electron wavefunctions  $\phi_{n,\vec{R}}(\vec{r}_i)$  is to remind us that they depend on the choice of fixed  $\vec{R}_I$ . The

wavefunctions also depend simultaneously on all of the electronic coordinates  $\vec{r}_i$ . As noted above, since the nuclear masses are not involved in eq. (19.1.9), the eigenvalues  $\mathcal{E}_{n,\vec{R}}$  should have values roughly of the order of magnitude of a Rydberg, and the wavefunctions  $\phi_{n,\vec{R}}(\vec{r}_i)$  should vary appreciably over length scales comparable to the Bohr radius. Finding these solutions may be quite difficult, but with sufficient computing power they can be obtained using the variational principle. Note that the last term in eq. (19.1.7) proportional to  $Z_I Z_J$  is simply a constant in this part of the calculation.

Since  $\phi_{n,\vec{R}}(\vec{r}_i)$  are the eigenstates of the Hermitian operator  $H_e$ , they can be chosen as an orthonormal basis for the electronic bound state part of the Hilbert space. This means that we can always write the full (multi-electron, multi-nucleus) wavefunction as

$$\Psi(\vec{r}_i, \vec{R}_I) = \sum_n \psi_n(\vec{R}_I) \phi_{n,\vec{R}}(\vec{r}_i), \quad (19.1.10)$$

for some coefficients  $\psi_n(\vec{R}_I)$  that will serve as the wavefunctions for the nuclear position degrees of freedom. The next step is to look for eigenstates of this form for the full Hamiltonian  $H$  with eigenvalue  $E$ . Using eqs. (19.1.9) and (19.1.8) in (19.1.6), we have

$$\sum_n \left( - \sum_I \frac{\hbar^2 \nabla_I^2}{2M_I} + \mathcal{E}_{n,\vec{R}} - E \right) \left[ \psi_n(\vec{R}_I) \phi_{n,\vec{R}}(\vec{r}_i) \right] = 0. \quad (19.1.11)$$

Now using the product rule for Laplacians,  $\nabla^2(fg) = f\nabla^2g + g\nabla^2f + 2(\vec{\nabla}f) \cdot (\vec{\nabla}g)$ , this becomes

$$\begin{aligned} \sum_n \phi_{n,\vec{R}}(\vec{r}_i) \left( - \sum_I \frac{\hbar^2 \nabla_I^2}{2M_I} + \mathcal{E}_{n,\vec{R}} - E \right) \psi_n(\vec{R}_I) = \\ \sum_n \sum_I \frac{\hbar^2}{M_I} \left[ (\vec{\nabla}_I \psi_n) \cdot (\vec{\nabla}_I \phi_{n,\vec{R}}) + \frac{1}{2} \psi_n \nabla_I^2 \phi_{n,\vec{R}} \right]. \end{aligned} \quad (19.1.12)$$

The Born-Oppenheimer approximation entails neglecting the right-hand side of this equation, with the partial justification that the amplitude of nuclear vibrations is smaller than the distances between nuclei by a factor of order  $(m/M)^{1/4}$ , as noted in eq. (19.1.3). To see how this works, recall that the wavefunction for an electronic state in a hydrogen atom, for example, involves a factor  $e^{-r/a_0}$ , where  $r$  is the separation between the nucleus and an electron. Therefore, considering the effect of changing a nuclear position slightly, we expect

$$\frac{|\vec{\nabla}_I \phi_{n,\vec{R}}|}{\phi_{n,\vec{R}}} \sim \frac{1}{a_0}. \quad (19.1.13)$$

In comparison, we can approximate the wavefunction for each nucleus  $I$  by that of a harmonic oscillator with angular frequency  $\omega_I = (m/M_I)^{1/2} e^2 / \hbar a_0$  as in eq. (19.1.2), so that  $\psi_n$  is proportional to  $e^{-M_I \omega_I |\vec{R}_I - \vec{R}_{I0}|^2 / 2\hbar}$ , where  $\vec{R}_{I0}$  is the equilibrium position of the nucleus. We then

estimate  $|\vec{\nabla}_I \psi_n|/\psi_n \sim |\vec{R}_I - \vec{R}_{I0}| M_I \omega_I / \hbar$ . Inserting the estimates for  $\omega_I$  and  $\Delta x_N = |\vec{R}_I - \vec{R}_{I0}|$  from eqs. (19.1.2) and (19.1.3), we obtain

$$\frac{|\vec{\nabla}_I \psi_n|}{\psi_n} \sim \left( \frac{M_I}{m} \right)^{1/4} \frac{1}{a_0} \quad (19.1.14)$$

Thus, each transfer of a derivative  $\vec{\nabla}_I$  from the nuclear wavefunction  $\psi_n$  to the electronic wavefunction  $\phi_n$  costs a relative suppression factor of  $(m/M_I)^{1/4}$ , in accord with the intuition that the heavy nuclei are more sharply localized.

Replacing the right side of eq. (19.1.12) by 0, we realize the added benefit that since the wavefunctions  $\phi_{n,\vec{R}}(\vec{r}_i)$  form an orthonormal basis, each term in the sum over  $n$  must vanish separately, and so we can write

$$\left( -\sum_I \frac{\hbar^2 \nabla_I^2}{2M_I} + \mathcal{E}_{n,\vec{R}} - E \right) \psi_n(\vec{R}_I) = 0 \quad (19.1.15)$$

as the energy eigenstate equation for the nuclear excitations. For each electronic state labeled by  $n$ , there is a family of nuclear vibrational and rotational states and energies associated with the solutions to eq. (19.1.15). In this Schrödinger-like equation, the  $\mathcal{E}_{n,\vec{R}}$ , obtained by solving for the electronic state energies, serve as the potential for the nuclear wavefunction.

To recap, the Born-Oppenheimer approximation separates the molecular problem into solving eq. (19.1.9) for the electronic states with fixed nuclei, and eq. (19.1.15) for the nuclear degrees of freedom with the electronic states providing the potential. While it is tempting to say that we have treated the nuclear kinetic term  $H_N$  in eq. (19.1.8) as a perturbation, it should be recognized that this is not the same sort of perturbation theory as in Chapter 15. As we have seen, the Born-Oppenheimer approach is an expansion in  $(m/M)^{1/4}$ , while standard perturbation theory would instead imply an expansion in powers of  $m/M$ . (The latter type of perturbative expansion does not work here, because the dynamics of the nuclear degrees of freedom cannot be well-approximated by an unperturbed Hamiltonian without any kinetic energy term.)

In the remaining sections of this chapter, we will discuss only a few minimal aspects of molecular physics, restricted to simple cases. First, in section 19.2, we will find an approximation for the solution of eq. (19.1.9) for the electronic state of the minimal example of a molecule, the  $\text{H}_2^+$  ion. In section 19.3, we will then treat the vibrational nuclear state determination of eq. (19.1.15) for more general diatomic molecules, based on a widely used and empirically successful potential. The general subject of quantized rigid body rotations, and the application to the rotational energy levels of molecules, is given in section 19.4. Section 19.5 discusses the vibrational and rotational spectra for diatomic molecules, including numerical parameters for some common ones. Section 19.6 contains a brief account of the rotational and vibrational excitations for the water molecule  $\text{H}_2\text{O}$ .

## 19.2 The simplest molecule: the $\text{H}_2^+$ ion

The  $\text{H}_2^+$  ion is the simplest molecule, consisting of one electron and two protons. In the Born-Oppenheimer approximation, we first fix the positions of the protons at  $\vec{R}_A$  and  $\vec{R}_B$ , with the distance between them denoted by

$$R = |\vec{R}_A - \vec{R}_B|, \quad (19.2.1)$$

and describe the electron's position with a vector  $\vec{r}$  with corresponding Laplacian  $\nabla^2$ . The Hamiltonian for the electron's state is, in the position representation,

$$H_e = -\frac{\hbar^2 \nabla^2}{2m} + e^2 \left( -\frac{1}{|\vec{r} - \vec{R}_A|} - \frac{1}{|\vec{r} - \vec{R}_B|} + \frac{1}{R} \right), \quad (19.2.2)$$

with the corresponding geometry shown in Figure 19.2.1. Our first goal is to estimate the ground-state energy  $\mathcal{E}_R$  and wavefunction  $\phi_R(\vec{r})$ , using a variational principle. The resulting  $\mathcal{E}_R$  then serves as the potential energy for the nuclear separation, as we discussed in eq. (19.1.15), the second part of the Born-Oppenheimer approximation. From this, we will determine the binding energy and the nuclear separation of the  $\text{H}_2^+$  ion.




Figure 19.2.1: The  $\text{H}_2^+$  ion molecule consists of two protons with charges  $+e$  at positions  $\vec{R}_A$  and  $\vec{R}_B$ , and an electron with charge  $-e$  at position  $\vec{r}$ . In the Born-Oppenheimer approximation, we first treat the proton positions  $\vec{R}_A$  and  $\vec{R}_B$  as fixed, to find the binding potential energy as a function of  $R = |\vec{R}_A - \vec{R}_B|$ .

To construct a trial state, let us start with unit-norm atomic orbital wavefunctions

$$\varphi_A(\vec{r}) = \left( \frac{Z}{a_0} \right)^{3/2} \frac{1}{\sqrt{\pi}} e^{-Z|\vec{r} - \vec{R}_A|/a_0}, \quad \varphi_B(\vec{r}) = \left( \frac{Z}{a_0} \right)^{3/2} \frac{1}{\sqrt{\pi}} e^{-Z|\vec{r} - \vec{R}_B|/a_0}. \quad (19.2.3)$$

These have the same form as the ground state of a hydrogen atom centered at each of the two proton positions, except that we have included a variational parameter  $Z$ , which governs the size of the orbitals. Our trial wavefunction  $\phi(\vec{r})$  will be a linear combination of  $\varphi_A$  and  $\varphi_B$ . Without loss of generality, we can take

$$\vec{R}_B = -\vec{R}_A = (R/2)\hat{z}, \quad (19.2.4)$$

so that the Hamiltonian has even parity,  $[\Pi, H] = 0$ . This ensures that the energy eigenstates can be chosen to have well-defined parity  $= \pm 1$ , so we try un-normalized variational states

$$|\phi_{\pm}\rangle = |\varphi_A\rangle \pm |\varphi_B\rangle, \quad (19.2.5)$$



with  $\Pi|\phi_{\pm}\rangle = \pm|\phi_{\pm}\rangle$ . From here on, the trial state and energy will be denoted just  $|\phi\rangle$  and  $\mathcal{E}_{Z,R}$  (without the subscript  $\pm$ ) to avoid clutter, with the understanding that we are trying both parity possibilities simultaneously.

Applying the variational principle of Chapter 16, the ground-state energy estimate is

$$\mathcal{E}_{Z,R} = \frac{\langle\phi|H_e|\phi\rangle}{\langle\phi|\phi\rangle} = \frac{\langle\varphi_A|H_e|\varphi_A\rangle \pm \langle\varphi_B|H_e|\varphi_A\rangle}{1 \pm \langle\varphi_B|\varphi_A\rangle}. \quad (19.2.6)$$

In writing the last expression, we have used the facts that  $\langle\varphi_A|\varphi_A\rangle = \langle\varphi_B|\varphi_B\rangle = 1$ , and  $\langle\varphi_A|\varphi_B\rangle = \langle\varphi_B|\varphi_A\rangle$ , since the atomic orbital wavefunctions are normalized and real, and  $\langle\varphi_A|H_e|\varphi_B\rangle = \langle\varphi_B|H_e|\varphi_A\rangle$ , since  $H_e$  is Hermitian, and  $\langle\varphi_B|H_e|\varphi_B\rangle = \langle\varphi_A|H_e|\varphi_A\rangle$  by symmetry, since the orbitals have the same form up to a displacement by distance  $R$ .

The electronic state depends on two parameters:  $R$  (the distance between the protons, which is temporarily fixed as part of the Born-Oppenheimer approximation) and  $Z$  (the variational parameter). Before doing the detailed calculations, let us consider two extreme special cases of  $R$  in which the problem simplifies and we know what to expect. First, if  $R$  is very large compared to  $a_0$ , then the electron will essentially be forced to choose between the two protons, and the ground state can be thought of as a symmetrized tensor product between a hydrogen atom and a lonely distant proton. In that limit, we expect that  $Z$  will be very close to 1, and the eigenvalue of  $H_e$  will be almost exactly  $-1$  Rydberg  $= -e^2/2a_0$ . In the opposite extreme case that  $R$  is very small compared to  $a_0$ , the situation is the same as if there were only a single nucleus with charge  $+2e$ , so we expect that  $Z$  will approach 2 in that limit. This will be accompanied by a diverging positive contribution to the  $H_e$  eigenvalue, due to the repulsive energy between the two protons (the last term in eq. (19.2.2)).

For intermediate  $R$ , the minimum  $\mathcal{E}_R$  should be achieved for  $1 < Z < 2$ . To minimize the energy, the preferred position for the electron is between the two protons, where it can benefit from its attraction to them both. However, for the odd-parity case, the wavefunction  $\phi(\vec{r})$  vanishes at the midpoint  $\vec{r} = 0$ , and therefore it must be small everywhere nearby, by continuity. In the even-parity case, the wavefunction is enhanced in the good region between the protons due to constructive interference. We therefore expect the ground state with strongest binding to have even parity. The essential reason for the bond is that the electron can be shared by the two nuclei to which it is attracted, lowering the potential energy.

To proceed, we need to find the matrix elements  $\langle\varphi_B|\varphi_A\rangle$ ,  $\langle\varphi_A|H_e|\varphi_A\rangle$ , and  $\langle\varphi_B|H_e|\varphi_A\rangle$  in eq. (19.2.6). In each case, the evaluation in the position representation involves integrals with integrands that depend on  $|\vec{r} - \vec{R}_A|$  and  $|\vec{r} - \vec{R}_B|$ , which at first might appear severely daunting. Fortunately, things simplify if we carry out the integrations using **elliptical coordinates**  $(\mu, \nu, \phi)$ , of which the last is the usual spherical coordinate angle measured around the  $z$ -axis,



and the first two are defined by

$$|\vec{r} - \vec{R}_A| = (\mu + \nu)R/2, \quad |\vec{r} - \vec{R}_B| = (\mu - \nu)R/2. \quad (19.2.7)$$

This is convenient because the integrands for the needed matrix elements are simple functions of  $\mu$  and  $\nu$ . Using  $\vec{R}_B = -\vec{R}_A = (R/2)\hat{z}$  one finds the ranges  $1 \leq \mu < \infty$  and  $-1 \leq \nu \leq 1$ , and

$$r = \sqrt{\mu^2 + \nu^2 - 1} R/2, \quad \cos \theta = \mu\nu / \sqrt{\mu^2 + \nu^2 - 1}, \quad (19.2.8)$$

from which the volume element can be found, by computing the Jacobian determinant, to be

$$d^3\vec{r} = r^2 dr d(\cos \theta) d\phi = \frac{R^3}{8}(\mu^2 - \nu^2) d\mu d\nu d\phi. \quad (19.2.9)$$

It is also convenient to define a rescaled constant distance between the protons,

$$\rho = ZR/a_0. \quad (19.2.10)$$

With these variable changes, the needed matrix elements are expressed as the integrals

$$\langle \varphi_B | \varphi_A \rangle = \frac{\rho^3}{4} \int_1^\infty d\mu \int_{-1}^1 d\nu (\mu^2 - \nu^2) e^{-\rho\mu}, \quad (19.2.11)$$

$$\langle \varphi_A | H_e | \varphi_A \rangle = \frac{e^2}{2a_0} Z \rho^2 \int_1^\infty d\mu \int_{-1}^1 d\nu f(\mu, \nu) e^{-\rho(\mu+\nu)}, \quad (19.2.12)$$

$$\langle \varphi_B | H_e | \varphi_A \rangle = \frac{e^2}{2a_0} Z \rho^2 \int_1^\infty d\mu \int_{-1}^1 d\nu f(\mu, \nu) e^{-\rho\mu}, \quad (19.2.13)$$

with the last two involving a common polynomial factor

$$f(\mu, \nu) = \left( \frac{1}{2} - \frac{Z\rho}{4} \right) (\mu^2 - \nu^2) + (Z - 2)\mu - Z\nu \quad (19.2.14)$$

coming from evaluating  $H_e \varphi_A(\vec{r})$  in the position representation. Doing the integrations gives

$$\langle \varphi_B | \varphi_A \rangle = (1 + \rho + \rho^2/3) e^{-\rho}, \quad (19.2.15)$$

$$\langle \varphi_A | H_e | \varphi_A \rangle = \frac{e^2}{2a_0} Z [Z - 2 + 2(1 + 1/\rho)e^{-2\rho}], \quad (19.2.16)$$

$$\langle \varphi_B | H_e | \varphi_A \rangle = \frac{e^2}{2a_0} Z [2/\rho - 2 - 10\rho/3 + Z(1 + \rho - \rho^2/3)] e^{-\rho}. \quad (19.2.17)$$

The resulting energy as a function of the variational parameter  $Z$  is therefore

$$\mathcal{E}_{Z,R} = \frac{e^2}{2a_0} \frac{Z \{ [Z - 2 + 2(1 + 1/\rho)e^{-2\rho}] \pm [2/\rho - 2 - 10\rho/3 + Z(1 + \rho - \rho^2/3)] e^{-\rho} \}}{1 \pm (1 + \rho + \rho^2/3)e^{-\rho}}. \quad (19.2.18)$$

Now, for each  $R$ , one can minimize  $\mathcal{E}_{Z,R}$  with respect to  $Z$  to obtain the energy  $\mathcal{E}_R$ , which according to eq. (19.1.15) serves as the Schrödinger equation potential for determining the nuclear separation and vibrational states.




Figure 19.2.2: The energy potential of the  $\text{H}_2^+$  ion ground state,  $\mathcal{E}_R + e^2/2a_0$ , as a function of the nuclear separation  $R$  in units of the Bohr radius, as estimated by the variational principle for the even-parity binding (solid curve) and odd-parity anti-binding (dashed curve) solutions. For each  $R$ , the energy is obtained by minimizing eq. (19.2.18) with respect to the variational parameter  $Z$ . The minimum is  $\mathcal{E}_{R,\min} + e^2/2a_0 = -2.35$  eV, obtained for  $R/a_0 = 2.003$  and  $Z = 1.238$ . The actual experimental binding energy of the  $\text{H}_2^+$  ion is  $E = -2.79$  eV.

The minimization is best done by computer; note that there is a dependence on  $Z$  hidden in the definition of  $\rho$ . The resulting  $\mathcal{E}_R + e^2/2a_0$  for each of the two parity choices is shown in Figure 19.2.2. The odd-parity state has its minimum at  $R = \infty$ , and so is rejected, as an anti-binding solution. For the even-parity binding state, for very large  $R$  one finds  $Z = 1$  and  $\mathcal{E}_R = -e^2/2a_0$ , and for very small  $R$  one finds  $Z \rightarrow 2$  and a large positive energy from the proton-proton repulsion term, as expected. The minimum occurs for

$$R/a_0 = 2.003, \quad Z = 1.238, \quad \mathcal{E}_{R,\min} + e^2/2a_0 = -2.35 \text{ eV}. \quad (19.2.19)$$

This is about 16% higher than the experimental binding energy of the  $\text{H}_2^+$  ion,  $E = -2.79$  eV. As usual, the variational method gives an upper bound on the energy, and using a trial wavefunction with more parameters and a more general shape would get us even closer to the true value. However, our simple try is already sufficient to get a reasonable approximation with the correct features of the exact answer.

A similar approach can be used to find the binding energy and the nuclear separation potential for the neutral  $\text{H}_2$  molecule with two electrons and two protons, which proceeds similarly but with slightly more involved integrals. This also holds for more complicated molecules with more complicated wavefunctions. Discussions can be found in more specialized books.

### 19.3 Diatomic bonds and vibrations from the Morse potential

The example in the preceding section illustrates the calculation of the potential energy that determines the properties of the wavefunctions associated with the nuclear-position wavefunction, in the Born-Oppenheimer approximation. In principle, we can now use this sort of potential to find the expectation value of the nuclear separation  $R$ , and the excitation energies associated

with nuclear vibrations and rotations. However, in practice the potential we found in Figure 19.2.2 is only known approximately and numerically after minimizing with respect to the variational parameter  $Z$ , even in the relatively easy case of  $\text{H}_2^+$ . It is useful to have a simpler analytic potential  $V(R)$  with the same qualitative features, containing parameters that can either be estimated from the variational principle applied to the electronic wavefunction, or determined empirically by experiment, for general diatomic molecules. The appropriate qualitative features are that the potential should:

- diverge positively for  $R \rightarrow 0$ , due to the electrostatic repulsion of the nuclei,
- have a stable minimum for some  $R$  comparable to the Bohr radius,
- approach a constant for  $R \rightarrow \infty$ , where the molecule separates into two atoms,
- have a large, but finite, number of vibrational states with nearly equal energy differences, in accord with experimental observations.

The last two points are consistent with the intuition that if we try to put too much energy into the molecule, we will break it into its atomic constituents. All of these qualitative features are realized by the approximate potential that we found for the  $\text{H}_2^+$  ion in Figure 19.2.2.

For general diatomic molecules, there is a simple three-parameter form, called the **Morse potential** after its invention by Philip M. Morse in 1929, which is both tractable for calculations and empirically successful. It is, using  $r$  from here on to denote the nuclear separation distance,

$$V(r) = V_0 [1 - e^{-\beta(r-a)}]^2. \quad (19.3.1)$$

Here  $V_0$  is the depth of the potential,  $a$  is the position of its minimum where  $V(r) = 0$ , and  $\beta$  is the parameter governing the restoring force for small oscillations about the minimum. For large  $r$  the potential approaches  $V_0$ , which is therefore the classical approximation to the energy needed to break up the molecule. For small  $r$ , the potential approaches  $V_0(e^{\beta a} - 1)^2$ , which is positive, but not infinite. However, this slight deviation from the first of our desired features is of little practical consequence, because experience shows that  $e^{\beta a} \gg 1$  for molecules, and thus the potential is so large at  $r = 0$  that it does not matter exactly how large it is; the probability density for small  $r$  is tiny in any case.

For small deviations about the minimum position, the Morse potential is close to that of a harmonic oscillator,

$$V(r) \approx V_0 \beta^2 (r - a)^2. \quad (19.3.2)$$

If we denote the two nuclear masses as  $M_1$  and  $M_2$ , then the reduced mass for the relative two-body problem (as discussed in section 4.2) can be defined as

$$M = M_1 M_2 / (M_1 + M_2). \quad (19.3.3)$$

Then, by setting eq. (19.3.2) equal to  $\frac{1}{2}M\omega^2(r-a)^2$ , we see that low-level vibrational excitations should be associated with a harmonic oscillator angular frequency

$$\omega = \sqrt{\frac{2V_0}{M}}\beta. \quad (19.3.4)$$

However, because of the requirement that the molecule can be broken up into its constituent atoms by adding finite energy  $V_0$ , the potential necessarily includes an anharmonicity which will become increasingly important for higher energy vibrational states.

Let us now solve for the energy eigenvalues and wavefunctions of the Morse potential, taking the orbital angular momentum to be 0 so that the wavefunction depends only on  $r$ . (Rotational excitations have relatively small energies, suppressed by a factor  $(m/M)^{1/2}$  as noted in section 19.1, and will be discussed in the next section.) Writing  $\psi(r) = U(r)/r$  as in section 10.1, the time-independent Schrödinger equation becomes

$$\left(-\frac{\hbar^2}{2M} \frac{d^2}{dr^2} + V_0 [1 - e^{-\beta(r-a)}]^2\right) U = EU, \quad (19.3.5)$$

for energy eigenvalues  $E$ . We now define dimensionless quantities

$$x = \beta(r - a), \quad \kappa = \sqrt{2MV_0}/\hbar\beta, \quad \mathcal{E} = E/V_0, \quad (19.3.6)$$

which transforms the eigenvalue equation into

$$\frac{d^2 U}{dx^2} + \kappa^2 [\mathcal{E} - (1 - e^{-x})^2] U = 0, \quad (19.3.7)$$

to be solved in the domain  $-\beta a < x < \infty$ . The boundary conditions are that  $U = 0$  for  $x = -\beta a$  (so that  $\psi$  is finite at  $r = 0$ ) and at  $x = \infty$  (so that we have a bound state). However, in order to find exact solutions, we cheat slightly by extending the domain to  $-\infty < x < \infty$ , and impose boundary conditions  $U(-\infty) = U(\infty) = 0$ , the same as for a 1-dimensional bound state. As mentioned above, this modification makes no practical difference for  $e^{\beta a} \gg 1$ , and it will allow us to solve the problem exactly.

To get rid of the pesky exponential, now make the further change of variable

$$z = \kappa e^{-x}, \quad \frac{d}{dx} = -z \frac{d}{dz}, \quad (19.3.8)$$

so that the differential equation becomes

$$\frac{d^2 U}{dz^2} + \frac{1}{z} \frac{dU}{dz} + \left[ \frac{\kappa^2 \mathcal{E}}{z^2} - \left(1 - \frac{\kappa}{z}\right)^2 \right] U = 0. \quad (19.3.9)$$

The domain boundaries, where we require that  $U$  vanishes, are at  $z = 0$  (corresponding to  $r \rightarrow \infty$ ) and  $z = \infty$  (corresponding, approximately, to  $r = 0$ ).

The remaining process of finding the exact eigensolutions  $U$  and  $\mathcal{E}$  is quite reminiscent of the methods we used to find the bound states of the 1-d harmonic oscillator in section 7.2, the isotropic 3-d harmonic oscillator in section 10.5, so we will be brief. First, we note that for large  $z$ , the differential equation (19.3.9) becomes just  $U'' \approx U$ , with solutions proportional to  $e^{\pm z}$ . Choosing the negative exponent in order to have a normalizable bound state with good behavior at large  $z$ , we therefore write

$$U(z) = e^{-z}S(z), \quad (19.3.10)$$

where  $S(z)$  must not grow exponentially at large  $z$  and must satisfy the differential equation

$$\frac{d^2S}{dz^2} + \left(\frac{1}{z} - 2\right) \frac{dS}{dz} + \left(\frac{\kappa^2(\mathcal{E} - 1)}{z^2} + \frac{2\kappa - 1}{z}\right) S = 0. \quad (19.3.11)$$

For small  $z$ , trying a power law  $S \approx c_0 z^p$  gives  $p^2 = \kappa^2(1 - \mathcal{E})$ . The positive root for  $p$  must be the correct one, in order to satisfy the boundary condition  $S = 0$  at  $z = 0$ , so

$$p = \kappa\sqrt{1 - \mathcal{E}}, \quad (19.3.12)$$

which is consistent with the requirement that  $\mathcal{E} = E/V_0 < 1$  in order to have a bound state. Therefore, let us try a power series solution

$$S(z) = z^p \sum_{n=0}^{\infty} c_n z^n. \quad (19.3.13)$$

Substituting this into eq. (19.3.11), we find the recurrence relation

$$c_{n+1}/c_n = \frac{1 + 2n + 2p - 2\kappa}{(1 + n + p)^2 - p^2}. \quad (19.3.14)$$

Now we can make the standard argument that the series must terminate so that  $S(z)$  is a polynomial; otherwise  $c_{n+1}/c_n \approx 2/n$  for large  $n$ , and we would find  $S(z) \sim e^{2z}$ . Therefore, the allowed energy eigenvalues  $\mathcal{E}_n$  correspond to the vanishing of the numerator of eq. (19.3.14) for some nonnegative integer  $n = 0, 1, 2, 3, \dots$ . Plugging in eq. (19.3.12) gives

$$\sqrt{1 - \mathcal{E}_n} = 1 - \frac{n + 1/2}{\kappa}. \quad (19.3.15)$$

Since the left side is positive, there can be only a finite number of bound state energy levels, with

$$n < \kappa - 1/2. \quad (19.3.16)$$

Solving eq. (19.3.15) for  $\mathcal{E}_n$  and multiplying by  $V_0$ , we finally obtain the remarkably simple result for the vibrational energy levels of the Morse potential,

$$E_n = \hbar\omega(n + 1/2) - \hbar\omega\chi(n + 1/2)^2, \quad (19.3.17)$$

where  $\omega$  was defined in eq. (19.3.4), and the dimensionless number

$$\chi = \frac{1}{2\kappa} = \frac{\hbar\beta}{2\sqrt{2MV_0}} \quad (19.3.18)$$

is called the **anharmonicity parameter**. For molecules,  $\chi$  is generally quite small. Thus, for small  $n$  so that the  $\chi$  term can be neglected, the energy levels  $E_n$  are nearly equally spaced, with behavior close to that of a 1-d harmonic oscillator corresponding to vibrations of the nuclei along their common axis. The effect of the anharmonicity is to decrease the energy spacings as  $n$  increases to its maximum allowed value, with  $E_n$  approaching  $V_0$  from below. The total number of vibrational states is the largest integer less than  $\kappa + 1/2$ . Data from fits to  $\hbar\omega$  and  $\chi$  for a few common molecules will be shown in section 19.5, after we discuss the much smaller rotational contributions to the molecular energy, in the next section.

## 19.4 Rigid bodies and molecular rotational states

In the preceding section, we considered molecular vibrations of diatomic molecules corresponding to nuclear oscillations along their common axis. We now turn to rotational excitations of molecules. This could be done by adding an angular momentum term  $\hbar^2 l(l+1)/2Mr^2$  to the Morse potential Schrödinger equation (19.3.5), and then finding solutions with non-zero total angular momentum quantum number  $l$ . However, in keeping with the Born-Oppenheimer philosophy, we will instead follow the somewhat simpler path of treating the molecule as a rigid body with fixed electronic state and nuclear separation, and moments of inertia of order  $Ma_0^2$ . We start with a general discussion of the quantum mechanics of the rotations of rigid bodies.

Consider a rigid body with moments of inertia  $I_A$ ,  $I_B$ , and  $I_C$  along three principal axes with  $\hat{A}$ ,  $\hat{B}$ , and  $\hat{C}$ , as in Figure 19.4.1. These are orthogonal unit vectors that are body-fixed (constant in the rotating frame of the body), and satisfy  $\hat{A} \cdot \hat{A} = \hat{B} \cdot \hat{B} = \hat{C} \cdot \hat{C} = 1$ , and  $\hat{A} \cdot \hat{B} = \hat{A} \cdot \hat{C} = \hat{B} \cdot \hat{C} = 0$ , and  $\hat{A} \times \hat{B} = \hat{C}$ , etc. In quantum mechanics, we treat  $\hat{A}$ ,  $\hat{B}$ ,  $\hat{C}$




Figure 19.4.1: The orientation of a rigid body with respect to a space-fixed coordinate system frame  $(\hat{x}, \hat{y}, \hat{z})$  can be specified by two orthogonal unit body-fixed vectors  $\hat{A}$ ,  $\hat{B}$ , and a third  $\hat{C} = \hat{A} \times \hat{B}$ . We choose them to be the principal axis directions, which means that the moment-of-inertia tensor is diagonal with components  $I_A$ ,  $I_B$ , and  $I_C$ .

as operators. Since they are each vectors, we know that their commutation relations with the components of the angular momentum operator  $\vec{L}$  are, as in eq. (13.1.12),

$$[L_a, \hat{A}_b] = i\hbar\epsilon_{abc}\hat{A}_c, \quad [L_a, \hat{B}_b] = i\hbar\epsilon_{abc}\hat{B}_c, \quad [L_a, \hat{C}_b] = i\hbar\epsilon_{abc}\hat{C}_c, \quad (19.4.1)$$

for  $a, b, c = x, y, z$ . In terms of the total angular momentum operator  $\vec{L}$  for the body, we define the principal-axis angular momentum components

$$\mathcal{L}_A = \hat{A} \cdot \vec{L}, \quad \mathcal{L}_B = \hat{B} \cdot \vec{L}, \quad \mathcal{L}_C = \hat{C} \cdot \vec{L}. \quad (19.4.2)$$

Referring to any good classical mechanics textbook, one finds that the rotational kinetic energy of the rigid body is then

$$H = \frac{\mathcal{L}_A^2}{2I_A} + \frac{\mathcal{L}_B^2}{2I_B} + \frac{\mathcal{L}_C^2}{2I_C}, \quad (19.4.3)$$

whose quantum mechanical eigenvalues we wish to find.

It is important to note that  $\mathcal{L}_{A,B,C}$  are not the components of a vector operator, but are instead three separate scalar operators, because they satisfy

$$[\vec{L}, \mathcal{L}_A] = [\vec{L}, \mathcal{L}_B] = [\vec{L}, \mathcal{L}_C] = 0. \quad (19.4.4)$$

It is left to Exercise 19.2 to verify this, using eq. (19.4.1) and  $[L_a, L_b] = i\hbar\epsilon_{abc}L_c$ , and to check that

$$[\mathcal{L}_A, \mathcal{L}_B] = -i\hbar\mathcal{L}_C, \quad [\mathcal{L}_B, \mathcal{L}_C] = -i\hbar\mathcal{L}_A, \quad [\mathcal{L}_C, \mathcal{L}_A] = -i\hbar\mathcal{L}_B, \quad (19.4.5)$$

where the minus signs are notable, and not errors. Thus, the principal-axis components  $-\mathcal{L}_A$ ,  $-\mathcal{L}_B$  and  $-\mathcal{L}_C$  satisfy the same commutation algebra as the ordinary fixed-frame rectangular components  $L_x, L_y, L_z$ , and therefore, together with

$$\mathcal{L}^2 = \mathcal{L}_A^2 + \mathcal{L}_B^2 + \mathcal{L}_C^2 = L^2, \quad (19.4.6)$$

have the same solutions for their eigenvalue problem. In particular, if  $L^2$  has eigenvalue  $\hbar^2 l(l+1)$ , then the allowed eigenvalues of  $-\mathcal{L}_C$  are  $-\hbar k$ , where  $k$  is an integer with magnitude less than or equal to  $l$ . We can choose as a CSCO for the rigid body the mutually commuting operators

$$L^2, L_z, \mathcal{L}_C, \quad (19.4.7)$$

with simultaneous eigenvalues, respectively,

$$\hbar^2 l(l+1), \quad \hbar m, \quad \hbar k, \quad (19.4.8)$$

where  $l$ ,  $m$ , and  $k$  are integers, with  $-l \leq m, k \leq l$ . Since  $L^2$  and  $L_z$  both commute with the Hamiltonian  $H$  in eq. (19.4.3), each energy level can be assigned one or more quantum numbers  $l$ , and there is always a degeneracy of at least  $2l + 1$  associated with the quantum number  $m$ .

As a particularly simple example, consider the **spherical top** with all three moments of inertia the same,  $I_A = I_B = I_C = I$ . In that case, the Hamiltonian is proportional to  $L^2$  and so commutes with all three CSCO operators, and the energy eigenvalues of the Hamiltonian in eq. (19.4.3) for a given  $l$  are simply

$$E_l = \frac{\hbar^2 l(l+1)}{2I}. \quad (19.4.9)$$

There is a degeneracy  $2l + 1$  due to the  $m$  quantum number (the orientation of the angular momentum with respect to fixed space) and another degeneracy  $2l + 1$  due to the  $k$  quantum number (the orientation of the angular momentum with respect to the body-fixed axes), giving a total degeneracy of  $(2l + 1)^2$  for the energy level  $E_l$ . The molecule  $\text{CH}_4$  (methane) is an example of a spherical top; although it is of course not spherical in shape, its tetrahedral symmetry dictates that its three moments of inertia are identical.

In the opposite extreme, suppose that none of the moments of inertia  $I_{A,B,C}$  coincide, an **asymmetric top**. A typical example is the  $\text{H}_2\text{O}$  molecule to be considered in section 19.6. In that case,  $H$  commutes with  $L^2$  and  $L_z$  but not with  $\mathcal{L}_C$ . There is no simple way of writing the energy eigenvalues for general  $l, m$ , but for any specific  $l$  one can diagonalize the Hamiltonian as a  $(2l + 1) \times (2l + 1)$  matrix. For example, if  $l = 1$ , then with the help of eqs. (8.3.2) and (8.3.4) with  $J_{x,y,z} \rightarrow \mathcal{L}_{A,B,C}$ , one finds the matrix representation in the basis of eigenstates of  $\mathcal{L}_C$ ,

$$H = \frac{\hbar^2}{4I_A} \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 1 \end{pmatrix} + \frac{\hbar^2}{4I_B} \begin{pmatrix} 1 & 0 & -1 \\ 0 & 2 & 0 \\ -1 & 0 & 1 \end{pmatrix} + \frac{\hbar^2}{2I_C} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (19.4.10)$$

with the energy eigenvalues

$$E_1 = \frac{\hbar^2}{2} \left( \frac{1}{I_A} + \frac{1}{I_B} \right), \quad E_2 = \frac{\hbar^2}{2} \left( \frac{1}{I_A} + \frac{1}{I_C} \right), \quad E_3 = \frac{\hbar^2}{2} \left( \frac{1}{I_B} + \frac{1}{I_C} \right). \quad (19.4.11)$$

Each of these three energies has degeneracy  $2l + 1 = 3$ , associated with the  $m$  quantum number. For larger  $l$ , the non-linear nature of the energy eigenvalue equation leads to irregular spacings between energies. The  $l = 2$  case is left to Exercise 19.3.

Now consider the case of a **symmetric top**, with  $I_A = I_B \neq I_C$ . This allows us to rewrite the Hamiltonian eq. (19.4.3) as

$$H = \frac{L^2}{2I_A} + \left( \frac{1}{2I_C} - \frac{1}{2I_A} \right) \mathcal{L}_C^2, \quad (19.4.12)$$



which is diagonal in our CSCO of eq. (19.4.7). So, in this case the energy levels are

$$E_{l,k} = \frac{\hbar^2}{2I_A} l(l+1) + \left( \frac{1}{2I_C} - \frac{1}{2I_A} \right) \hbar^2 k^2, \quad (19.4.13)$$

each with degeneracy  $2l+1$ , again due to the  $m$  quantum number. However, there is an additional degeneracy factor of 2 for  $k \neq 0$ , due to the symmetry under  $k \rightarrow -k$ .

Finally we consider the case of a diatomic molecule, with the axis of symmetry containing both nuclei along  $\hat{C}$ . This is a special case of the symmetric top, with  $I_C \ll I_A = I_B$ . This means that the contribution to the energy in eq. (19.4.13) proportional to  $k^2$  is very large, and one often only needs to consider the states with  $k = 0$  and degeneracy  $2l+1$ , because the energy cost to excite states with  $k \neq 0$  is too great. In the idealized limit  $I_C = 0$ , this is a **linear rigid rotor** (epitomized by two point masses attached with a thin rod), and  $k = 0$  is forced in order to have finite energy, so that

$$E_l = \frac{\hbar^2}{2I_A} l(l+1). \quad (19.4.14)$$

Despite the similarity to the formula for the spherical top in eq. (19.4.9), this case is quite different, as the degeneracy of  $E_l$  is only  $2l+1$ . For a diatomic molecule with reduced mass  $M$  and expectation value  $R$  for the nuclear separation, it is traditional to define

$$B = \frac{\hbar^2}{2I_A} = \frac{\hbar^2}{2MR^2}, \quad (19.4.15)$$

so that the rotational contribution to the molecular energy is  $Bl(l+1)$ . In a first approximation, we may treat the electronic state as fixed when computing the rotational excitations. However, if the molecule is in an excited vibrational state like the Morse potential states labeled by  $n$  in section 19.3, then it will have a slightly increased moment of inertia. To parameterize this effect, one may subtract a contribution proportional to  $n+1/2$  from  $1/I_A$ . Thus

$$E_{\text{rot}} = l(l+1) [B - \alpha(n+1/2)], \quad (19.4.16)$$

where  $\alpha$  is a very small positive constant (not to be confused with the fine structure constant), which can be calculated with some effort, or obtained empirically, for each diatomic molecule.

## 19.5 Vibrational and rotational spectra for diatomic molecules

Combining the rotational contributions of the previous section with the larger vibrational contributions found in eq. (19.3.17), we obtain the energy eigenvalues for diatomic molecules,

$$E_{n,l} = \hbar\omega(n+1/2) - \hbar\omega\chi(n+1/2)^2 + l(l+1) [B - \alpha(n+1/2)], \quad (19.5.1)$$

molecule	$M/m_p$	$\hbar\omega$	$\hbar\omega\chi$	$B$	$\alpha$
H <sub>2</sub>	0.50	0.546	0.0151	0.00755	$3.80 \times 10^{-4}$
D <sub>2</sub>	1.00	0.387	0.00767	0.00378	$1.34 \times 10^{-4}$
HCl	0.97	0.371	0.00655	0.00131	$3.81 \times 10^{-5}$
N <sub>2</sub>	7.00	0.293	0.00178	$2.48 \times 10^{-4}$	$2.13 \times 10^{-6}$
CO	6.86	0.269	0.00165	$2.40 \times 10^{-4}$	$2.17 \times 10^{-6}$
O <sub>2</sub>	8.00	0.196	0.00148	$1.79 \times 10^{-4}$	$1.98 \times 10^{-6}$
Cl <sub>2</sub>	17.7	0.0694	$3.34 \times 10^{-4}$	$3.03 \times 10^{-5}$	$1.88 \times 10^{-7}$
NaCl	13.9	0.0452	$2.20 \times 10^{-4}$	$2.71 \times 10^{-5}$	$2.02 \times 10^{-7}$

Table 19.5.4: Empirical values for vibrational ( $\hbar\omega$  and  $\hbar\omega\chi$ ) and rotational ( $B$  and  $\alpha$ ) energy parameters for some common diatomic molecules, in units of eV. Here D is the deuterium isotope of hydrogen. In the specialized literature, these energy parameters are usually given as equivalent photon wavenumbers with units  $\text{cm}^{-1}$ , obtained by dividing the numbers above by  $2\pi\hbar c = 1.2407 \times 10^{-4} \text{ eV}\cdot\text{cm}$ . Also shown in the second column is the reduced nuclear mass  $M = M_1 M_2 / (M_1 + M_2)$  in units of the proton mass.

with degeneracy  $2l + 1$ . This parameterization has proven empirically successful, and fits to data quite generally obey

$$\hbar\omega \gg \hbar\omega\chi \gg B \gg \alpha > 0. \quad (19.5.2)$$

For illustration, the observed numerical parameters for some common diatomic molecules are shown in Table 19.5.4. This confirms the argument in section 19.1 that the vibrational energy scales are larger than the rotational energies, and both tend to be smaller for larger  $M$ , but not absolutely so because of differences in electronic wavefunctions. Comparing ordinary hydrogen and deuterium, which have similar electronic states but different nuclear masses, the ratios are close to the predictions of eq. (19.1.5),  $\hbar\omega_{H_2}/\hbar\omega_{D_2} \approx (M_{D_2}/M_{H_2})^{1/2} = \sqrt{2}$  for vibrations, and  $B_{H_2}/B_{D_2} \approx M_{D_2}/M_{H_2} = 2$  for rotational excitations.

Molecules can absorb and emit photons by jumping from one energy level to another. For **heteronuclear** diatomic molecules (those with different nuclei), there is always an asymmetry in the electronic state wavefunction, which gives rise to a permanent electric dipole moment pointing along the axis common to the nuclei. This electric dipole moment couples to the electromagnetic field. Such molecules therefore are **infrared active**, with efficient transitions between adjacent vibrational levels,  $n \rightarrow n + 1$  for absorption of a photon and  $n \rightarrow n - 1$  for emission. In a first approximation relevant for low resolution experiments, one can neglect the anharmonicity of the vibrational modes and the rotational contributions, implying a single spectral line corresponding to photons with energies near  $\hbar\omega$ , which is usually in the infrared.

At a higher level of resolution, it is necessary to take into account not only the anharmonicity  $\chi$  but also the fact that there is a selection rule  $\Delta l = \pm 1$  for electric dipole transitions, similar




Figure 19.5.1: Infrared vibrational-rotational absorption spectrum for a heteronuclear diatomic molecule undergoing transitions  $n \rightarrow n + 1$ , with either  $l \rightarrow l + 1$  (right) or  $l + 1 \rightarrow l$  (left) as labeled. The central energy is at  $\hbar\omega_n = \hbar\omega(1 - 2(n + 1)\chi)$ , where  $\omega$  is the harmonic angular frequency and  $\chi$  is the anharmonicity. However, the would-be line with  $E_{\text{photon}} = \hbar\omega_n$  is absent, due to the selection rule  $\Delta l = \pm 1$ . The energy differences between lines are  $2B$  and  $4B$  as shown, with  $B$  the rotational energy parameter. The relative occupancies of initial states depend on temperature, and as shown here assume  $k_B T = 10B$ . Due to Boltzmann suppression of higher vibrational modes, the most prominent absorption lines have  $n = 0$ , with  $\omega_0 = \omega(1 - 2\chi)$  called the fundamental angular frequency. At low resolution, the rotational lines shown and the lines for different  $n$  coalesce into a single one with energy  $E \approx \hbar\omega_0$ .

to the one we will derive for single-electron transitions in atoms in section 22.4. The absorbed or emitted photon energies can therefore be obtained by taking differences between the levels in eq. (19.5.1), with the results (neglecting the small correction  $\alpha$  for simplicity)

$$\Delta E_{n \rightarrow n+1}^{l \rightarrow l+1} = \hbar\omega(1 - 2(n + 1)\chi) + 2(l + 1)B, \quad (19.5.3)$$

$$\Delta E_{n \rightarrow n+1}^{l+1 \rightarrow l} = \hbar\omega(1 - 2(n + 1)\chi) - 2(l + 1)B. \quad (19.5.4)$$

This shows that there is actually no photon absorption or emission line at  $\hbar\omega_n$ , where  $\omega_n = \omega(1 - 2(n + 1)\chi)$  is the angular frequency difference corrected for the anharmonicity. Instead, the spectrum for each  $n$  is as shown in Figure 19.5.1, and features a series of lines equally spaced by  $2B$  above  $\hbar\omega_n$  for  $l \rightarrow l + 1$ , and another series of lines equally spaced by  $2B$  below  $\hbar\omega_n$  for  $l + 1 \rightarrow l$ , with a gap of  $4B$  between them due to the selection rule that  $l$  must change by 1 unit. Measuring the energy gaps reveals the moment of inertia of the molecule. The relative occupancies of the initial states with eigenvalues  $l$  are governed by the statistical factor

$$(2l + 1)e^{-l(l+1)B/k_B T}. \quad (19.5.5)$$

For small  $l$ , this factor rises with  $l$  due to the rotational degeneracy factor  $2l + 1$ , and for large  $l$  it is suppressed by the exponential Boltzmann factor. The relative initial-state occupancies shown in Figure 19.5.1 correspond to  $k_B T = 10B$ . The number of prominent lines on either side of the central value  $\hbar\omega_n$  increases with the temperature.

For **homonuclear** molecules (those with identical nuclei), there are two special considerations. First, according to the spin-statistics principle, the interchange of the two nuclei must multiply the state ket by a minus sign if they are fermions, or a plus sign if they are bosons. For a nucleus with spin  $s$ , this sign equals  $(-1)^{2s}$ . To match this, recall that the interchange of nuclei in the relative coordinates just coincides with a parity transformation (replacing each nuclear coordinate vector by minus itself), and therefore is associated with a factor of  $(-1)^l$  for orbital angular momentum  $l$ , as in eq. (8.7.5). Also, if the two spins  $s$  combine to form a total spin  $S$ , then the exchange of the two nuclear spin states results in a factor of  $(-1)^{2s-S}$ , as can be seen from eq. (12.4.25), or from the symmetric and antisymmetric labels in eq. (12.4.44). Therefore, combining the wavefunction and spin-state exchange factors, we must have

$$(-1)^{2s} = (-1)^l (-1)^{2s-S}. \quad (19.5.6)$$

It follows that  $l$  must be even if the total nuclear spin state has even  $S$ , and  $l$  must be odd if  $S$  is odd. The molecular states with symmetric nuclear spins and  $(-1)^{2s-S} = 1$  are referred to as “ortho-”, while those with antisymmetric nuclear spin states are called “para-”.<sup>†</sup>

Counting states using eq. (12.4.44) for spin  $s$ , the total number of ortho- states with  $S = 2s, 2s-2, \dots$  is  $(s+1)(2s+1)$ , while the total number of para- states with  $S = 2s-1, 2s-3, \dots$  is  $s(2s+1)$ . To get this counting correct, one includes  $2S+1$  states for each  $S$ ; the grand total number of nuclear spin states is  $(2s+1)^2$ . For molecules in thermal equilibrium at lab temperatures,  $k_B T = 0.0255$  eV is larger than the rotational energy differences, so each nuclear spin state has close to the same occupancy. Therefore, for diatomic molecules with identical nuclei, the ratio of the total number of para- states to the total number of ortho- states is

$$\frac{N_{\text{para}}}{N_{\text{ortho}}} = \frac{s}{s+1}. \quad (19.5.7)$$

This can be used to experimentally determine the spins of nuclei that form diatomic molecules. For example, for  $\text{H}_2$  gas at lab temperature with  $s = 1/2$ , the abundance of ortho- $\text{H}_2$  with  $S = 1$  and therefore odd  $l$  is about 3 times that of para- $\text{H}_2$  with  $S = 0$  and even  $l$ . In contrast, for deuterium with  $s = 1$ , the abundance at lab temperature of ortho- $\text{D}_2$  with  $S = 2$  or  $S = 0$  and even  $l$  is about 2 times that of para- $\text{D}_2$  with  $S = 1$  and odd  $l$ . For lower temperatures, the right side of eq. (19.5.7) must be corrected by a degeneracy-and-Boltzmann statistics factor

$$\frac{\sum_{l=1,3,5,\dots} (2l+1) e^{-l(l+1)B/k_B T}}{\sum_{l=0,2,4,\dots} (2l+1) e^{-l(l+1)B/k_B T}} \quad (19.5.8)$$

---

<sup>†</sup>The terminology is similar to that for atomic helium as discussed in section 18.3, in the sense that symmetric spin states are ortho- and antisymmetric spin states are para-. However, for atomic helium this refers to the electronic spins, while here the spins referred to are the nuclear ones.

when  $2s$  is even, or its reciprocal when  $2s$  is odd. This factor is close to 1 at high temperatures, and decreases from more than 0.99 when  $k_B T/B > 3$  (which includes lab temperature for all molecules) to 0 at lower temperatures. For example, at very low temperatures, so that most of the molecules are in the  $l = 0$  level,  $\text{H}_2$  is almost all para-, while  $\text{D}_2$  is mostly ortho-. For nuclei with  $s = 0$ , like the main oxygen isotope, the para- states with odd  $l$  do not exist at all.

The second special consideration for homonuclear diatomic molecules is that they cannot have an electric dipole moment, since by symmetry there is no special direction along which it could point. Such molecules therefore have a much smaller coupling to electromagnetic fields, and are called “inactive” in the infrared and microwave ranges.

However, the vibrational and rotational energy levels of homonuclear molecules can still be observed by inelastic scattering of photons, called the **Raman effect**. In Raman scattering, a photon with known energy  $\hbar\Omega$  impacts the molecule, sometimes changing its vibrational or rotational state. The molecule’s change in energy is then seen as an opposite change in energy of the scattered photon. The Raman effect also occurs in heteronuclear molecules.

When the molecular state does not change, the final state photon energy is also  $\hbar\Omega$ ; this is called Rayleigh scattering. In a first approximation in which anharmonicity and rotational excitations are neglected, if the molecule absorbs some energy by transitioning to the next higher vibrational state  $n \rightarrow n + 1$ , the scattered photon will have energy  $\hbar(\Omega - \omega)$ , called Raman–Stokes scattering, and if the molecule gives up energy by transitioning to the next lower vibrational state by  $n + 1 \rightarrow n$ , then the scattered photon will have energy  $\hbar(\Omega + \omega)$ , called Raman–anti-Stokes scattering. The main Rayleigh scattering line is therefore accompanied by two equally spaced lines, as shown in Figure 19.5.2. Except at very high temperatures, the  $n = 0$  state has the highest initial occupancy, and the anti-Stokes line is much less intense than the Stokes line for vibrational Raman scattering, because the initial states for anti-Stokes have higher energies and thus a relative Boltzmann suppression  $e^{-\hbar\omega/k_B T}$ .

The Raman effect can also result from inelastic scattering with transitions between states with different rotational energies. Here, it is important that there is a selection rule (which we will not attempt to derive) for Raman scattering,  $\Delta l = 0$  or  $\pm 2$ . Using the approximation  $E_l = Bl(l + 1)$ , one therefore finds for the allowed scattered photon energies

$$E_{\text{scattered}}^{l \rightarrow l+2} = \hbar\Omega - (4l + 6)B, \quad (\text{Stokes}), \quad (19.5.9)$$

$$E_{\text{scattered}}^{l+2 \rightarrow l} = \hbar\Omega + (4l + 6)B, \quad (\text{anti-Stokes}), \quad (19.5.10)$$

along with the Rayleigh line  $E_{\text{scattered}}^{l \rightarrow l} = \hbar\Omega$ . The positions of the spectral lines for the rotational Raman effect are shown in Figure 19.5.3. There is an energy gap  $6B$  on either side of the Rayleigh line to the nearest rotational Raman lines, and gaps  $4B$  between different rotational Raman–Stokes and Raman–anti-Stokes lines. The relative intensities depend on the occupancies of the




Figure 19.5.2: The vibrational Raman effect spectrum of scattered photon energies for diatomic molecules. The incident photon has energy  $\hbar\Omega$ . If the molecule remains in the same state after the scattering, the scattered photon energy is also  $\hbar\Omega$ , giving the Rayleigh scattering line at center. The Stokes line results from the molecule transitioning to the next-higher energy vibrational state by  $n \rightarrow n+1$ , and the anti-Stokes line is due to the molecule moving to the next-lower vibrational state by  $n+1 \rightarrow n$ . The anti-Stokes line is less intense than the Stokes line, as its initial occupancies have a relative Boltzmann suppression  $e^{-\hbar\omega/k_B T}$ .




Figure 19.5.3: The rotational Raman effect spectrum of scattered photon energies for diatomic molecules. The incident photon has energy  $\hbar\Omega$ . When the molecule remains in the same state, the scattered photon energy is also  $\hbar\Omega$ , giving the Rayleigh scattering line in the center. The left series of Stokes lines result when the molecule transitions to a higher energy rotational state via  $l \rightarrow l+2$  as labeled for various  $l$ , and the right series of anti-Stokes lines are due to the molecule transitioning to lower energy states by  $l+2 \rightarrow l$ . The energy separations between lines are  $4B$  or  $6B$  as shown, where  $B$  is the rotational energy parameter, inversely proportional to the moment of inertia. The relative intensities of the lines are not shown, and depend on the temperature and whether the molecule is homonuclear or heteronuclear.

initial states  $l$ , which for homonuclear molecules are proportional to

$$(s+1)(2l+1)e^{-l(l+1)B/k_B T} \quad (\text{homonuclear ortho- states}), \quad (19.5.11)$$

$$s(2l+1)e^{-l(l+1)B/k_B T} \quad (\text{homonuclear para- states}), \quad (19.5.12)$$

alternating between odd and even  $l$ . For heteronuclear molecules, the initial-state occupancies are simply proportional to the factor already given in eq. (19.5.5).

## 19.6 The most important molecule: H<sub>2</sub>O

Polyatomic molecules are obviously more complicated than diatomic ones, and mostly beyond our scope. However, given the importance of water to life and happiness on Earth, it seems worthwhile to give a brief phenomenological account of its vibrational and rotational quantum excitations.

The H<sub>2</sub>O molecule has a bent (non-linear) shape, with the geometry of the three nuclei shown in Figure 19.6.1. The bond angle measured from the oxygen nucleus is famously about 104.5°. The electrons on the hydrogen atoms are attracted to the oxygen nucleus, leaving them with a net positive charge and ensuring that the molecule has a permanent electric dipole moment in its ground state, pointing in the direction of the principal axis labeled  $\hat{B}$ . The reasons for this odd geometry and the nature of the bonding are discussed in more specialized books.




Figure 19.6.1: The equilibrium geometry of the nuclei in the H<sub>2</sub>O molecule is shown on the left. The rectangular coordinates of the nuclear positions, with respect to the center of mass, are O at (0, −0.6563) and H at (7.572, 5.209) and (−7.572, 5.209), in units of 10<sup>−11</sup> meters. The angle between the OH bonds is 104.5°. The principal axes in which the moment-of-inertia tensor is diagonal are shown from the center of mass position as  $\hat{A}$  and  $\hat{B}$ , with  $\hat{C}$  out of the plane of this page. The three diagrams on the right illustrate the three vibrational modes: bending (b), symmetric stretching (s), and asymmetric stretching (a).

There are three<sup>†</sup> vibrational modes, called bending (b), symmetric stretching (s), and asymmetric stretching (a). The classical depictions of these modes are also shown in Figure 19.6.1. The bending mode can be thought of as an oscillation of the bond angle about its equilibrium value. In the symmetric stretching mode, the two hydrogen nuclei (protons) move farther and closer to the center of mass in unison, while in the asymmetric stretching mode one moves closer as the other moves farther. Considering each vibrational mode separately, for simplicity, one can parameterize the energies in terms of a Morse-potential-inspired spectrum,

$$E_n^{(m)} = \hbar\omega_m(n + 1/2) - \hbar\omega_m\chi_m(n + 1/2)^2, \quad (m = b, s, a). \quad (19.6.1)$$

For each mode,  $\omega_m$  is often called the harmonic angular frequency, and the slightly smaller

<sup>†</sup>In general, a molecule has  $3N$  degrees of freedom associated with the position coordinates of the  $N$  nuclei. Of these, 3 degrees of freedom correspond to translations of the molecule as a whole, and 3 more are associated with the rotations of the molecule if it is non-linear, or 2 more if it is linear with  $I_C = 0$ . Therefore, in the approximation of harmonic oscillations about equilibrium, there are  $3N - 6$  vibrational modes for a non-linear molecule like H<sub>2</sub>O, and  $3N - 5$  vibrational modes for a linear molecule like CO<sub>2</sub> or any diatomic molecule.



difference in angular frequencies between the  $n = 0$  and  $n = 1$  energies,  $\omega_m(1 - 2\chi_m)$ , is referred to as the fundamental angular frequency. The experimental values in this parameterization are

$$\hbar\omega_b = 0.2046 \text{ eV}, \quad \hbar\omega_b\chi_b = 0.00335 \text{ eV}, \quad (19.6.2)$$

$$\hbar\omega_s = 0.4754 \text{ eV}, \quad \hbar\omega_s\chi_s = 0.0109 \text{ eV}, \quad (19.6.3)$$

$$\hbar\omega_a = 0.4892 \text{ eV}, \quad \hbar\omega_a\chi_a = 0.0116 \text{ eV}. \quad (19.6.4)$$

These are all in the IR spectrum, as argued on general grounds for molecular vibrations in section 19.1. The fundamental frequencies are  $\nu = \omega(1 - 2\chi)/2\pi = 4.785 \times 10^{13}$ ,  $1.097 \times 10^{14}$ , and  $1.127 \times 10^{14}$  Hz, with vacuum wavelengths 6.27, 2.73, and 2.66 microns. These modes are easily excited by electromagnetic radiation due to the polar nature of the molecule, leading to strong IR absorption features. The anharmonicities help to effectively broaden the absorption spectrum to longer wavelengths, with an even stronger broadening effect in both directions from interactions between neighboring molecules in liquid water. Fortunately for us, water remains mostly transparent to the higher energy photons in the visible range. This is because the lowest electronic state excitations have much larger energies corresponding to UV photons, leaving a window in the absorption spectrum for liquid water for frequencies from roughly  $\nu = 3.75 \times 10^{14}$  to  $1.5 \times 10^{15}$  Hz, corresponding to the visible and the very near IR and UV.

At lower energies, rotational modes are important. Since  $\text{H}_2\text{O}$  is not a linear molecule, it behaves as a quantum asymmetric top for rotational excitations. Given the geometry in Figure 19.6.1 and its caption, it is a short exercise in arithmetic to compute the moments of inertia for the principal axes  $\hat{A}$  and  $\hat{B}$  in the plane common to the nuclei, and  $\hat{C}$  perpendicular to that plane, using oxygen's nuclear mass  $M_O = 15.875m_p$ . This successfully predicts results within a few per cent of the empirical rotational energy parameters for  $\text{H}_2\text{O}$ ,

$$e_A = \frac{\hbar^2}{2I_A} = 0.003459 \text{ eV}, \quad e_B = \frac{\hbar^2}{2I_B} = 0.001801 \text{ eV}, \quad e_C = \frac{\hbar^2}{2I_C} = 0.001152 \text{ eV}, \quad (19.6.5)$$

in terms of which the asymmetric top Hamiltonian is

$$H = e_A(\mathcal{L}_A/\hbar)^2 + e_B(\mathcal{L}_B/\hbar)^2 + e_C(\mathcal{L}_C/\hbar)^2. \quad (19.6.6)$$

As discussed in section 19.4, there is no simple formula for the energies of an asymmetric top that works for all  $l$ , but for the  $l = 1$  states the three energy eigenvalues are, from eq. (19.4.11),

$$E_1^{(1)} = e_A + e_B = 0.00526 \text{ eV}, \quad E_2^{(1)} = e_A + e_C = 0.00461 \text{ eV}, \quad (19.6.7)$$

$$E_3^{(1)} = e_B + e_C = 0.00295 \text{ eV}, \quad (19.6.8)$$

each with degeneracy 3. For the  $l = 2$  states, the five energy eigenvalues are, numerically,

$$E_1^{(2)} = 0.00987 \text{ eV}, \quad E_2^{(2)} = 0.01181 \text{ eV}, \quad E_3^{(2)} = 0.01679 \text{ eV}, \quad (19.6.9)$$

$$E_4^{(2)} = 0.01694 \text{ eV}, \quad E_5^{(2)} = 0.00870 \text{ eV}, \quad (19.6.10)$$



each with degeneracy 5. It is left to Exercise 19.3 to derive the symbolic versions of these  $l = 2$  energy eigenvalues in terms of the moments of inertia for a general asymmetric top.

Transitions between the above rotational states, and many others with higher  $l$  relevant for lab temperature  $k_B T = 0.0255$  eV, correspond to photons with wavelengths in the near microwave spectrum. However, they are all much shorter than the 12.2 cm wavelength used by modern consumer microwave ovens to heat food. Microwave ovens efficiently heat liquid water, but do not operate by resonantly exciting rotational states (or the vibrational states described above, which have even higher energies). Instead, they produce time dependence in the superpositions of the rotational states, corresponding semi-classically to oscillatory changes in orientation rather than complete rotations. Microwave ovens are much less efficient at heating ice, which has nearly locked-in molecular orientations.

## 19.7 Exercises

**Exercise 19.1.** The radial wavefunctions of the Morse potential discussed in section 19.3 can be written exactly in terms of the associated Laguerre polynomials  $L_N^\alpha$  defined in eq. (10.5.36). Prove this, by using eq. (19.3.15) and the associated Laguerre differential equation (10.5.37) to show that  $S_n(z) = z^{\kappa-n-1/2} L_n^{2\kappa-2n-1}(2z)$  solves the differential equation (19.3.11) for the rescaled energy eigenvalue  $\mathcal{E}_n = 1 - (1 - (n + 1/2)/\kappa)^2$ .

**Exercise 19.2.** Prove that the principal-axis angular momentum components of a rigid body, defined in eq. (19.4.2), commute with the ordinary angular momentum components, as in eq. (19.4.4). Also show that they obey the commutation relations in eq. (19.4.5), with the notable minus signs.

**Exercise 19.3.** In this problem, we will find the five  $l = 2$  rotational energy levels, each with degeneracy 5, for a general asymmetric top.

(a) Start by writing the  $5 \times 5$  matrix representations for generic angular momentum components  $J_x$ ,  $J_y$ , and  $J_z$ , making use of eqs. (8.4.4) and (8.4.5). Then substitute  $J_{x,y,z} \rightarrow \mathcal{L}_{A,B,C}$ , and use them in eq. (19.4.3) to find the  $5 \times 5$  Hamiltonian matrix for  $l = 2$ . For convenience, write it in terms of  $e_A = \hbar^2/2I_A$  and  $e_B = \hbar^2/2I_B$  and  $e_C = \hbar^2/2I_C$ .

(b) Find the energy eigenvalues of the  $l = 2$  Hamiltonian matrix obtained in part (a). This should involve three linear equations with solutions

$$E_1 = e_A + e_B + 4e_C, \quad E_2 = e_A + e_C + 4e_B, \quad E_3 = e_B + e_C + 4e_A, \quad (19.7.1)$$

and one quadratic equation with solutions

$$E_{4,5} = 2 \left( e_A + e_B + e_C \pm \sqrt{e_A^2 + e_B^2 + e_C^2 - e_A e_B - e_A e_C - e_B e_C} \right). \quad (19.7.2)$$

## 20 Heisenberg and interaction representations

### 20.1 The Heisenberg picture and equations of motion

Everything we have done so far uses a formulation that is sometimes called the **Schrödinger picture** of quantum mechanics. This just means that the state of the system evolves according to Postulate 6 of section 3.1,

$$i\hbar \frac{d}{dt} |\psi(t)\rangle = H |\psi(t)\rangle. \quad (20.1.1)$$

However, there are other, equivalent, formulations of quantum mechanics, called “pictures”, which are related to the Schrödinger picture by doing a time-dependent unitary transformation on the state of the system and a corresponding transformation to all operators. Done consistently, this cannot affect the predictions for physically measurable quantities, but the intermediate steps for solving problems will differ. The main reason for considering other pictures is to make some calculations easier to do and/or understand.

Recall from the discussion in section 3.4 that the solution to eq. (20.1.1), starting from an initial state  $|\psi(t_0)\rangle$ , is

$$|\psi(t)\rangle = U(t, t_0) |\psi(t_0)\rangle, \quad (20.1.2)$$

where  $U(t, t_0)$  is a unitary operator satisfying the differential equation

$$i\hbar \frac{d}{dt} U(t, t_0) = H U(t, t_0), \quad (20.1.3)$$

with  $U(t, t_0)^{-1} = U(t, t_0)^\dagger = U(t_0, t)$ . If  $H$  has no explicit time dependence, then the solution to eq. (20.1.3) is relatively easy, just  $U(t, t_0) = e^{-i(t-t_0)H/\hbar}$ . In that case, there is no particular advantage to the Heisenberg or interaction pictures described in the following.

In the **Heisenberg picture**, all time dependence is transferred from the state of the system to the operators. To accomplish this, define

$$|\psi_H\rangle = U(t, t_0)^\dagger |\psi(t)\rangle, \quad (20.1.4)$$

$$A_H(t) = U(t, t_0)^\dagger A(t) U(t, t_0), \quad (20.1.5)$$

where  $|\psi_H\rangle$  is the state ket in the Heisenberg picture, and for each Schrödinger picture operator  $A(t)$ , the corresponding Heisenberg picture operator is denoted  $A_H(t)$ . Combining eqs. (20.1.2) and (20.1.4) gives

$$|\psi_H\rangle = |\psi(t_0)\rangle. \quad (20.1.6)$$

Thus the Heisenberg picture state ket  $|\psi_H\rangle$  is independent of the time  $t$ ; it only depends on the initial condition at  $t_0$ . Note that there are really infinitely many Heisenberg pictures, one for

each choice of the reference initial time  $t_0$ . We include a subscript  $H$  to denote the state of the system or an operator as being in the Heisenberg picture. If an object is not written with the  $H$  subscript, it can be assumed to be in the Schrödinger picture. For convenience and simplicity, we will often write  $U = U(t, t_0)$  in the rest of this section.

A possible point of confusion is that it is only the state of the system that we are defining differently in the Heisenberg and Schrödinger pictures via the unitary transformation in eq. (20.1.4). Both the Schrödinger picture state ket  $|\psi(t)\rangle$  and the Heisenberg picture state ket  $|\psi_H\rangle$  can be expressed as (different!) linear combinations of a set of *fixed* orthobasis kets, which are defined independently of the choice of picture. A convenient way to choose the fixed orthobasis kets is as time-independent eigenstates of some appropriate time-independent CSCO in the Schrödinger picture. Once so chosen, the orthobasis kets do not depend on  $t$ , by definition.

Inner products and matrix elements are the same in both pictures, since they are related by a unitary transformation. To see this, consider a matrix element of an arbitrary observable  $A$  between two possible states of the system  $|\psi\rangle$  and  $|\chi\rangle$  that both obey Schrödinger's equation in the Schrödinger picture. We have

$$\langle\chi(t)|A(t)|\psi(t)\rangle = \langle\chi(t)|UU^\dagger A(t)UU^\dagger|\psi(t)\rangle = \langle\chi_H|A_H(t)|\psi_H\rangle. \quad (20.1.7)$$

This shows the physical equivalence of the two pictures.

Even though many important observable operators (for example,  $A$  = position, momentum, or spin) are time-independent in the Schrödinger picture, the corresponding Heisenberg picture operators  $A_H(t)$  will generally depend on  $t$ , and often in a complicated way, depending on the Hamiltonian. This is the price to be paid in the Heisenberg picture for the convenience that the state of the system has no  $t$  dependence.

However, there is an important special case in which  $A_H(t)$  does not depend on  $t$ . Suppose that in the Schrödinger picture,  $H$  and  $A$  do not depend on  $t$ , and  $[A, H] = 0$ . Then  $A$  commutes with  $U(t, t_0)$  as well, since the latter is built out of  $H$ . In that case,

$$A_H = U^\dagger A U = U^\dagger U A = A, \quad (20.1.8)$$

and we say that  $A$  is a **constant of motion**, following classical physics terminology. Note that this corresponds to a conserved quantity, as defined in section 5.1 in the Schrödinger picture. In particular, if  $H$  does not depend explicitly on  $t$ , then  $H_H = H$ , and the Schrödinger and Heisenberg picture Hamiltonians are equal.

More generally, we can ask how  $A_H(t)$  evolves in time. To find out, we calculate, using the product rule for derivatives,

$$\frac{dA_H}{dt} = \left[ \frac{d}{dt} U(t, t_0)^\dagger \right] A U + U^\dagger A \left[ \frac{d}{dt} U(t, t_0) \right] + U^\dagger \frac{\partial A}{\partial t} U. \quad (20.1.9)$$

Evaluating the derivatives in square brackets using the differential equation (20.1.3) gives

$$\frac{dA_H}{dt} = \frac{i}{\hbar}U^\dagger H A U - \frac{i}{\hbar}U^\dagger A H U + U^\dagger \frac{\partial A}{\partial t} U, \quad (20.1.10)$$

which can be rewritten as

$$\frac{dA_H}{dt} = \frac{i}{\hbar}[H_H, A_H] + \left(\frac{\partial A}{\partial t}\right)_H. \quad (20.1.11)$$

The meaning of the last term is that we take the derivative of  $A$  with respect to its explicit dependence on  $t$  in the Schrödinger picture, and then convert the result to the Heisenberg picture. Equation (20.1.11) is called the **Heisenberg equation of motion** for the operator  $A_H$ . Since the state of the system does not change in the Heisenberg picture in the absence of a measurement, this equation captures all physical effects of unitary time evolution. It is a direct analog in quantum mechanics of the classical equations of motion.

The form of **equal-time commutation relations** does not change if one switches pictures. If one has a commutation relation

$$[A, B] = C, \quad (20.1.12)$$

in the Schrödinger picture, then

$$[A_H(t), B_H(t)] = [U^\dagger A U, U^\dagger B U] = U^\dagger [A, B] U = U^\dagger C U = C_H(t). \quad (20.1.13)$$

So, for example,

$$[X_H(t), P_H(t)] = [X, P] = i\hbar, \quad (20.1.14)$$

and if  $\vec{J}$  is an angular momentum operator, then

$$[J_{Ha}(t), J_{Hb}(t)] = i\hbar\epsilon_{abc}J_{Hc}(t), \quad (a, b, c = x, y, z). \quad (20.1.15)$$

It is crucial that we specified equal-time commutation relations, because in general the commutator  $[A_H(t), B_H(t')]$  does not have a simple relationship to  $[A, B]$  unless  $t' = t$ . For example,

$$[X_H(t), P_H(t')] \neq i\hbar \quad (20.1.16)$$

in general, unless  $t' = t$ .

To illustrate how the Heisenberg picture works in the simple case that the Hamiltonian does not explicitly depend on time, suppose that in the Schrödinger picture we have the usual Hamiltonian for a spinless particle of mass  $m$  moving in a potential,

$$H = \frac{P^2}{2m} + V(X), \quad (20.1.17)$$

where  $V(x) = \sum_n v_n x^n$ , with  $v_n$  a set of fixed real numbers. Then

$$P_H^2 = U^\dagger P U U^\dagger P U = U^\dagger P^2 U, \quad (20.1.18)$$

and similarly

$$(X_H)^n = U^\dagger X^n U. \quad (20.1.19)$$

Therefore, the Heisenberg picture Hamiltonian is

$$H_H = U^\dagger H U = \frac{P_H^2}{2m} + V(X_H). \quad (20.1.20)$$

The Schrödinger picture position and momentum operators have no explicit time dependence,

$$\frac{\partial X}{\partial t} = 0, \quad \frac{\partial P}{\partial t} = 0, \quad (20.1.21)$$

and so we obtain from eq. (20.1.11), by using eq. (20.1.14),

$$\frac{dX_H}{dt} = \frac{i}{\hbar} [H_H, X_H] = \frac{P_H}{m}, \quad (20.1.22)$$

$$\frac{dP_H}{dt} = \frac{i}{\hbar} [H_H, P_H] = -V'(X_H). \quad (20.1.23)$$

The Heisenberg equations of motion (20.1.22) and (20.1.23) have the same form as the classical equations of motion, but with the classical position and momentum promoted to the corresponding Heisenberg-picture operators. Using them, one can check that

$$\frac{d}{dt} H_H = 0, \quad (20.1.24)$$

which verifies that  $H_H = H$  is indeed a constant of motion.

## 20.2 The interaction picture and transition amplitudes

Suppose that we have a Hamiltonian (in the Schrödinger picture) of the form

$$H(t) = H_0 + W(t), \quad (20.2.1)$$

where  $W(t)$  may depend explicitly on time, but  $H_0$  does not. Then one can define another picture called the **interaction picture**, also known as the **Dirac picture**.

The idea of the interaction picture is to transfer from the state ket to the operators only the part of the time dependence that  $H_0$  is responsible for. This is done by defining, for the ket describing the Schrödinger picture state of the system  $|\psi(t)\rangle$  and for all observables  $A$ ,

$$|\psi_I(t)\rangle = e^{itH_0/\hbar} |\psi(t)\rangle, \quad (20.2.2)$$

$$A_I(t) = e^{itH_0/\hbar} A e^{-itH_0/\hbar}. \quad (20.2.3)$$

The subscript  $I$  is used to denote the interaction picture state and operators; when it is not present, the state or operator should be assumed to be in the Schrödinger picture. The interaction picture states and operators coincide with their Schrödinger picture counterparts at time  $t = 0$ . (One could instead choose some other time to do the matching, but there is no particular advantage to doing so.) Note that  $(H_0)_I = e^{itH_0/\hbar} H_0 e^{-itH_0/\hbar} = H_0$ , so a distinguishing subscript for  $H_0$  is not needed. However, in general,  $W_I(t) \neq W(t)$ , so we must be careful to distinguish them from now on. Note that eqs. (20.2.2) and (20.2.3) can be considered a change of basis, since  $e^{-itH_0/\hbar}$  is a unitary operator.

Clearly, if  $W(t) = 0$ , then the interaction and Heisenberg pictures are the same, while if  $H_0 = 0$ , then the interaction and Schrödinger pictures are the same. More generally, the interaction picture is intermediate between the Schrödinger and Heisenberg pictures. At first, it might seem that the interaction picture combines the worst features of both, since both the state and observable operators evolve non-trivially in time. However, it turns out to be useful for an elegant formulation of time-dependent perturbation theory, when the effects of  $H_0$  are large but easily calculable, and the effects of  $W(t)$  can be treated as the perturbation.

By an argument exactly analogous to the derivation of eq. (20.1.11), one can find the equation giving the time evolution of operators in the interaction picture,

$$\frac{dA_I}{dt} = \frac{i}{\hbar} [H_0, A_I] + \left( \frac{\partial A}{\partial t} \right)_I. \quad (20.2.4)$$

Here, the meaning of the last term is that one should take the derivative of  $A$  with respect to its explicit time dependence in the Schrödinger picture, and then convert the result to the interaction picture using eq. (20.2.3).

Let us now define the unitary operator  $\mathcal{U}_I(t, t_0)$  which gives the time evolution of  $|\psi_I(t)\rangle$  in terms of its initial condition at an arbitrary time  $t = t_0$ , according to

$$|\psi_I(t)\rangle = \mathcal{U}_I(t, t_0) |\psi_I(t_0)\rangle. \quad (20.2.5)$$

To avoid confusion, it is important to note that  $\mathcal{U}_I(t, t_0)$  is not the same as  $U_I(t, t_0)$ , which is defined to be the interaction-picture version of the Schrödinger picture unitary time-evolution operator  $U(t, t_0)$ . In fact, the relation between these three unitary operators is

$$\mathcal{U}_I(t, t_0) = e^{itH_0/\hbar} U(t, t_0) e^{-it_0H_0/\hbar} = U_I(t, t_0) e^{i(t-t_0)H_0/\hbar}. \quad (20.2.6)$$

The first equality follows from eqs. (20.1.2), (20.2.2), and (20.2.5), and the last equality then follows directly from the definition eq. (20.2.3). In the special case that  $H$  has no explicit time dependence, we can now apply the known solution for  $U(t, t_0)$  in eq. (3.4.3) to learn that

$$\mathcal{U}_I(t, t_0) = e^{itH_0/\hbar} e^{i(t_0-t)H/\hbar} e^{-it_0H_0/\hbar}, \quad (\text{for time-independent } H). \quad (20.2.7)$$

Our goal now is to solve for  $\mathcal{U}_I(t, t_0)$  more generally. To accomplish this, we calculate

$$i\hbar \frac{d}{dt} |\psi_I(t)\rangle = i\hbar \frac{d}{dt} (e^{itH_0/\hbar} |\psi(t)\rangle) = -e^{itH_0/\hbar} H_0 |\psi(t)\rangle + i\hbar e^{itH_0/\hbar} \frac{d}{dt} |\psi(t)\rangle. \quad (20.2.8)$$

Using the Schrödinger equation  $i\hbar \frac{d}{dt} |\psi\rangle = (H_0 + W) |\psi\rangle$ , this becomes

$$i\hbar \frac{d}{dt} |\psi_I(t)\rangle = e^{itH_0/\hbar} W |\psi\rangle, \quad (20.2.9)$$

which we can rewrite in a form involving only interaction picture objects, as

$$i\hbar \frac{d}{dt} |\psi_I(t)\rangle = W_I(t) |\psi_I(t)\rangle. \quad (20.2.10)$$

Plugging the definition eq. (20.2.5) into eq. (20.2.10), we get the operator equation

$$i\hbar \frac{d}{dt} \mathcal{U}_I(t, t_0) = W_I(t) \mathcal{U}_I(t, t_0). \quad (20.2.11)$$

Integrating both sides gives

$$i\hbar [\mathcal{U}_I(t, t_0) - \mathcal{U}_I(t_0, t_0)] = \int_{t_0}^t dt' W_I(t') \mathcal{U}_I(t', t_0). \quad (20.2.12)$$

Rewriting this using  $\mathcal{U}_I(t_0, t_0) = I$ , we find the useful integral equation

$$\mathcal{U}_I(t, t_0) = I - \frac{i}{\hbar} \int_{t_0}^t dt' W_I(t') \mathcal{U}_I(t', t_0). \quad (20.2.13)$$

Taking the Hermitian adjoint of this, and using the unitarity relation  $\mathcal{U}_I(t, t_0)^\dagger = \mathcal{U}_I(t, t_0)^{-1} = \mathcal{U}_I(t_0, t)$  and the Hermiticity of  $W_I(t')$ , one finds  $\mathcal{U}_I(t_0, t) = I + \frac{i}{\hbar} \int_{t_0}^t dt' \mathcal{U}_I(t_0, t') W_I(t')$ . Then, interchanging the roles of  $t$  and  $t_0$  gives an alternative version of the integral equation,

$$\mathcal{U}_I(t, t_0) = I - \frac{i}{\hbar} \int_{t_0}^t dt' \mathcal{U}_I(t, t') W_I(t'). \quad (20.2.14)$$

Although eqs. (20.2.13) and (20.2.14) contain the same information, the latter will be the more useful form in the context of scattering, as we will see in section 23.2.

Solving the integral equation (20.2.13) by iteration gives

$$\mathcal{U}_I(t, t_0) = I - \frac{i}{\hbar} \int_{t_0}^t dt' W_I(t') + \left(-\frac{i}{\hbar}\right)^2 \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' W_I(t') W_I(t'') + \dots \quad (20.2.15)$$

or, rewriting more systematically,

$$\mathcal{U}_I(t, t_0) = \sum_{N=0}^{\infty} \left(-\frac{i}{\hbar}\right)^N \int_{t_0}^t dt_N \int_{t_0}^{t_N} dt_{N-1} \dots \int_{t_0}^{t_2} dt_1 W_I(t_N) W_I(t_{N-1}) \dots W_I(t_1). \quad (20.2.16)$$

Equation (20.2.16) is the **Dyson series** for the time evolution of the state in the interaction picture; compare to the similar structure of eq. (3.4.32). This solution can be checked by plugging it directly into eq. (20.2.11). In general,  $W_I(t)$  may not commute with  $W_I(t')$  when  $t \neq t'$ . Therefore, we are careful to put the operators in increasing time order, from right to left, as specified by the nested integrations with  $t_0 < t_1 < t_2 < \cdots < t_{N-1} < t_N < t$ . In practice, eq. (20.2.16) is usually approximated by truncating it to the first few terms, treating  $W_I$  as small. A more compact (but less directly useful) way of writing eq. (20.2.16) is

$$\mathcal{U}_I(t, t_0) = T \exp \left[ -\frac{i}{\hbar} \int_{t_0}^t dt' W_I(t') \right], \quad (20.2.17)$$

where  $T$  is the time-ordering symbol introduced in the discussion surrounding eq. (3.4.31), to which eq. (20.2.17) can be compared. The symbol  $T$  is an instruction to re-order the operators following it, so that after expanding the exponential,  $W_I(t_j)$  appears to the right of  $W_I(t_k)$  whenever  $t_j < t_k$ . The time-ordering rule is that earlier operators act first on kets.

Given our solution for  $\mathcal{U}_I(t, t_0)$  in eq. (20.2.16), one can express the time dependence of the state back in the Schrödinger picture. Using eqs. (20.1.2) and (20.2.6),

$$|\psi(t)\rangle = U(t, t_0)|\psi(t_0)\rangle = e^{-itH_0/\hbar} \mathcal{U}_I(t, t_0) e^{it_0H_0/\hbar} |\psi(t_0)\rangle. \quad (20.2.18)$$

It is convenient to express this in terms of an orthobasis of eigenstates  $|n\rangle$  of  $H_0$  with eigenvalues  $\mathcal{E}_n$ . Using completeness twice, we have

$$|\psi(t)\rangle = \sum_m \sum_n e^{-i\mathcal{E}_m t/\hbar} e^{i\mathcal{E}_n t_0/\hbar} |m\rangle \langle m | \mathcal{U}_I(t, t_0) | n \rangle \langle n | \psi(t_0) \rangle. \quad (20.2.19)$$

Now, define the **transition amplitudes** for the eigenstates of  $H_0$  as

$$a_{n \rightarrow m}(t, t_0) \equiv \langle m | \mathcal{U}_I(t, t_0) | n \rangle. \quad (20.2.20)$$

Then, given the initial state coefficients

$$c_n(t_0) = \langle n | \psi(t_0) \rangle, \quad (20.2.21)$$

it follows that the Schrödinger picture state of the system at time  $t$  is

$$|\psi(t)\rangle = \sum_m c_m(t) |m\rangle, \quad (20.2.22)$$

where, from eq. (20.2.19),

$$c_m(t) = \sum_n e^{i(\mathcal{E}_n t_0 - \mathcal{E}_m t)/\hbar} a_{n \rightarrow m}(t, t_0) c_n(t_0). \quad (20.2.23)$$



Note that  $a_{n \rightarrow m}(t_0, t_0) = \langle m | n \rangle = \delta_{nm}$ , so that eq. (20.2.23) is consistent for  $t = t_0$ . The non-trivial part of the time evolution of the system due to  $W(t)$  is encoded in the transition amplitudes  $a_{n \rightarrow m}(t, t_0)$ , which in turn rely on our solution for  $\mathcal{U}_I(t, t_0)$  in eq. (20.2.16).

As a special case, suppose we are only interested in transitions between eigenstates of  $H_0$ , but with time evolution governed by the full Hamiltonian  $H = H_0 + W(t)$ . The probability to start at time  $t_0$  in one of the  $H_0$  eigenstates  $|n\rangle$  and find the system at time  $t$  in another  $H_0$  eigenstate  $|m\rangle$  is

$$\mathcal{P}_{n \rightarrow m} = |a_{n \rightarrow m}|^2. \quad (20.2.24)$$

More generally, the transition amplitudes  $a_{n \rightarrow m}$  defined in eq. (20.2.20) can be used to find the time evolution of an arbitrary state  $|\psi(t)\rangle$ , through eqs. (20.2.22) and (20.2.23).

## 20.3 Exercises

**Exercise 20.1.** Consider the 1-d harmonic oscillator for a particle of mass  $m$  and frequency  $\omega$ .

- (a) Use Theorem 2.4.3 to find the Heisenberg-picture operators  $a_H(t)$  and  $a_H^\dagger(t)$  in terms of the corresponding Schrödinger picture operators, taking them to coincide at  $t = t_0 = 0$ . Use the results to find  $X_H(t)$  and  $P_H(t)$ . Check that your results satisfy the differential equations from eq. (20.1.11).
- (b) Find the commutators  $[X_H(t), P_H(t')]$  and  $[X_H(t), X_H(t')]$  and  $[P_H(t), P_H(t')]$ .
- (c) Find the ground state correlation functions,  $\langle 0 | X_H(t) X_H(t') | 0 \rangle$  and  $\langle 0 | P_H(t) P_H(t') | 0 \rangle$ .

**Exercise 20.2.** A particle of mass  $m$  has a Hamiltonian  $H = H_0 + W$ , where  $H_0 = P^2/2m$  and  $W = V(X)$  does not depend on time.

- (a) Find the interaction picture operators  $X_I(t)$  and  $P_I(t)$  in terms of the corresponding Schrödinger picture operators, with which they coincide at  $t = 0$ .
- (b) Find the commutators  $[X_I(t), P_I(t')]$  and  $[P_I(t), P_I(t')]$  and  $[X_I(t), X_I(t')]$ . In particular, show that the last one does not vanish except when  $t' = t$ .
- (c) Repeat parts (a) and (b) with the roles of  $H_0$  and  $W$  reversed, so that  $H_0 = V(X)$  and  $W = P^2/2m$ .

## 21 Time-dependent perturbation theory

### 21.1 The short-time and sudden approximations

As a warm-up for a more general treatment of time-dependent perturbation theory, suppose that the Hamiltonian for a system is

$$H(t) = \begin{cases} H_0 & (t < 0), \\ H_{\text{int}} & (0 < t < \delta), \\ H'_0 & (t > \delta). \end{cases} \quad (21.1.1)$$

The starting and ending Hamiltonians are assumed to have no time dependence, and to have known eigenstates and eigenvalues,  $|n\rangle$  and  $\mathcal{E}_n$  for  $H_0$ , and  $|m'\rangle$  and  $\mathcal{E}_{m'}$  for  $H'_0$ . In the interregnum  $0 < t < \delta$ , the Hamiltonian is  $H_{\text{int}}$ . Consider the **short-time approximation**, in which  $\delta$  is small compared to the time scales set by  $\hbar$  divided by characteristic energy differences. The idea is to take advantage of the smallness of  $\delta$  to develop an approximation for the state of the system at late times, despite the possibly complicated form of  $H_{\text{int}}$ , and in particular even if we do not know its eigenstates and eigenvalues. We assume that  $\delta$  is so small that any explicit time dependence of  $H_{\text{int}}$  over that time interval can be ignored.

For each of  $t \leq 0$  and  $t \geq \delta$ , we can use the Schrödinger equation to write the form of the state as

$$|\psi(t)\rangle = \sum_n b_n e^{-it\mathcal{E}_n/\hbar} |n\rangle \quad (t \leq 0), \quad (21.1.2)$$

$$|\psi(t)\rangle = \sum_{m'} c_{m'} e^{-i(t-\delta)\mathcal{E}_{m'}/\hbar} |m'\rangle \quad (t \geq \delta), \quad (21.1.3)$$

for some constants  $b_n$  and  $c_{m'}$ . For later convenience, a constant phase  $e^{i\mathcal{E}_{m'}\delta/\hbar}$  has been included in eq. (21.1.3), by choice of the definition of  $c_{m'}$ . For  $0 < t < \delta$ , the Schrödinger equation is

$$i\hbar \frac{d}{dt} |\psi(t)\rangle = H_{\text{int}} |\psi(t)\rangle. \quad (21.1.4)$$

So, to first order in  $\delta$ ,

$$|\psi(\delta)\rangle = \left( I - i\frac{\delta}{\hbar} H_{\text{int}} \right) |\psi(0)\rangle. \quad (21.1.5)$$

Comparing this to eq. (21.1.2) with  $t = 0$ , and eq. (21.1.3) with  $t = \delta$ , we find

$$\sum_{m'} c_{m'} |m'\rangle = \left( I - i\frac{\delta}{\hbar} H_{\text{int}} \right) \sum_n b_n |n\rangle. \quad (21.1.6)$$

Now taking the inner product with another  $H'_0$  eigenstate  $\langle k'|$ , and then relabeling  $k' \rightarrow m'$ , we get the solution

$$c_{m'} = \sum_n b_n a_{n \rightarrow m'}, \quad (21.1.7)$$

where the transition amplitude is

$$a_{n \rightarrow m'} = \langle m' | n \rangle - i \frac{\delta}{\hbar} \langle m' | H_{\text{int}} | n \rangle. \quad (21.1.8)$$

Given the initial state, specified by the coefficients  $b_n$  of the  $H_0$  eigenstates, the final state is determined by the coefficients  $c_{m'}$  of the  $H'_0$  eigenstates.

One application of eq. (21.1.8) is to the special case  $H'_0 = H_0$ , so that  $H_{\text{int}} - H_0$  acts for a short time as a perturbation to an otherwise constant Hamiltonian. Then  $\langle m | n \rangle = \delta_{nm}$ , and

$$a_{n \rightarrow m} = \delta_{nm} - i \frac{\delta}{\hbar} \sum_n \langle m | H_{\text{int}} | n \rangle. \quad (21.1.9)$$

For example, if the initial state is an eigenstate  $|n\rangle$  of  $H_0$ , so that only one of the  $b_n$  is non-zero, then the probability of finding the system in a different  $H_0$  eigenstate  $|m\rangle$  after  $t > \delta$  is

$$\mathcal{P}_{n \rightarrow m} = \frac{\delta^2}{\hbar^2} |\langle m | H_{\text{int}} | n \rangle|^2. \quad (21.1.10)$$

In the next section, we will see that this result can be obtained as a special case application of a general treatment that we will develop for time-dependent perturbation theory; see eqs. (21.2.8) and (21.2.9) with  $W = H_{\text{int}} - H_0$  for  $0 < t < \delta$ .

The **sudden approximation** is obtained if we assume that  $\delta$  is so small that  $H_{\text{int}}$  is irrelevant, but  $H'_0 \neq H_0$ . Then, if the system was originally in an  $H_0$  eigenstate  $|n\rangle$  for  $t < 0$ , the probability of finding the system in an  $H'_0$  eigenstate  $|m'\rangle$  for  $t > 0$  is

$$\mathcal{P}_{n \rightarrow m'} = |c_{m'}|^2 = |\langle m' | n \rangle|^2. \quad (21.1.11)$$

This follows from the  $\delta \rightarrow 0$  limit of eq. (21.1.8).

As an example, suppose that a one-dimensional harmonic oscillator is in its ground state, when the angular frequency in the Hamiltonian suddenly doubles, due to some outside agent. Let us compute the probability that the oscillator will be found in the ground state of the new Hamiltonian. Working in the position representation, the normalized wavefunctions of the ground states are

$$\langle x | 0 \rangle_\omega = \left( \frac{m\omega}{\pi\hbar} \right)^{1/4} \exp(-m\omega x^2 / 2\hbar) \quad (\text{before}), \quad (21.1.12)$$

$$\langle x | 0 \rangle_{2\omega} = \left( \frac{2m\omega}{\pi\hbar} \right)^{1/4} \exp(-m\omega x^2 / \hbar) \quad (\text{after}). \quad (21.1.13)$$

Then, in the sudden approximation, we compute

$${}_{2\omega} \langle 0 | 0 \rangle_\omega = \int_{-\infty}^{\infty} dx {}_{2\omega} \langle 0 | x \rangle \langle x | 0 \rangle_\omega = \sqrt{\frac{\sqrt{2}m\omega}{\pi\hbar}} \int_{-\infty}^{\infty} dx \exp(-3m\omega x^2 / 2\hbar) = \frac{2^{3/4}}{3^{1/2}}, \quad (21.1.14)$$

which results in  $\mathcal{P}_{0_\omega \rightarrow 0_{2\omega}} = 2\sqrt{2}/3 \approx 0.9428$ .

As a more practical example, consider the effect of the decay of the tritium ( $^3\text{H}$ ) nucleus on the atomic electron state. The tritium nucleus consists of 1 proton and 2 neutrons, so  $Z = 1$ . It undergoes a  $\beta$  decay to a helium isotope and an electron and antineutrino,

$$^3\text{H} \rightarrow ^3\text{He} + e^- + \bar{\nu}_e. \quad (21.1.15)$$

The antineutrino has only weak interactions, and so can be ignored for the present purposes. The  $\beta$ -decay electron is emitted with average (maximum) kinetic energy 0.0057 MeV (0.0186 MeV), so that its average (maximum) speed can be computed to be  $0.15c$  ( $0.26c$ ). These are much larger than the typical semi-classical speeds of electrons in a  $Z = 1$  atom,  $v = \alpha c \approx c/137$ . Therefore, it is a good approximation to also ignore the  $\beta$ -decay electron, as it leaves the scene very quickly.

The remaining  $^3\text{He}$  atom consists of the nucleus with 2 protons and 1 neutron, so  $Z = 2$ , together with the left-over atomic electron from the  $^3\text{H}$  atom. Therefore, in the sudden approximation the probability of the atomic electron starting in the ground state of  $^3\text{H}$  and ending up in the ground state of the  $^3\text{He}^+$  ion is  $\mathcal{P}_{1,0,0 \rightarrow 1,0,0} = |a_{1,0,0 \rightarrow 1,0,0}|^2$ , where

$$a_{1,0,0 \rightarrow 1,0,0} = \int d^3\vec{r} [\psi_{1,0,0}^{Z=2}(\vec{r})]^* \psi_{1,0,0}^{Z=1}(\vec{r}). \quad (21.1.16)$$

Recalling that

$$\psi_{1,0,0}^Z(\vec{r}) = \sqrt{\frac{Z^3}{\pi a_0^3}} e^{-Zr/a_0}, \quad (21.1.17)$$

we have

$$a_{1,0,0 \rightarrow 1,0,0} = 4\pi \int_0^\infty dr r^2 \frac{2^{3/2}}{\pi a_0^3} e^{-3r/a_0} = 2^{9/2}/3^3, \quad (21.1.18)$$

and the probability of the electron remaining in the atomic ground state after the decay is

$$\mathcal{P}_{1,0,0 \rightarrow 1,0,0} = 2^9/3^6 = 512/729 \approx 0.702. \quad (21.1.19)$$

## 21.2 Transition amplitudes and probabilities in perturbation theory

In section 3.4, we learned how to solve for the time-dependence of quantum state. If the Hamiltonian does not depend on time, and one can solve its eigenvalue problem, then the time evolution can always be found by writing the unitary operator  $U(t, t_0)$  in terms of the stationary states with appropriate phases factors that depend on the energy; see eqs. (3.4.1) and (3.4.7). However, if the Hamiltonian does depend on time, then as we saw in section 20.2, the time

evolution is given in terms of a Dyson series that typically cannot be obtained exactly in closed form. The approximation that follows from truncating the Dyson series in the interaction picture is known as time-dependent perturbation theory.

Just as in our discussion of the interaction picture, we suppose that the full Hamiltonian is

$$H(t) = H_0 + W(t), \quad (21.2.1)$$

where  $H_0$  does not depend explicitly on time, and we assume that we know (at least approximately) the solutions to its eigenvalue problem,

$$H_0|n\rangle = \mathcal{E}_n|n\rangle. \quad (21.2.2)$$

Since  $H_0$  is an observable, the kets  $|n\rangle$  form a complete orthobasis. The remaining part of the full Hamiltonian,  $W(t)$ , contains the explicit time dependence. Then, given the state of the system at time  $t_0$ , the solution for the state at a later time  $t$  is found in terms of the transition amplitudes  $a_{n \rightarrow m}$  defined by eq. (20.2.20), which relies on eq. (20.2.16), where

$$W_I(t) = e^{iH_0t/\hbar} W(t) e^{-iH_0t/\hbar} \quad (21.2.3)$$

is the perturbation part of the Hamiltonian, translated into the interaction picture.

The time-dependent perturbative expansion simply amounts to assuming that  $W(t)$  is small, so that we can expand the transition amplitude in eq. (20.2.20) as

$$a_{n \rightarrow m}(t, t_0) = a_{n \rightarrow m}^{(0)} + a_{n \rightarrow m}^{(1)} + a_{n \rightarrow m}^{(2)} + a_{n \rightarrow m}^{(3)} + \cdots, \quad (21.2.4)$$

where each term  $a_{n \rightarrow m}^{(N)}$  contains  $N$  factors of  $W$ , following from the corresponding terms in eq. (20.2.16). In practice, eq. (21.2.4) is then truncated at some finite  $N$ , usually at  $N = 1$  or  $N = 2$ . Note that the zeroth order contribution is just

$$a_{n \rightarrow m}^{(0)} = \langle m|n\rangle = \delta_{nm}, \quad (21.2.5)$$

as follows from the identity operator ( $N = 0$ ) part of  $\mathcal{U}_I$ .

The preceding tells us that the first-order contribution to the transition amplitude is

$$a_{n \rightarrow m}^{(1)} = -\frac{i}{\hbar} \int_{t_0}^t dt_1 \langle m| e^{iH_0t_1/\hbar} W(t_1) e^{-iH_0t_1/\hbar} |n\rangle. \quad (21.2.6)$$

Fortunately, each  $H_0$  can be evaluated acting on a neighboring eigenket or eigenbra. So, defining

$$\omega_{mn} = (\mathcal{E}_m - \mathcal{E}_n)/\hbar, \quad (21.2.7)$$

we obtain

$$a_{n \rightarrow m}^{(1)} = -\frac{i}{\hbar} \int_{t_0}^t dt_1 e^{i\omega_{mn}t_1} \langle m| W(t_1) |n\rangle. \quad (21.2.8)$$

This result provides the time evolution of a general state from  $t_0$  to  $t$ , at the leading order in time-dependent perturbation theory, using eqs. (20.2.22)–(20.2.23) with  $a_{n \rightarrow m} = \delta_{nm} + a_{n \rightarrow m}^{(1)}$ .

Before moving on to consider the higher-order contributions, consider the special case that the system starts at time  $t_0$  in an eigenstate  $|n\rangle$  of  $H_0$ , and we then measure  $H_0$  at time  $t$ . The transition probability to transfer from the stationary state  $|n\rangle$  to a *different* eigenstate  $|m\rangle$  of  $H_0$ , at the leading non-trivial order in the perturbation, is simply

$$\mathcal{P}_{n \rightarrow m} = |a_{n \rightarrow m}^{(1)}|^2, \quad (\text{for } m \neq n). \quad (21.2.9)$$

The validity of the approximation requires that we find  $|a_{n \rightarrow m}^{(1)}| \leq 1$ , so that the probability does not exceed 1. This is a necessary but not sufficient condition, because the sum of  $|a_{n \rightarrow m}^{(1)}|^2$  for multiple final states  $|m\rangle$  could still exceed 1. And, even if that does not happen, the series expansion in eq. (21.2.4) could still fail to converge when higher-order terms are included. The perturbation expansion is often, but not always, more reliable if  $|a_{n \rightarrow m}^{(1)}| \ll 1$ .

It is also instructive to consider what happens if we try to use the first-order approximation to directly compute the probability to remain in the initial stationary state  $|n\rangle$ . Since  $a_{n \rightarrow n}^{(0)} = 1$ , we get, instead of eq. (21.2.9):

$$\mathcal{P}_{n \rightarrow n} = |1 + a_{n \rightarrow n}^{(1)} + a_{n \rightarrow n}^{(2)} + \cdots|^2. \quad (21.2.10)$$

Now, as part of the Hamiltonian,  $W(t_1)$  must be Hermitian, so its expectation values are always real, and

$$a_{n \rightarrow n}^{(1)} = -\frac{i}{\hbar} \int_{t_0}^t dt_1 \langle n | W(t_1) | n \rangle \quad (21.2.11)$$

is a pure imaginary number. Therefore, although one might naively neglect  $a_{n \rightarrow n}^{(2)}$  as being of higher order, eq. (21.2.10) would then give  $1 + |a_{n \rightarrow n}^{(1)}|^2$ , which is clearly unacceptable for a probability unless the first-order correction vanishes. To get a meaningful result from this direct calculation, we must work consistently to at least second order, by also including the contribution from  $a_{n \rightarrow n}^{(2)}$  in the transition amplitude, with the result [dropping contributions of third order in  $W(t)$ ]

$$\mathcal{P}_{n \rightarrow n} = 1 + |a_{n \rightarrow n}^{(1)}|^2 + 2\text{Re}[a_{n \rightarrow n}^{(2)}]. \quad (21.2.12)$$

Alternatively, the probability to remain in the initial eigenstate of  $H_0$  can be obtained indirectly, from the complementary probability,

$$\mathcal{P}_{n \rightarrow n} = 1 - \sum_{m \neq n} \mathcal{P}_{n \rightarrow m} = 1 - \sum_{m \neq n} |a_{n \rightarrow m}^{(1)}|^2, \quad (21.2.13)$$

which is also second order in the perturbation. Comparing these, we learn that

$$2\text{Re}[a_{n \rightarrow n}^{(2)}] = - \sum_m |a_{n \rightarrow m}^{(1)}|^2, \quad (21.2.14)$$

with the sum over  $m$  now including  $n$ .

Now consider the second-order contribution to the transition amplitude, obtained from eqs. (20.2.16), (21.2.3), and eq. (20.2.20),

$$a_{n \rightarrow m}^{(2)} = \left(-\frac{i}{\hbar}\right)^2 \int_{t_0}^t dt_2 \int_{t_0}^{t_2} dt_1 \langle m | e^{iH_0 t_2 / \hbar} W(t_2) e^{iH_0(t_1 - t_2) / \hbar} W(t_1) e^{-iH_0 t_1 / \hbar} | n \rangle. \quad (21.2.15)$$

To evaluate this, we use completeness, by inserting a sum over all eigenstates of  $H_0$ ,

$$I = \sum_{k_1} |k_1\rangle \langle k_1|, \quad (21.2.16)$$

immediately before  $W(t_1)$ . This puts every  $H_0$  adjacent to one of its eigenstate bras or kets, allowing it to be replaced by  $\mathcal{E}_n$ ,  $\mathcal{E}_m$ , or  $\mathcal{E}_{k_1}$ , with the result

$$a_{n \rightarrow m}^{(2)} = \left(-\frac{i}{\hbar}\right)^2 \int_{t_0}^t dt_2 \int_{t_0}^{t_2} dt_1 \sum_{k_1} e^{i\omega_{mk_1} t_2} \langle m | W(t_2) | k_1 \rangle e^{i\omega_{k_1 n} t_1} \langle k_1 | W(t_1) | n \rangle. \quad (21.2.17)$$

It is left to Exercise 21.3 to perform the check of taking the special case  $|m\rangle = |n\rangle$  and manipulating the result to verify that eq. (21.2.14) indeed holds.

In the same way, the third-order contribution is found to be

$$a_{n \rightarrow m}^{(3)} = \left(-\frac{i}{\hbar}\right)^3 \int_{t_0}^t dt_3 \int_{t_0}^{t_3} dt_2 \int_{t_0}^{t_2} dt_1 \sum_{k_1} \sum_{k_2} e^{i\omega_{mk_2} t_3} \langle m | W(t_3) | k_2 \rangle e^{i\omega_{k_2 k_1} t_2} \langle k_2 | W(t_2) | k_1 \rangle e^{i\omega_{k_1 n} t_1} \langle k_1 | W(t_1) | n \rangle. \quad (21.2.18)$$

A diagrammatic representation of the contributions  $a_{n \rightarrow m}^{(N)}$  is shown in Figure 21.2.1. Each line with an arrow represents an eigenstate of  $H_0$ , as labeled. For each vertex with incoming state line  $k$ , outgoing state line  $k'$ , and time label  $t_j$ , one writes a factor

$$-\frac{i}{\hbar} e^{i\omega_{k'k} t_j} \langle k' | W(t_j) | k \rangle. \quad (21.2.19)$$

Then one integrates  $\int_{t_0}^{t_{j+1}} dt_j$  for each of the intermediate times  $t_1, \dots, t_N$  in the diagram for  $a_{n \rightarrow m}^{(N)}$ , with the identification  $t_{N+1} = t$ . All internal state labels  $k_j$  are summed over. This diagrammatic mnemonic allows us to quickly write down the formula for  $a_{n \rightarrow m}^{(N)}$ , corresponding to each sub-diagram starting with  $n$  and ending with  $m$ .

$$\begin{aligned}
\begin{array}{c} n \\ \hline \longrightarrow \\ t_0 \end{array} \begin{array}{c} m \\ \hline \longrightarrow \\ t \end{array} &= \begin{array}{c} n \\ \longrightarrow \\ t_0 \end{array} \begin{array}{c} m \\ \longrightarrow \\ t \end{array} + \begin{array}{c} n \\ \longrightarrow \\ t_0 \end{array} \begin{array}{c} \bullet \\ t_1 \end{array} \begin{array}{c} m \\ \longrightarrow \\ t \end{array} + \begin{array}{c} n \\ \longrightarrow \\ t_0 \end{array} \begin{array}{c} \bullet \\ t_1 \end{array} \begin{array}{c} k_1 \\ \longrightarrow \\ t_2 \end{array} \begin{array}{c} m \\ \longrightarrow \\ t \end{array} \\
&+ \begin{array}{c} n \\ \longrightarrow \\ t_0 \end{array} \begin{array}{c} \bullet \\ t_1 \end{array} \begin{array}{c} k_1 \\ \longrightarrow \\ t_2 \end{array} \begin{array}{c} \bullet \\ t_2 \end{array} \begin{array}{c} k_2 \\ \longrightarrow \\ t_3 \end{array} \begin{array}{c} m \\ \longrightarrow \\ t \end{array} + \begin{array}{c} n \\ \longrightarrow \\ t_0 \end{array} \begin{array}{c} \bullet \\ t_1 \end{array} \begin{array}{c} k_1 \\ \longrightarrow \\ t_2 \end{array} \begin{array}{c} \bullet \\ t_2 \end{array} \begin{array}{c} k_2 \\ \longrightarrow \\ t_3 \end{array} \begin{array}{c} \bullet \\ t_3 \end{array} \begin{array}{c} k_3 \\ \longrightarrow \\ t_4 \end{array} \begin{array}{c} m \\ \longrightarrow \\ t \end{array} + \dots
\end{aligned}$$

Figure 21.2.1: A diagrammatic representation of the time-dependent perturbation theory expansion for the transition amplitude  $a_{n \rightarrow m}(t, t_0)$ , represented by the double line with arrow, on the left. In the sum on the right side of the equality, each contribution with  $N$  vertices represents  $a_{n \rightarrow m}^{(N)}(t, t_0)$ . Each single line with arrow represents an  $H_0$  eigenstate, as labeled. Each vertex labeled by a time  $t_j$  and incoming line  $k$  and outgoing line  $k'$  represents a factor of  $-\frac{i}{\hbar} e^{i\omega_{k'k}t_j} \langle k' | W(t_j) | k \rangle$ , with an integration  $\int_{t_0}^{t_{j+1}} dt_j$ . This is done for each of the intermediate times  $t_1, \dots, t_N$ , with the identification  $t_{N+1} = t$ . All internal  $H_0$  eigenstate labels  $k_j$  are summed over.

### 21.3 Applying first-order time-dependent perturbation theory, and Fermi's golden rule

In this section and the next, we consider the application of time-dependent perturbation theory in the approximation of keeping only the first-order contribution to the transition amplitude, eq. (21.2.8).

As a first example, consider a one-dimensional harmonic oscillator with the usual unperturbed Hamiltonian  $H_0 = \frac{P^2}{2m} + \frac{1}{2}m\omega^2 X^2$ , taken to be in its ground state  $|0\rangle$  in the far past, at time  $t = -\infty$ . A perturbation

$$W(t) = -qE_0 e^{-t^2/\tau^2} X \quad (21.3.1)$$

is applied, corresponding to a continuous turning on and then off of a uniform electric field  $E_0$  that couples to the particle's charge  $q$ . The constant  $\tau$  controls the time scale of the turn-on and turn-off. Our goal is to find the probability that the particle is found in a given excited state  $|n\rangle$  when the perturbation has become negligible in the far future as  $t \rightarrow \infty$ .

Applying eq. (21.2.8) with  $\mathcal{E}_n = \hbar\omega(n + 1/2)$ , the amplitude for the transition in the far future, when the perturbation has effectively turned off, is

$$a_{0 \rightarrow n}^{(1)}(t = \infty, t_0) = -\frac{i}{\hbar} \int_{t_0}^{\infty} dt_1 e^{i\omega t_1} (-qE_0 e^{-t_1^2/\tau^2}) \langle n | X | 0 \rangle. \quad (21.3.2)$$

The necessary matrix element is

$$\langle n | X | 0 \rangle = \sqrt{\frac{\hbar}{2m\omega}} \delta_{n1}, \quad (21.3.3)$$



so that within the first-order approximation, there is a selection rule that only the first excited state can be reached. Taking the limit  $t_0 \rightarrow -\infty$ ,

$$a_{0 \rightarrow 1}^{(1)} = \frac{iqE_0}{\sqrt{2m\hbar\omega}} \int_{-\infty}^{\infty} dt_1 \exp(i\omega t_1 - t_1^2/\tau^2). \quad (21.3.4)$$

The integral can be done by completing the square, using the change of variables  $u = t_1/\tau - i\omega\tau/2$ , resulting in a transition probability

$$\mathcal{P}(0 \rightarrow 1) = \frac{\pi q^2 E_0^2 \tau^2}{2m\hbar\omega} e^{-\omega^2 \tau^2/2}, \quad (21.3.5)$$

while  $a_{0 \rightarrow n}^{(1)}$  and  $\mathcal{P}(0 \rightarrow n)$  vanish for all  $n > 1$ . Since  $\mathcal{P}$  cannot exceed 1, the approximation leading to eq. (21.3.5) must fail if the applied field is too large, that is if  $q^2 E_0^2 \tau^2$  is too large in comparison to  $m\hbar\omega e^{\omega^2 \tau^2/2}$ .

As a second example with many applications, we consider a generic unspecified  $H_0$ , and a perturbation that turns on at  $t = 0$  and is constant thereafter,

$$W(t) = \begin{cases} 0 & (\text{for } t < 0), \\ V & (\text{for } t > 0). \end{cases} \quad (21.3.6)$$

To be as general as possible, let  $V$  be an arbitrary Hermitian operator, with the only restriction that it has no time dependence. Now consider the probability that the perturbation will induce a transition from an initial  $H_0$  eigenstate  $|i\rangle$  to a different final  $H_0$  eigenstate  $|f\rangle$ . From eq. (21.2.8), the amplitude for the transition is

$$a_{i \rightarrow f}^{(1)} = -\frac{i}{\hbar} \langle f|V|i\rangle \int_0^t dt_1 e^{i\omega_{fi}t_1}, \quad (21.3.7)$$

where we have exploited the fact that the matrix element of  $V$  does not depend on time to pull it out of the integral, and defined  $\omega_{fi} = (\mathcal{E}_f - \mathcal{E}_i)/\hbar$ . Evaluating the integral, and squaring the magnitude of the result, we obtain the transition probability for  $t > 0$ ,

$$\mathcal{P}_{i \rightarrow f}(t) = |a_{i \rightarrow f}^{(1)}|^2 = \frac{4 \sin^2(t\omega_{fi}/2)}{\hbar^2 \omega_{fi}^2} |\langle f|V|i\rangle|^2, \quad (21.3.8)$$

valid at first order in perturbation theory. For future convenience, we define the function

$$F(\mathcal{E}_f - \mathcal{E}_i, t) \equiv \frac{4}{(\mathcal{E}_f - \mathcal{E}_i)^2} \sin^2\left(\frac{t(\mathcal{E}_f - \mathcal{E}_i)}{2\hbar}\right), \quad (21.3.9)$$

so that

$$\mathcal{P}_{i \rightarrow f}(t) = F(\mathcal{E}_f - \mathcal{E}_i, t) |\langle f|V|i\rangle|^2. \quad (21.3.10)$$

The transitions at first order in perturbation theory are limited to those that satisfy

$$\langle f|V|i\rangle \neq 0, \quad (21.3.11)$$




Figure 21.3.1: The transition probability at first order in time-dependent perturbation theory,  $\mathcal{P}_{i \rightarrow f} = |a_{i \rightarrow f}^{(1)}|^2$ , as a function of the time  $t$ , for a constant perturbation  $V$  that turns on at time  $t = 0$ . For short times, the probability grows quadratically like  $t^2 |\langle f|V|i \rangle|^2 / \hbar^2$ , and for longer times oscillates between 0 and a maximum value  $4 |\langle f|V|i \rangle|^2 / (\mathcal{E}_f - \mathcal{E}_i)^2$ .

an example of a matrix element selection rule.

When applying eq. (21.3.10) in practice, it is important to distinguish the cases for which  $|f\rangle$  is a discrete state with an isolated  $H_0$  eigenvalue  $\mathcal{E}_f$  (for example, a bound state of an electron in an atom) from those in which  $|f\rangle$  is a state in a continuum (for example, a free-particle plane wave, or an unbound state of an electron ionized from an atom but still under the influence of the Coulomb potential of the nucleus).

First, suppose that  $|f\rangle$  is discrete. In that case, eq. (21.3.10) can be directly applied. For very small  $t$ , the transition probability grows quadratically, because for small  $t$ ,

$$F(\mathcal{E}_f - \mathcal{E}_i, t) \approx t^2 / \hbar^2, \quad (21.3.12)$$

which is independent of the unperturbed energies. Therefore, eq.(21.3.10) becomes

$$\mathcal{P}_{i \rightarrow f} \approx \frac{t^2}{\hbar^2} |\langle f|V|i \rangle|^2 \quad (\text{small } t). \quad (21.3.13)$$

Of course, this must fail for sufficiently large  $t$ , as the probability cannot exceed 1. Equation (21.3.13) agrees with the result we had already found in the short-time approximation in eq. (21.1.10). For longer time scales, the transition probability for a given discrete final state oscillates between 0 and a maximum value  $4 |\langle f|V|i \rangle|^2 / (\mathcal{E}_f - \mathcal{E}_i)^2$ , as illustrated<sup>†</sup> in Figure 21.3.1.

Due to the denominator, the final states  $|f\rangle$  that have  $\mathcal{E}_f \approx \mathcal{E}_i$  will tend to have the largest probabilities, provided that the matrix element  $\langle f|V|i \rangle$  is not too small in magnitude. Also, the probability for a given final state  $|f\rangle$  vanishes whenever  $t = 2\pi n / \omega_{fi} = 2\pi \hbar n / (\mathcal{E}_f - \mathcal{E}_i)$  for integer  $n$ . These features are illustrated by the graph of  $F(\mathcal{E}_f - \mathcal{E}_i, t)$  as a function of  $\mathcal{E}_f - \mathcal{E}_i$ ,

<sup>†</sup>As a function of the time  $t$ , the probability oscillates with a similar appearance to the Rabi oscillations found for a magnetic spin resonance system in section 9.2; compare to Figure 9.2.1. However, the physical reason behind the oscillations is quite different, because in the present case the excitation is driven by a constant interaction rather than a harmonic driving frequency.




Figure 21.3.2: The function  $F(\mathcal{E}_f - \mathcal{E}_i, t)$  defined by eq. (21.3.9) and appearing in the transition probability for first-order time-dependent perturbation theory in eq. (21.3.10), as a function of  $\mathcal{E}_f - \mathcal{E}_i$ . The main peak height  $t^2/\hbar^2$  grows quadratically with time, while its width shrinks proportionally to  $1/t$ . The total area under the curve is  $2\pi t/\hbar$ .

shown in Figure 21.3.2. The final states  $|f\rangle$  most likely to be reached at a given time  $t$  are those with energies that lie under the main peak around  $\mathcal{E}_i$ , which is bounded by

$$|\mathcal{E}_f - \mathcal{E}_i| < 2\pi\hbar/t. \quad (21.3.14)$$

The main peak gets narrower and taller for larger  $t$ , so that in the limit of long times, the most likely transitions are those for which the difference in unperturbed energies is constrained, increasingly strongly, by eq. (21.3.14).

Now consider the case that  $|f\rangle$  is not discrete, so that the unperturbed energy eigenvalues  $\mathcal{E}_f$  and the matrix elements  $\langle f|V|i\rangle$  each form a continuum. In that case, we write

$$|f\rangle = |\mathcal{E}_f, u_{\mathcal{E}_f}\rangle, \quad (21.3.15)$$

where  $u_{\mathcal{E}_f}$  represents the degeneracy label (or labels) for the unperturbed energy. (For example, the degeneracy labels could include the angular direction of a final-state momentum vector, and/or an unobserved spin.) We also define the density of states  $\rho(\mathcal{E}_f, u_{\mathcal{E}_f})$  according to

$$\rho(\mathcal{E}_f, u_{\mathcal{E}_f}) d\mathcal{E}_f du_{\mathcal{E}_f} = \left( \begin{array}{c} \text{number of states between } \mathcal{E}_f \text{ and } \mathcal{E}_f + d\mathcal{E}_f, \\ \text{and between } u_{\mathcal{E}_f} \text{ and } u_{\mathcal{E}_f} + du_{\mathcal{E}_f} \end{array} \right), \quad (21.3.16)$$

where, for now, the degeneracy labels are also assumed continuous. The corresponding contribution to the transition probability is, from eq. (21.3.10),

$$d\mathcal{P}_{i \rightarrow f} = |\langle f|V|i\rangle|^2 F(\mathcal{E}_f - \mathcal{E}_i, t) \rho(\mathcal{E}_f, u_{\mathcal{E}_f}) d\mathcal{E}_f du_{\mathcal{E}_f}, \quad (21.3.17)$$

in terms of the function  $F$  defined in eq.(21.3.9). In practice, eq. (21.3.17) must always be integrated over some ranges of  $\mathcal{E}_f$  and  $u_{\mathcal{E}_f}$  in order to give the probability of transition to that range of states,

$$\Delta\mathcal{P} = \int d\mathcal{P}_{i \rightarrow f}. \quad (21.3.18)$$

If we want only the transition probability for some of the final states (for example, those with energy in some range, or momentum direction inside some cone), this can be enforced either by integrating only over the appropriate ranges of energy and degeneracy labels, or by putting a projection operator before  $V$  within the matrix element  $\langle f|V|i\rangle$ .

In the limit of large time  $t$ , we can take advantage of the fact that  $F(\mathcal{E}_f - \mathcal{E}_i, t)$  as a function of  $\mathcal{E}_f$  becomes increasingly sharply peaked near  $\mathcal{E}_i$ , and so approaches proportionality to a delta function. To understand the proportionality, we note that

$$\int_{-\infty}^{\infty} d\mathcal{E}_f F(\mathcal{E}_f - \mathcal{E}_i, t) = 2\pi t/\hbar, \quad (21.3.19)$$

so that we can make the replacement

$$F(\mathcal{E}_f - \mathcal{E}_i, t) \rightarrow \frac{2\pi t}{\hbar} \delta(\mathcal{E}_f - \mathcal{E}_i) \quad (21.3.20)$$

for large  $t$ . The transition probability eq. (21.3.17) is then directly proportional to  $t$ ,

$$d\mathcal{P}_{i \rightarrow f} = t dR_{i \rightarrow f}, \quad (21.3.21)$$

where the proportionality can therefore be interpreted as a transition rate per unit time,

$$dR_{i \rightarrow f} = \frac{2\pi}{\hbar} |\langle f|V|i\rangle|^2 \delta(\mathcal{E}_f - \mathcal{E}_i) \rho(\mathcal{E}_f, u_{\mathcal{E}_f}) d\mathcal{E}_f du_{\mathcal{E}_f}. \quad (21.3.22)$$

For a group of states with  $\mathcal{E}_f \approx \mathcal{E}_i$ , the total transition rate to those states is obtained by integration over  $\mathcal{E}_f$ , and integrating (or summing) over the degeneracy label(s)  $u_{\mathcal{E}_f}$  for it. Doing the energy integration, and using the notation of eq. (21.3.15), we get the transition rate

$$R = \frac{2\pi}{\hbar} \int du_{\mathcal{E}_f} \rho(\mathcal{E}_f, u_{\mathcal{E}_f}) |\langle \mathcal{E}_f, u_{\mathcal{E}_f} | V | i \rangle|^2, \quad (21.3.23)$$

where it is now understood that only final states with  $\mathcal{E}_f = \mathcal{E}_i$  are included, and again projection operators can be used inside the matrix element to restrict to particular classes of final states with some desired properties. Either eq. (21.3.23) [or its equivalent (21.3.22)] is one version of **Fermi's golden rule**. (It was actually originally obtained by Dirac, but Fermi popularized it.) If the energy degeneracy labels  $u_{\mathcal{E}_f}$  are discrete rather than continuous, then one makes the replacement  $\int du_{\mathcal{E}_f} \rightarrow \sum_{u_{\mathcal{E}_f}}$ .

Although the derivation of the golden rule turned  $F$  into a delta function by assuming that  $t$  is large, we have to be careful if it is *too* large. This is simply because interpretation of eq. (21.3.23) as a constant rate clearly requires that  $t < 1/R$ , so that the probability does not exceed 1. More generally, for times that are not short compared to  $1/R$ , the transition rate decreases as the initial state  $|i\rangle$  is depleted. If we write  $\mathcal{P}_{i \rightarrow i}(t) = e^{-R_{\text{tot}} t}$  for the probability of

remaining in the initial state, including the effects of changing to *all* possible final states, then the transition rate to a particular class of final states approaches  $Re^{-R_{\text{tot}}t}$  at late times.

The golden rule can also be applied as an approximation in the case of a group of many discrete but closely spaced final states with approximately the same matrix elements  $\langle \mathcal{E}_f, u_{\mathcal{E}_f} | V | i \rangle$ . This works if the unperturbed energy differences  $\Delta \mathcal{E}_f$  between neighboring discrete final states are small enough that there are many states under the main peak in the function  $F(\mathcal{E}_f - \mathcal{E}_i, t)$ . This provision requires that

$$t \Delta \mathcal{E}_f \ll 2\pi\hbar, \quad (21.3.24)$$

so that the elapsed time is short compared to the inverse energy spacing between adjacent states.

## 21.4 Harmonic time-dependent perturbations

An even more common application of time-dependent perturbation theory is the case of a **harmonic time-dependent perturbation**, in which  $W(t)$  is proportional to a sine or cosine function of time. For example, this includes the case of electromagnetic waves interacting with a charged particle. Suppose that the Hamiltonian perturbation turns on at time  $t = 0$ , so

$$W(t) = \begin{cases} 0 & (\text{for } t < 0), \\ Ve^{-i\omega t} + V^\dagger e^{i\omega t} & (\text{for } t > 0), \end{cases} \quad (21.4.1)$$

where  $V$  is an operator that does not depend on time. Note that  $W(t)$  is Hermitian, but  $V$  need not be. This perturbation can also be written as  $(V + V^\dagger) \cos(\omega t) + i(V^\dagger - V) \sin(\omega t)$  for  $t > 0$ , with  $(V + V^\dagger)$  and  $i(V^\dagger - V)$  both Hermitian operators.

Applying first-order time-dependent perturbation theory in the form of eq. (21.2.8), the transition amplitude is

$$a_{i \rightarrow f}^{(1), \omega}(t) = -\frac{i}{\hbar} \int_0^t dt' \left( e^{i(\omega_f - \omega_i - \omega)t'} \langle f | V | i \rangle + e^{i(\omega_f - \omega_i + \omega)t'} \langle f | V^\dagger | i \rangle \right) \quad (21.4.2)$$

$$= \frac{1 - e^{i(\omega_f - \omega_i - \omega)t}}{\hbar(\omega_f - \omega_i - \omega)} V_{fi} + \frac{1 - e^{i(\omega_f - \omega_i + \omega)t}}{\hbar(\omega_f - \omega_i + \omega)} V_{fi}^\dagger, \quad (21.4.3)$$

where we have used a short-hand notation for the time-independent matrix elements,

$$V_{fi} = \langle f | V | i \rangle, \quad V_{fi}^\dagger = \langle f | V^\dagger | i \rangle. \quad (21.4.4)$$

Requiring that these do not vanish provides a selection rule for first-order time-dependent perturbation theory.

The first term in eq. (21.4.3) is resonant only if  $\omega_f \approx \omega_i + \omega$ , and the second term only if  $\omega_f \approx \omega_i - \omega$ . For  $\omega \neq 0$ , at most one of these conditions can be satisfied for any particular  $\mathcal{E}_f$ , so

only one term will be important within the resonance approximation. Therefore, we can make the approximation of ignoring the cross terms in  $|a_{i \rightarrow f}^{(1), \omega}(t)|^2$ . Then the probability of transition between eigenstates of  $H_0$  is

$$\mathcal{P}_{i \rightarrow f} = F(\mathcal{E}_f - \mathcal{E}_i - \hbar\omega, t) |V_{fi}|^2 + F(\mathcal{E}_f - \mathcal{E}_i + \hbar\omega, t) |V_{fi}^\dagger|^2, \quad (21.4.5)$$

where the same function  $F$  defined by eq. (21.3.9) appears again. The application of eq. (21.4.5) is slightly different depending on whether the final state is discrete or part of a continuum, although the final results will have a similar appearance.

First, consider the continuum case, for which we write

$$|f\rangle = |\mathcal{E}_f, u_{\mathcal{E}_f}\rangle \quad (21.4.6)$$

for the group of final states with  $H_0$  eigenvalue  $\mathcal{E}_f$  and degeneracy label  $u_{\mathcal{E}_f}$ . For large  $t$ , the same argument that led to eq. (21.3.22) now gives

$$d\mathcal{P}_{i \rightarrow f} = t dR_{i \rightarrow f}, \quad (21.4.7)$$

where the differential contribution to the transition rate is

$$dR_{i \rightarrow f} = \frac{2\pi}{\hbar} \left( \delta(\mathcal{E}_f - \mathcal{E}_i - \hbar\omega) |V_{fi}|^2 + \delta(\mathcal{E}_f - \mathcal{E}_i + \hbar\omega) |V_{fi}^\dagger|^2 \right) \rho(\mathcal{E}_f, u_{\mathcal{E}_f}) d\mathcal{E}_f du_{\mathcal{E}_f}. \quad (21.4.8)$$

The effect of the first term can be referred to as **absorption**; since  $\mathcal{E}_f = \mathcal{E}_i + \hbar\omega$ , the Hamiltonian perturbation  $V$  causes the initial state to gain energy  $\hbar\omega$  and jump up to the final state. The second term is **emission**, as the perturbation  $V^\dagger$  cause the initial state to lose energy  $\hbar\omega$  and jump down to the final state. Since we ignored the cross-terms, and the final state energies must be different in the two cases anyway, it makes sense to separate these, and write

$$dR_{i \rightarrow f}^{\text{abs}} = \frac{2\pi}{\hbar} |V_{fi}|^2 \delta(\mathcal{E}_f - \mathcal{E}_i - \hbar\omega) \rho(\mathcal{E}_f, u_{\mathcal{E}_f}) d\mathcal{E}_f du_{\mathcal{E}_f}, \quad (21.4.9)$$

$$dR_{i \rightarrow f}^{\text{em}} = \frac{2\pi}{\hbar} |V_{fi}^\dagger|^2 \delta(\mathcal{E}_f - \mathcal{E}_i + \hbar\omega) \rho(\mathcal{E}_f, u_{\mathcal{E}_f}) d\mathcal{E}_f du_{\mathcal{E}_f}. \quad (21.4.10)$$

After integration over  $\mathcal{E}_f$ ,

$$dR_{i \rightarrow f}^{\text{abs}} = \frac{2\pi}{\hbar} |\langle \mathcal{E}_f, u_{\mathcal{E}_f} | V | i \rangle|^2 \rho(\mathcal{E}_f, u_{\mathcal{E}_f}) du_{\mathcal{E}_f}, \quad (21.4.11)$$

$$dR_{i \rightarrow f}^{\text{em}} = \frac{2\pi}{\hbar} |\langle \mathcal{E}_f, u_{\mathcal{E}_f} | V^\dagger | i \rangle|^2 \rho(\mathcal{E}_f, u_{\mathcal{E}_f}) du_{\mathcal{E}_f}, \quad (21.4.12)$$

where it is now understood that the energy conservation conditions

$$\mathcal{E}_f = \mathcal{E}_i + \hbar\omega, \quad (\text{absorption}), \quad (21.4.13)$$

$$\mathcal{E}_f = \mathcal{E}_i - \hbar\omega, \quad (\text{emission}), \quad (21.4.14)$$

are enforced, due to the delta functions. Equations (21.4.11) and (21.4.12), or their equivalents (21.4.9) and (21.4.10), are **Fermi's golden rule for harmonic perturbations**. If the energy degeneracy labels  $u_{\mathcal{E}_f}$  are continuous, then they should be integrated over, and if they are discrete then the differential  $du_{\mathcal{E}_f}$  is removed and they can be summed over.

If the transition from  $|i\rangle$  to  $|f\rangle$  by absorbing energy  $\hbar\omega$  from  $V$  is allowed, then it follows that for the same  $\omega$  one can also have the reverse process of  $|f\rangle$  to  $|i\rangle$  by emitting energy  $\hbar\omega$  to  $V^\dagger$ ; both the energy conservation and matrix element selection rules work. For both absorption and emission, we can obtain the differential rate for the opposite process  $dR_{f \rightarrow i}$  by interchanging  $\mathcal{E}_f \leftrightarrow \mathcal{E}_i$  and  $|V_{fi}|^2 \leftrightarrow |V_{if}^\dagger|^2$ . Note that the relevant matrix elements are simply related by

$$V_{if}^\dagger = \langle i|V^\dagger|f\rangle = (\langle f|V|i\rangle)^* = (V_{fi})^*, \quad (21.4.15)$$

so  $|V_{if}^\dagger|^2 = |V_{fi}|^2$ . Let us now change the  $i$  and  $f$  labels to a more neutral notation  $a$  and  $b$ , since each can play the role of initial and final state. If we multiply eq. (21.4.12) with  $i = a$  and  $f = b$  by  $\rho(\mathcal{E}_a, u_{\mathcal{E}_a})du_{\mathcal{E}_a}$ , the result is the same as multiplying eq. (21.4.11) with  $i = b$  and  $f = a$  by  $\rho(\mathcal{E}_b, u_{\mathcal{E}_b})du_{\mathcal{E}_b}$ , after integrating over the same intervals in the degeneracy labels in each case. This shows that

$$\rho(\mathcal{E}_a, u_{\mathcal{E}_a}) \frac{dR_{a \rightarrow b + \hbar\omega}^{\text{em}}}{du_{\mathcal{E}_b}} = \rho(\mathcal{E}_b, u_{\mathcal{E}_b}) \frac{dR_{b + \hbar\omega \rightarrow a}^{\text{abs}}}{du_{\mathcal{E}_a}}. \quad (21.4.16)$$

This is known as the **detailed balance** relation between emission and absorption rates and the densities of states.

Let us now reconsider eq. (21.4.5), this time assuming that the final state  $|f\rangle$  is discrete. We now suppose that the perturbations occur as an incoherent distribution with a range of angular frequencies. By “incoherent”, it is implied that interference contributions will be negligible, so that to a good approximation the probabilities, not the amplitudes, add for different driving frequencies. To see how this works, consider instead of eq. (21.4.1), a perturbation that is a superposition of many such terms with infinitesimally spaced  $\omega$ 's with smoothly varying real amplitudes  $c_\omega$  and, crucially, phases  $\varphi_\omega$  that are assumed to be random, with no relation (coherence) whatsoever between neighboring values of  $\omega$ ,

$$W(t) = \sum_{\omega} c_{\omega} (e^{i\varphi_{\omega}} V e^{-i\omega t} + e^{-i\varphi_{\omega}} V^\dagger e^{i\omega t}) \quad (\text{for } t > 0). \quad (21.4.17)$$

After isolating the emission and absorption resonant pieces as before, the square of the transition amplitude  $a_{i \rightarrow f}^{(1)}$  will contain terms of the form

$$|a_{i \rightarrow f}^{(1)}|^2 = \sum_{\omega} \sum_{\omega'} c_{\omega} c_{\omega'} e^{i(\varphi_{\omega} - \varphi_{\omega'})} \dots, \quad (21.4.18)$$

where the  $\dots$  represents quantities that are relatively slowly varying functions of  $\omega$  and  $\omega'$ . Because of the random phases and the slow variation of the summand, all terms with  $\omega' \neq \omega$  now cancel in the limit of infinitesimal spacing of the sums, leaving behind a single sum of the form  $\sum_{\omega} c_{\omega}^2 \dots$ , coming from the subset of terms with  $\omega' = \omega$ , which add with constructive interference. This sum can now be converted to an integral

$$\sum_{\omega} c_{\omega}^2 \rightarrow \int d\omega \rho(\omega), \quad (21.4.19)$$

described by some density per unit frequency  $\rho(\omega)$ . This means that the transition probabilities for continuous  $\omega$  are assumed<sup>†</sup> to be weighted by a function  $\rho(\omega)$  that characterizes the perturbation. For example, if the perturbations take the form of electromagnetic radiation, one can relate  $\rho(\omega)$  to the intensity spectrum of the incoherent light, as we will do in section 22.2. In the presence of the integral over continuous  $\omega$ , even though  $\mathcal{E}_f$  is discrete, we can again replace

$$F(\mathcal{E}_f - \mathcal{E}_i \pm \hbar\omega, t) \rightarrow \frac{2\pi t}{\hbar} \delta(\mathcal{E}_f - \mathcal{E}_i \pm \hbar\omega), \quad (21.4.20)$$

and so obtain the differential transition rates

$$dR_{i+\hbar\omega \rightarrow f}^{\text{abs}} = \frac{2\pi}{\hbar} |V_{fi}|^2 \delta(\mathcal{E}_f - \mathcal{E}_i - \hbar\omega) \rho(\omega) d\omega, \quad (21.4.21)$$

$$dR_{i \rightarrow f+\hbar\omega}^{\text{em}} = \frac{2\pi}{\hbar} |V_{fi}^{\dagger}|^2 \delta(\mathcal{E}_f - \mathcal{E}_i + \hbar\omega) \rho(\omega) d\omega. \quad (21.4.22)$$

After integrating over  $\omega$  according to  $\int d\omega \delta(\mathcal{E}_f - \mathcal{E}_i \pm \hbar\omega) = 1/\hbar$ , we have

$$R_{i+\hbar\omega \rightarrow f}^{\text{abs}} = \frac{2\pi}{\hbar^2} |\langle f|V|i\rangle|^2 \rho(\omega), \quad (21.4.23)$$

$$R_{i \rightarrow f+\hbar\omega}^{\text{em}} = \frac{2\pi}{\hbar^2} |\langle f|V^{\dagger}|i\rangle|^2 \rho(\omega), \quad (21.4.24)$$

where it is understood now that  $\omega$  is fixed in terms of  $\mathcal{E}_f$  and  $\mathcal{E}_i$  by eqs. (21.4.14) and (21.4.13). The energy degeneracies of the final state  $|f\rangle$  are assumed here to be also discrete, and so are not indicated explicitly, but can be summed over.

Although eqs. (21.4.11)–(21.4.12) and (21.4.23)–(21.4.24) look similar, it bears emphasis that they apply in slightly different circumstances. In eqs. (21.4.23)–(21.4.24), the final states are discrete, and the driving angular frequency is selected (by energy conservation) from an incoherent continuum of perturbations to the Hamiltonian with some assumed density function with respect to angular frequency,  $\rho(\omega)$ . In eqs. (21.4.11)–(21.4.12), it is the final state that

---

<sup>†</sup>It is also possible to take the distribution of perturbations to be coherent, which means that the phases are slowly varying functions of  $\omega$ . In that case one cannot neglect interference cross-terms in the transition amplitude  $a_{i \rightarrow f}^{(1)}$  for given initial and final states. We will not consider that case here.



is selected by energy conservation from among the continuous final states with density with respect to energy,  $\rho(\mathcal{E}_f, u_{\mathcal{E}_f})$ , for a driving angular frequency that can be treated as a single monochromatic Hamiltonian perturbation. In the latter case, the results could also be summed over an incoherent continuous distribution. Both of these situations arise in the interaction of matter with electromagnetic waves, depending on whether the final electronic state is bound or unbound. This is the subject of the next chapter.

## 21.5 Exercises

**Exercise 21.1.** At the end of section 21.1, we found  $\mathcal{P} = 512/729$  for the probability that a  ${}^3\text{H}$  (tritium) atom in its ground state would end up in the ground state of  ${}^3\text{He}$  after beta decay. Find the probability that the atom ends up in each of the  $n = 2$  or  $n = 3$  levels instead. Check that the total probability for ending up in the  $n = 1, 2$ , and  $3$  levels does not exceed 1.

**Exercise 21.2.** A particle of mass  $m$  moves freely in a 1-dimensional box of length  $a$ , with the left edge at  $x = 0$  and the right edge at  $x = a$ . At time  $t = 0$ , the particle is in the ground state, and the box is suddenly lengthened by expanding the right edge out to  $x = b$ .

- (a) Find the probability that the particle ends up in each of the energy eigenstates of the final box, labeled by positive integers  $n$ .
- (b) Use your results from part (a), and your knowledge of the laws of probability, to evaluate the following mathematical infinite sum

$$S(a, b) = \sum_{n=1}^{\infty} \frac{\sin^2(n\pi a/b)}{(n^2 a^2 - b^2)^2}. \quad (21.5.1)$$

Quantum mechanics knows how to do non-trivial sums!

- (c) As a check, for the special case  $b = 2a$ , evaluate the individual probabilities numerically for each integer up to  $n = 10$ , and their total with 6 significant digits. (Be careful for  $n = 2$ ; the contribution is non-zero and finite.)

**Exercise 21.3.** Show that the real part of the second-order transition amplitude for a state to itself,  $a_{n \rightarrow n}^{(2)}$ , as found in eq. (21.2.17) with  $|m\rangle = |n\rangle$ , agrees with the right side of the sum rule eq. (21.2.14).

**Exercise 21.4.** A particle of mass  $m$  is bound in a 1-dimensional simple harmonic oscillator of natural frequency  $\omega$ . At  $t = 0$ , it is in the ground state, and a perturbation to the Hamiltonian is turned on, of the form  $W(X, t) = \lambda X^2 e^{-t/T}$ , where  $\lambda$  and  $T$  are constants.

- (a) Use first-order time-dependent perturbation theory to find the probabilities that, after a very long time ( $t \gg T$ ), the particle will have jumped to each of the excited states.

- (b) Using your answers for part (a), infer the probability that after a long time the particle will be found in the ground state.
- (c) Use second-order time-dependent perturbation theory to directly calculate the probability of finding the particle in the ground state. Check that your answer agrees with part (b).

**Exercise 21.5.** A hydrogen atom is in its ground state at time  $t = 0$ , when a decaying electric field  $\vec{E} = E_0 \hat{z} e^{-t/T}$  is applied. Use first-order time-dependent perturbation theory to find the probability for the atom to be found at  $t \gg T$  in each of the first excited states  $[(n, l, m) = (2, 0, 0) \text{ and } (2, 1, 1) \text{ and } (2, 1, 0) \text{ and } (2, 1, -1)]$ .

**Exercise 21.6.** Consider the magnetic spin resonance problem discussed in section 9.2.

- (a) Treating  $\tilde{B}$  (and therefore  $\Gamma$ ) as small, use first-order time-dependent perturbation theory to calculate the probability for transition from the state  $|\downarrow\rangle$  to the state  $|\uparrow\rangle$ , as a function of  $t$ . How does this compare to the exact answer found in eq. (9.2.16)?
- (b) When  $\omega$  is very close to  $\omega_B$ , show that the perturbative answer you got in part (a) can be interpreted in terms of a rate  $\approx N\gamma^2 \tilde{B}^2 \delta(\omega - \omega_B)$ , where  $N$  is a certain number that you will find. [Hint: note the relationship between eqs. (21.3.9) and (21.3.20).] To apply this in practice, you would integrate over a spectrum of driving frequencies  $\omega$ .

## 22 Absorption and emission of light

### 22.1 Electrons in the presence of electromagnetic waves

In this chapter, we will be discussing the interactions of electrons with a classical electromagnetic wave. The Hamiltonian is obtained from the Coulomb gauge result in eq. (4.3.35), with  $q = -e$  and  $\gamma = -g_e e / 2m_e c \approx -e / m_e c$  for electrons,

$$H = H_0 + \frac{e}{m_e c} \vec{A} \cdot \vec{P} + \frac{e^2}{2m_e c^2} A^2 - e\Phi + \frac{e}{m_e c} \vec{S} \cdot \vec{B}, \quad (22.1.1)$$

where the electron's Hamiltonian in the absence of the electromagnetic wave is

$$H_0 = \frac{P^2}{2m_e} + U(\vec{R}). \quad (22.1.2)$$

We have included here an additional potential  $U$ , which includes all effects not associated with the wave, for example the Coulomb attraction of the electron to a fixed heavy nucleus, or repulsion by other electrons. Let the wave have unit polarization vector  $\hat{\varepsilon}$ , and wavevector

$$\vec{k} = \hat{k} \omega / c, \quad (22.1.3)$$

so that the classical electric and magnetic fields are

$$\vec{E}(\vec{r}, t) = \hat{\varepsilon} E_0 \sin(\vec{k} \cdot \vec{r} - \omega t), \quad (22.1.4)$$

$$\vec{B}(\vec{r}, t) = (\hat{k} \times \hat{\varepsilon}) E_0 \sin(\vec{k} \cdot \vec{r} - \omega t), \quad (22.1.5)$$

which satisfy Maxwell's equations provided that  $\hat{\varepsilon} \cdot \hat{k} = 0$ . In Coulomb gauge, the potentials describing this classical wave are

$$\Phi(\vec{r}, t) = 0, \quad (22.1.6)$$

$$\vec{A}(\vec{r}, t) = -\hat{\varepsilon} \frac{c}{\omega} E_0 \cos(\vec{k} \cdot \vec{r} - \omega t), \quad (22.1.7)$$

as one can check using eqs. (4.3.8)–(4.3.9) and (4.3.34).

If the wave is not too intense, it is a good approximation to drop the non-linear term proportional to  $A^2$ , so that our Hamiltonian simplifies to  $H = H_0 + W$ , where

$$W(t) = \frac{e}{m_e c} (\vec{A} \cdot \vec{P} + \vec{B} \cdot \vec{S}) \quad (22.1.8)$$

$$= \frac{e E_0}{m_e \omega} \left[ -\cos(\vec{k} \cdot \vec{R} - \omega t) \hat{\varepsilon} \cdot \vec{P} + \sin(\vec{k} \cdot \vec{R} - \omega t) (\vec{k} \times \hat{\varepsilon}) \cdot \vec{S} \right]. \quad (22.1.9)$$

This can be treated as a perturbation using the formalism of section 21.4, specifically by writing

$$W(t) = V e^{-i\omega t} + V^\dagger e^{i\omega t}, \quad (22.1.10)$$

as in eq. (21.4.1), with

$$V = -\frac{eE_0}{2m_e\omega} e^{i\vec{k}\cdot\vec{R}} \left[ \hat{\varepsilon} \cdot \vec{P} + i(\vec{k} \times \hat{\varepsilon}) \cdot \vec{S} \right]. \quad (22.1.11)$$

To put this into a more compact form, we use the identity  $(\vec{k} \times \hat{\varepsilon}) \cdot \vec{S} = -\hat{\varepsilon} \cdot (\vec{k} \times \vec{S})$  to obtain

$$V = -\frac{eE_0}{2m_e\omega} \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}}, \quad (22.1.12)$$

where we have defined a vector operator

$$\vec{\mathcal{P}}_{\vec{k}} \equiv e^{i\vec{k}\cdot\vec{R}} (\vec{P} - i\vec{k} \times \vec{S}). \quad (22.1.13)$$

Note that  $\vec{\mathcal{P}}_{\vec{k}}$  has the same units as momentum, but it is not Hermitian. Its matrix elements will be used in our discussion of absorption and emission of electromagnetic waves in the next few sections. The polarization vector  $\hat{\varepsilon}$  can now be taken to be complex, for example to describe circularly polarized waves, but must satisfy  $\hat{\varepsilon} \cdot \hat{k} = 0$  and  $\hat{\varepsilon}^* \cdot \hat{\varepsilon} = 1$ .

The Hamiltonian for a system of  $N_e$  electrons in an external classical electromagnetic wave can be similarly treated using an appropriate  $H_0$  (which will include the repulsive potentials between each pair of electrons, their attractive potentials due to the heavy nucleus, and spin-orbit and spin-spin couplings) together with a harmonic perturbation

$$V = -\frac{eE_0}{2m_e\omega} \sum_{i=1}^{N_e} \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k},i}, \quad (22.1.14)$$

Here, each  $\vec{\mathcal{P}}_{\vec{k},i}$  is defined in terms of the position, momentum, and spin operators  $\vec{R}_i$ ,  $\vec{P}_i$ , and  $\vec{S}_i$  for the individual electrons, as in eq. (22.1.13). This is the linearized approximation of a Hamiltonian perturbation describing the interaction with an external electromagnetic wave.

## 22.2 Absorption of electromagnetic waves

Consider an electron described by some Hamiltonian  $H_0$ , which then encounters a classical electromagnetic wave. As described in the previous section, we can now apply time-dependent perturbation theory as discussed in section 21.4, with

$$V = -\frac{eE_0}{2m_e\omega} \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} \quad (22.2.1)$$

in eq. (21.4.1), where  $E_0$  and  $\hat{\varepsilon}$  are the electric field amplitude and unit polarization vector of the wave, and the operator  $\vec{\mathcal{P}}_{\vec{k}}$  was defined in eq. (22.1.13) in terms of the wavevector. We will compute results in terms of matrix elements of  $\vec{\mathcal{P}}_{\vec{k}}$ , and then work out the simpler electric dipole and higher multipole approximations in sections 22.4 and 22.5, using an expansion in the angular frequency  $\omega$ .

Consider the rate for an initial  $H_0$  eigenstate  $|i\rangle$  to absorb energy from the wave and end up in a group of final  $H_0$  eigenstates

$$|f\rangle = |\mathcal{E}_f, u_{\mathcal{E}_f}\rangle. \quad (22.2.2)$$

For now, we take these to have continuous eigenvalues  $\mathcal{E}_f$  and degeneracy labels  $u_{\mathcal{E}_f}$  and density of states  $\rho(\mathcal{E}_f, u_{\mathcal{E}_f})$  as defined in eq. (21.3.16). Fermi's golden rule, eq. (21.4.9), then gives the rate for absorption transitions,

$$dR_{i+\hbar\omega\rightarrow f}^{\text{abs}} = \frac{2\pi}{\hbar} \left( \frac{eE_0}{2m_e\omega} \right)^2 |\langle f | \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} | i \rangle|^2 \rho(\mathcal{E}_f, u_{\mathcal{E}_f}) du_{\mathcal{E}_f} \delta(\mathcal{E}_f - \mathcal{E}_i - \hbar\omega) d\mathcal{E}_f. \quad (22.2.3)$$

The time-averaged flux of energy in the incident wave that is causing these transitions is

$$\text{energy flux in wave} = \frac{\text{energy}}{(\text{time})(\text{area})} = \frac{cE_0^2}{8\pi}. \quad (22.2.4)$$

This can be obtained by time-averaging either  $cu_{\text{EM}}$  in eq. (4.3.6), or the magnitude of the Poynting vector  $|\vec{S}|$  in eq. (4.3.7). Since both eq. (22.2.3) and eq. (22.2.4) are proportional to  $E_0^2$ , it make sense to define a measurable quantity proportional to their ratio, in which the incident wave intensity cancels out. The appropriate ratio is called the **absorption cross-section**:

$$\sigma^{\text{abs}} = \frac{\text{absorbed energy/time}}{\text{energy flux in incident wave}}. \quad (22.2.5)$$

Note that  $\sigma^{\text{abs}}$  has units of area. The energy absorbed in each  $i \rightarrow f$  transition is  $\hbar\omega$ , so

$$d\sigma_{i+\hbar\omega\rightarrow f}^{\text{abs}} = \hbar\omega dR_{i+\hbar\omega\rightarrow f}^{\text{abs}} / (cE_0^2/8\pi), \quad (22.2.6)$$

or, in terms of the fine-structure constant  $\alpha = e^2/\hbar c$ ,

$$d\sigma_{i+\hbar\omega\rightarrow f}^{\text{abs}} = \frac{4\pi^2\alpha\hbar}{m_e^2\omega} |\langle f | \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} | i \rangle|^2 \rho(\mathcal{E}_f, u_{\mathcal{E}_f}) du_{\mathcal{E}_f} \delta(\mathcal{E}_f - \mathcal{E}_i - \hbar\omega) d\mathcal{E}_f. \quad (22.2.7)$$

If the final state energies  $\mathcal{E}_f$  are continuous near  $\mathcal{E}_i + \hbar\omega$ , one can immediately integrate over them, with the result

$$d\sigma_{i+\hbar\omega\rightarrow f}^{\text{abs}} = \frac{4\pi^2\alpha\hbar}{m_e^2\omega} |\langle f | \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} | i \rangle|^2 \rho(\mathcal{E}_f, u_{\mathcal{E}_f}) du_{\mathcal{E}_f}, \quad (22.2.8)$$

where it is now implicitly required that  $\mathcal{E}_f = \mathcal{E}_i + \hbar\omega$ . Equation (22.2.8) also applies as an approximation if the relevant final states are discrete but very closely spaced. If some of the degeneracy labels are discrete rather than continuous, then one should remove  $du_{\mathcal{E}_f}$  and sum over them rather than integrating.

Now suppose instead that the final states selected by energy conservation have discrete  $H_0$  eigenvalues  $\mathcal{E}_f$ . We will discuss this situation in three different ways, appropriate for different circumstances.

First, suppose that the electromagnetic fields are plane waves with fixed propagation direction  $\hat{k}$  and polarization  $\hat{\varepsilon}$ , but with incoherent phases and a range of angular frequencies wide enough to cover the final state in question. Then, as discussed in section 21.4, the incoherent phases mean that we can sum the probabilities (not the amplitudes) to obtain a transition rate. From using eq. (22.2.1) in eq. (21.4.21),

$$R_{i+\hbar\omega \rightarrow f}^{\text{abs}} = \int_0^\infty d\omega \rho(\omega) \frac{2\pi}{\hbar} \delta(\mathcal{E}_f - \mathcal{E}_i - \hbar\omega) \left( \frac{eE_0}{2m_e\omega} \right)^2 |\langle f | \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} | i \rangle|^2, \quad (22.2.9)$$

where  $\rho(\omega)$  is the density per unit angular frequency of incoherent electromagnetic wave perturbations of the form eq. (21.4.1). Let  $I$  be the total intensity (power per unit area) in the electromagnetic radiation, so that

$$dI = \frac{cE_0^2}{8\pi} \rho(\omega) d\omega. \quad (22.2.10)$$

We thus obtain, in terms of the intensity per unit angular frequency,  $dI/d\omega$ ,

$$R_{i+\hbar\omega \rightarrow f}^{\text{abs}} = \int_0^\infty d\omega \frac{dI}{d\omega} \delta(\mathcal{E}_f - \mathcal{E}_i - \hbar\omega) \frac{4\pi^2\alpha}{m_e^2\omega^2} |\langle f | \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} | i \rangle|^2. \quad (22.2.11)$$

Doing the integration over  $\omega$ , we find<sup>†</sup> an absorption rate

$$R_{i+\hbar\omega \rightarrow f}^{\text{abs}} = \frac{4\pi^2\alpha}{m_e^2\hbar\omega^2} \frac{dI}{d\omega} |\langle f | \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} | i \rangle|^2, \quad (22.2.12)$$

where  $\omega = (\mathcal{E}_f - \mathcal{E}_i)/\hbar$  has now been implicitly fixed.

As a second scenario, suppose that the system is bathed in electromagnetic radiation that is not only phase-incoherent and broad-band in  $\omega$ , but may be coming from all directions and with all possible polarization vectors. Although we have been discussing the electromagnetic field as a classical wave, in this case we choose to frame our discussion in terms of the numbers of photon quanta with energy  $\hbar\omega$  and particular wavenumbers and polarizations. To do so, we take the system to be in a large cubic box of side  $L$ , with periodic boundary conditions giving allowed wavevectors

$$\vec{k} = \hat{k}\omega/c = \frac{2\pi}{L}(\hat{x}n_x + \hat{y}n_y + \hat{z}n_z) \quad (22.2.13)$$

where  $(n_x, n_y, n_z)$  are integers. In the large-volume continuum limit, a discrete sum over these wavevectors will correspond to the integration

$$\sum_{\vec{k}} \rightarrow \left( \frac{L}{2\pi} \right)^3 \int d^3\vec{k} = \left( \frac{L}{2\pi c} \right)^3 \int_0^\infty d\omega \omega^2 \int d\Omega_{\vec{k}}. \quad (22.2.14)$$

---

<sup>†</sup>This could also be obtained directly from eq. (21.4.23).

Denote by  $N_{\vec{k},\hat{\varepsilon}}$  the occupation number of photons in a mode with wavevector  $\vec{k}$  and polarization vector  $\hat{\varepsilon}$ . Then the energy associated with that mode is

$$\hbar\omega N_{\vec{k},\hat{\varepsilon}} = (\text{volume}) \left( \frac{\text{energy}}{\text{volume}} \right) = L^3 (E_0^{\vec{k},\hat{\varepsilon}})^2 / 8\pi, \quad (22.2.15)$$

where  $E_0^{\vec{k},\hat{\varepsilon}}$  is the corresponding classical electric field amplitude.

Going back to eq. (21.4.5), and using the assumption of incoherence to sum the probabilities rather than amplitudes, the total probability for absorption is

$$\mathcal{P}_{i+\hbar\omega \rightarrow f}^{\text{abs}} = \sum_{\vec{k}} \sum_{\hat{\varepsilon}} \left( \frac{eE_0^{\vec{k},\hat{\varepsilon}}}{2m_e\omega} \right)^2 |\langle f | \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} | i \rangle|^2 F(\mathcal{E}_f - \mathcal{E}_i - \hbar\omega, t). \quad (22.2.16)$$

Now, using eqs. (22.2.14) and (22.2.15), this becomes

$$\mathcal{P}_{i+\hbar\omega \rightarrow f}^{\text{abs}} = \int_0^\infty d\omega \int d\Omega_{\vec{k}} \sum_{\hat{\varepsilon}} \frac{e^2 \hbar \omega}{4\pi^2 m_e^2 c^3} N_{\vec{k},\hat{\varepsilon}} |\langle f | \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} | i \rangle|^2 F(\mathcal{E}_f - \mathcal{E}_i - \hbar\omega, t). \quad (22.2.17)$$

Note that the polarizations  $\hat{\varepsilon}$  that are being summed over depend on the direction of  $\vec{k}$ , because of the transversality constraint  $\hat{\varepsilon} \cdot \vec{k} = 0$  for electromagnetic waves. As usual, we now apply  $F(\mathcal{E}_f - \mathcal{E}_i - \hbar\omega, t) = (2\pi t/\hbar) \delta(\mathcal{E}_f - \mathcal{E}_i - \hbar\omega)$ , and do the integral over  $\omega$  to obtain  $\mathcal{P}_{i+\hbar\omega \rightarrow f}^{\text{abs}} = t R_{i+\hbar\omega \rightarrow f}^{\text{abs}}$ , where the absorption rate is

$$R_{i+\hbar\omega \rightarrow f}^{\text{abs}} = \frac{\alpha\omega}{2\pi m_e^2 c^2} \int d\Omega_{\vec{k}} \sum_{\hat{\varepsilon}} N_{\vec{k},\hat{\varepsilon}} |\langle f | \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} | i \rangle|^2, \quad (22.2.18)$$

in terms of the photon occupation numbers  $N_{\vec{k},\hat{\varepsilon}}$  for modes with  $\omega = c|\vec{k}| = (\mathcal{E}_f - \mathcal{E}_i)/\hbar$ . As one might expect, the contribution to the rate from each mode is proportional to the number of photons available in that mode. Comparing eq. (22.2.12) to eq. (22.2.18), we see that

$$\frac{dI}{d\omega} \rightarrow \frac{\hbar\omega^3}{8\pi^3 c^2} \int d\Omega_{\vec{k}} \sum_{\hat{\varepsilon}} N_{\vec{k},\hat{\varepsilon}} \quad (22.2.19)$$

is the effective intensity per unit angular frequency in the bath of photons.

The third way of treating the absorption to a discrete final state is to compute a cross-section, as we have already done for the case of absorption by a continuous group of final states, in eq. (22.2.8). To do so, it is necessary to realize that there will always be at least a small effective broadening of the discrete energy level of the final-state, for two reasons. First,  $|f\rangle$  must be able to decay, at least back to the lower-energy initial state  $|i\rangle$  if not others, leading to a finite lifetime. The second reason, which is often more important in practice, is that there are environmental effects on the  $H_0$  system from perturbations due to nearby atoms with a thermal distribution of velocities, and from fluctuations in the electromagnetic fields.

Both types of line-broadening effects imply that the energy level of the final state is not perfectly sharp, but is described by a Breit–Wigner resonance, with a small finite width that may need to be obtained empirically. Therefore, when using eq. (22.2.7), the density of final states, which naively would be a delta function in the case of discrete  $|f\rangle$ , should be

$$\rho(\mathcal{E}_f, u_{\mathcal{E}_f}) = f_{\text{BW}}(\mathcal{E}_f - \mathcal{E}_{f0}, \hbar\gamma) \equiv \frac{\hbar\gamma}{2\pi} \frac{1}{(\mathcal{E}_f - \mathcal{E}_{f0})^2 + \hbar^2\gamma^2/4}. \quad (22.2.20)$$

Here  $\mathcal{E}_{f0}$  is the nominal discrete  $H_0$  eigenvalue before the line-broadening is taken into account, and  $\hbar\gamma$  is the full width at half maximum (FWHM) of the **Breit–Wigner line-shape** (named after Gregory Breit and Wigner, and also known as the **Lorentzian line-shape**) distribution for the energy  $\mathcal{E}_f$ , which is now continuous. Note that the normalization is

$$\int_{-\infty}^{\infty} d\mathcal{E}_f f_{\text{BW}}(\mathcal{E}_f - \mathcal{E}_{f0}, \hbar\gamma) = 1, \quad (22.2.21)$$

so that in the narrow-width approximation,

$$\lim_{\gamma \rightarrow 0} f_{\text{BW}}(\mathcal{E}_f - \mathcal{E}_{f0}, \hbar\gamma) = \delta(\mathcal{E}_f - \mathcal{E}_{f0}), \quad (22.2.22)$$

as expected. The absorption cross-section for a nominally discrete final state  $|f\rangle$  becomes, after using eq. (22.2.20) in eq. (22.2.7) and integrating over  $d\mathcal{E}_f$ ,

$$\sigma_{i \rightarrow f}^{\text{abs}} = \frac{2\pi\alpha\hbar^2\gamma}{m_e^2\omega} \frac{|\langle f | \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} | i \rangle|^2}{(\mathcal{E}_{f0} - \mathcal{E}_i - \hbar\omega)^2 + \hbar^2\gamma^2/4}. \quad (22.2.23)$$

Here we have removed the  $du_{\mathcal{E}_f}$  under the assumption that the degeneracy labels for the final state, such as angular momentum quantum numbers, are also discrete and can be summed over. This cross-section is a strongly peaked function of  $\omega$ , becoming sharper in the limit  $\gamma \rightarrow 0$ .

## 22.3 Induced and spontaneous emission of light

We now turn to the case of transitions between electronic states involving emission of energy in the form of electromagnetic radiation. The presence of a classical electromagnetic wave with angular frequency  $\omega$  will cause an initial  $H_0$  eigenstate  $|i\rangle$  to fall to a state  $|f\rangle$  of lower energy, provided that the  $H_0$  eigenvalues satisfy  $\mathcal{E}_i - \mathcal{E}_f = \hbar\omega$ . If  $dI/d\omega$  is the intensity of incoherent electromagnetic waves per unit  $\omega$ , then we can apply exactly the same reasoning we used to arrive at eq. (22.2.12), except that we use eq. (21.4.22) for emission rather than eq. (21.4.21) for absorption. The result is the rate for **induced emission** (or **stimulated emission**),

$$R_{i \rightarrow f + \hbar\omega}^{\text{ind. em.}} = \frac{4\pi^2\alpha}{m_e^2\hbar\omega^2} \frac{dI}{d\omega} |\langle f | \hat{\varepsilon}^* \cdot \vec{\mathcal{P}}_{\vec{k}}^\dagger | i \rangle|^2, \quad (22.3.1)$$



where now  $\omega = (\mathcal{E}_i - \mathcal{E}_f)/\hbar$  is implicit. This assumes that the wavevector direction  $\hat{k}$  is fixed. For the case of light incoming from all directions, the transition rate can be expressed in terms of the photon occupation numbers using eq. (22.2.19), again just as we did for the absorption case in arriving at eq. (22.2.18),

$$R_{i \rightarrow f + \hbar\omega}^{\text{ind. em.}} = \frac{\alpha\omega}{2\pi m_e^2 c^2} \int d\Omega_{\vec{k}} \sum_{\vec{\varepsilon}} N_{\vec{k}, \vec{\varepsilon}} |\langle f | \hat{\varepsilon}^* \cdot \vec{\mathcal{P}}_{\vec{k}}^\dagger | i \rangle|^2. \quad (22.3.2)$$

The reason for referring to this as “induced” is that it is proportional to the intensity of incident light, or equivalently to the numbers of photons already present.

Despite the fact that our calculations so far do not show it, the transition from  $|i\rangle$  to  $|f\rangle$  with the emission of a photon can also occur even if there are no incident classical waves or photons present. That process is called **spontaneous emission**, and we will now show that it yields a result equal to eq. (22.3.2) but with the factor  $N_{\vec{k}, \vec{\varepsilon}}$  replaced by 1. The phenomenon of spontaneous emission, in which an electronic state decays to a lower energy state by emitting a photon, is the quantum mechanical analog of the classical Larmor radiation of an accelerating charge. However, it can only occur if there is a quantum state with lower energy, while classical Larmor radiation would always occur; compare to the discussion in section 1.1.

To obtain the rate for spontaneous emission in the most direct way, the electromagnetic fields should be treated quantum mechanically, rather than classically as we have done. Recall that our method so far was to treat the potentials  $\Phi$  and  $\vec{A}$  and their derivatives  $\vec{E}$  and  $\vec{B}$  as classical quantities, which are then evaluated in terms of the position and momentum operators for the electron that witnesses them. In a more logical and complete approach, both the electronic and radiation degrees of freedom should be treated as a unified quantum system. Then  $\Phi$  and  $\vec{A}$  will be operators in their own right, rather than just functions of the position operator, and will act on a Hilbert space that describes not just the electronic state, but the photons as well. This allows a direct computation of spontaneous emission (and more complicated phenomena). In this section, instead of pursuing that direct method, we will derive the results for spontaneous emission using an indirect argument, based on inference from requiring the consistency of statistical mechanics, called the **Einstein A and B coefficients** method.

Consider the thought experiment of a cavity kept in thermal equilibrium at temperature  $T$  with the electromagnetic radiation inside of it. The walls of the cavity are made of a very large number of atoms that include two states  $|a\rangle$  and  $|b\rangle$  with energy difference

$$\mathcal{E}_b - \mathcal{E}_a = \hbar\omega, \quad (22.3.3)$$

so that the photon absorption process  $a + \hbar\omega \rightarrow b$  and the emission process  $b \rightarrow a + \hbar\omega$  are both constantly occurring to ensure the equilibrium. According to statistical mechanics, and

our discussion in section 3.5, the relative probability for a state with energy  $E$  is proportional to the Boltzmann factor  $e^{-E/k_B T}$ , where  $k_B$  is Boltzmann's constant. Therefore, since  $\hbar\omega N_{\vec{k},\hat{\varepsilon}}$  is the energy of a state with  $N_{\vec{k},\hat{\varepsilon}}$  photons in the mode with wavevector  $\vec{k}$  (with  $k = \omega/c$ ) and polarization  $\hat{\varepsilon}$ , the average number of photons in that mode is

$$\overline{N}_{\vec{k},\hat{\varepsilon}} = \left( \sum_{n=0}^{\infty} n e^{-n\hbar\omega/k_B T} \right) / \sum_{n=0}^{\infty} e^{-n\hbar\omega/k_B T} = \frac{1}{e^{\hbar\omega/k_B T} - 1}. \quad (22.3.4)$$

In equilibrium, each  $N_{\vec{k},\hat{\varepsilon}}$ , which we will refer to as  $N$  for short in the following, is constant and equal to the mean number given by eq. (22.3.4). Therefore, we must have

$$\left( \frac{dN}{dt} \right)_{\text{spon. em.}} + \left( \frac{dN}{dt} \right)_{\text{ind. em.}} + \left( \frac{dN}{dt} \right)_{\text{absorption}} = 0, \quad (22.3.5)$$

with contributions from spontaneous emission, induced emission, and absorption rates that are respectively of the forms

$$\left( \frac{dN}{dt} \right)_{\text{spon. em.}} = n_b A, \quad (22.3.6)$$

$$\left( \frac{dN}{dt} \right)_{\text{ind. em.}} = N n_b B_{ba}, \quad (22.3.7)$$

$$\left( \frac{dN}{dt} \right)_{\text{absorption}} = -N n_a B_{ab}. \quad (22.3.8)$$

Here, we have used the fact that the spontaneous and induced emission contributions to the rate for creation of photons must both be proportional to the number of atoms  $n_b$  in the state  $|b\rangle$ , while the absorption rate contribution to the rate for subtraction of photons is proportional to the number of atoms  $n_a$  in the state  $|a\rangle$ . Also, the induced emission and absorption rates are both proportional to the number of photons already present  $N$ , but by definition the spontaneous emission rate has no such factor. Our goal is to determine the ratios of the coefficients<sup>†</sup> of proportionality,  $A$ ,  $B_{ba}$ , and  $B_{ab}$ . These are in turn proportional to the transition rates  $R_{b \rightarrow a + \hbar\omega}^{\text{spon. em.}}$ ,  $R_{b \rightarrow a + \hbar\omega}^{\text{ind. em.}}$ , and  $R_{a + \hbar\omega \rightarrow b}^{\text{abs}}$ , respectively, but with the corresponding factors of  $N$  extracted from the last two, since we have put them explicitly in eqs. (22.3.7) and (22.3.8).

The numbers  $n_a$  and  $n_b$  are unknown, but since the system is in thermal equilibrium, we know their ratio, as they are each proportional to the corresponding Boltzmann factor. So,

$$n_a/n_b = e^{-(\mathcal{E}_a - \mathcal{E}_b)/k_B T} = e^{\hbar\omega/k_B T}. \quad (22.3.9)$$

Using eqs. (22.3.6)–(22.3.9) in eq. (22.3.5), we can solve for the photon occupation number,

$$N = \frac{A}{B_{ab} e^{\hbar\omega/k_B T} - B_{ba}}. \quad (22.3.10)$$

---

<sup>†</sup>Einstein's original coefficients were normalized differently.

Now the key point is that  $A$ ,  $B_{ab}$ , and  $B_{ba}$  should depend only on the properties of the states  $|a\rangle$  and  $|b\rangle$ , but not on the temperature. Therefore, the only way to reconcile the functional dependences on  $T$  of eqs. (22.3.4) and (22.3.10) is

$$A = B_{ab} = B_{ba}, \quad (22.3.11)$$

and, reassuringly, this indeed does give eq. (22.3.4).

The equality  $B_{ab} = B_{ba}$  is just a check of the detailed balance equality between  $R_{a+\hbar\omega\rightarrow b}^{\text{abs}}$  and  $R_{b\rightarrow a+\hbar\omega}^{\text{ind. em.}}$ , which we had already found by direct computation in eqs. (22.2.18) and (22.3.2). The new requirement that  $A = B_{ba}$  says that, because  $N$  was already factored out of  $B_{ba}$ , the rate we calculated for induced emission alone is related to the total emission rate by

$$\text{induced} \rightarrow \text{induced} + \text{spontaneous} \quad (22.3.12)$$

$$N_{\vec{k},\hat{\varepsilon}} \rightarrow N_{\vec{k},\hat{\varepsilon}} + 1. \quad (22.3.13)$$

Thus, the rate for spontaneous emission is obtained by simply removing the factor of  $N_{\vec{k},\hat{\varepsilon}}$  from the formula eq. (22.3.2) for induced emission,

$$R_{i\rightarrow f+\hbar\omega}^{\text{spont. em.}} = \frac{\alpha\omega}{2\pi m_e^2 c^2} \int d\Omega_{\vec{k}} \sum_{\hat{\varepsilon}} |\langle f | \hat{\varepsilon}^* \cdot \vec{\mathcal{P}}_{\vec{k}}^\dagger | i \rangle|^2. \quad (22.3.14)$$

A more direct calculation that treats the electromagnetic fields quantum mechanically gives the same result.

## 22.4 Electric dipole approximation

The case in which the wavelength of the electromagnetic wave is much larger than the spatial extent of the initial-state wavefunction is called the **electric dipole approximation**. To see why this might apply as a valid approximation, suppose that the electron is in a hydrogen-like atomic orbital state with principal quantum number  $n$  and atomic number  $Z$ . Then the minimum photon energy needed to ionize the state obeys  $\hbar\omega = Z^2 e^2 / 2a_0 n^2$  from eq. (11.1.16), while the characteristic spatial extent of the initial wavefunction is  $\langle R \rangle = 3n^2 a_0 / 2Z$ , using eq. (11.1.65) with  $l = 0$ . For larger  $l$  in the initial state, the characteristic size  $\langle R \rangle$  is even smaller (for fixed  $n$ ), according to eq. (11.1.65). To compare the atomic size scale to the inverse wavenumber of the light (with minimum ionizing energy),

$$k\langle R \rangle = \frac{\omega\langle R \rangle}{c} \sim \frac{3\alpha Z}{4}. \quad (22.4.1)$$

This is smaller than 1, even when  $Z$  is large, since  $\alpha = e^2/\hbar c \approx 1/137$ . For transitions between bound states, the relevant wavelength is even longer because the change in energy is smaller.

As an extreme case, the wavelength is macroscopic for transitions between states with the same principal quantum number, which have very nearly the same energy. This means that in transitions between bound states we can make the long-wavelength limit approximation

$$e^{i\vec{k}\cdot\vec{R}} \approx 1 \quad (22.4.2)$$

over the relevant spatial extent of the atom. In the same limit, one can also neglect the  $\vec{k} \times \vec{S}$  contribution compared to the momentum operator  $\vec{P}$  in  $\vec{\mathcal{P}}_{\vec{k}}$  of eq. (22.1.13). This is made plausible by the rough estimate

$$\frac{|\vec{k} \times \vec{S}|}{|\vec{P}|} \sim \frac{(\omega/c)\hbar}{\hbar/\langle R \rangle} \quad (22.4.3)$$

(using the uncertainty relation for position and momentum in the denominator), which is parametrically the same as eq. (22.4.1), and so again suppressed by a factor  $\alpha Z$ .

Therefore, the electric dipole (long-wavelength) approximation is defined by taking

$$\langle f | \vec{\mathcal{P}}_{\vec{k}} | i \rangle \approx \langle f | \vec{P} | i \rangle \quad (22.4.4)$$

in the absorption and emission rates. Now we can use a trick, by noticing that if the only momentum dependence of  $H_0$  is in the kinetic term, then

$$[H_0, \vec{R}] = \frac{1}{2m_e} [P^2, \vec{R}] = -i\frac{\hbar}{m_e} \vec{P}, \quad (22.4.5)$$

so that in the case of absorption,

$$\langle f | \hat{\varepsilon} \cdot \vec{P} | i \rangle = \frac{im_e}{\hbar} \hat{\varepsilon} \cdot \langle f | [H_0, \vec{R}] | i \rangle = \frac{im_e}{\hbar} (\mathcal{E}_f - \mathcal{E}_i) \langle f | \hat{\varepsilon} \cdot \vec{R} | i \rangle = im_e \omega \langle f | \hat{\varepsilon} \cdot \vec{R} | i \rangle. \quad (22.4.6)$$

For the emission case, the only difference is that  $\hat{\varepsilon} \rightarrow \hat{\varepsilon}^*$  and  $\omega \rightarrow -\omega$ , so

$$\langle f | \hat{\varepsilon}^* \cdot \vec{P} | i \rangle = -im_e \omega \langle f | \hat{\varepsilon}^* \cdot \vec{R} | i \rangle. \quad (22.4.7)$$

The sign change makes no difference, and the complex conjugation of the polarization vector has no effect for linear polarizations, but makes a difference for circular polarization vectors [such as  $\hat{\varepsilon} = (\hat{x} \pm i\hat{y})/\sqrt{2}$  if  $\hat{k} = \hat{z}$ ].

For example, using eq. (22.4.6) in eq. (22.2.8), we have for the absorption cross-section in the case of continuous final state energies

$$d\sigma_{i \rightarrow f}^{\text{abs, dipole}} = 4\pi^2 \alpha \hbar \omega |\langle f | \hat{\varepsilon} \cdot \vec{R} | i \rangle|^2 \rho(\mathcal{E}_f, u_{\mathcal{E}_f}) du_{\mathcal{E}_f}. \quad (22.4.8)$$

Recalling that  $\hat{\varepsilon}$  is the direction of the polarization of the absorbed electromagnetic wave, we see that the relevant matrix element in the electric dipole approximation is proportional to the position operator along the polarization direction. More generally, the electric dipole approximation amounts to making the substitutions

$$|\langle f | \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} | i \rangle|^2 \rightarrow m_e^2 \omega^2 |\langle f | \hat{\varepsilon} \cdot \vec{R} | i \rangle|^2 \quad (\text{electric dipole absorption}) \quad (22.4.9)$$

in the absorption rates for discrete final states with incoherent plane-wave light in eq. (22.2.12), incoherent light from all directions in (22.2.18), and in the cross-section for a line-broadened discrete final state in eq. (22.2.23). Similarly, one makes the replacement

$$|\langle f | \hat{\varepsilon}^* \cdot \vec{\mathcal{P}}_k^\dagger | i \rangle|^2 \rightarrow m_e^2 \omega^2 |\langle f | \hat{\varepsilon}^* \cdot \vec{R} | i \rangle|^2 \quad (\text{electric dipole emission}) \quad (22.4.10)$$

in the induced emission rates eq. (22.3.1) for plane waves with a fixed direction of propagation and eq. (22.3.2) for light from all directions in terms of the photon occupation numbers, and in eq. (22.3.14) for spontaneous emission.

In the case of spontaneous emission in eq. (22.3.14), the matrix element  $\langle f | \hat{\varepsilon}^* \cdot \vec{R} | i \rangle$  in the electric dipole approximation does not depend directly on the wavevector magnitude, but does depend indirectly on the wavevector direction because of the requirement  $\hat{\varepsilon}^* \cdot \hat{k} = 0$ . Temporarily fixing  $\hat{k} = \hat{z}$ , the two independent polarization directions to be summed over can be taken along  $\hat{x}$  and  $\hat{y}$ , so that

$$\sum_{\hat{\varepsilon}} |\langle f | \hat{\varepsilon}^* \cdot \vec{R} | i \rangle|^2 = |\langle f | X | i \rangle|^2 + |\langle f | Y | i \rangle|^2. \quad (22.4.11)$$

Then, integrating over all possible directions for  $\hat{k}$  will effectively give an average of the two terms over three possible directions, so

$$\int d\Omega_{\vec{k}} \sum_{\hat{\varepsilon}} |\langle f | \hat{\varepsilon}^* \cdot \vec{R} | i \rangle|^2 = (4\pi) \frac{2}{3} \left( |\langle f | X | i \rangle|^2 + |\langle f | Y | i \rangle|^2 + |\langle f | Z | i \rangle|^2 \right) \quad (22.4.12)$$

$$= \frac{8\pi}{3} \langle f | \vec{R} | i \rangle \cdot \langle i | \vec{R} | f \rangle. \quad (22.4.13)$$

This reduces eq. (22.3.14) to the simple formula

$$R_{i \rightarrow f}^{\text{spon. em.}} = \frac{4\alpha\omega^3}{3c^2} |\langle f | \vec{R} | i \rangle|^2 \quad (22.4.14)$$

for the rate for an initial state  $|i\rangle$  to decay to a final state  $|f\rangle$  by the spontaneous emission of a photon. Note that the right side involves

$$|\langle f | \vec{R} | i \rangle|^2 \equiv \langle f | \vec{R} | i \rangle \cdot \langle i | \vec{R} | f \rangle, \quad (22.4.15)$$

the square of the magnitude of a complex vector matrix element.

Suppose that there is no background electromagnetic radiation to induce emission or absorption, and let the number of atoms in the state  $|i\rangle$  be  $N_i(t)$  at time  $t$ . Then at time  $t + \Delta t$ ,

$$N_i(t + \Delta t) = N_i(t) - \Delta t R_i N_i(t), \quad (22.4.16)$$

where the total decay rate of the initial state is

$$R_i = \sum_f R_{i \rightarrow f}^{\text{spon. em.}}. \quad (22.4.17)$$

Then

$$\frac{N_i(t + \Delta t) - N_i(t)}{\Delta t} = -R_i N_i(t), \quad (22.4.18)$$

which in the limit  $\Delta t \rightarrow 0$  becomes the differential equation

$$\frac{dN_i}{dt} = -R_i N_i, \quad (22.4.19)$$

with the solution

$$N_i(t) = N_i(t_0) e^{-(t-t_0)R_i}. \quad (22.4.20)$$

Therefore  $\tau_i = 1/R_i$  is the lifetime of the state  $|i\rangle$ , defined as the amount of time needed for the number of atoms in that state to decrease by a factor of  $1/e \approx 0.367879$  in the absence of other perturbations. The half-life, defined as the amount of time required for the number to decrease by half, is related to this by  $t_{1/2} = \tau_i \ln 2 = 0.693147/R_i$ .

The preceding results imply matrix element selection rules that govern both absorption and emission in the electric dipole approximation when  $|i\rangle$  and  $|f\rangle$  are angular momentum eigenstates. These follow immediately from eqs. (13.2.15)–(13.2.18). First, eq. (13.2.15) tells us that the change in the orbital angular momentum quantum number  $l$  must be exactly 1 unit,  $\Delta l = l_f - l_i = \pm 1$ . As a corollary, the parities of the initial and final states must be opposite, since the parity of a state with orbital angular momentum  $l$  is  $(-1)^l$  from eq. (8.7.5); this can also be seen directly from the parity selection rule, since the operator  $\vec{R}$  has odd parity. Also, eqs. (13.2.16)–(13.2.18) tell us that the  $z$ -component of the orbital angular momentum changes by 1 unit or less:  $\Delta m_l = m_{l,f} - m_{l,i} = 0, \pm 1$ . As special cases, if the polarization of the wave is along  $\hat{\varepsilon} = \hat{z}$ , then eq. (13.2.16) says  $\Delta m_l = 0$ , and if it is in the  $x, y$  plane, then eqs. (13.2.17)–(13.2.18) demand  $\Delta m_l = \pm 1$ . The spin is not affected by the position operator, so that  $m_{s,f} = m_{s,i}$ .

We therefore have the **electric dipole selection rules** for single-electron transitions,

$$\Delta l = \pm 1, \quad \Delta m_l = 0, \pm 1 \quad \Delta m_s = 0, \quad (\text{electric dipole}). \quad (22.4.21)$$

These can be generalized to the case of multi-electron transitions (where more than one electron changes orbitals in going from the initial to the final state), for which the electric dipole selection rules are

$$\pi_i \pi_f = -1, \quad \Delta J = 0, \pm 1, \quad \Delta m_J = 0, \pm 1, \quad J_i + J_f > 0, \quad (\text{electric dipole}). \quad (22.4.22)$$

The first three of these follow from requiring the non-vanishing of the matrix element of the parity-odd electric dipole operator  $\hat{\varepsilon} \cdot \vec{R}$ . The  $\Delta J$  rule follows from the fact that  $\vec{R}$  is a vector

operator, and therefore carries total angular momentum 1 in the sense explained in Chapter 13. In terms of addition of angular momentum,  $J \otimes 1 = (J-1) \oplus J \oplus (J+1)$  for  $J \geq 1$ .

The last selection rule in eq. (22.4.22) is completely general, even if we do not make the electric dipole approximation. It says just this: a total angular momentum  $J = 0$  state can never go to another  $J = 0$  state by absorbing or emitting a single photon. The reason is that a single photon can always be written as a linear combination of transverse circular polarizations, carrying intrinsic angular momentum component along the propagation direction with eigenvalue either  $+\hbar$  or  $-\hbar$ , which cannot be matched by the initial and final electronic states if they both have no angular momentum. Therefore, we have a general rule

$$J_i + J_f > 0, \quad (\text{all single-photon transitions}). \quad (22.4.23)$$

However, this rule (and all other selection rules listed here) can be violated if one considers emission or absorption of multiple photons, either as distinct sequential events involving intermediate states, or by going beyond first order in time-dependent perturbation theory or through perturbations caused by the environment of the atom. Multi-photon transition rates at higher order in perturbation theory are suppressed by additional powers of the fine-structure constant, however, and so can often be neglected when they compete with single-photon transitions.

In the approximation of the  $LS$  coupling scheme, the total spin, total orbital, and total angular momentum orthobasis quantum numbers  $L, S, J, m_J$  are good quantum numbers for the initial and final stationary states. Then, because the total spin is not involved at all in the electric dipole moment operator, one has the additional rules

$$\Delta S = 0, \quad \Delta L = 0, \pm 1, \quad L_i + L_f > 0, \quad (\text{electric dipole, } LS \text{ coupling}) \quad (22.4.24)$$

Recall, from the discussion in section 18.4, that the  $LS$  coupling scheme holds to a good approximation for atoms with atomic number  $Z$  not too large.

## 22.5 Magnetic dipole, electric quadrupole, and higher orders

Transitions that violate the electric dipole selection rules of eqs. (22.4.21) and (22.4.22) have rates that are typically suppressed by extra powers of  $Z\alpha$ , or by extra inverse powers of the wavelength of the photon emitted or absorbed. For this reason, they are often referred to as “forbidden”. Traditionally, this terminology does not mean that they are literally forbidden, but rather that they take place at highly suppressed rates. By the same terminology, transitions that take place through the electric dipole operator are often called “allowed”.

To go beyond the electric dipole approximation and its selection rules for absorption and emission of electromagnetic radiation, consider the expansion in small  $k$  of eq. (22.1.13). To

make things a little simpler, let us temporarily choose the wavevector of the light as  $\vec{k} = k\hat{z}$  and the polarization as  $\hat{\varepsilon} = \hat{x}$ , so

$$\hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} = e^{i\vec{k} \cdot \vec{R}} \hat{\varepsilon} \cdot (\vec{P} - i\vec{k} \times \vec{S}) = e^{ikZ} (P_x + ikS_y). \quad (22.5.1)$$

Now, expanding  $e^{ikZ} \approx 1 + ikZ + \dots$ , and using  $ZP_x = (ZP_x - XP_z)/2 + (ZP_x + XP_z)/2 = L_y/2 + (ZP_x + P_zX)/2$ , we have

$$\hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} = P_x + i\frac{\omega}{2c} [L_y + 2S_y + (ZP_x + P_zX)] + \dots. \quad (22.5.2)$$

Recasting this back into a form valid for general orthogonal  $\hat{k}$  and  $\hat{\varepsilon}$  gives

$$\hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} = \hat{\varepsilon} \cdot \vec{P} + i\frac{\omega}{2c} \left[ (\hat{k} \times \hat{\varepsilon}) \cdot (\vec{L} + 2\vec{S}) + (\hat{k} \cdot \vec{R} \hat{\varepsilon} \cdot \vec{P} + \hat{k} \cdot \vec{P} \hat{\varepsilon} \cdot \vec{R}) \right] + \dots, \quad (22.5.3)$$

which is the result to next-to-leading order in the long-wavelength expansion. In the last term, the ordering of operators  $\vec{R}$  and  $\vec{P}$  does not matter; they commute because of the transversality condition  $\hat{\varepsilon} \cdot \hat{k} = 0$

Therefore, for the harmonic perturbation to be used in eq. (21.4.1) and subsequent equations, we can write

$$V = V_{E1} + V_{M1} + V_{E2} + \dots, \quad (22.5.4)$$

where the separated contributions,

$$V_{E1} = -\frac{eE_0}{2m_e\omega} \hat{\varepsilon} \cdot \vec{P}, \quad (22.5.5)$$

$$V_{M1} = -i\frac{eE_0}{4m_e c} (\hat{k} \times \hat{\varepsilon}) \cdot (\vec{L} + 2\vec{S}), \quad (22.5.6)$$

$$V_{E2} = -i\frac{eE_0}{4m_e c} (\hat{k} \cdot \vec{R} \hat{\varepsilon} \cdot \vec{P} + \hat{k} \cdot \vec{P} \hat{\varepsilon} \cdot \vec{R}) \quad (22.5.7)$$

are called electric dipole (E1), magnetic dipole (M1), and electric quadrupole (E2), respectively. Note that in the magnetic dipole operator, the vector  $(\hat{k} \times \hat{\varepsilon})$  is the unit vector along the magnetic field direction in the wave; the magnetic field couples to the combination  $\vec{L} + 2\vec{S}$ .

Single-photon transitions that are dominantly magnetic dipole in character can now be evaluated from the general formulas in sections 22.2-22.3, by using

$$|\langle f | \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} | i \rangle|^2 \rightarrow \frac{\omega^2}{4c^2} |\langle f | (\hat{k} \times \hat{\varepsilon}) \cdot (\vec{L} + 2\vec{S}) | i \rangle|^2 \quad (\text{M1 absorption}), \quad (22.5.8)$$

$$|\langle f | \hat{\varepsilon}^* \cdot \vec{\mathcal{P}}_{\vec{k}}^\dagger | i \rangle|^2 \rightarrow \frac{\omega^2}{4c^2} |\langle f | (\hat{k} \times \hat{\varepsilon}^*) \cdot (\vec{L} + 2\vec{S}) | i \rangle|^2 \quad (\text{M1 emission}). \quad (22.5.9)$$

For the electric quadrupole term, we can do a trick similar to the one we used in the electric dipole case to get eqs. (22.4.9) and (22.4.10). Assuming that  $H_0$  does not depend on the momentum operator except through the kinetic term  $P^2/2m_e$ , we find from eq. (22.4.5) that

$$[H_0, \hat{k} \cdot \vec{R} \hat{\varepsilon} \cdot \vec{R}] = -i\frac{\hbar}{m_e} (\hat{k} \cdot \vec{R} \hat{\varepsilon} \cdot \vec{P} + \hat{k} \cdot \vec{P} \hat{\varepsilon} \cdot \vec{R}), \quad (22.5.10)$$



where we have made use of  $\hat{\varepsilon} \cdot \hat{k} = 0$ . Therefore,

$$V_{E2} = \frac{eE_0}{4\hbar c} [H_0, \hat{k} \cdot \vec{R} \hat{\varepsilon} \cdot \vec{R}], \quad (22.5.11)$$

and in the matrix elements, one can evaluate  $H_0$  on the adjacent ket  $|i\rangle$  or bra  $\langle f|$ , giving a factor of  $\mathcal{E}_f - \mathcal{E}_i = \hbar\omega$  for absorption and  $-\hbar\omega$  for emission. It follows that in matrix elements for single-photon absorption, one can make the replacement

$$V_{E2} \rightarrow \frac{eE_0\omega}{4c} \hat{k} \cdot \overleftrightarrow{\mathcal{Q}} \cdot \hat{\varepsilon}, \quad (22.5.12)$$

where<sup>†</sup> the symmetric, traceless rank-2 tensor quadrupole moment operator is

$$\mathcal{Q}_{ab} = R_a R_b - \delta_{ab} R^2/3, \quad (22.5.13)$$

for  $a, b = x, y, z$ . This uses the fact that the  $\delta_{ab}$  part of  $\mathcal{Q}_{ab}$  does not contribute to  $V_{E2}$  because of  $\hat{\varepsilon} \cdot \hat{k} = 0$ . Within matrix elements for single-photon emission,

$$V_{E2}^\dagger \rightarrow -\frac{eE_0\omega}{4c} \hat{k} \cdot \overleftrightarrow{\mathcal{Q}} \cdot \hat{\varepsilon}^* \quad (22.5.14)$$

similarly applies. Comparing eqs. (22.5.12) and (22.5.14) to eq. (22.2.1), we see that for dominantly electric quadrupole transitions, the general formulas in sections 22.2-22.3 apply with

$$|\langle f | \hat{\varepsilon} \cdot \vec{\mathcal{P}}_{\vec{k}} | i \rangle|^2 \rightarrow \frac{m_e^2 \omega^4}{4c^2} |\langle f | \hat{k} \cdot \overleftrightarrow{\mathcal{Q}} \cdot \hat{\varepsilon} | i \rangle|^2 \quad (\text{E2 absorption}), \quad (22.5.15)$$

$$|\langle f | \hat{\varepsilon}^* \cdot \vec{\mathcal{P}}_{\vec{k}}^\dagger | i \rangle|^2 \rightarrow \frac{m_e^2 \omega^4}{4c^2} |\langle f | \hat{k} \cdot \overleftrightarrow{\mathcal{Q}} \cdot \hat{\varepsilon}^* | i \rangle|^2 \quad (\text{E2 emission}), \quad (22.5.16)$$

for single-photon absorption and emission, respectively.

We now turn to the selection rules for M1 and E2 transitions. First, let us use eqs. (22.5.8) and (22.5.9) to find the requirements on the initial and final states in order for a magnetic dipole transition to occur. In transitions involving a single electron, neither  $\vec{L}$  nor  $\vec{S}$  changes the orbital quantum number  $l$ , and the spin quantum number  $s$  is of course fixed at  $1/2$ . However, the components of  $\vec{L}$  are linear combinations of  $L_z, L_\pm$ , and the components of  $\vec{S}$  are linear combinations of  $S_z, S_\pm$ , so they can change  $m_l$  and  $m_s$  by zero or one unit. Furthermore, magnetic dipole transitions cannot occur between states with different principal (radial) quantum numbers, because  $\vec{L}$  and  $\vec{S}$  do not operate on the radial wavefunctions and do not change  $l$ , so the matrix element calculation in the position representation will be proportional to

$$\int_0^\infty dr r^2 R_{nl}(r) R_{n'l}(r) = \delta_{nn'}. \quad (22.5.17)$$

---

<sup>†</sup>For two vectors  $\vec{v}$  and  $\vec{w}$  with components  $v_a$  and  $w_a$ , and a rank-2 tensor  $\overleftrightarrow{T}$  with components  $T_{ab}$ , the notation  $\vec{v} \cdot \overleftrightarrow{T} \cdot \vec{w}$  means  $\sum_{a=x,y,z} \sum_{b=x,y,z} v_a T_{ab} w_b$ . We have already encountered the quadrupole moment operator  $\mathcal{Q}_{ab}$  in eqs. (13.1.30)-(13.1.33).

Therefore, the **magnetic dipole selection rules** are, for single-electron transitions,

$$\Delta n = 0, \quad \Delta l = 0, \quad \Delta m_l = 0, \pm 1, \quad \Delta m_s = 0, \pm 1, \quad (\text{magnetic dipole}). \quad (22.5.18)$$

For multi-electron transitions, the selection rules from eqs. (22.5.8) and (22.5.9) are

$$\pi_i \pi_f = +1, \quad \Delta J = 0, \pm 1, \quad \Delta m_J = 0, \pm 1, \quad J_i + J_f > 0, \quad (\text{magnetic dipole}). \quad (22.5.19)$$

The parity selection rule follows from the fact that the operator  $\vec{L} + 2\vec{S}$  has even parity, and the  $\Delta J$  and  $\Delta m_J$  selection rules follow from the fact that it is a vector, and therefore as explained in Chapter 13 it carries total angular momentum 1. The rule  $J_i + J_f > 0$  is completely general, as we already noted in our discussion of the electric dipole case. In the  $LS$  coupling scheme approximation, where initial and final stationary states can be assigned total angular momentum basis quantum numbers  $L, S, J, m_J$ , there are additional rules

$$\Delta L = 0, \quad \Delta S = 0 \quad (\text{magnetic dipole, } LS \text{ coupling}). \quad (22.5.20)$$

Note that the electric dipole and magnetic dipole contributions cannot both be relevant for a given initial and final state. Magnetic dipole transitions never change the parity, while electric dipole transitions always do.

The famous 21 centimeter hyperfine transition line within the ground state of hydrogen is a magnetic dipole transition, as it involved a spin flip but does not change the orbital angular momentum or principal quantum number, which are  $l = 0$  and  $n = 1$  in both the initial and final state. Since the resulting photon has an extremely long wavelength compared to the atomic size, the hyperfine transition occurs at a highly suppressed rate (compared to what one might expect if it were an electric dipole transition), because  $k = \omega/c \ll 1/a_0$ .

Now consider the matrix element selection rules that apply to the electric quadrupole term. Depending on the choices for  $\hat{k}$  and  $\hat{\varepsilon}$ , the matrix elements in eqs. (22.5.15) and (22.5.16) can involve linear combinations of the operators  $XY, XZ, YZ, X^2 - Y^2$ , and  $X^2 + Y^2 - 2Z^2$ , which are the components of the quadrupole tensor  $\vec{\vec{Q}}$ . These operators all have parity  $\pi_Q = +1$ , so electric quadrupole transitions can only connect states with the same parity. Acting on single-electron orbital angular momentum eigenstates, these operators can change  $l$  by 0 or 2 units (but not 1 unit, as that would not satisfy the parity selection rule), and  $m_l$  by 0, 1, or 2 units. Unlike the magnetic dipole interaction, the electric quadrupole operator does act non-trivially on the radial wavefunction, so there is no restriction on changes in the principal quantum number. However, it does not change the spin state. So, we have the **electric quadrupole selection rules** for single-electron transitions,

$$\Delta l = 0, \pm 2, \quad \Delta m_l = 0, \pm 1, \pm 2, \quad \Delta m_s = 0, \quad (\text{electric quadrupole}). \quad (22.5.21)$$

For transitions involving multiple electrons, the selection rules are

$$\pi_i \pi_f = +1, \quad |\Delta J| \leq 2, \quad |\Delta m_J| \leq 2, \quad J_i + J_f > 1, \quad (\text{electric quadrupole}), \quad (22.5.22)$$

and in the  $LS$  coupling scheme approximation in which the initial and final multi-electron stationary states have good quantum numbers  $L, S, J, m_J$ , there are additional selection rules

$$\Delta S = 0, \quad |\Delta L| \leq 2, \quad L_i + L_f > 1 \quad (\text{electric quadrupole, } LS \text{ coupling}). \quad (22.5.23)$$

The general constraint that angular momentum can change by at most two units in electric quadrupole transitions follows from the Wigner–Eckart Theorem 13.3.1 and the fact that the quadrupole operator is an irreducible tensor of order  $j = 2$ , as we noted in section 13.1. The electric quadrupole transitions can only occur between states that have the same parity, so they can never occur if electric dipole transitions are allowed. If the initial and final states in question have  $\Delta l = 0$  or  $\Delta L = 0$ , then the transition can be both magnetic dipole and electric quadrupole in character, but if  $\Delta L = 2$  or  $\Delta J = 2$  or it changes  $m_l$  or  $m_J$  by two units, then it will be electric quadrupole at leading order in the long-wavelength approximation.

If one continues the expansion in  $\vec{k} \cdot \vec{R}$  of  $\vec{\mathcal{P}}_{\vec{k}}$ , the resulting electric and magnetic multipole contributions are called  $Eq$  and  $Mq$  with  $q = 1, 2, 3, \dots$ , where M2 is magnetic quadrupole, E3 is electric octopole, etc. The general selection rules for these contributions are

$$\pi_i \pi_f = (-1)^q, \quad (Eq), \quad (22.5.24)$$

$$\pi_i \pi_f = (-1)^{q-1}, \quad (Mq), \quad (22.5.25)$$

$$|\Delta J| \leq q \leq J_i + J_f, \quad |\Delta m_J| \leq q, \quad (Eq) \text{ and } (Mq). \quad (22.5.26)$$

In the long wavelength limit, the leading  $Eq$  or  $Mq$  contributions for a transition between two states are those with the smallest  $q$  that satisfies these selection rules. For a possible transition between an odd-parity state with  $J = 2$  and an even-parity state with  $J = 0$ , the single-photon transition would be uniquely determined to be M2, magnetic quadrupole. For two states with the same parity and both with  $J = 1$ , the transition could be either M1 or E2. For two states both with  $J = 0$ , there is no allowed single-photon transition as we have already discussed, and there will instead be a highly suppressed double-photon emission.

## 22.6 Photo-electric effect for atoms

In this section, we will compute the photo-electric effect cross-section for atoms, as an application of the results obtained in section 22.2. Instead of considering the photo-electric effect for a metal surface, we will take an initial hydrogen-like atomic bound state  $|i\rangle$  for an electron. The light to be absorbed has angular frequency  $\omega$ , polarization  $\hat{\epsilon}$ , and wavenumber  $\vec{k} = \hat{k}\omega/c$ . The final

state  $|f\rangle$  is taken to be a continuum momentum eigenstate with momentum eigenvalue  $\vec{p}_f = \hbar\vec{K}$ . This final state is an approximation, as we are neglecting the influence on its wavefunction of the nuclear electric field; this is to make the problem more tractable, and works best in the limit that the final-state electron energy  $\hbar^2 K^2/2m_e$  is large compared to the binding energy of the initial state. Since we are not assuming that the wavelength of the electromagnetic wave is large compared to the de Broglie wavelength  $2\pi/K$  of the final state, we will not use the multipole expansion of the previous two sections in the following derivation.

To be specific, let the initial state of the electron be a hydrogen-like atomic ground-state wavefunction

$$\langle \vec{r} | i \rangle = \psi_{1,0,0}(\vec{r}) = \sqrt{\frac{Z^3}{\pi a_0^3}} e^{-Zr/a_0}, \quad (22.6.1)$$

which also applies approximately for 1s electrons in multi-electron atoms, with

$$\mathcal{E}_i = -Z^2 e^2 / 2a_0. \quad (22.6.2)$$

The final state electronic wavefunction is approximated by

$$\langle \vec{r} | f \rangle = \psi_{\vec{K}}(\vec{r}) = A e^{i\vec{K} \cdot \vec{r}}, \quad (22.6.3)$$

where  $A$  is a normalization constant and the momentum and energy are

$$\vec{p}_f = \hbar\vec{K}, \quad \text{and} \quad \mathcal{E}_f = \frac{\hbar^2 K^2}{2m_e}. \quad (22.6.4)$$

The delta function in eq. (22.2.7) then reveals that the photon energy is

$$\hbar\omega = \hbar ck = \frac{\hbar^2 K^2}{2m_e} + \frac{Z^2 e^2}{2a_0}. \quad (22.6.5)$$

This shows that there is a minimum possible value of  $\hbar\omega_{\min} = Z^2 e^2 / 2a_0$  in order for the process to take place, achieved when the kinetic energy of the final-state electron is small. This is in agreement with the observation that the photo-electric effect requires a minimum incident photon energy quantum  $\hbar\omega$ , regardless of the intensity of the light, as noted in section 1.3.

However, our calculation will work best in the opposite limit of large  $K$  compared to the binding energy, so that the plane-wave approximation of eq. (22.6.3) is more nearly valid. In that limit, we have  $\hbar^2 K^2 / 2m_e \gg Z^2 e^2 / 2a_0$ , which can be rewritten as  $K \gg Z/a_0$  by using  $a_0 = \hbar^2 / m_e e^2$ . This also implies  $\hbar\omega \gg Z^2 e^2 / 2a_0$ , so  $\omega/c \gg Z^2 e^2 / 2a_0 \hbar c$ , or  $\omega/c \gg Z^2 \alpha / 2a_0$ . The velocity of the ejected electron is then

$$v_f = \frac{p_f}{m_e} = \frac{\hbar K}{m_e} \approx \frac{2\omega}{K}, \quad (22.6.6)$$

where we used eq. (22.6.5) with the approximation that the binding energy is small compared to the photon energy. Therefore, since we will be treating the electron as nonrelativistic, we need  $\omega \ll cK$  so that  $v_f \ll c$ . So, the following results will be most reliable when

$$K \gg \omega/c \gg Z^2\alpha/2a_0, \quad \text{and} \quad K \gg Z/a_0. \quad (22.6.7)$$

The first condition allows us to neglect the spin contribution  $\vec{k} \times \vec{S} = (\omega/c)\hat{k} \times \vec{S}$  in eq. (22.1.13) compared to the  $\vec{P}$  term.

In order to have a well-defined normalization factor  $A$  in eq. (22.6.3), the final state is taken to be confined to a very large cubic box with side  $L \gg a_0/Z$ ; as a check, we will show that the dependence on  $L$  cancels in the final result. Then the normalization condition is

$$1 = \int_{\text{box}} d^3\vec{r} |\psi_{\vec{K}}(\vec{r})|^2 = |A|^2 \int_{\text{box}} d^3\vec{r} = |A|^2 L^3, \quad (22.6.8)$$

so  $A = L^{-3/2}$ . Imposing periodic boundary conditions on the wavefunction implies that

$$\psi_{\vec{K}}(\vec{r}) = \psi_{\vec{K}}(\vec{r} + L[n_x\hat{x} + n_y\hat{y} + n_z\hat{z}]), \quad (22.6.9)$$

for integers  $n_x, n_y, n_z$ , so we obtain the allowed wavenumber components

$$K_x = \frac{2\pi}{L}n_x, \quad K_y = \frac{2\pi}{L}n_y, \quad K_z = \frac{2\pi}{L}n_z. \quad (22.6.10)$$

We will need the density of final states,  $\rho(\mathcal{E}_f, \cos\theta_f, \phi_f)$ , as defined by eq. (21.3.16), where  $(\theta_f, \phi_f)$  are the spherical coordinate angles of the final-state electron's momentum vector  $\vec{K}$ , and serve as the degeneracy labels for the energy. For a volume of  $\vec{K}$ -space  $dK_x dK_y dK_z$ , the number of allowed final states is

$$\left(\frac{L}{2\pi}\right)^3 dK_x dK_y dK_z = \left(\frac{L}{2\pi}\right)^3 d^3\vec{K} = \left(\frac{L}{2\pi}\right)^3 K^2 dK d(\cos\theta_f) d\phi_f. \quad (22.6.11)$$

Now, since

$$d\mathcal{E}_f = \frac{\hbar^2}{m_e} K dK, \quad (22.6.12)$$

we have

$$\left(\text{number of states in the range } d\mathcal{E}_f d\phi_f d(\cos\theta_f)\right) = \left(\frac{L}{2\pi}\right)^3 \frac{m_e}{\hbar^2} K d\mathcal{E}_f d\phi_f d(\cos\theta_f), \quad (22.6.13)$$

from which we obtain, by comparison with eq. (21.3.16), the density of final electron states

$$\rho(\mathcal{E}_f, \cos\theta_f, \phi_f) = \left(\frac{L}{2\pi}\right)^3 \frac{m_e K}{\hbar^2} \quad (22.6.14)$$

in the plane-wave approximation.

We are assuming that the spin can be neglected due to eq. (22.6.7), so eq. (22.1.13) becomes  $\vec{\mathcal{P}}_{\vec{k}} = e^{i\vec{k}\cdot\vec{R}}\hat{\varepsilon}\cdot\vec{P}$ . Applying eq. (22.2.8), we now obtain the differential cross-section for absorption,

$$\frac{d\sigma}{d\Omega_f} = \frac{4\pi^2\alpha K}{m_e\hbar\omega} \left(\frac{L}{2\pi}\right)^3 |\langle f|e^{i\vec{k}\cdot\vec{R}}\hat{\varepsilon}\cdot\vec{P}|i\rangle|^2, \quad (22.6.15)$$

where  $d\Omega_f = d(\cos\theta_f) d\phi_f$ . It remains to evaluate the matrix element. Because of the transverse nature of electromagnetic waves  $\hat{\varepsilon}\cdot\vec{k} = 0$ , we have the commutator  $[\vec{k}\cdot\vec{R}, \hat{\varepsilon}\cdot\vec{P}] = 0$ , so

$$\langle f|e^{i\vec{k}\cdot\vec{R}}\hat{\varepsilon}\cdot\vec{P}|i\rangle = \langle f|\hat{\varepsilon}\cdot\vec{P}e^{i\vec{k}\cdot\vec{R}}|i\rangle = \hbar\hat{\varepsilon}\cdot\vec{K}\langle f|e^{i\vec{k}\cdot\vec{R}}|i\rangle, \quad (22.6.16)$$

where we used  $\vec{p}_f = \hbar\vec{K}$ . The last matrix element is evaluated in the position representation as

$$\langle f|e^{i\vec{k}\cdot\vec{R}}|i\rangle = \int d^3\vec{r} \left(\frac{1}{\sqrt{L^3}}e^{-i\vec{K}\cdot\vec{r}}\right) e^{i\vec{k}\cdot\vec{r}} \left(\sqrt{\frac{Z^3}{\pi a_0^3}}e^{-Zr/a_0}\right) = \sqrt{\frac{Z^3}{\pi a_0^3 L^3}} I(q), \quad (22.6.17)$$

where we have defined the integral

$$I(q) = \int d^3\vec{r} e^{-Zr/a_0} e^{i\vec{q}\cdot\vec{r}} \quad \text{with} \quad \vec{q} \equiv \vec{k} - \vec{K}, \quad (22.6.18)$$

so that  $\hbar\vec{q}$  is the momentum transferred from the photon to the atom.

Due to the spherical symmetry of the initial-state wavefunction, we see that  $I(q)$  only depends on the magnitude of  $\vec{q}$ , not its direction. So, for the purpose of evaluating  $I(q)$ , we can temporarily choose  $\vec{q} = q\hat{z}$ , so that  $\vec{q}\cdot\vec{r} = qr\cos\theta$ , giving

$$I(q) = \int_0^\infty dr r^2 \int_{-1}^1 d(\cos\theta) \int_0^{2\pi} d\phi e^{-r(Z/a_0 - iq\cos\theta)}. \quad (22.6.19)$$

The  $\phi$  integration trivially gives a factor of  $2\pi$ . Doing the  $\cos\theta$  integration next, and the  $r$  integration last,

$$I(q) = -\frac{2\pi i}{q} \int_0^\infty dr r (e^{-r(Z/a_0 - iq)} - e^{-r(Z/a_0 + iq)}) = \frac{8\pi a_0^3}{Z^3(1 + q^2 a_0^2/Z^2)^2}. \quad (22.6.20)$$

Putting together eqs. (22.6.15)-(22.6.17) and (22.6.20) we arrive at

$$\frac{d\sigma}{d\Omega_f} = \frac{32\alpha\hbar K}{m_e\omega} |\hat{\varepsilon}\cdot\vec{K}|^2 \frac{a_0^3}{Z^3(1 + q^2 a_0^2/Z^2)^4}, \quad (22.6.21)$$

in which the dependences on the size of the box  $L$  (from the density of states, and from the matrix element) have canceled, as promised. Note that the differential cross-section is largest when the electron is ejected in the direction of the polarization of the electric field of the wave.

It is now convenient to choose a coordinate system in which (without loss of generality, since the initial electron wavefunction is spherically symmetric) the incident photon momentum is in the  $\hat{z}$  direction, and has polarization along the  $\hat{x}$  direction, so

$$\vec{k} = \hat{z}k = \hat{z}\omega/c, \quad \hat{\varepsilon} = \hat{x}. \quad (22.6.22)$$

Writing, from here on,  $\theta = \theta_f$  and  $\phi = \phi_f$  for the ejected electron's momentum spherical coordinate direction angles, with  $\vec{K} = \hat{r}K$ , we have

$$\hat{\varepsilon} \cdot \vec{K} = K \sin \theta \cos \phi, \quad (22.6.23)$$

$$q^2 = (\vec{k} - \vec{K})^2 = K^2 + (\omega/c)^2 - 2K(\omega/c) \cos \theta, \quad (22.6.24)$$

so that eq. (22.6.21) becomes

$$\frac{d\sigma}{d\Omega} = \frac{32\alpha\hbar}{m_e\omega} \frac{a_0^3 K^3}{Z^3(N_1 - N_2 \cos \theta)^4} \sin^2 \theta \cos^2 \phi, \quad (22.6.25)$$

in which we have introduced dimensionless quantities

$$N_1 = 1 + a_0^2(K^2 + \omega^2/c^2)/Z^2, \quad (22.6.26)$$

$$N_2 = 2a_0^2 K \omega / c Z^2. \quad (22.6.27)$$

To find the total cross-section, we use the integrals

$$\int_0^{2\pi} d\phi \cos^2 \phi = \pi, \quad (22.6.28)$$

$$\int_{-1}^1 d(\cos \theta) \frac{\sin^2 \theta}{(N_1 - N_2 \cos \theta)^4} = \frac{4}{3(N_1^2 - N_2^2)^2}, \quad (22.6.29)$$

to obtain

$$\sigma = \int d\Omega \frac{d\sigma}{d\Omega} = \frac{128\pi\alpha\hbar}{3m_e\omega} \frac{a_0^3 K^3}{Z^3(N_1^2 - N_2^2)^2}. \quad (22.6.30)$$

If the light is polarized along  $\hat{y}$  rather than  $\hat{x}$ , then  $\cos^2 \phi$  should be replaced by  $\sin^2 \phi$  in the differential cross-section eq. (22.6.25). If the light is unpolarized we should instead make the replacement based on the average of the two transverse polarizations,  $\cos^2 \phi \rightarrow (\cos^2 \phi + \sin^2 \phi)/2 = 1/2$ , which gives, after integrating over  $\phi$ ,

$$\frac{d\sigma}{d(\cos \theta)} = \frac{32\pi\alpha\hbar}{m_e\omega} \frac{a_0^3 K^3}{Z^3(N_1 - N_2 \cos \theta)^4} \sin^2 \theta. \quad (22.6.31)$$

The total cross-section is independent of the polarization, because the angle  $\phi$  is integrated over.

Now consider the high-energy limit  $K \gg \omega/c$  and  $K \gg Z/a_0$  of eq. (22.6.7), which we showed was appropriate for our plane-wave approximation for the final state. Expanding to next-to-leading order in large  $K$ , we have

$$\frac{1}{(N_1 - N_2 \cos \theta)^4} = \left( \frac{Z}{Ka_0} \right)^8 \left( 1 + \frac{8\omega}{cK} \cos \theta + \dots \right), \quad (22.6.32)$$

so that eqs. (22.6.25) and (22.6.30) become

$$\frac{d\sigma}{d\Omega} = \frac{32\alpha\hbar}{m_e\omega} \left( \frac{Z}{Ka_0} \right)^5 \left( 1 + \frac{8\omega}{cK} \cos \theta \right) \sin^2 \theta \cos^2 \phi, \quad (22.6.33)$$

$$\sigma = \frac{128\pi\alpha\hbar}{3m_e\omega} \left( \frac{Z}{Ka_0} \right)^5. \quad (22.6.34)$$

This can be rewritten to eliminate  $K$  in favor of  $\omega$  using the energy conservation condition  $K = \sqrt{2m_e\omega/\hbar}$ , which follows from eq. (22.6.5) in the limit of small binding energy compared to the photon energy. The result for the total cross-section is

$$\sigma = \frac{256\pi}{3} \alpha \left( \frac{a_0}{Z} \right)^2 \left( \frac{|E_i|}{\hbar\omega} \right)^{7/2}, \quad (22.6.35)$$

where  $a_0/Z$  is the characteristic length scale of the initial state, and  $|E_i| = Z^2 e^2 / 2a_0$  is the binding energy, and  $\hbar\omega$  is the incident photon energy.

The preceding results apply not just to single-electron wavefunctions for hydrogen-like ions, but to the ejection of electrons from the  $K$ -shell<sup>†</sup> of heavier atoms. There are two  $K$ -shell electrons (one for each spin), so the cross-section we obtained should be multiplied by 2 to find the total  $K$ -shell ejection cross-section for multi-electron atoms. The most important qualitative features are that the cross-section grows with increasing  $Z$  like  $Z^5$ , and falls sharply with increasing incident photon energy like  $1/(\text{energy})^{7/2}$ . The ejected electrons are preferentially emitted along the photon's electric field polarization, which is perpendicular to the photon's propagation direction, but with a slight preference for the forward direction ( $\cos \theta > 0$ ) as a subleading effect in the high-energy expansion in eq. (22.6.33).

In the opposite limit of small  $K$ , eqs. (22.6.25) and (22.6.30) appear to show a threshold behavior proportional to  $K^3$ . However, in that regime, the calculation we did is not to be trusted, because the plane-wave approximation for the final state is quite bad.

## 22.7 Exercises

**Exercise 22.1.** In this problem you will find the lifetime of the  $|n, l, m\rangle = |3, 2, 2\rangle$  state of an isolated hydrogen atom, due to spontaneous emission of a single photon. Ignore fine, hyperfine,

---

<sup>†</sup>The execrable but traditional jargon “ $K$ -shell” just means the innermost shell with principal quantum number  $n = 1$ , called  $1s$  in section 18.4. The  $2s$ ,  $2p$  electrons are called  $L$ -shell, and  $3s$ ,  $3p$  and  $3d$  are called  $M$ -shell, etc. For each shell with principal quantum number  $n$ , there can be up to  $2 \sum_{l=0}^{n-1} (2l+1) = 2n^2$  electrons.



and Lamb shift splitting effects, and use the electric dipole approximation.

(a) According to the electric dipole selection rules, what state or states could be the final state? What are the angular frequency  $\omega$ , the wavelength  $\lambda$ , and the color of the emitted light? Note that  $\lambda \gg a_0$ , so that the electric dipole approximation is indeed valid.

(b) Use eq. (22.4.14) to evaluate the emission rate, and compute  $\tau = 1/\Gamma$  to find the lifetime, both symbolically in terms of quantities like  $\alpha$ ,  $c$ ,  $\hbar$ ,  $a_0$ , and  $\omega$ , and then numerically in seconds.

**Exercise 22.2.** Consider an electron trapped in a 3-dimensional isotropic harmonic oscillator potential with angular frequency  $\Omega$ . The electron is initially in the first excited state with  $l = 1$ ,  $m = 0$ , and spontaneously emits a photon to decay to the ground state.

(a) How is the angular frequency  $\omega$  of the emitted photon related to the angular frequency  $\Omega$  of the harmonic oscillator potential?

(b) Compute the lifetime of the first excited state.

## 23 Scattering in three dimensions

### 23.1 Cross-sections and scattering amplitudes

Consider the problem of scattering of free particles from a localized target in three dimensions. Suppose that the incoming particles are in a mono-energetic beam, all moving in the same direction, which we will take to be  $\hat{z}$  by a choice of coordinate system. If the beam is uniform over some area large compared to the size of the target, it can be described by a flux

$$n = \frac{\text{number of incident particles}}{(\text{area})(\text{time})}. \quad (23.1.1)$$

Here, the area is measured in the  $xy$  plane, perpendicular to the beam propagation direction. The experiment then counts (or infers, perhaps from measurements of energy deposited in a detector) the number of particles scattered in a differential of solid angle

$$d\Omega = d\phi d(\cos\theta). \quad (23.1.2)$$

Let the number of scattered particles as a function of direction be described by

$$ds = \frac{\text{number of scattered particles in } d\Omega}{\text{time}}. \quad (23.1.3)$$

Clearly,  $ds$  must be proportional to  $n$ , and to  $d\Omega$ . We therefore define the proportionality factor to be the scattering differential **cross-section**,

$$\frac{d\sigma}{d\Omega} = \frac{ds}{n d\Omega}. \quad (23.1.4)$$

Note that this has units of area. The total cross-section is then defined as

$$\sigma = \int d\Omega \frac{d\sigma}{d\Omega} = \int_0^{2\pi} d\phi \int_{-1}^1 d(\cos\theta) \frac{d\sigma}{d\Omega}, \quad (23.1.5)$$

also with units of area. Intuitively, the cross-section is the effective area of the target for the type of scattering under consideration.

It is often true that the differential cross-section has **azimuthal symmetry**, which means that it does not depend on  $\phi$ . This occurs if the target and the properties of the scattering process are invariant under rotations about the initial beam propagation direction. In that case, one can give the differential cross-section in terms of

$$\frac{d\sigma}{d(\cos\theta)} = \int_0^{2\pi} d\phi \frac{d\sigma}{d\Omega} = 2\pi \frac{d\sigma}{d\Omega}. \quad (23.1.6)$$

In particular, this will occur if the target is spherically symmetric and the incident particles are either spinless or have randomly oriented spins.




Figure 23.1.1: A classical trajectory for scattering from a hard sphere of radius  $R$ , with impact parameter  $b$  and scattering angle  $\theta$  measured from the forward direction  $\hat{z}$  (the direction pointing to the right). The geometry gives the relation  $b = R \cos(\theta/2)$ .

For classical scattering problems, the cross-section can be computed in terms of specific trajectories followed by the incident particles. It is useful to define the **impact parameter**  $b$  as the distance between a particular initial particle trajectory line (when it is still very far from the target) and the line parallel to it through a suitably chosen central point in the target, which we take to be the origin of our coordinate system.

Consider first a classical<sup>†</sup> example, **hard-sphere scattering**. Particles are fired at a target ball (of radius  $R$ , and heavy enough that it doesn't move when hit). Gravitational effects are neglected, so that the incident particles move in straight lines. For a particular scattering event, the impact parameter  $b$  is the distance between the initial particle trajectory line and the line parallel to it through the center of the target ball, as shown in Figure 23.1.1. To find the differential cross-section, the general strategy is to relate  $b$  to the scattering angle  $\theta$ . Using elementary geometry, the angles  $\alpha$  and  $\theta$  are related to each other and the impact parameter by

$$\alpha = \arcsin(b/R), \quad \theta = \pi - 2\alpha, \quad (23.1.7)$$

which implies

$$b = R \sin(\pi/2 - \theta/2) = R \cos(\theta/2). \quad (23.1.8)$$

Assuming a uniform flux  $n$  of incident particles moving in the  $\hat{z}$  direction, the number of them per unit time with impact parameter between  $b$  and  $b + db$  is equal to the product of  $n$  with the area of the annulus in the  $xy$  plane with inner radius  $b$  and outer radius  $b + db$ ,

$$n(2\pi b db) = n2\pi R \cos(\theta/2) R d(\cos(\theta/2)) = \frac{1}{2}n\pi R^2 d(\cos \theta). \quad (23.1.9)$$

This means that the rate of scattered particles in solid angle  $d\Omega$  in eq. (23.1.3) is

$$ds = \frac{1}{2}n\pi R^2 d(\cos \theta) \frac{d\phi}{2\pi} = \frac{1}{4}nR^2 d\Omega. \quad (23.1.10)$$

---

<sup>†</sup>We will later obtain the result for the corresponding quantum mechanical problem, in section 23.5.

We therefore obtain, from the definition eq. (23.1.4), the differential cross-section

$$\frac{d\sigma}{d\Omega} = R^2/4. \quad (23.1.11)$$

Because of the azimuthal symmetry, this is equivalently

$$\frac{d\sigma}{d(\cos\theta)} = \pi R^2/2, \quad (23.1.12)$$

which integrates to give the total cross-section

$$\sigma = \int_{-1}^1 d(\cos\theta) \frac{d\sigma}{d(\cos\theta)} = \pi R^2. \quad (23.1.13)$$

This example has two rather special features. First, the differential cross-section is isotropic; it does not depend on  $\theta$ . Second, the total cross-section is simply equal to the geometrical cross-sectional area of the target; the scattering takes place if and only if the impact parameter does not exceed the radius  $R$ .

Another classical (and classic) example is Rutherford scattering of a light spinless particle with charge  $Z_1 e$  (for example, an  $\alpha$  particle with  $Z_1 = 2$ ) and initial energy  $E$  from a static Coulomb potential

$$V(r) = Z_2 e/r, \quad (23.1.14)$$

for example, caused by a heavy nucleus containing  $Z_2$  protons. From the classical hyperbolic trajectories, one can show (again by finding  $\theta$  in terms of the impact parameter distance defined when the scattering particle is very far away) that

$$\frac{d\sigma}{d\Omega} = \frac{Z_1^2 Z_2^2 e^4}{16E^2 \sin^4(\theta/2)}. \quad (23.1.15)$$

This differential cross-section is certainly not constant, and in fact blows up with a non-integrable divergence for small  $\theta$ . To get a finite value for the total cross-section, one must modify the question slightly and count only those events in which the electron is scattered by some minimal cut-off angle  $\theta_{\min}$ . (In real-world experiments, this is justified by the finite resolution of detectors, and by the difficulties encountered if one tries to put a detector too close to the beam.) Then

$$\sigma(\theta > \theta_{\min}) = \int_0^{2\pi} d\phi \int_{-1}^{\cos\theta_{\min}} d(\cos\theta) \frac{Z_1^2 Z_2^2 e^4}{16E^2 \sin^4(\theta/2)} = \frac{\pi Z_1^2 Z_2^2 e^4}{4E^2} \left( \frac{1 + \cos\theta_{\min}}{1 - \cos\theta_{\min}} \right). \quad (23.1.16)$$

The divergence is recovered in the limit  $\theta_{\min} \rightarrow 0$ . The interpretation is that in scattering from a  $1/r$  potential, the initial charged particle is always scattered at least a little, with sufficient strength that the unrestricted total cross-section is infinite; we say that the Coulomb potential has infinite range.

Most scattering problems are intermediate between the extremes of hard-sphere scattering (with a sharp cut-off limited range of interaction) and Coulomb scattering (with an infinite range). Even if scattering occurs for every impact parameter or scattering angle, the integrated total cross-section  $\sigma$  can still be finite, if the differential cross-section rises less quickly than  $1/(1 - \cos \theta)$  for small  $\theta$ . If so, then the potential giving rise to it is said to have finite range.

Enough classical warm-ups; we now turn to the quantum theory of scattering. Consider a beam of particles with fixed momentum  $\vec{p} = \hbar \vec{k}$  that encounter a target in the form of a potential  $V(\vec{r})$  that has support within a finite volume near the origin. If the potential were completely absent, the beam could be described by a plane wavefunction

$$\psi(\vec{r}) = \langle \vec{r} | \vec{k} \rangle = \frac{1}{(2\pi)^{3/2}} e^{i\vec{k} \cdot \vec{r}} = \frac{1}{(2\pi)^{3/2}} e^{ikz}. \quad (23.1.17)$$

In writing the last equality, we chose our coordinate system so that the initial beam momentum is along the  $z$  direction, so

$$\vec{k} = \hat{z}k. \quad (23.1.18)$$

The wavefunction eq. (23.1.17) cannot be normalized to unity, so we have chosen to instead normalize it so that it satisfies orthonormality and completeness relations

$$\langle \vec{k}' | \vec{k} \rangle = \delta^{(3)}(\vec{k}' - \vec{k}), \quad \int d^3k |\vec{k}\rangle \langle \vec{k}| = I, \quad (23.1.19)$$

and the probability of finding a particle per unit volume is

$$\frac{\text{probability}}{\text{volume}} = |\psi(\vec{r}, t)|^2 = |e^{-iEt/\hbar} \psi(\vec{r})|^2 = \frac{1}{(2\pi)^3}, \quad (23.1.20)$$

a constant in both time and position. This normalization is rather arbitrary, but will cancel in the differential cross-section. The total flux of particles (probability for a particle in the incident beam to pass through a unit area, per unit time) described by the wavefunction is therefore

$$n = \frac{\text{probability}}{(\text{area})(\text{time})} = \left( \frac{\text{probability}}{\text{volume}} \right) (\text{speed}) = \frac{1}{(2\pi)^3} \frac{\hbar k}{m}. \quad (23.1.21)$$

Let us next anticipate the form of a stationary-state wavefunction for large  $r$ , taking into account the scattering potential,

$$\psi(\vec{r}) = \frac{1}{(2\pi)^{3/2}} \left( e^{ikz} + \frac{e^{ikr}}{r} f_k(\theta, \phi) \right), \quad (r \rightarrow \infty). \quad (23.1.22)$$

We will justify this form before the end of section 23.2. For now, we just note that the first term represents the incident wave, while the quantity

$$f_k(\theta, \phi) \quad (23.1.23)$$

characterizes the outgoing flux of probability density, and is called the **scattering amplitude**. It multiplies a factor  $e^{ikr}$ , which corresponds to a wave traveling away from the origin, and a factor  $1/r$  which indicates that the probability amplitude falls linearly with distance. This gives a probability density that falls off like  $1/r^2$ , which is the requisite behavior so that the probability to detect a particle scattering within a given solid angle  $d\Omega$  with area  $r^2 d\Omega$  is independent of the distance to the detector  $r$ , when  $r$  is large. Due to energy conservation and the assumption that the scattering potential vanishes at large distances, the magnitude of the wavenumber is the same  $k$  for both incoming and outgoing waves. The scattering amplitude  $f_k(\theta, \phi)$  has units of [length] and depends only on the magnitude  $k$  and the spherical coordinate angles, interpreted as those of the propagation direction of a scattered particle. In cases with azimuthal symmetry, we will write the scattering amplitude as  $f_k(\theta)$ .

A small detector at  $(\theta, \phi)$  covering a solid angle  $d\Omega$ , at a large distance  $r$  from the target, will see a total probability flux

$$ds = \left( \frac{\text{probability}}{\text{volume}} \right) (\text{area})(\text{speed}) = \left| \frac{1}{(2\pi)^{3/2}} f_k(\theta, \phi) \frac{e^{ikr}}{r} \right|^2 (r^2 d\Omega) \left( \frac{\hbar k}{m} \right) \quad (23.1.24)$$

$$= \frac{|f_k(\theta, \phi)|^2}{(2\pi)^3} \frac{\hbar k}{m} d\Omega. \quad (23.1.25)$$

Here we have neglected the interference with the incident part of the wavefunction proportional to  $e^{ikz}$ , with the justification that in the real world the incoming particle beam has only a finite extent that will not overlap with sufficiently distant detectors measuring the scattered particles. (To summarize the assumptions made: the beam size perpendicular to its direction is large compared to the target, but small compared to the distance to the detector.) Using eqs. (23.1.21) and (23.1.25) in eq. (23.1.4), we obtain the important result

$$\frac{d\sigma}{d\Omega} = |f_k(\theta, \phi)|^2. \quad (23.1.26)$$

The differential cross-section is equal to the squared magnitude of the scattering amplitude.

## 23.2 The scattering matrix ( $S$ -matrix) operator and the Lippmann–Schwinger equation

A general scattering problem involves an initial state of well-separated free particles denoted  $|a\rangle$ . “Free” means that they are described by plane wave momentum eigenstates, and the symbol “ $a$ ” is a shorthand for all of the data about the momenta and spins and identities of the incoming particles. Interactions then occur in a localized scattering region, described by a wavefunction that might be quite complicated. Finally, one observes particles moving far away from the scattering region as well-separated plane-waves, in a state denoted  $|b\rangle$ .

If the incoming and outgoing particles were really pure plane waves, then they would fill all space at all times. In realistic scattering events, the states are actually wave packets, consisting of superpositions of plane wave states with some spread of momenta, like the Gaussian superpositions of section 6.2. Instead of dealing directly with wave packets, the problem can be finessed by taking our scattering potential to be localized in time as well as position. We imagine that at some times in both the far past and the far future the non-kinetic Hamiltonian terms are turned off. Thus, particles in our wave packet that are far away are free in those two extremes, with a purely kinetic-energy Hamiltonian  $H_0$ . It is convenient to work in the interaction picture of section 20.2, so that the plane wave states have no time dependence at all at those times in the far past and future.

Suppose that the state is a simple plane-wave ket  $|a\rangle$  in the interaction picture at some very early time, which we can write (somewhat loosely for now) as  $t = -\infty$ . As we learned in section 20.2, the interaction-picture state at later times  $t$  will be

$$\mathcal{U}_I(t, -\infty) |a\rangle, \quad (23.2.1)$$

where  $\mathcal{U}_I$  is the unitary time-evolution operator. This motivates the definition of the **scattering matrix** or **S-matrix** operator as its very late-time limit,

$$S = \mathcal{U}_I(+\infty, -\infty), \quad (23.2.2)$$

where again  $+\infty$  is understood to represent some particular large positive time. It follows that the probability amplitude to find the particle in the far future in another plane wave state  $|b\rangle$  is the  $S$ -matrix element

$$\langle b|S|a\rangle. \quad (23.2.3)$$

By its definition, the  $S$ -matrix operator is unitary, which is a way of saying that probability is conserved. Since  $\mathcal{U}_I$  is built out of the interaction (non-kinetic) Hamiltonian, the  $S$ -matrix is also invariant under any symmetries that the system might have. For example,  $S$  will commute with the angular momentum operator if the interaction Hamiltonian is rotationally invariant. This means that  $S$  will be diagonal in a suitably chosen basis of eigenstates of those symmetries. The  $S$ -matrix also encodes further information about the interaction Hamiltonian, including its characteristic length and energy scales, and any bound states or interesting substructure that it may have. Understanding the  $S$ -matrix through scattering is therefore a useful probe of physics at short distance scales that might otherwise be difficult to access directly.

One way to evaluate the  $S$ -matrix elements in eq. (23.2.3) is to work at a fixed intermediate time, which we can choose to be  $t = 0$ . To do this, let us define two scattering states  $|\psi_a^+\rangle$  and

$|\psi_b^-\rangle$  according to

$$|\psi_a^+\rangle = \mathcal{U}_I(0, -\infty) |a\rangle, \quad (23.2.4)$$

$$\langle\psi_b^-| = \langle b|\mathcal{U}_I(\infty, 0). \quad (23.2.5)$$

In words,  $|\psi_a^+\rangle$  is the (generally quite complicated) result at time  $t = 0$  of evolving the plane-wave state  $|a\rangle$  from the distant past  $t = -\infty$ , in the interaction picture. The interpretation of  $|\psi_b^-\rangle$  is perhaps somewhat less intuitive; it is the complicated state at time  $t = 0$  that is obtained by evolving the simple plane-wave state  $|b\rangle$  backwards in time from the far future  $t = \infty$ . This can be seen by using  $\mathcal{U}_I(\infty, 0)^\dagger = \mathcal{U}_I(0, \infty)$  in the adjoint of eq. (23.2.5), which implies

$$|\psi_b^-\rangle = \mathcal{U}_I(0, \infty) |b\rangle. \quad (23.2.6)$$

The point of making these definitions is that we can now write the  $S$ -matrix elements between plane-wave states as the inner product

$$\langle b|S|a\rangle = \langle\psi_b^-|\psi_a^+\rangle, \quad (23.2.7)$$

evaluated at the fixed time  $t = 0$ .

Suppose that the interaction Hamiltonian has no explicit time-dependence, so that energy is conserved. Then the initial and final states have the same energy  $E_a = E_b$ , and we can write

$$S = I - 2\pi i \delta(E_a - E_b) T, \quad (23.2.8)$$

which defines an operator  $T$ , called the **transition operator**. It is the part of the  $S$ -matrix associated with genuinely non-trivial scattering. The factor of  $-2\pi i$  is for future convenience, as we will see soon when we identify  $T$  more explicitly.

The preceding discussion has been very general, as we have not specified whether the free particles in the initial and final configurations are governed by non-relativistic or relativistic kinematics, or whether their numbers and identities change in the scattering process. Quantum field theory is the natural framework for addressing the numerous difficult issues that can arise, including changes in the particle numbers, infrared divergences associated with massless particles, and ultraviolet divergences associated with very high energies of intermediate states. In the rest of this book, we will be content to specialize to the simpler case that the scattering interaction is a potential  $V(\vec{R})$  that depends on a single coordinate vector operator. This  $\vec{R}$  could either describe a single particle that appears in both the initial and final states and interacts with an external potential, or it could be the relative coordinate between two particles that interact with each other through  $V(\vec{R})$ . In both cases, we assume that the scattering is



elastic, so that particles do not change into particles of another type, and that the kinematics conserves the non-relativistic energy, so that the Hamiltonian has the time-independent form

$$H = H_0 + V(\vec{R}), \quad H_0 = P^2/2m. \quad (23.2.9)$$

The potential is assumed to be short-range, which in practice usually means weaker than a Coulomb potential at large distances, so that  $rV(\vec{r}) \rightarrow 0$  for  $r \rightarrow \infty$ . From here on, we will replace the plane-wave labels  $a$  and  $b$  by wavenumbers  $\vec{k}$  and  $\vec{k}'$  associated with momentum eigenvalues  $\hbar\vec{k}$  and  $\hbar\vec{k}'$ , with

$$H_0 |\vec{k}\rangle = E |\vec{k}\rangle, \quad E = \hbar^2 k^2 / 2m. \quad (23.2.10)$$

Spin labels, if any, are suppressed, so that the  $S$ -matrix elements are denoted  $\langle \vec{k}' | S | \vec{k} \rangle$ .

With these assumptions, in the interaction picture we have

$$V_I(t) = e^{iH_0 t/\hbar} V e^{-iH_0 t/\hbar}, \quad (23.2.11)$$

and recall from eq. (20.2.7) that the unitary time-evolution operator for *finite* times is

$$\mathcal{U}_I(t, t') = e^{iH_0 t/\hbar} e^{-iH(t-t')/\hbar} e^{-iH_0 t'/\hbar}. \quad (23.2.12)$$

However, the latter formula has problematic limits for  $t$  or  $t' \rightarrow \pm\infty$ , due to its oscillatory nature. To get around this, we define modified versions of  $\mathcal{U}_I(0, -\infty)$  and  $\mathcal{U}_I(0, +\infty)$ , by inserting exponential damping factors  $e^{\epsilon t'/\hbar}$  and  $e^{-\epsilon t'/\hbar}$  in the integral equation (20.2.14), to turn off the interactions in the far past and the far future, respectively,

$$\Omega_+ \equiv I - \frac{i}{\hbar} \int_{-\infty}^0 dt' e^{\epsilon t'/\hbar} \mathcal{U}_I(0, t') V_I(t'), \quad [\text{modified } \mathcal{U}_I(0, -\infty)], \quad (23.2.13)$$

$$\Omega_- \equiv I + \frac{i}{\hbar} \int_0^{\infty} dt' e^{-\epsilon t'/\hbar} \mathcal{U}_I(0, t') V_I(t'), \quad [\text{modified } \mathcal{U}_I(0, \infty)], \quad (23.2.14)$$

with positive infinitesimal  $\epsilon$ . These are called **Møller operators**, after Christian Møller. Inserting eqs. (23.2.11) and (23.2.12) with  $t = 0$  in eqs. (23.2.13) and (23.2.14), we get

$$|\psi_{\vec{k}}^+\rangle = \Omega_+ |\vec{k}\rangle = \left( I - \frac{i}{\hbar} \int_{-\infty}^0 dt' e^{i(H-E-i\epsilon)t'/\hbar} V \right) |\vec{k}\rangle, \quad (23.2.15)$$

$$|\psi_{\vec{k}}^-\rangle = \Omega_- |\vec{k}\rangle = \left( I + \frac{i}{\hbar} \int_0^{\infty} dt' e^{i(H-E+i\epsilon)t'/\hbar} V \right) |\vec{k}\rangle, \quad (23.2.16)$$

where we used  $e^{-iH_0 t'/\hbar} |\vec{k}\rangle = e^{-iE t'/\hbar} |\vec{k}\rangle$ . Doing the  $t'$  integrals, we find

$$|\psi_{\vec{k}}^{\pm}\rangle = \left( I + \frac{1}{E - H \pm i\epsilon} V \right) |\vec{k}\rangle. \quad (23.2.17)$$

The states  $|\psi_k^\pm\rangle$  are seen to differ in how they avoid the possible singularity in the formal limit  $H = E$ , through opposite  $\pm i\epsilon$  prescriptions in the denominator. This equation contains an inverse operator involving the full Hamiltonian, which is not so convenient for practical evaluation. It is therefore useful to rearrange it, making use of  $H = H_0 + V$ , to arrive at

$$|\psi_k^\pm\rangle = |\vec{k}\rangle + \frac{1}{E - H_0 \pm i\epsilon} V |\psi_k^\pm\rangle. \quad (23.2.18)$$

This is called the **Lippmann–Schwinger equation**, after Bernard Lippmann and Julian Schwinger. It will play an essential role in the following.

Acting on the Lippmann–Schwinger equation with the operator  $E - H_0 \pm i\epsilon$ , and making use of  $H_0 |\vec{k}\rangle = E |\vec{k}\rangle$  and  $H = H_0 + V$  again, one finds  $(H - E) |\psi_k^\pm\rangle = \pm i\epsilon (|\psi_k^\pm\rangle - |\vec{k}\rangle)$ , which in the limit  $\epsilon \rightarrow 0$  reduces to

$$H |\psi_k^\pm\rangle = E |\psi_k^\pm\rangle. \quad (23.2.19)$$

Thus we have the nice result that  $|\psi_k^+\rangle$  and  $|\psi_k^-\rangle$  are both stationary states of the full Hamiltonian  $H$ , each with exactly the same energy eigenvalue  $E = \hbar^2 k^2 / 2m$  as the  $H_0$  eigenvalue of the corresponding plane-wave state. As we will see,  $|\psi_k^+\rangle$  in particular has a wavefunction that has the large- $r$  form that we anticipated in eq. (23.1.22).

Another remarkable result is that the scattering states have the same orthonormality properties as the corresponding plane wave state,

$$\langle \psi_{\vec{k}'}^+ | \psi_{\vec{k}}^+ \rangle = \langle \psi_{\vec{k}'}^- | \psi_{\vec{k}}^- \rangle = \langle \vec{k}' | \vec{k} \rangle = \delta^{(3)}(\vec{k}' - \vec{k}), \quad (23.2.20)$$

where the last equality reflects the assumed normalization in eq. (23.1.19). To prove it, start by considering the matrix element  $\langle \psi_{\vec{k}'}^+ | V | \psi_{\vec{k}}^+ \rangle$ , which can be rewritten in two ways, using the Lippmann–Schwinger equation for  $|\psi_{\vec{k}}^+\rangle$ , and its adjoint for  $\langle \psi_{\vec{k}'}^+ |$ ,

$$\langle \psi_{\vec{k}'}^+ | V | \psi_{\vec{k}}^+ \rangle = \langle \psi_{\vec{k}'}^+ | V \left( |\vec{k}\rangle + \frac{1}{E - H_0 + i\epsilon} V | \psi_{\vec{k}}^+ \right), \quad (23.2.21)$$

$$\langle \psi_{\vec{k}'}^+ | V | \psi_{\vec{k}}^+ \rangle = \left( \langle \vec{k}' | + \langle \psi_{\vec{k}'}^+ | V \frac{1}{E' - H_0 - i\epsilon} \right) V | \psi_{\vec{k}}^+ \rangle, \quad (23.2.22)$$

where we have used the Hermiticity of  $H_0$  and  $V$ , and  $E = \hbar^2 k^2 / 2m$  and  $E' = \hbar^2 k'^2 / 2m$ . Taking the difference of these two equations, we find

$$\langle \psi_{\vec{k}'}^+ | V | \vec{k} \rangle - \langle \vec{k}' | V | \psi_{\vec{k}}^+ \rangle + \langle \psi_{\vec{k}'}^+ | V \left( \frac{1}{E - H_0 + i\epsilon} - \frac{1}{E' - H_0 - i\epsilon} \right) V | \psi_{\vec{k}}^+ \rangle = 0. \quad (23.2.23)$$

Now we write

$$\frac{1}{E - H_0 + i\epsilon} - \frac{1}{E' - H_0 - i\epsilon} = \frac{E' - E - 2i\epsilon}{(E' - H_0 - i\epsilon)(E - H_0 + i\epsilon)}, \quad (23.2.24)$$

and then divide eq. (23.2.23) by  $E' - E - 2i\epsilon$ . In doing so, we can change the  $-2i\epsilon$  to  $-i\epsilon$ , since  $\epsilon$  is infinitesimal and its only purpose is to avoid singular denominators. The result can be expressed, using  $E |\vec{k}\rangle = H_0 |\vec{k}\rangle$  in the first term and  $E' |\vec{k}'\rangle = \langle \vec{k}' | H_0$  in the second term, as

$$\begin{aligned} \langle \psi_{\vec{k}'}^+ | V \frac{1}{E' - H_0 - i\epsilon} |\vec{k}\rangle + \langle \vec{k}' | \frac{1}{E - H_0 + i\epsilon} V |\psi_{\vec{k}}^+\rangle + \\ \langle \psi_{\vec{k}'}^+ | V \frac{1}{E' - H_0 - i\epsilon} \frac{1}{E - H_0 + i\epsilon} V |\psi_{\vec{k}}^+\rangle = 0. \end{aligned} \quad (23.2.25)$$

Again applying the Lippmann–Schwinger equation and its adjoint, this becomes

$$\left( \langle \psi_{\vec{k}'}^+ | - \langle \vec{k}' | \right) |\vec{k}\rangle + \langle \vec{k}' | \left( |\psi_{\vec{k}}^+\rangle - |\vec{k}\rangle \right) + \left( \langle \psi_{\vec{k}'}^+ | - \langle \vec{k}' | \right) \left( |\psi_{\vec{k}}^+\rangle - |\vec{k}\rangle \right) = 0, \quad (23.2.26)$$

which finally simplifies to  $\langle \psi_{\vec{k}'}^+ | \psi_{\vec{k}}^+ \rangle = \langle \vec{k}' | \vec{k} \rangle$ . The proof of  $\langle \psi_{\vec{k}'}^- | \psi_{\vec{k}}^- \rangle = \langle \vec{k}' | \vec{k} \rangle$  is identical except for  $i\epsilon \rightarrow -i\epsilon$  everywhere.

Equation (23.2.20) means that the operators  $\Omega_+$  and  $\Omega_-$  preserve the norms of states, as they obey

$$\Omega_+^\dagger \Omega_+ = \Omega_-^\dagger \Omega_- = I. \quad (23.2.27)$$

It may therefore come as a surprise to learn that these operators are usually *not* unitary. In a finite-dimensional space, an operator  $A$  that satisfies  $A^\dagger A = I$  is invertible, with  $A^{-1} = A^\dagger$ , and also satisfies  $AA^\dagger = I$ , and so is unitary. However, in an infinite-dimensional Hilbert space, one can have  $A^\dagger A = I$  even if  $A$  is not invertible, in which case  $AA^\dagger \neq I$ . Such an operator that is not unitary, but preserves inner products, is called an **isometric operator**. In our present situation, the modification by inserting the convergence factors  $e^{\pm\epsilon t'/\hbar}$  causes  $\Omega_\pm$  to be isometric but not necessarily unitary, unlike the unitary operators  $\mathcal{U}_I(t, t')$  for finite times  $t, t'$ .

To better understand this, note that applying completeness of the plane wave states  $|\vec{k}\rangle$  to eqs. (23.2.15) and (23.2.16) yields

$$\Omega_\pm = \int d^3\vec{k} |\psi_{\vec{k}}^\pm\rangle \langle \vec{k}|, \quad \Omega_\pm^\dagger = \int d^3\vec{k} |\vec{k}\rangle \langle \psi_{\vec{k}}^\pm|. \quad (23.2.28)$$

Now, as a check of eq. (23.2.27),

$$\begin{aligned} \Omega_\pm^\dagger \Omega_\pm &= \int d^3\vec{k} \int d^3\vec{q} |\vec{k}\rangle \langle \psi_{\vec{k}}^\pm | \psi_{\vec{q}}^\pm \rangle \langle \vec{q}| = \int d^3\vec{k} \int d^3\vec{q} \delta^{(3)}(\vec{k} - \vec{q}) |\vec{k}\rangle \langle \vec{q}| \\ &= \int d^3\vec{k} |\vec{k}\rangle \langle \vec{k}| = I \end{aligned} \quad (23.2.29)$$

However, while the scattering states  $|\psi_{\vec{k}}^+\rangle$  satisfy Dirac orthonormality, they are not necessarily a complete orthobasis, because we know that they have positive energy, while the potential  $V$  may have bound states  $|\chi_n\rangle$  with negative energy. Therefore, completeness for them reads

$$\int d^3\vec{k} |\psi_{\vec{k}}^+\rangle \langle \psi_{\vec{k}}^+| + \sum_n |\chi_n\rangle \langle \chi_n| = I, \quad (23.2.30)$$

and similarly for  $|\psi_{\vec{k}}^-\rangle$ . It follows from eqs. (23.2.28) and (23.2.30) that

$$\Omega_+ \Omega_+^\dagger = \Omega_- \Omega_-^\dagger = I - \sum_n |\chi_n\rangle \langle \chi_n|. \quad (23.2.31)$$

So, the Møller operators are unitary (and invertible) only if the potential  $V$  has no bound states.

The  $S$ -matrix operator and its adjoint are given by

$$S = \Omega_-^\dagger \Omega_+, \quad S^\dagger = \Omega_+^\dagger \Omega_-. \quad (23.2.32)$$

The unitarity of  $S$  defined in this way may not be immediately obvious, but it is true. To prove it, note first that since the scattering states  $|\psi_{\vec{k}}^\pm\rangle$  and the bound states  $|\chi_n\rangle$  are eigenstates of the same Hermitian operator  $H$  with different eigenvalues, Theorem 2.6.4 guarantees us that they are orthogonal. It follows immediately from eq. (23.2.28) that

$$\Omega_\pm^\dagger |\chi_n\rangle = 0, \quad \langle \chi_n | \Omega_\pm = 0. \quad (23.2.33)$$

So, we can now compute

$$S^\dagger S = \Omega_+^\dagger \Omega_- \Omega_-^\dagger \Omega_+ = \Omega_+^\dagger \left( I - \sum_n |\chi_n\rangle \langle \chi_n| \right) \Omega_+ = \Omega_+^\dagger \Omega_+ = I, \quad (23.2.34)$$

and similarly

$$S S^\dagger = \Omega_-^\dagger \Omega_+ \Omega_+^\dagger \Omega_- = \Omega_-^\dagger \left( I - \sum_n |\chi_n\rangle \langle \chi_n| \right) \Omega_- = \Omega_-^\dagger \Omega_- = I. \quad (23.2.35)$$

Thus the unitarity of the  $S$  matrix survives the exponential time damping modification for infinitesimal  $\epsilon$ , even though the Møller operators that it is built out of are not unitary.

Let us investigate the  $S$ -matrix element between two arbitrary plane-wave states:

$$\langle \vec{k} | S | \vec{k}' \rangle = \langle \psi_{\vec{k}}^- | \psi_{\vec{k}'}^+ \rangle = \langle \psi_{\vec{k}}^+ | \psi_{\vec{k}'}^+ \rangle + \left( \langle \psi_{\vec{k}}^- | - \langle \psi_{\vec{k}}^+ | \right) | \psi_{\vec{k}'}^+ \rangle \quad (23.2.36)$$

$$= \delta^{(3)}(\vec{k} - \vec{k}') + \langle \vec{k} | V \left( \frac{1}{E - H + i\epsilon} - \frac{1}{E - H - i\epsilon} \right) | \psi_{\vec{k}'}^+ \rangle, \quad (23.2.37)$$

where we used eq. (23.2.20) and the adjoint of eq. (23.2.17) to get the last equality. Now, thanks to eq. (23.2.19), we can replace  $H$  by  $E' = \hbar^2 k'^2 / 2m$ , to get

$$\langle \vec{k} | S | \vec{k}' \rangle = \delta^{(3)}(\vec{k} - \vec{k}') - 2i \frac{\epsilon}{(E - E')^2 + \epsilon^2} \langle \vec{k} | V | \psi_{\vec{k}'}^+ \rangle. \quad (23.2.38)$$

It might be tempting to suppose that the last term should vanish in the limit  $\epsilon \rightarrow 0$ , but special care is needed when  $E = E'$ . Recalling that

$$\lim_{\epsilon \rightarrow 0} \frac{\epsilon}{x^2 + \epsilon^2} = \pi \delta(x), \quad (23.2.39)$$

we actually find

$$\langle \vec{k} | S | \vec{k}' \rangle = \delta^{(3)}(\vec{k} - \vec{k}') - 2\pi i \delta(E - E') \langle \vec{k} | V | \psi_{\vec{k}'}^+ \rangle. \quad (23.2.40)$$

Comparing with eq. (23.2.8), we see that the transition operator  $T$  must satisfy

$$T | \vec{k} \rangle = V | \psi_{\vec{k}}^+ \rangle, \quad (23.2.41)$$

so that

$$\langle \vec{k} | S | \vec{k}' \rangle = \delta^{(3)}(\vec{k} - \vec{k}') - 2\pi i \delta(E - E') \langle \vec{k} | T | \vec{k}' \rangle. \quad (23.2.42)$$

Since the plane-wave kets  $| \vec{k} \rangle$  form an orthobasis, by superposition eq. (23.2.41) provides a complete definition of  $T$  acting on any state.

Our next goal is to relate the matrix elements of  $T$  to the scattering amplitude  $f_k(\theta, \phi)$  introduced in eqs. (23.1.22)-(23.1.26) of the previous section. To do this, we will work with the Lippmann–Schwinger equation (23.2.18) in the position representation. From here on, we drop the subscript  $\vec{k}$  for the scattering state kets and wavefunctions to reduce clutter, so that  $|\psi^\pm\rangle$  are understood to be the scattering states related to  $| \vec{k} \rangle$  by time evolution from  $t = \mp\infty$  to  $t = 0$ , with wavefunctions  $\psi^\pm(\vec{r}) = \langle \vec{r} | \psi^\pm \rangle$ . The wavefunction form of the Lippmann–Schwinger equation is then

$$\langle \vec{r} | \psi^\pm \rangle = \langle \vec{r} | \vec{k} \rangle + \int d^3\vec{r}' \langle \vec{r} | \frac{1}{E - H_0 \pm i\epsilon} | \vec{r}' \rangle \langle \vec{r}' | V | \psi^\pm \rangle, \quad (23.2.43)$$

where we have used completeness with respect to position eigenstates  $| \vec{r}' \rangle$ . The last term in eq. (23.2.43) involves a matrix element that we write as

$$\langle \vec{r} | \frac{1}{E - H_0 \pm i\epsilon} | \vec{r}' \rangle = \frac{2m}{\hbar^2} G_\pm(\vec{r}, \vec{r}'), \quad (23.2.44)$$

which defines  $G_\pm(\vec{r}, \vec{r}')$  as a function independent of the scattering potential  $V$ .

To evaluate the function  $G_\pm(\vec{r}, \vec{r}')$ , we will now use completeness twice more, this time with respect to plane-wave states with momenta  $\hbar\vec{q}$  and  $\hbar\vec{q}'$ , normalized just as in eq. (23.1.17), so

$$\langle \vec{r} | \vec{q} \rangle = \frac{1}{(2\pi)^{3/2}} e^{i\vec{q}\cdot\vec{r}}, \quad (23.2.45)$$

with orthonormality and completeness relations

$$\langle \vec{q}' | \vec{q} \rangle = \delta^{(3)}(\vec{q}' - \vec{q}), \quad \int d^3\vec{q} | \vec{q} \rangle \langle \vec{q} | = I. \quad (23.2.46)$$

Thus eq. (23.2.44) becomes

$$G_\pm(\vec{r}, \vec{r}') = \frac{\hbar^2}{2m} \int d^3\vec{q}' \int d^3\vec{q} \langle \vec{r} | \vec{q}' \rangle \langle \vec{q}' | \frac{1}{E - H_0 \pm i\epsilon} | \vec{q} \rangle \langle \vec{q} | \vec{r}' \rangle. \quad (23.2.47)$$

The reason for doing this is that now the operator  $H_0$  can be evaluated as a number, using

$$H_0|\vec{q}\rangle = \frac{\hbar^2 q^2}{2m}|\vec{q}\rangle, \quad (23.2.48)$$

so that

$$\langle\vec{q}'|\frac{1}{E-H_0\pm i\epsilon}|\vec{q}\rangle = \delta^{(3)}(\vec{q}'-\vec{q})\frac{1}{E-\hbar^2 q^2/2m\pm i\epsilon}. \quad (23.2.49)$$

Doing the  $\int d^3\vec{q}'$  integration in eq. (23.2.47) using the delta function, and remembering that  $E = \hbar^2 k^2/2m$  is the energy of the incident beam particles, we find an integral expression for the function defined by eq. (23.2.44),

$$G_{\pm}(\vec{r}, \vec{r}') = \int \frac{d^3\vec{q}}{(2\pi)^3} \frac{e^{i\vec{q}\cdot(\vec{r}-\vec{r}')}}{k^2 - q^2 \pm i\epsilon}. \quad (23.2.50)$$

Before doing this integral, as an aside we observe that  $G_{\pm}(\vec{r}, \vec{r}')$  could also be defined mathematically as the solution to a differential equation. To see this, note that

$$\nabla^2 G_{\pm}(\vec{r}, \vec{r}') = \int \frac{d^3\vec{q}}{(2\pi)^3} \frac{(-q^2)}{k^2 - q^2 \pm i\epsilon} e^{i\vec{q}\cdot(\vec{r}-\vec{r}')}, \quad (23.2.51)$$

where the Laplacian was taken with respect to the coordinate  $\vec{r}$ , taking advantage of the simple exponential dependence. Therefore,

$$(\nabla^2 + k^2)G_{\pm}(\vec{r}, \vec{r}') = \int \frac{d^3\vec{q}}{(2\pi)^3} e^{i\vec{q}\cdot(\vec{r}-\vec{r}')} = \delta^{(3)}(\vec{r}-\vec{r}'), \quad (23.2.52)$$

which shows that  $G_{\pm}(\vec{r}, \vec{r}')$  is a **Green function** (named after mathematician George Green) for the Helmholtz differential operator  $\nabla^2 + k^2$ , with a delta-function source at  $\vec{r}'$ .

To evaluate the Green function, go back to eq. (23.2.50) and put the integral  $\int d^3\vec{q}$  into spherical coordinates,

$$G_{\pm}(\vec{r}, \vec{r}') = \frac{1}{(2\pi)^3} \int_0^\infty dq q^2 \int_0^{2\pi} d\phi \int_{-1}^1 d(\cos\theta) \frac{e^{iq|\vec{r}-\vec{r}'|\cos\theta}}{k^2 - q^2 \pm i\epsilon}, \quad (23.2.53)$$

where the angles  $\theta$  and  $\phi$  of  $\vec{q}$  are measured from the direction  $\vec{r} - \vec{r}'$ . Now do the angular integrals to get

$$G_{\pm}(\vec{r}, \vec{r}') = \frac{i}{4\pi^2|\vec{r}-\vec{r}'|} \int_0^\infty dq \frac{q}{q^2 - k^2 \mp i\epsilon} \left( e^{iq|\vec{r}-\vec{r}'|} - e^{-iq|\vec{r}-\vec{r}'|} \right). \quad (23.2.54)$$

Let  $x = |\vec{r} - \vec{r}'|$ , and call the integral  $I(x, k)$ . It can be evaluated (using the method of complex variable contour integration, for example) as  $I(x, k) = i\pi e^{\pm ikx}$ . Therefore, the Green function is

$$G_{\pm}(\vec{r}, \vec{r}') = -\frac{e^{\pm ik|\vec{r}-\vec{r}'|}}{4\pi|\vec{r}-\vec{r}'|}. \quad (23.2.55)$$

Plugging this result into the Lippmann–Schwinger equation (23.2.43), using eq. (23.2.44), gives

$$\langle \vec{r} | \psi^\pm \rangle = \langle \vec{r} | \vec{k} \rangle - \frac{2m}{\hbar^2} \int d^3 \vec{r}' \frac{e^{\pm i k |\vec{r} - \vec{r}'|}}{4\pi |\vec{r} - \vec{r}'|} \langle \vec{r}' | V | \psi^\pm \rangle. \quad (23.2.56)$$

We now complete the translation into the position representation by noticing that

$$\langle \vec{r}' | V | \psi^\pm \rangle = \int d^3 \vec{r}'' \langle \vec{r}' | V | \vec{r}'' \rangle \langle \vec{r}'' | \psi^\pm \rangle = V(\vec{r}') \langle \vec{r}' | \psi^\pm \rangle, \quad (23.2.57)$$

where we have used  $\langle \vec{r}' | V | \vec{r}'' \rangle = \delta^{(3)}(\vec{r}' - \vec{r}'') V(\vec{r}')$ . Therefore, the Lippmann–Schwinger equation in the position representation is

$$\psi^\pm(\vec{r}) = \frac{e^{ikz}}{(2\pi)^{3/2}} - \frac{2m}{\hbar^2} \int d^3 \vec{r}' \frac{e^{\pm i k |\vec{r} - \vec{r}'|}}{4\pi |\vec{r} - \vec{r}'|} V(\vec{r}') \psi^\pm(\vec{r}'). \quad (23.2.58)$$

Here,  $\psi^\pm(\vec{r})$  is the total wavefunction, split into the part describing the incident particle beam, and the integral coming from the effect of scattering by the potential.

We now specialize to points  $\vec{r}$  that are very far away from the points  $\vec{r}'$  in the neighborhood of the origin where  $V(\vec{r}')$  has its support. We do this by expanding for  $r' \ll r$ , so that

$$|\vec{r} - \vec{r}'| = \sqrt{r^2 - 2\vec{r} \cdot \vec{r}' + r'^2} = r \sqrt{1 - 2\hat{r} \cdot \vec{r}'/r + \dots} = r - \hat{r} \cdot \vec{r}' + \dots \quad (23.2.59)$$

Let us define a wavevector  $\vec{k}'$ , with the same magnitude as the incoming wavevector  $\vec{k} = k\hat{z}$ , but directed away from the scattering region (in the radial direction),

$$\vec{k}' = k\hat{r}. \quad (23.2.60)$$

Then eq. (23.2.59) implies

$$e^{\pm i k |\vec{r} - \vec{r}'|} \approx e^{\pm i k r} e^{\mp i \vec{k}' \cdot \vec{r}'}. \quad (23.2.61)$$

and eq. (23.2.58) becomes, for large  $r$ ,

$$\psi^\pm(\vec{r}) = \frac{1}{(2\pi)^{3/2}} e^{ikz} - \frac{m}{2\pi\hbar^2} \frac{e^{\pm i k r}}{r} \int d^3 \vec{r}' e^{-i \vec{k}' \cdot \vec{r}'} V(\vec{r}') \psi^\pm(\vec{r}'). \quad (23.2.62)$$

We now see that the factor  $e^{+ikr}$  in  $\psi^+(\vec{r})$  corresponds to a scattered wave moving *away* from the target potential. The factor  $e^{-ikr}$  in  $\psi^-(\vec{r})$  corresponds to the counterintuitive situation of an incoming radial wave converging on a target from all directions and cleverly matching itself onto a plane wave. Therefore we concentrate on

$$\psi^+(\vec{r}) = \frac{1}{(2\pi)^{3/2}} \left( e^{ikz} + \frac{e^{ikr}}{r} f_k(\theta, \phi) \right), \quad (r \rightarrow \infty), \quad (23.2.63)$$

where we have now identified an expression for the scattering amplitude,

$$f_k(\theta, \phi) = -\frac{m}{2\pi\hbar^2} (2\pi)^{3/2} \int d^3\vec{r}' e^{-i\vec{k}' \cdot \vec{r}'} V(\vec{r}') \psi^+(\vec{r}'), \quad (23.2.64)$$

with the angles  $(\theta, \phi)$  defined to be the spherical coordinate angles of  $\vec{k}'$  when the coordinate system has been chosen so that  $\vec{k} = \hat{z}k$ . Equation (23.2.63) indeed has the form promised in eq. (23.1.22). However, eqs. (23.2.63) and (23.2.64) do not yet constitute a solution, because eq. (23.2.63) writes  $\psi^+$  in terms of  $f_k(\theta, \phi)$ , which in turn depends on  $\psi^+$ . Our remaining task in this section is to see how to solve these equations for  $f_k(\theta, \phi)$  by eliminating  $\psi^+$ .

Since the ket  $|\vec{k}'\rangle$  is the state with position wavefunction  $e^{i\vec{k}' \cdot \vec{r}}/(2\pi)^{3/2}$ , an equivalent way of writing eq. (23.2.64) is

$$f_k(\theta, \phi) = -\frac{4\pi^2 m}{\hbar^2} \langle \vec{k}' | V | \psi^+ \rangle, \quad (23.2.65)$$

Now, using the definition of the transition operator in eq. (23.2.41), this becomes a matrix element between completely known (plane-wave) states,

$$f_k(\theta, \phi) = -\frac{4\pi^2 m}{\hbar^2} \langle \vec{k}' | T | \vec{k} \rangle. \quad (23.2.66)$$

The problem of finding the scattering amplitude has therefore been reduced to solving for the matrix elements of  $T$ .

From the Lippmann–Schwinger equation (23.2.18) with the  $+$  sign selected, and again using the definition of the transition operator in eq. (23.2.41), we find

$$T|\vec{k}\rangle = V \left( |\vec{k}\rangle + \frac{1}{E - H_0 + i\epsilon} V |\psi^+\rangle \right) = \left( V + V \frac{1}{E - H_0 + i\epsilon} T \right) |\vec{k}\rangle. \quad (23.2.67)$$

Since this is true for every plane-wave ket  $|\vec{k}\rangle$ , we have an operator relation between  $T$ ,  $V$ ,  $H_0$ , and the incident particle energy  $E$ ,

$$T = V + V \frac{1}{E - H_0 + i\epsilon} T. \quad (23.2.68)$$

This can be formally solved for  $T$ , with the result

$$T = \left( I - V \frac{1}{E - H_0 + i\epsilon} \right)^{-1} V. \quad (23.2.69)$$

Together with eq. (23.2.66), we can finally say that we have a solution for the scattering amplitude. However, practical evaluation of it is still a non-trivial task that typically requires approximation. Different approximation schemes amount to practical methods for evaluating the formal inverse operator on the right side of eq. (23.2.69).



### 23.3 The Optical Theorem

The **Optical Theorem** is an identity relating the total integrated cross-section to the imaginary part of the forward scattering amplitude  $f_k(\theta = 0)$ , as follows:

$$\sigma = \frac{4\pi}{k} \text{Im} [f_k(0)]. \quad (23.3.1)$$

This is useful for at least two reasons. First, it can provide a check of the consistency of a calculation. Second, it is sometimes easier to obtain  $\text{Im}[f_k(0)]$  than it is to find the total cross-section directly. Note that we write  $f_k(0)$  for  $\theta = 0$  even though we are *not* assuming azimuthal symmetry, because the coordinate  $\phi$  is both meaningless and irrelevant when  $\theta = 0$ .

To prove the Optical Theorem, start with

$$\text{Im} [f_k(0)] = -\frac{4\pi^2 m}{\hbar^2} \text{Im} \left( \langle \vec{k} | V | \psi^+ \rangle \right), \quad (23.3.2)$$

which is obtained from eq. (23.2.65) by setting  $\langle \vec{k}' | = \langle \vec{k} |$  for forward scattering, and then taking the imaginary part. To evaluate the right side, we can apply the Hermitian adjoint of the Lippmann–Schwinger equation (23.2.18) with the  $+$  sign selected,

$$\langle \vec{k} | = \langle \psi^+ | - \langle \psi^+ | V \frac{1}{E - H_0 - i\epsilon}, \quad (23.3.3)$$

to obtain

$$\text{Im} \left( \langle \vec{k} | V | \psi^+ \rangle \right) = \text{Im} \left( \langle \psi^+ | V | \psi^+ \rangle \right) - \text{Im} \left( \langle \psi^+ | V \frac{1}{E - H_0 - i\epsilon} V | \psi^+ \rangle \right). \quad (23.3.4)$$

The first term on the right side is 0, because  $V$  is Hermitian, and therefore always has real expectation values. In contrast, the second term does not vanish, because the operator  $1/(E - H_0 - i\epsilon)$  is not Hermitian, due to the  $i\epsilon$ . Using the definition of the transition operator in eq. (23.2.41),

$$\text{Im} \left( \langle \vec{k} | V | \psi^+ \rangle \right) = -\text{Im} \left( \langle \vec{k} | T^\dagger \frac{1}{E - H_0 - i\epsilon} T | \vec{k} \rangle \right) \quad (23.3.5)$$

$$= -\frac{1}{2i} \langle \vec{k} | T^\dagger \left( \frac{1}{E - H_0 - i\epsilon} - \frac{1}{E - H_0 + i\epsilon} \right) T | \vec{k} \rangle \quad (23.3.6)$$

$$= -\langle \vec{k} | T^\dagger \frac{\epsilon}{(E - H_0)^2 + \epsilon^2} T | \vec{k} \rangle. \quad (23.3.7)$$

Inserting a completeness relation  $\int d^3 \vec{k}' |\vec{k}' \rangle \langle \vec{k}'| = I$  just before the  $T$  operator, this becomes

$$\text{Im} \left( \langle \vec{k} | V | \psi^+ \rangle \right) = -\int d^3 \vec{k}' |\langle \vec{k} | T | \vec{k}' \rangle|^2 \pi \delta(E - E'), \quad (23.3.8)$$

where we made use of  $\lim_{\epsilon \rightarrow 0} \frac{\epsilon}{x^2 + \epsilon^2} = \pi \delta(x)$ . Separating the integration over  $\vec{k}'$  into an angular and radial part, and using  $\delta(E - E') = (mk/\hbar^2)\delta(k - k')$  to do the radial integration over  $k'$ , eq. (23.3.8) turns into

$$\text{Im} \left( \langle \vec{k} | V | \psi^+ \rangle \right) = -\frac{\pi m k}{\hbar^2} \int d\Omega_{\vec{k}'} |\langle \vec{k} | T | \vec{k}' \rangle|^2. \quad (23.3.9)$$

Putting this into eq. (23.3.2) gives the imaginary part of the forward scattering amplitude in terms of the matrix elements of the transition operator,

$$\text{Im} [f_k(0)] = \frac{4\pi^3 m^2 k}{\hbar^4} \int d\Omega_{\vec{k}'} |\langle \vec{k} | T | \vec{k}' \rangle|^2. \quad (23.3.10)$$

Since  $|\langle \vec{k} | T | \vec{k}' \rangle|^2$  depends only on the common magnitude of  $\vec{k}$  and  $\vec{k}'$  and the angle between them, we can take it to be a function of  $\vec{k}'$  with  $\vec{k}$  held fixed, and use eqs. (23.1.26) and (23.2.66) to write

$$|\langle \vec{k} | T | \vec{k}' \rangle|^2 = \frac{\hbar^4}{16\pi^4 m^2} |f_k(\theta', \phi')|^2 = \frac{\hbar^4}{16\pi^4 m^2} \frac{d\sigma}{d\Omega'}, \quad (23.3.11)$$

where the angles  $(\theta', \phi')$  of  $\vec{k}'$  are now measured with respect to the  $\vec{k}$  direction. Inserting this in eq. (23.3.10) and doing the angular integration gives

$$\text{Im} [f_k(0)] = \frac{k}{4\pi} \sigma, \quad (23.3.12)$$

which is the Optical Theorem.

## 23.4 Born approximation

The **Born approximation** is a type of perturbative expansion applied to scattering problems, in which the wavefunction within the target region is assumed to be only mildly altered by the potential. In the first-order (or “leading”) Born approximation, for the purposes of eq. (23.2.64) we simply take the wavefunction inside the target to be the same as the incident plane wave,

$$\psi^+(\vec{r}') \approx \frac{1}{(2\pi)^{3/2}} e^{ikz'}. \quad (23.4.1)$$

Recall that the wavevectors of the incident and scattered wavefunctions are  $\vec{k} = k\hat{z}$  and  $\vec{k}' = k\hat{r}$  respectively, so that  $kz' = \vec{k} \cdot \vec{r}'$ . Equation (23.2.64) then reduces to an integral that can be evaluated for any given potential,

$$f_k(\theta, \phi) \approx -\frac{m}{2\pi\hbar^2} \int d^3\vec{r}' e^{i(\vec{k} - \vec{k}') \cdot \vec{r}'} V(\vec{r}'). \quad (23.4.2)$$

Note that in this first-order Born approximation, the sign of the potential does not make any difference, since it enters linearly in the scattering amplitude, which then gets squared to give

$\frac{d\sigma}{d\Omega} = |f_k(\theta, \phi)|^2$ . In particular, within the leading Born approximation, an attractive potential gives the same differential cross-section as a repulsive potential with the same magnitude.

Before proceeding to use it, let us first address the question of the validity of the Born approximation. Intuitively, this relies on the potential not being too strong. Consider the wavefunction form of the Lippmann–Schwinger equation valid inside the target region, as it appeared in eq. (23.2.58) before we took the large  $r$  limit. To make a rough estimate, let us call  $a$  the characteristic range of the potential, meaning that  $V(\vec{r})$  is significant only for  $r < a$ . Then we can require that in the right side of eq. (23.2.58) the correction term is much smaller in magnitude than the plane wave term,

$$\frac{2m}{\hbar^2} \left| \int d^3\vec{r}' \frac{e^{ik|\vec{r}-\vec{r}'|}}{4\pi|\vec{r}-\vec{r}'|} V(\vec{r}') e^{ikz'} \right| \ll 1, \quad (23.4.3)$$

for  $r < a$ , where we have presumptively replaced the wavefunction  $\psi^+$  by its putative approximation eq. (23.4.1). To make things easier, we can just consider  $\vec{r}$  close to the origin, where the correction might be expected to be largest, or at least not much smaller than at other points. Then a rough condition for validity of the Born approximation is

$$\frac{m}{2\pi\hbar^2} \left| \int d^3\vec{r}' \frac{1}{r'} e^{ik(r'+z')} V(\vec{r}') \right| \ll 1. \quad (23.4.4)$$

Now, we make the further approximation of taking  $k \approx 0$  and replacing the potential by its average over  $r' < a$ , a constant factor that we will call  $\bar{V}$ . The rest of the integral is then

$$\int d^3\vec{r}' \frac{1}{r'} = 4\pi \int_0^a dr' r' = 2\pi a^2. \quad (23.4.5)$$

Thus our estimate for the validity condition of the Born approximation is

$$\frac{ma^2}{\hbar^2} |\bar{V}| \ll 1, \quad (23.4.6)$$

where  $a$  and  $\bar{V}$  are the characteristic range and strength of the potential. This could have been guessed merely on dimensional grounds, at least in the low-energy limit of small  $k$ . However, this estimate for the range of validity of the Born approximation can often be too conservative, particularly for larger  $k$ , because then the oscillation of the integrand can produce efficient cancellation in the left side of eq. (23.4.3), not accounted for in our rough estimate. In that case, dimensional analysis suggests that the weaker condition  $ma|\bar{V}|/\hbar^2 k \ll 1$  could suffice.

It is also an amusing fact that the Optical Theorem completely fails in the leading Born approximation. This is immediately obvious from the fact that eq. (23.4.2) is real when  $\vec{k}' = \vec{k}$ , so that  $\text{Im}[f_k(0)] = 0$ , but the total cross-section does not vanish. Going beyond the leading order Born approximation, as we will do before the end of this section, is necessary to restore the validity of the Optical Theorem.




Figure 23.4.1: Geometry of vectors involved in elastic scattering from a potential. The initial and final wavevectors  $\vec{k}$  and  $\vec{k}'$  have equal length  $k$ . The vector  $\vec{q} = \vec{k} - \vec{k}'$ , where  $\hbar\vec{q}$  is the momentum transferred from the initial particle to the scattering potential, has length  $q = 2k \sin(\theta/2)$ , where  $\theta$  is the angle between  $\vec{k}$  and  $\vec{k}'$ .

Let us now consider examples of the leading Born approximation. An important special case is that of a spherically symmetric potential, so that the scattering amplitude has azimuthal symmetry (no dependence on  $\phi$ ). Define

$$\vec{q} = \vec{k} - \vec{k}', \quad (23.4.7)$$

so that  $\hbar\vec{q}$  is the momentum transferred from the initial particle to the scattering potential. (See Figure 23.4.1.) Then  $f_k(\theta)$  only depends on  $k$  and  $\theta$  through the combination

$$q = |\vec{k} - \vec{k}'| = 2k \sin(\theta/2). \quad (23.4.8)$$

Indeed, eq. (23.4.2) becomes

$$f_k(\theta) = -\frac{m}{2\pi\hbar^2} \int d^3\vec{r}' V(r') e^{i\vec{q}\vec{r}' \cos \theta'}, \quad (23.4.9)$$

where we have taken advantage of the spherical symmetry to choose our  $\vec{r}'$  coordinate system in the integral so that  $\vec{q}$  is in the  $\cos \theta' = 0$  direction. It follows that

$$f_k(\theta) = -\frac{m}{2\pi\hbar^2} \int_0^\infty dr' r'^2 \int_0^{2\pi} d\phi' \int_{-1}^1 d(\cos \theta') V(r') e^{iqr' \cos \theta'} \quad (23.4.10)$$

$$= \frac{im}{\hbar^2 q} \int_0^\infty dr' r' V(r') (e^{iqr'} - e^{-iqr'}), \quad (23.4.11)$$

which we can rewrite, without the distracting primes, as

$$f_k(\theta) = -\frac{2m}{\hbar^2 q} \int_0^\infty dr r V(r) \sin(qr). \quad (23.4.12)$$

This is the first-order Born approximation result for the scattering amplitude with an arbitrary spherically symmetric potential. In this case,  $f_k(\theta)$  is always real. Note that the large distance contribution to the integral converges if  $V(r)$  decreases faster than  $1/r$  for large  $r$ .

In the low-energy approximation  $q \rightarrow 0$ , one can use  $\sin(qr) \approx qr$  to find that

$$f_k(\theta) = -\frac{2m}{\hbar^2} \int_0^\infty dr r^2 V(r), \quad (\text{small } q \text{ limit}), \quad (23.4.13)$$

which is independent of  $q$ , and therefore independent of  $\theta$ . In the opposite limit of very large  $q$ , eq. (23.4.12) gives  $f_k(\theta) \rightarrow 0$  because of the  $q$  in the denominator together with the cancellation due to the very rapid oscillation of  $\sin(qr)$  in the integrand.

As an example, consider the **Yukawa potential**, named after Hideki Yukawa, who used it in 1935 to model the strong interactions between nucleons due to the exchange of mesons. It is

$$V(r) = -\frac{g}{r}e^{-r/a}, \quad (23.4.14)$$

where  $g$  and  $a$  are constants with units of [(energy)(length)] and [length], respectively. For  $r \ll a$ , this has the same form as a Coulomb potential. However, for large  $r \gg a$ , the potential approaches 0 exponentially faster than the Coulomb potential does. For that reason it is sometimes also known as the **screened Coulomb potential**, and  $a$  can be thought of as the range of the potential. Applying eq. (23.4.12) gives

$$f_k(\theta) = \frac{2mg}{\hbar^2 q} \int_0^\infty dr e^{-r/a} \sin(qr) = \frac{2mg}{\hbar^2} \frac{1}{q^2 + 1/a^2}. \quad (23.4.15)$$

Now we can use  $q^2 = 4k^2 \sin^2(\theta/2) = 2k^2(1 - \cos \theta)$ , to find

$$\frac{d\sigma}{d\Omega} = |f_k(\theta)|^2 = \left( \frac{2mg}{\hbar^2} \right)^2 \frac{1}{[2k^2(1 - \cos \theta) + 1/a^2]^2} \quad (23.4.16)$$

in the Born approximation. If we take the long-range limit  $a \rightarrow \infty$  and let  $-g = Z_1 Z_2 e^2$ , then

$$\frac{d\sigma}{d\Omega} = \frac{Z_1^2 Z_2^2 e^4 m^2}{4\hbar^4 k^4 \sin^4(\theta/2)} = \frac{Z_1^2 Z_2^2 e^4}{16E^2 \sin^4(\theta/2)}, \quad (23.4.17)$$

where in the last expression we used  $E = \hbar^2 k^2 / 2m$ . This is the same differential scattering cross-section as we found for the corresponding classical Rutherford scattering problem in section 23.1.

In the limit of low-energy scattering from the Yukawa potential, the rough condition for validity of the Born approximation is obtained by taking  $\bar{V} = g/a$  in eq. (23.4.6), so

$$|g| \ll \hbar^2 / ma. \quad (23.4.18)$$

This can be compared to the condition necessary for a bound state to exist, which can be shown (for example, applying the variational method with a carefully chosen trial wavefunction) to be approximately  $g > 0.8399\hbar^2 / ma$ . The general lesson is that if the potential allows a bound state, then it is too strong for the Born approximation to be valid for low  $E$ . However, it can be valid for larger  $E$ , such that  $k \gg 1/a$ .

Returning to the general scattering problem, let us consider how to extend the Born approximation beyond leading order. We can start with the expression eq. (23.2.69) for the transition operator, and use the expansion  $(I - X)^{-1} = I + X + X^2 + X^3 + \dots$ . The result is

$$T = V + V \frac{1}{E - H_0 + i\epsilon} V + V \frac{1}{E - H_0 + i\epsilon} V \frac{1}{E - H_0 + i\epsilon} V + \dots \quad (23.4.19)$$

The ordering of operators matters, because  $H_0$  and  $V$  do not commute unless the latter is a trivial (constant) potential. Recalling, from eq. (23.2.66), that

$$f_k(\theta, \phi) = -\frac{4\pi^2 m}{\hbar^2} \langle \vec{k}' | T | \vec{k} \rangle, \quad (23.4.20)$$

we obtain an expansion in powers of the potential,

$$f_k(\theta, \phi) = f_k^{(1)}(\theta, \phi) + f_k^{(2)}(\theta, \phi) + f_k^{(3)}(\theta, \phi) + \cdots, \quad (23.4.21)$$

where  $f_k^{(N)}(\theta, \phi)$  has  $N$  factors of the potential  $V$ . In particular,

$$f_k^{(1)}(\theta, \phi) = -\frac{4\pi^2 m}{\hbar^2} \langle \vec{k}' | V | \vec{k} \rangle \quad (23.4.22)$$

is just the leading Born approximation that we have already studied, and the second-order and third-order contributions in the Born expansion are now seen to be

$$f_k^{(2)}(\theta, \phi) = -\frac{4\pi^2 m}{\hbar^2} \langle \vec{k}' | V \frac{1}{E - H_0 + i\epsilon} V | \vec{k} \rangle, \quad (23.4.23)$$

$$f_k^{(3)}(\theta, \phi) = -\frac{4\pi^2 m}{\hbar^2} \langle \vec{k}' | V \frac{1}{E - H_0 + i\epsilon} V \frac{1}{E - H_0 + i\epsilon} V | \vec{k} \rangle, \quad (23.4.24)$$

etc. Note that the leading Born approximation just amounts to setting  $T = V$  in eq. (23.4.20).

We can now work out these contributions in the wavefunction representation, in terms of the Green function, by using completeness with respect to position and eqs. (23.2.45) and (23.2.44) for the plane wave and Green function. The results are

$$f_k^{(1)}(\theta, \phi) = -\frac{1}{4\pi} \left( \frac{2m}{\hbar^2} \right) \int d^3 \vec{r}_1 e^{i\vec{k} \cdot \vec{r}_1} V(\vec{r}_1) e^{-i\vec{k}' \cdot \vec{r}_1}, \quad (23.4.25)$$

$$f_k^{(2)}(\theta, \phi) = -\frac{1}{4\pi} \left( \frac{2m}{\hbar^2} \right)^2 \int d^3 \vec{r}_1 \int d^3 \vec{r}_2 e^{i\vec{k} \cdot \vec{r}_1} V(\vec{r}_1) G_+(\vec{r}_1, \vec{r}_2) V(\vec{r}_2) e^{-i\vec{k}' \cdot \vec{r}_2}, \quad (23.4.26)$$

$$f_k^{(3)}(\theta, \phi) = -\frac{1}{4\pi} \left( \frac{2m}{\hbar^2} \right)^3 \int d^3 \vec{r}_1 \int d^3 \vec{r}_2 \int d^3 \vec{r}_3 e^{i\vec{k} \cdot \vec{r}_1} V(\vec{r}_1) G_+(\vec{r}_1, \vec{r}_2) V(\vec{r}_2) G_+(\vec{r}_2, \vec{r}_3) V(\vec{r}_3) e^{-i\vec{k}' \cdot \vec{r}_3}. \quad (23.4.27)$$

A diagrammatic version of these contributions is shown in Figure 23.4.2. The formula for  $f_k^{(N)}(\theta, \phi)$  contains  $N$  vertices where the scattering potential acts. At each scattering vertex, labeled  $n = 1, \dots, N$ , there is an integration

$$\text{vertex} \leftrightarrow \frac{2m}{\hbar^2} \int d^3 \vec{r}_n V(\vec{r}_n), \quad (23.4.28)$$

and we assign factors

$$\text{internal lines} \leftrightarrow G_+(\vec{r}_n, \vec{r}_{n+1}) \quad (23.4.29)$$




Figure 23.4.2: Diagrammatic representation of the first three terms in the Born approximation expansion for the scattering amplitude,  $f_k^{(1)}$ ,  $f_k^{(2)}$ , and  $f_k^{(3)}$  from eqs. (23.4.25)–(23.4.27). Each of the vertex positions  $\vec{r}_n$  is integrated over, with an associated factor of  $2m/\hbar^2$ .

to the lines between consecutive vertices  $\vec{r}_n$  and  $\vec{r}_{n+1}$ . There are also two factors

$$\text{external lines} \leftrightarrow e^{i\vec{k}\cdot\vec{r}_1} e^{-i\vec{k}'\cdot\vec{r}_N} \quad (23.4.30)$$

associated with the initial and final momenta, and a single factor of  $-1/4\pi$ .

An intuitive interpretation for the Born expansion is that the particle is described by plane waves except for interactions with the potential. Between potential interactions, it propagates according to the Green function of the Helmholtz operator, and the initial and final wavefunctions are free plane-waves. All possible numbers of interactions are summed over, and then all possible interaction positions are integrated over, to give the total scattering amplitude.

## 23.5 Spherical potential scattering and the partial wave expansion

Consider scattering from a spherically symmetric potential  $V(\vec{r}) = V(r)$ . As usual, we take the initial particle wavenumber to be  $\vec{k} = \hat{z}k$ , so that we have azimuthal symmetry and the scattering amplitude can only depend on  $\theta$ , not  $\phi$ . This implies that we can write a **partial wave expansion** in the orbital angular momentum quantum number  $l$ ,

$$f_k(\theta) = \sum_{l=0}^{\infty} (2l+1) P_l(\cos \theta) f_l(k). \quad (23.5.1)$$

The  $P_l(\cos \theta)$  are the Legendre polynomials, and the factor of  $2l+1$  is a convention. This equation defines the quantities  $f_l(k)$ , called **partial wave amplitudes**, that appear on the right side. They can equivalently be viewed as functions  $f_l(E)$  of the energy  $E = \hbar^2 k^2 / 2m$ .

To better understand the partial wave expansion, recall the expansion of a plane wave in the absence of scattering, worked out at the end of section 10.2,

$$e^{ikz} = e^{ikr \cos \theta} = \sum_{l=0}^{\infty} (2l+1) P_l(\cos \theta) i^l j_l(kr). \quad (23.5.2)$$

The  $j_l(kr)$  are spherical Bessel functions. For large  $r$ , we find from eq. (10.2.17) that

$$i^l j_l(kr) \approx \frac{e^{ikr} - e^{-ikr + i\pi l}}{2ikr}, \quad (r \gg 1/k). \quad (23.5.3)$$

Plugging eqs. (23.5.1)-(23.5.3) into eq. (23.2.63)

$$\psi^+(\vec{r}) = \frac{1}{(2\pi)^{3/2}} \left[ e^{ikz} + f_k(\theta) \frac{e^{ikr}}{r} \right] \quad (23.5.4)$$

$$= \frac{1}{(2\pi)^{3/2}} \left[ \sum_{l=0}^{\infty} (2l+1) P_l(\cos \theta) \left( \frac{e^{ikr} - e^{-ikr+i\pi l}}{2ikr} \right) + \sum_{l=0}^{\infty} (2l+1) P_l(\cos \theta) f_l(k) \frac{e^{ikr}}{r} \right], \quad (23.5.5)$$

where the large  $r$  condition is implicit. Collecting like terms, we find

$$\psi^+(\vec{r}) = \frac{1}{(2\pi)^{3/2} 2ik} \sum_{l=0}^{\infty} (2l+1) P_l(\cos \theta) \left( S_l \frac{e^{ikr}}{r} + (-1)^{l+1} \frac{e^{-ikr}}{r} \right) \quad (23.5.6)$$

for  $r \rightarrow \infty$ , where we have defined

$$S_l = 1 + i2kf_l(k). \quad (23.5.7)$$

The terms proportional to  $e^{ikr}$  and  $e^{-ikr}$  are outgoing and incoming spherical waves, respectively; compare to the discussion around eqs. (10.2.24)–(10.2.27). They are called the **partial waves** of angular momentum  $l$ . Note that the outgoing spherical waves depend on the potential  $V(r)$  through the coefficients  $S_l$ . In contrast, the incoming spherical waves do not depend on the  $S_l$  and therefore do not depend on  $V(r)$ ; they are just a re-writing of the incoming beam.

The quantities  $S_l$  have a simple interpretation: they are the  $S$ -matrix elements in an angular momentum basis. To see this, consider general spherical coordinate angles for the directions of  $\vec{k}$  and  $\vec{k}'$ , which are  $\Omega_{\vec{k}} = (\theta_{\vec{k}}, \phi_{\vec{k}})$  and  $\Omega_{\vec{k}'} = (\theta_{\vec{k}'}, \phi_{\vec{k}'})$  respectively. The angle between these two vectors is the  $\theta$  appearing in the preceding equations. Therefore, the spherical harmonics addition formula (8.6.71) allows us to rewrite the scattering amplitude in eq. (23.5.1) as

$$f_k(\theta) = 4\pi \sum_{l=0}^{\infty} \sum_{m=-l}^l Y_l^m(\theta_{\vec{k}'}, \phi_{\vec{k}'})^* Y_l^m(\theta_{\vec{k}}, \phi_{\vec{k}}) \frac{S_l - 1}{i2k}. \quad (23.5.8)$$

So, eq. (23.2.66) tells us that the transition operator matrix elements are

$$\langle \vec{k}' | T | \vec{k} \rangle = \frac{i\hbar^2}{2\pi m k} \sum_{l,m} Y_l^m(\theta_{\vec{k}'}, \phi_{\vec{k}'})^* Y_l^m(\theta_{\vec{k}}, \phi_{\vec{k}}) (S_l - 1), \quad (23.5.9)$$

and so the  $S$ -matrix elements are, from eq. (23.2.42),

$$\langle \vec{k}' | S | \vec{k} \rangle = \delta^{(3)}(\vec{k} - \vec{k}') + \frac{\hbar^2}{mk} \delta(E - E') \sum_{l,m} Y_l^m(\theta_{\vec{k}'}, \phi_{\vec{k}'})^* Y_l^m(\theta_{\vec{k}}, \phi_{\vec{k}}) (S_l - 1). \quad (23.5.10)$$

This can be simplified using  $\delta(E - E') = (m/\hbar^2 k) \delta(k - k')$ , and the definition

$$\delta^{(3)}(\vec{k} - \vec{k}') = \frac{1}{k^2} \delta(k - k') \delta(\phi_{\vec{k}} - \phi_{\vec{k}'}) \delta(\cos \theta_{\vec{k}} - \cos \theta_{\vec{k}'}), \quad (23.5.11)$$



and the spherical harmonic completeness formula (8.6.32), reproduced here for convenience,  $\delta(\phi - \phi')\delta(\cos \theta - \cos \theta') = \sum_{l=0}^{\infty} \sum_{m=-l}^l Y_l^m(\theta', \phi')^* Y_l^m(\theta, \phi)$ , to give

$$\langle \vec{k}' | S | \vec{k} \rangle = \frac{1}{k^2} \delta(k - k') \sum_{l=0}^{\infty} \sum_{m=-l}^l Y_l^m(\theta_{\vec{k}'}, \phi_{\vec{k}'})^* Y_l^m(\theta_{\vec{k}}, \phi_{\vec{k}}) S_l. \quad (23.5.12)$$

Now consider a basis of free-particle angular momentum eigenstates  $|k, l, m\rangle$ , of the type discussed in section 10.2. These states are related to the plane-wave states by

$$|k, l, m\rangle = \int d\Omega_{\vec{k}} Y_l^m(\theta_{\vec{k}}, \phi_{\vec{k}}) |\vec{k}\rangle, \quad (23.5.13)$$

$$|\vec{k}\rangle = \sum_{l=0}^{\infty} \sum_{m=-l}^l Y_l^m(\theta_{\vec{k}}, \phi_{\vec{k}}) |k, l, m\rangle, \quad (23.5.14)$$

and satisfy orthonormality and completeness relations

$$\langle k', l', m' | k, l, m \rangle = \frac{1}{k^2} \delta(k - k') \delta_{ll'} \delta_{mm'}, \quad (23.5.15)$$

$$\int_0^{\infty} dk k^2 \sum_{l=0}^{\infty} \sum_{m=-l}^l |k, l, m\rangle \langle k, l, m| = I. \quad (23.5.16)$$

It is a short exercise to show that the content of eq. (23.5.12) can now be expressed as

$$\langle k', l', m' | S | k, l, m \rangle = \frac{1}{k^2} \delta(k - k') \delta_{ll'} \delta_{mm'} S_l, \quad (23.5.17)$$

or, in yet another way, as

$$S = \int_0^{\infty} dk k^2 \sum_{l=0}^{\infty} \sum_{m=-l}^l S_l |k, l, m\rangle \langle k, l, m| \quad (23.5.18)$$

The fact that the  $S$ -matrix is diagonal in a basis of orbital angular momentum eigenstates is due to the assumed rotational symmetry invariance of the potential.

The unitarity of the  $S$ -matrix now implies

$$\begin{aligned} I &= S^\dagger S = \int_0^{\infty} dk' k'^2 \sum_{l', m'} \int_0^{\infty} dk k^2 \sum_{l, m} S_l^* S_l |k', l', m'\rangle \langle k', l', m' | k, l, m\rangle \langle k, l, m| \\ &= \int_0^{\infty} dk k^2 \sum_{l, m} |S_l|^2 |k, l, m\rangle \langle k, l, m| \end{aligned} \quad (23.5.19)$$

where the last equality made use of the orthonormality relation (23.5.15) to eliminate the  $k'$  integral and the  $l'$  and  $m'$  sums. Comparison with eq. (23.5.16) establishes that  $|S_l|^2 = 1$  for each  $l$ . This crucial fact is called **partial wave unitarity**, and it is a way of restating the

assumption that the potential is only scattering the particles, not creating or destroying them. It means that we can write, for the complex factors  $S_l$  in eq. (23.5.6),

$$S_l = 1 + i2kf_l(k) = e^{i2\delta_l}, \quad (23.5.20)$$

which defines a set of real numbers  $\delta_l$  called the **phase shifts** of the angular momentum  $l$  partial waves. (The factor of 2 in the exponent is a traditional convention.) Equation (23.5.20) exposes that the effect of the scattering is just to give each outgoing partial wave of orbital angular momentum  $l$  an extra phase factor  $e^{i2\delta_l}$  relative to the corresponding incoming partial wave.

The general scattering problem for a spherically symmetric potential thus reduces to finding the phase shifts  $\delta_l$  appearing in the wavefunction for large  $r$  obtained by re-writing eq. (23.5.6),

$$\psi^+(\vec{r}) = \frac{1}{(2\pi)^{3/2}} \sum_{l=0}^{\infty} (2l+1) P_l(\cos \theta) [e^{i2\delta_l} e^{ikr} - e^{-i(kr-l\pi)}] / 2ikr, \quad (r \rightarrow \infty). \quad (23.5.21)$$

In the limit that the potential  $V(r)$  vanishes, then  $\delta_l = 0$ , and more generally the  $\delta_l$  parameterize the effect of  $V(r)$  on each  $l$  partial wave. We will show before the end of this section that it is a good approximation to use only the lowest few partial waves  $l = 0, 1, 2, \dots$ , except in the case of very high-energy scattering.

Some useful equivalent ways of writing the relation between the phase shifts and the partial wave amplitudes  $f_l(k)$  are, from eq. (23.5.20),

$$f_l(k) = \frac{e^{i2\delta_l} - 1}{2ik} = \frac{e^{i\delta_l} \sin \delta_l}{k} = \frac{1}{k(\cot \delta_l - i)}. \quad (23.5.22)$$

Using the next-to-last expression of eq. (23.5.22) in eq. (23.5.1), we have the scattering amplitude in terms of the phase shifts,

$$f_k(\theta) = \frac{1}{k} \sum_{l=0}^{\infty} (2l+1) e^{i\delta_l} \sin \delta_l P_l(\cos \theta). \quad (23.5.23)$$

This can be used to compute the total cross-section,

$$\sigma = 2\pi \int_{-1}^1 d(\cos \theta) |f_k(\theta)|^2 \quad (23.5.24)$$

$$= \frac{2\pi}{k^2} \sum_{l=0}^{\infty} \sum_{l'=0}^{\infty} (2l+1)(2l'+1) e^{i(\delta_l - \delta_{l'})} \sin(\delta_l) \sin(\delta_{l'}) \int_{-1}^1 d(\cos \theta) P_{l'}(\cos \theta) P_l(\cos \theta). \quad (23.5.25)$$

To simplify this, apply the Legendre polynomial orthogonality condition  $\int_{-1}^1 du P_{l'}(u) P_l(u) = \delta_{ll'} 2/(2l+1)$ . We arrive at

$$\sigma = \frac{4\pi}{k^2} \sum_{l=0}^{\infty} (2l+1) \sin^2 \delta_l. \quad (23.5.26)$$

Note that even though the differential cross-section  $d\sigma/d\Omega = |f_k(\theta)|^2$  certainly has interferences between partial wave amplitudes with different  $l$ , there is no such interference in the total cross-section. As a check, or an alternative derivation, the Optical Theorem tells us that

$$\sigma = \frac{4\pi}{k} \text{Im}[f_k(\theta = 0)] = \frac{4\pi}{k} \left( \frac{1}{k} \sum_{l=0}^{\infty} (2l+1) P_l(1) \sin(\delta_l) \text{Im}[e^{i\delta_l}] \right), \quad (23.5.27)$$

which reproduces eq. (23.5.26) after using  $\text{Im}[e^{i\delta_l}] = \sin \delta_l$  and the Legendre polynomial normalization condition  $P_l(1) = 1$ .

An important consequence of eq. (23.5.22), as encoded in eq. (23.5.26), follows if we write

$$\sigma = \sum_{l=0}^{\infty} \sigma_l, \quad (23.5.28)$$

where  $\sigma_l$  is the contribution from orbital angular momentum  $l$ . Then  $\sin^2 \delta_l \leq 1$  implies

$$\sigma_l \leq \frac{4\pi}{k^2} (2l+1). \quad (23.5.29)$$

This is known as the **partial wave unitarity bound**. Equality is achieved only if  $\sin^2 \delta_l = 1$ , which implies  $\delta_l = (n + 1/2)\pi$  for integer  $n$ . In that case,  $\sigma_l$  is said to saturate the partial wave unitarity bound. Remarkably, this bound on the contribution  $\sigma_l$  for each partial wave applies no matter how big the scattering potential is. It becomes stronger with increasing energy, as it can be rewritten

$$\sigma_l \leq \frac{2\pi\hbar^2}{mE} (2l+1), \quad (23.5.30)$$

and it should be viewed as a requirement imposed by the conservation of probability.

Partial wave unitarity also restricts the values that  $f_l(k)$  can take to a circle of radius  $1/2k$  in the complex plane, as given by the first equality of eq. (23.5.22) and illustrated in Figure 23.5.1. If  $\delta_l$  is small, then  $f_l(k)$  is also small in magnitude, and is almost purely real. This is realized in the leading Born approximation limit. In the opposite case that  $|f_l(k)|$  is maximal, so that the partial wave unitarity bound eq. (23.5.29) is saturated, then  $f_l(k)$  is at the top of the circle and is almost pure imaginary, equal to  $i/k$ .

Consider potentials that vanish completely<sup>†</sup> outside of a finite range, so that  $V(r) = 0$  for  $r > R$ . Then the exact (not just the large  $r$  limit) wavefunction for a stationary scattering state can be written as, taking into account the azimuthal symmetry,

$$\psi^+(\vec{r}) = \frac{1}{(2\pi)^{3/2}} \sum_{l=0}^{\infty} (2l+1) P_l(\cos \theta) F_l(r), \quad (23.5.31)$$

---

<sup>†</sup>This is done mostly for convenience; such potentials are not completely realistic. However, the results we are about to obtain are often a good approximation, provided that the potential  $V(r)$  is suitably weak for  $r > R$ .




Figure 23.5.1: Partial wave unitarity illustrated: the possible values of the partial wave amplitude  $f_l(k) = (e^{i2\delta_l} - 1)/2ik$  lie on a circle with radius  $1/2k$  in the upper-half complex plane, and are related to the phase shift angle  $\delta_l$  as shown. The Born approximation limit is realized near the bottom of the circle, for small  $\sin \delta_l$ , which gives small and nearly real  $f_l(k)$ . The partial wave unitarity bound eq. (23.5.29) is saturated for  $f_l(k) \approx i/k$ , near the top of the circle, with  $\sin^2 \delta_l = 1$ , so  $\delta_l = (n + 1/2)\pi$  for integer  $n$ .

where the radial wavefunction for the angular momentum  $l$  partial wave is

$$F_l(r) = \begin{cases} A_l(r) & (\text{for } r \leq R), \\ B_l(r) & (\text{for } r \geq R). \end{cases} \quad (23.5.32)$$

The exact form of the function  $A_l(r)$  may be quite difficult to obtain, since it depends on the scattering potential. However, since we are taking the potential to vanish for  $r \geq R$ , the function  $B_l(r)$  must be a superposition of the free-particle solutions that we found in section 10.2,

$$B_l(r) = c_l h_l^{(1)}(kr) + d_l h_l^{(2)}(kr), \quad (23.5.33)$$

where  $c_l$  and  $d_l$  are constant coefficients, and  $h_l^{(1)}$  and  $h_l^{(2)}$  are the spherical Hankel functions

$$h_l^{(1)}(kr) = j_l(kr) + in_l(kr) \sim \frac{e^{i(kr-l\pi/2)}}{ikr}, \quad (23.5.34)$$

$$h_l^{(2)}(kr) = j_l(kr) - in_l(kr) \sim -\frac{e^{-i(kr-l\pi/2)}}{ikr}, \quad (23.5.35)$$

where the large  $r$  asymptotic forms are shown. Matching to our scattering wavefunction for large  $r$ , eq. (23.5.21), we can solve for the constants  $c_l$  and  $d_l$  in terms of the phase shifts  $\delta_l$ . The result is

$$B_l(r) = \frac{1}{2}e^{i\pi l/2} \left[ e^{i2\delta_l} h_l^{(1)}(kr) + h_l^{(2)}(kr) \right], \quad (23.5.36)$$

or equivalently,

$$B_l(r) = e^{i\pi l/2} e^{i\delta_l} [\cos(\delta_l) j_l(kr) - \sin(\delta_l) n_l(kr)]. \quad (23.5.37)$$

To obtain scattering state solutions, one can now adopt the following strategy. First, the difficult part: solve the time-independent Schrödinger differential equation for  $A_l(r)$  in the region of non-zero potential,  $r \leq R$ . Because this is a second-order differential equation in  $r$ , in general

there will be two linearly independent solutions with arbitrary coefficients. To fix the coefficients, match these solutions to eq. (23.5.37) at  $r = R$ , using continuity of the wavefunction,

$$A_l(R) = B_l(R), \quad (23.5.38)$$

and, if the potential is finite at  $r = R$ , continuity of the derivative

$$A'_l(R) = B'_l(R). \quad (23.5.39)$$

Since the  $B_l(R)$  are given in eq. (23.5.37) in terms of known functions, one can solve for the  $\delta_l$ . To write the solution as compactly as possible, it is useful to define the quantities

$$\alpha_l \equiv A'_l(R)/A_l(R), \quad (23.5.40)$$

in which the overall normalization cancels. Then, since eqs. (23.5.38) and (23.5.39) imply

$$\alpha_l = B'_l(R)/B_l(R), \quad (23.5.41)$$

we obtain the general solution for the phase shift using eq. (23.5.37),

$$\tan \delta_l = \frac{\alpha_l j_l(kR) - k j'_l(kR)}{\alpha_l n_l(kR) - k n'_l(kR)}. \quad (23.5.42)$$

Summarizing: once the  $\alpha_l$  have been found using eq. (23.5.40), the phase shifts can be calculated from eq. (23.5.42), and then used in eq. (23.5.23) to obtain  $f_k(\theta)$ , which in turn gives the differential cross-section  $|f_k(\theta)|^2$ .

At leading order in an expansion in small  $kR$ , eqs. (10.2.14) and (10.2.15) for the spherical Bessel and Neumann functions can be used to find

$$\tan \delta_l = \frac{(kR)^{2l+1}(l - R\alpha_l)}{(2l-1)!!(2l+1)!!(l+1+R\alpha_l)}. \quad (23.5.43)$$

For  $l = 0, 1, 2, 3$ , and  $4$ , the denominator factor  $(2l-1)!!(2l+1)!!$  is respectively  $1, 3, 45, 1575$ , and  $99225$ , so the coefficient of  $(kR)^{2l+1}$  gets smaller rapidly as  $l$  increases. For low and intermediate energies (meaning as long as  $kR$  is not very large compared to  $1$ ), it is therefore justified to keep only the first few partial wave contributions, as we had promised to show earlier.

Let us now specialize to the low energy limit,  $kR \ll 1$ , where it is justified to neglect all but the  $s$ -wave phase shift  $\delta_0$  contribution, which behaves like  $\tan \delta_0 \propto k$  and provides an isotropic differential cross-section. Expanding the  $l = 0$  case of eq. (23.5.42) to the next-to-leading order in  $k$ , the result can be parameterized in the form

$$k \cot \delta_0 = -\frac{1}{a} + \frac{1}{2}r_0 k^2 + \cdots, \quad (23.5.44)$$

which defines length scales  $a$  and  $r_0$ , called the **scattering length** and the **effective range** respectively. In terms of  $R$  and  $\alpha_0 = A'_0(R)/A_0(R)$ , one finds

$$a = \frac{\alpha_0 R^2}{1 + \alpha_0 R} \quad (23.5.45)$$

and the determination of  $r_0$  is left to Exercise 23.8. The scattering length  $a$  can be either positive or negative depending on the sign of  $\alpha_0$ , and can be much larger in magnitude than the length scale  $R$  built into the potential, as we will see. In terms of these parameters, we can evaluate the low-energy, and therefore  $s$ -wave, cross-section from eq. (23.5.26),

$$\sigma = \frac{4\pi}{k^2} \frac{1}{1 + \cot^2 \delta_0} = \frac{4\pi a^2}{1 + (1 - r_0/a)a^2 k^2}, \quad (23.5.46)$$

where terms of higher order than  $k^2$  have been consistently neglected in the denominator. In the extreme low-energy approximation,  $\sigma$  has a simple dependence on the scattering length,

$$\sigma \approx 4\pi a^2. \quad (23.5.47)$$

In principle,  $a$ ,  $r_0$ , and higher order contributions to eq. (23.5.44) can be obtained experimentally from the energy dependence and interference effects in the low-energy limit.

## 23.6 Bound states, resonances, and poles in scattering amplitudes

The partial wave scattering amplitudes and cross-sections, viewed as functions of the energy  $E$  analytically continued to the complex domain, have an analytic structure that conveys information about the potential. As we will explain in this section, a pole (divergence) in the partial wave scattering amplitude corresponds either to a true bound state, or to a **resonance**, also known as a **quasi-bound state**, of the potential  $V(r)$ .

First, consider the case that the potential has one or more discrete bound states. We assume that the constant part of the potential has been fixed so that it vanishes at large distances,  $V(\infty) = 0$ , so that the condition for a bound state is an isolated eigenvalue  $E_{\text{bound}} < 0$  for the time-independent Schrödinger equation. To uncover the connection between such states and the scattering amplitude, consider the unbound scattering solutions for the wavefunction  $\psi_E(\vec{r})$  with positive  $E = \hbar^2 k^2/2m$ . Here  $k$  is the wavenumber far from the origin as discussed in section 23.1 and 23.2, with the asymptotic form of the wavefunction for large  $r$  as given in eq. (23.5.21). Suppose we try the trick of making the replacement  $k \rightarrow i\kappa$  in that wavefunction, where  $\kappa$  is now taken to be a real number. The solution so obtained, if it exists, will have energy  $E_{\text{bound}} = -\hbar^2 \kappa^2/2m < 0$  and therefore will be a bound state.

The only obstacle to constructing bound state wavefunction solutions in this way is that they will almost always be hopelessly non-normalizable, and therefore unphysical. To see this,

note that the asymptotic form for large  $r$  given in eq. (23.5.21) will be, after  $k \rightarrow i\kappa$ ,

$$\psi(\vec{r}) = \frac{1}{(2\pi)^{3/2}} \sum_{l=0}^{\infty} (2l+1) P_l(\cos \theta) [-S_l e^{-\kappa r} + (-1)^l e^{\kappa r}] / 2\kappa r, \quad (r \rightarrow \infty). \quad (23.6.1)$$

The term proportional to  $e^{-\kappa r}$  is sensible, but the term  $e^{\kappa r}$  evidently gives a probability density that diverges exponentially for large  $r$ . It appears that this wavefunction would describe a physically absurd state in which the probability to find the particle in any finite region is vanishingly small compared to the probability to find it farther away from the origin.

However, this solution can be salvaged for special, isolated values of the energy. Since we are considering a spherically symmetric potential, we can look for eigenstates of the total orbital angular momentum. Taking a term of fixed  $l$  in eq. (23.6.1), and re-normalizing the wavefunction by multiplying by a constant factor  $-2(2\pi)^{3/2}/(2l+1)S_l$ , we have

$$\psi_l(\vec{r}) = P_l(\cos \theta) \left( e^{-\kappa r} + (-1)^{l+1} \frac{e^{\kappa r}}{S_l} \right) / \kappa r, \quad (r \rightarrow \infty). \quad (23.6.2)$$

In order to re-interpret this wavefunction as a sensible (normalizable) energy and angular momentum eigenstate wavefunction at large  $r$ , it is necessary to eliminate the offending second term by requiring

$$S_l \rightarrow \infty. \quad (23.6.3)$$

We conclude that a true bound state for the potential  $V(r)$  with negative energy  $E_{\text{bound}}$  must correspond to a pole in the  $S$ -matrix element  $S_l = e^{i2\delta_l}$ , and thus in the scattering amplitude  $f_l(k) = (e^{i2\delta_l} - 1)/(2ik)$ , when viewed as a function of energy analytically continued to  $E = E_{\text{bound}} < 0$ . Due to the factor  $P_l(\cos \theta)$ , the orbital angular momentum quantum number of the bound state is  $l$ . So, one can relate the bound state energies, and even their angular momenta, to the poles in the partial-wave scattering amplitudes.

Since scattering solutions always have positive energy, and the bound states always have negative energy, one never actually hits the bound state pole directly in scattering experiments. Indeed, this is clear from the fact that for positive  $E$ , partial wave unitarity tells us that  $e^{i2\delta_l}$  has magnitude 1. Nevertheless, the presence of the bound states can be inferred from the functional dependence of  $e^{i2\delta_l}$  (and thus the cross-section) on  $E$ , particularly if  $-E_{\text{bound}}$  is small.

As an example, consider low-energy (small  $k$ ) scattering as discussed in equations (23.5.44)–(23.5.47) of the previous section. Taking  $k \rightarrow i\kappa$  in eq. (23.5.44), we see from eq. (23.5.22) that the  $s$ -wave scattering amplitude can have a pole at the solution to

$$\kappa = \frac{1}{a} + \frac{1}{2} r_0 \kappa^2, \quad (23.6.4)$$

with  $\kappa$  positive to ensure that the bound-state wavefunction is well-behaved at  $r \rightarrow \infty$ . Thus, if the scattering length  $a$  is found to be positive and not too small, one can predict a weakly bound state with  $\kappa \approx 1/a$  and energy close to

$$E_{\text{bound}} \approx -\frac{\hbar^2}{2ma^2}. \quad (23.6.5)$$

Including the effect of  $r_0$ , this is more precisely

$$E_{\text{bound}} \approx -\frac{\hbar^2 \kappa^2}{2m}, \quad (23.6.6)$$

where  $\kappa$  is the smaller solution to the quadratic equation (23.6.4), approximately

$$\kappa \approx \frac{1}{a} + \frac{r_0}{2a^2}. \quad (23.6.7)$$

The low-energy  $s$ -wave cross-section eq. (23.5.46) in the presence of a weakly bound state with energy  $E_{\text{bound}} < 0$  can now be written in the convenient form

$$\sigma_0 \approx \frac{4\pi a^2}{1 - E/E_{\text{bound}}}, \quad (23.6.8)$$

as dictated by the  $E = 0$  limit found in eq. (23.5.47), together with the requirement of a pole at  $E = E_{\text{bound}}$ .

Returning to the case of general (not necessarily small)  $k$ , the bound-state energies are not the only possible poles in the partial wave amplitude. To see this, let us look for peaks in the cross-section. The partial wave cross-section from eq. (23.5.26) can be rewritten as

$$\sigma_l = \frac{4\pi}{k^2} \frac{2l+1}{1 + \cot^2 \delta_l}, \quad (23.6.9)$$

where  $\cot^2 \delta_l$  is often quite large, as for example in the Born approximation limit. However, if it happens for some  $E$  that  $\cot \delta_l \approx 0$ , then  $\sigma_l$  will peak at that energy, saturating the partial wave unitarity bound. Suppose that  $E = E_{\text{res}}$  is a resonant energy that makes  $\cot \delta_l = 0$ . For energies that are close to this, we can expand  $\cot \delta_l$  in  $E - E_{\text{res}}$ , so that

$$\cot(\delta_l) = \cot(\delta_l) \Big|_{E=E_{\text{res}}} + (E - E_{\text{res}}) \frac{d}{dE} \cot(\delta_l) \Big|_{E=E_{\text{res}}} + \mathcal{O}(E - E_{\text{res}})^2. \quad (23.6.10)$$

The first term on the right vanishes by assumption, and we define the quantity  $\Gamma$ , with units of energy, according to

$$\frac{2}{\Gamma} = -\frac{d}{dE} \cot(\delta_l) \Big|_{E=E_{\text{res}}}, \quad (23.6.11)$$

so that, near the resonant energy,

$$\cot(\delta_l) \approx -2(E - E_{\text{res}})/\Gamma. \quad (23.6.12)$$






Figure 23.6.1: Behavior of the partial-wave cross-section  $\sigma_l$  and the phase shift  $\delta_l$  as a function of the scattering energy  $E$  near a resonance pole  $E_{\text{res}} - i\Gamma/2$ . In the left figure, the partial-wave cross-section is given by the Breit–Wigner lineshape. The phase shift  $\delta_l$  increases by  $\pi$  near each resonance, and for the lowest such resonance  $n$  in the right figure can be taken to be 0.

Using this in eq. (23.6.9) gives, for  $E$  close to  $E_{\text{res}}$ ,

$$\sigma_l \approx \frac{4\pi(2l+1)}{k^2} \frac{\Gamma^2/4}{(E - E_{\text{res}})^2 + \Gamma^2/4}. \quad (23.6.13)$$

Excluding the energy dependence of the  $1/k^2 = \hbar^2/2mE$  factor, this is the Breit–Wigner, or Lorentzian, lineshape with full width at half maximum (FWHM)  $\Gamma$  for the quasi-bound state resonance with angular momentum  $l$  and energy  $E_{\text{res}}$ . It is depicted in Figure 23.6.1.

Near  $E = E_{\text{res}}$ , eq. (23.6.12) tells us that the phase shift behaves like

$$\delta_l \approx \pi(n + 1/2) + \arctan\left(\frac{E - E_{\text{res}}}{\Gamma/2}\right), \quad (23.6.14)$$

for some integer  $n$ . This behavior is also sketched in Figure 23.6.1, showing that the phase shift rises through  $\pi(n + 1/2)$  near a resonant energy. These resonances correspond to poles in the partial wave scattering amplitude as a function of complex energy, at  $E = E_{\text{res}} - i\Gamma/2$ , because

$$f_l(k) = \frac{1}{k(\cot \delta_l - i)} \approx -\frac{\Gamma/2k}{E - E_{\text{res}} + i\Gamma/2}, \quad (23.6.15)$$

which follows from eq. (23.5.22). The assignment of complex energy  $E_{\text{res}} - i\Gamma/2$  is the tell-tale signature of an unstable state, as discussed at the end of section 5.6.

Figure 23.6.2 shows plausible potential shapes that could give rise to true bound state (left panel) and resonant quasi-bound state (right panel) poles in the scattering amplitude. Recall from our discussion at the end of section 10.4 that in three dimensions, the existence of a true bound state with energy  $E_{\text{bound}} < 0$  is not guaranteed unless the potential well is sufficiently




Figure 23.6.2: Sketches of example effective potentials  $V_{\text{eff}}(r)$  and energies for a true bound state with  $l = 0$  (left) and a quasi-bound state resonance (right), which will cause the scattering amplitude and the cross-section to have poles at  $E = E_{\text{bound}} < 0$  or at  $E = E_{\text{res}} - i\Gamma/2$ , respectively. In the latter case, the local minimum of the effective potential could be either negative (as shown) or positive, and could arise from the combination of an attractive potential  $V(r) < 0$  and the repulsive centrifugal core  $\hbar^2 l(l+1)/2mr^2$ .

deep. Unlike a true bound state, a quasi-bound state resonant energy  $E_{\text{res}}$  is positive. One way that such a quasi-bound state can arise is from the interplay between an attractive central potential  $V(r)$  and the orbital angular momentum contribution to the effective potential for the radial wavefunction in the Schrödinger equation (recall the discussion in section 10.1),

$$V_{\text{eff}}(r) = V(r) + \frac{\hbar^2 l(l+1)}{2mr^2}. \quad (23.6.16)$$

For  $l \neq 0$ , the centrifugal term is repulsive and grows stronger at small  $r$ . The potential shown in the right panel of Figure 23.6.2 has a finite barrier height between its minimum well and the asymptotic region at very large distances where it vanishes. Any corresponding state peaked inside the effective well with positive energy is not a true bound state; the wavefunction will always leak outside by tunneling. This is the reason for the terminology “quasi-bound state”. Since a quasi-bound state resonant pole occurs only for complex  $E$ , it again can never be hit directly in scattering experiments, but can be inferred from the complex analytic singularity structure of the scattering amplitudes. The smaller  $\Gamma$  is, the closer one can approach the resonance pole. However,  $e^{i2\delta_l}$  always has unit magnitude for physical (real positive) values of the scattering energy  $E$ .

A schematic map of the two kinds of poles in the complex  $E$  plane for  $e^{i2\delta_l}$ , and therefore for  $f_l(k)$ , are shown in Figure 23.6.3. They consist of true bound states on the negative real  $E$  axis and quasi-bound states below the positive real  $E$  axis.




Figure 23.6.3: Positions of poles of the partial wave amplitudes and  $S$ -matrix elements  $e^{i2\delta_l}$ , in the complex energy plane. Quasi-bound state resonances have poles in the analytic continuation to complex  $E$  below the real  $E$  axis, at  $E = E_{\text{res}} - i\Gamma/2$ . True bound states have poles on the negative real  $E$  axis, at  $E = E_{\text{bound}} < 0$ . Physical scattering energies are only on the positive real  $E$  axis. Therefore, these poles are never directly accessed in scattering, where  $e^{i2\delta_l}$  is always finite with unit magnitude, but their influence can be seen in the dependence of the cross-section on energy.

## 23.7 Examples of scattering from spherical potentials

Consider the scattering of a particle from a hard (impenetrable) sphere, so that the potential is

$$V(r) = \begin{cases} \infty & (\text{for } r \leq R), \\ 0 & (\text{for } r \geq R). \end{cases} \quad (23.7.1)$$

This is the quantum mechanical version of the classical scattering of a particle from a hard sphere, discussed in section 23.1. In this particular case, it is not necessary to use eqs. (23.5.39)–(23.5.42), because the solution for  $r \leq R$  is trivial,

$$\psi^+(\vec{r}) = 0 \quad (r \leq R), \quad (23.7.2)$$

corresponding to  $A_l(r) = 0$ . Meanwhile, for  $r \geq R$ , we write

$$\psi^+(\vec{r}) = \frac{1}{(2\pi)^{3/2}} \sum_{l=0}^{\infty} (2l+1) P_l(\cos \theta) B_l(r), \quad (23.7.3)$$

with  $B_l(r)$  given by eq. (23.5.37). To match to the vanishing solution at  $r = R$ , we need  $B_l(R) = 0$ , so

$$\tan \delta_l = j_l(kR)/n_l(kR), \quad (l = 0, 1, 2, \dots) \quad (23.7.4)$$

for each of the  $s$ -wave,  $p$ -wave,  $d$ -wave,  $\dots$  partial shifts. This is especially simple for the  $s$ -wave,

$$\tan \delta_0 = [\sin(kR)/kR] / [-\cos(kR)/kR] = -\tan(kR), \quad (23.7.5)$$

with the solution

$$\delta_0 = -kR. \quad (23.7.6)$$

Therefore, for  $l = 0$ ,

$$B_0(r) \propto \cos \delta_0 \frac{\sin(kr)}{kr} + \sin \delta_0 \frac{\cos(kr)}{kr} = \frac{1}{kr} \sin(kr + \delta_0) = \frac{1}{kr} \sin(k(r - R)). \quad (23.7.7)$$




Figure 23.7.1: The solid line shows the  $s$ -wave radial wavefunction  $rF_0(r)$  as a function of  $r$ , for scattering from a hard sphere with  $V = \infty$  in the shaded region  $r < R$ . The dashed line shows what  $rF_0(r)$  would be if the potential were absent, corresponding to free particle propagation. The potential pushes out the wavefunction by a distance  $R$ , giving a phase shift  $\delta_0 = -kR$ .

As illustrated in Figure 23.7.1, the outgoing  $s$ -wave is just shifted out by a distance  $R$ , compared to the free-particle solution.

More generally, consider all  $\delta_l$  in the low-energy limit  $kR \ll 1$  in which the wavenumber is much smaller than the inverse size of the hard sphere. In that case,

$$j_l(kR) \approx \frac{(kR)^l}{(2l+1)!!}, \quad n_l(kR) \approx -\frac{(2l-1)!!}{(kR)^{l+1}}, \quad (23.7.8)$$

so, from eq. (23.7.4),

$$\tan \delta_l = -\frac{(kR)^{2l+1}}{(2l-1)!!(2l+1)!!}, \quad (23.7.9)$$

which quickly becomes very small as  $l$  is increased. [Note that this result for  $\tan \delta_l$  could also be obtained simply as the  $\alpha_l \rightarrow \infty$  limit of eq. (23.5.43).]

In the extreme low energy limit  $kR \ll 1$ , only the  $s$ -wave contributes, resulting in

$$\frac{d\sigma}{d\Omega} \approx |f_0(k)|^2 = \frac{\sin^2 \delta_0}{k^2} = \frac{\sin^2(kR)}{k^2} \approx R^2, \quad (kR \ll 1). \quad (23.7.10)$$

Since this is isotropic (constant in both  $\phi$  and  $\cos \theta$ ), the total cross-section is

$$\sigma = \int d\Omega \frac{d\sigma}{d\Omega} = 4\pi R^2, \quad (kR \ll 1), \quad (23.7.11)$$

which is 4 times the classical result  $\sigma_{\text{classical}} = \pi R^2$  that we found in section eq (23.1.13).

In the opposite limit of very high scattering energy, many  $\delta_l$  can contribute. For  $kR \gg 1$ , one can show that  $\sigma \approx 2\pi R^2$ , which is still twice the classical result. This can be understood from the fact that since  $V(r)$  is discontinuous at  $r = R$ , no matter how big  $E$  is, the wavefunction varies sharply on length scales  $\ll 1/k$ . To get  $\sigma = \pi R^2 = \sigma_{\text{classical}}$ , it is necessary to take a

smoothed potential (a “softened hard sphere”), in which the the potential decreases to  $V = 0$  over a distance scale much smaller than  $R$  but much larger than  $1/k$ .

As another example, consider a finite well or barrier, with potential

$$V(r) = \begin{cases} V_0 & (\text{for } r \leq R), \\ 0 & (\text{for } r \geq R), \end{cases} \quad (23.7.12)$$

where the constant  $V_0$  can be either negative (an attractive well) or positive (a repulsive core potential). Let us only consider  $s$ -wave scattering, valid at low energy. Outside the potential’s range, the radial wavefunction is proportional to

$$F_0(r) = e^{i\delta_0} [\cos \delta_0 j_0(kr) - \sin \delta_0 n_0(kr)] = e^{i\delta_0} \frac{\sin(kr + \delta_0)}{kr}, \quad (r \geq R), \quad (23.7.13)$$

with the normalization chosen arbitrarily, and

$$\frac{\hbar^2 k^2}{2m} = E. \quad (23.7.14)$$

Inside the potential’s range, assuming  $E > V_0$ , the radial wavefunction is

$$F_0(r) = C j_0(k'r) = C \sin(k'r)/k'r, \quad (r \leq R), \quad (23.7.15)$$

with a relative normalization constant  $C$ , and

$$\frac{\hbar^2 k'^2}{2m} = E - V_0. \quad (23.7.16)$$

The solution  $n_0(k'r)$  is rejected here, because it is not normalizable at  $r = 0$ , as discussed in section 10.2. Now we match the solutions at  $r = R$ , to obtain an equation that determines  $C$ ,

$$C \sin(k'R)/k'R = e^{i\delta_0} \sin(kR + \delta_0)/kR. \quad (23.7.17)$$

We also match  $d\psi^+/dr$  at  $r = R$ , by specializing the convenient general results of eqs. (23.5.40) and (23.5.42),

$$\tan \delta_0 = \frac{(k/k') \sin(k'R) \cos(kR) - \cos(k'R) \sin(kR)}{(k/k') \sin(k'R) \sin(kR) + \cos(k'R) \cos(kR)}. \quad (23.7.18)$$

The preceding assumed  $E > V_0$ . If instead  $E < V_0$ , then  $k'$  will be imaginary, but eqs. (23.7.17) and (23.7.18) are valid after making the replacements  $\sin(k'R)/k' \rightarrow \sinh(\kappa R)/\kappa$  and  $\cos(k'R) \rightarrow \cosh(\kappa R)$ , where  $\hbar^2 \kappa^2/2m = V_0 - E$ .

Graphs of the  $l = 0$  wavefunction normalized by a factor of the radial coordinate are shown in Figure 23.7.2 for the attractive case  $V_0 < 0$  so that  $k' > k$ , and the repulsive case  $V_0 > 0$  so that  $k' < k$ , respectively. In the attractive case,  $\delta_0$  is positive, and the potential “pulls in” the




Figure 23.7.2: The solid lines show examples of the  $s$ -wave radial wavefunction  $rF_0(r)$  as a function of  $r$ , for scattering from a spherical-well potential in the shaded region, if the potential is attractive (top) or repulsive (bottom). The dashed line shows the corresponding result if the potential is absent. The potential pulls in (for the attractive case,  $\delta_0 > 0$ ) or pushes out (for the repulsive case,  $\delta_0 < 0$ ) the wavefunction by a distance  $|\delta_0|/k$ .

$l = 0$  wave by a shift  $\delta_0/k$  in the position of the nodes. In contrast, for the repulsive case,  $\delta_0$  is negative, and the potential can be thought of as “pushing out” the  $l = 0$  wave by an amount  $|\delta_0|/k$ . Things are a little more complicated for the partial waves with  $l > 0$ , and for potentials that are not piecewise constant. However, the essential qualitative feature remains that outgoing partial waves are pulled in by attractive potentials (because the Schrödinger equation dictates that the wavefunction oscillates faster in the region where the potential energy is smaller) and pushed out by repulsive potentials (because the wavefunction oscillates more slowly, or not at all, in regions with larger potential energy).

The phase shift  $\delta_0$  is small in magnitude for  $E \gg |V_0|$ . In the case of an attractive potential ( $V_0 < 0$ ), the phase shift increases as  $E$  decreases until it reaches  $\delta_0 = \pi/2$ , where the  $s$ -wave cross-section is as large as it can possibly be, for a given energy,

$$\sigma_{l=0} \Big|_{\max} = \frac{4\pi}{k^2} \sin^2(\pi/2) = \frac{2\pi\hbar^2}{mE}. \quad (23.7.19)$$

In this case, the scattering has saturated the partial wave unitarity bound. Decreasing  $E$  even more, eventually one may reach  $\delta_0 = \pi$ , for which

$$\sigma_{l=0} \Big|_{\min} = \frac{4\pi}{k^2} \sin^2(\pi) = 0. \quad (23.7.20)$$

This is the Ramsauer–Townsend effect; there is no scattering even though  $V(r)$  is non-negligible and attractive, because the de Broglie wavelength happens to match the characteristic length

scale of the potential. We have seen this type of behavior already in a one-dimensional scattering example, at the end of section 6.6, and in the real world it was first observed as suppressions in the cross-sections for electron scattering from inert gas atoms Ar, Kr, and Xe.

In the low-energy limit, one can find the scattering length and effective range defined in eq. (23.5.44), by expanding eq. (23.7.18) for small  $k$ . The results are

$$a = R - \tan(k_0 R)/k_0, \quad (23.7.21)$$

$$r_0 = R - \frac{R^3}{3a^2} - \frac{1}{ak_0^2}, \quad (23.7.22)$$

where we have defined

$$k_0 = \sqrt{-2mV_0}/\hbar. \quad (23.7.23)$$

If  $k_0 R$  happens to be slightly larger than  $\pi/2$ , then the scattering length can be large,  $a \gg R$ . This corresponds to the existence of an  $s$ -wave bound state with small binding energy  $E_{\text{bound}} \approx -\hbar^2/2ma^2$ , and then  $\sigma \approx 4\pi a^2/(1 - E/E_{\text{bound}})$  as discussed in the previous section.

## 23.8 Neutron-proton scattering and the deuteron

The low-energy scattering of neutrons and protons provides a practical illustration of some of the ideas discussed in sections 23.5–23.7. The masses of the proton and neutron are respectively  $m_p = 938.272 \text{ MeV}/c^2$  and  $m_n = 939.565 \text{ MeV}/c^2$ , with a reduced mass in the two-body problem (see section 4.2),

$$\mu = \frac{m_n m_p}{m_n + m_p} \approx 469.459 \text{ MeV}/c^2. \quad (23.8.1)$$

The potential between nucleons arises from the rather complicated strong nuclear force, but the spectrum of bound states for two nucleons is very simple. There are no  $pp$  or  $nn$  bound states, and there is only one bound state for the neutron and the proton. This is the deuteron ( $d$ ), a state with spin 1 and binding energy

$$-E_{\text{bound}}^d = 2.2246 \text{ MeV}, \quad (23.8.2)$$

which is determined by measuring the energy of the photon ( $\gamma$ ) emitted when a proton captures a neutron,  $n + p \rightarrow d + \gamma$ . The binding energy 2.2246 MeV is rather small in comparison to other nuclear binding energies per nucleon pair. Since the deuteron is weakly bound, there should be a pole in the neutron-proton scattering cross-section when extrapolated to negative energy.

Let us now understand the angular momentum and parity quantum numbers of the deuteron, as a prerequisite to figuring out its role in neutron-proton scattering. Since the full Hamiltonian

must be invariant under rotations, the energy eigenstates are also eigenstates of the square of the total angular momentum operator  $\vec{J}$ , obtained by combining the spins of both of the constituents  $\vec{S}_p$  and  $\vec{S}_n$  and their total orbital angular momentum  $\vec{L}$  in the center-of-mass frame. The statement that the deuteron has spin 1 means that the operator  $J^2$  has eigenvalue  $\hbar^2 J(J+1)$ , where  $J = 1$ . The neutron and proton each have spin  $1/2$ , so the possible total constituent spin combinations for the deuteron are  $S = 0$  and  $S = 1$ . The  $J = 1$  spin of the deuteron is then obtained by combining this with the total orbital angular momentum quantum number  $L$  in the center-of-mass frame, which can therefore only be  $L = 0, 1$ , or  $2$ . Now, the magnetic dipole moment of the deuteron is known experimentally to be  $\mu_d = 0.85744\mu_N$ , which is very close to the sum of the dipole moments of the neutron and proton,  $\mu_p + \mu_n = 2.79285\mu_N - 1.91304\mu_N = 0.87981\mu_N$ . This indicates that the deuteron magnetic moment is very nearly realized when the spins are aligned in the  $S = 1$  state, without much contribution from the orbital motion of the charged proton. Therefore, the deuteron must have predominantly  $L = 0$ . However, there are also two clear experimental indications that it cannot be a pure  $L = 0$  eigenstate. First, there is the small but significant discrepancy in the magnetic moment sum noted above. Second, the deuteron is found experimentally to have a non-zero electric quadrupole moment, which would be inconsistent with the perfect spherical symmetry of a pure  $L = 0$  eigenstate. The strong interaction Hamiltonian responsible for binding the deuteron is known (from its more fundamental formulation, quantum chromodynamics, or QCD) to commute with parity. Therefore, eq. (8.7.8) applies, and it must be possible to assign the deuteron a definite parity eigenvalue  $\eta_d = (-1)^L \eta_p \eta_n$ . Here,  $\eta_p$  and  $\eta_n$  are the intrinsic parities of the proton and neutron, which are conventionally taken to be  $\eta_p = \eta_n = 1$ . To have a definite parity eigenvalue  $\eta_d = \pm 1$ , the deuteron must be a superposition of states with either all  $L$  even, or all  $L$  odd. Since we already know that the state is mostly  $L = 0$ , this implies  $\eta_d = +1$  and rules out the possibility of any  $L = 1$  component. Therefore, the deuteron must be a linear combination of  $L = 0$  and  $L = 2$ . In order to give  $J = 1$ , it follows that the combination of constituent spins can only be pure  $S = 1$ . In spectroscopic notation, the deuteron state is predominantly  ${}^{2S+1}L_J = {}^3S_1$ , but with a few percent  ${}^3D_1$  component.

For our present purposes, the most important feature of the previous paragraph is that the deuteron is a pure  $S = 1$  combination. The absence of a bound state in the  $S = 0$  combination shows that the strong nuclear force is spin-dependent, which means that we should also keep track of the  $S = 0$  and  $S = 1$  combinations separately in scattering. The data for low-energy scattering of neutrons and protons in the  $S = 0$  and  $S = 1$  combinations yield scattering lengths



and effective ranges, as defined in eq. (23.5.44), of

$$a^{S=1} = 5.42 \text{ fm}, \quad r_0^{S=1} = 1.75 \text{ fm}, \quad (23.8.3)$$

$$a^{S=0} = -23.7 \text{ fm}, \quad r_0^{S=0} = 2.7 \text{ fm}, \quad (23.8.4)$$

in units of  $1 \text{ fm} = 10^{-15} \text{ meters}$ . The positive scattering length in the  $S = 1$  channel supports the hypothesis of a bound state. Solving eq. (23.6.4) gives a prediction for the bound state of

$$\kappa = 0.231 \text{ fm}^{-1}, \quad (23.8.5)$$

which then yields the estimate

$$E_{\text{bound}} = -\frac{\hbar^2 \kappa^2}{2\mu} = -2.22 \text{ MeV}, \quad (23.8.6)$$

in good agreement<sup>†</sup> with the experimental deuteron binding energy quoted in eq. (23.8.2). Now, using the facts that the  $E = 0$  cross-section for  $s$ -wave scattering is  $4\pi a^2$ , and that it must have a pole at  $E = E_{\text{bound}}^d$ , we have as in eq. (23.6.8),

$$\sigma^{S=1} = \frac{4\pi (a^{S=1})^2}{1 - E/E_{\text{bound}}^d} \approx \frac{3.69 \text{ b}}{1 + E/2.2246 \text{ MeV}}, \quad (23.8.7)$$

where we have used the traditional nuclear physics unit of cross-section

$$1 \text{ b} = 100 \text{ fm}^2 = 10^{-28} \text{ meters}^2, \quad (23.8.8)$$

called<sup>‡</sup> a **barn**.

In the  $S = 0$  channel for neutron-proton scattering, the fact that the scattering length  $a^{S=0}$  is negative confirms that there is no bound state. Still, we can write the cross-section in a form similar to eq. (23.8.7), by using eq. (23.5.46),

$$\sigma^{S=0} = \frac{4\pi (a^{S=0})^2}{1 + (ka^{S=0})^2(1 - r_0^{S=0}/a^{S=0})} \approx \frac{70.6 \text{ b}}{1 + E/(0.066 \text{ MeV})}. \quad (23.8.9)$$

However, it should be emphasized that there really is no neutron-proton bound state with energy near  $-0.066 \text{ MeV}$ . Such a pole in the scattering cross-section at negative real  $E$ , but with a negative scattering length  $a$  and therefore not associated with an actual bound state, is

---

<sup>†</sup>Neglecting  $r_0^{S=1}$  here would give much worse agreement. However, it must be admitted that the excellence of the agreement as presented here is partly accidental. Since eq. (23.5.44) was truncated beyond order  $k^2$ , rather than solving the quadratic equation (23.6.4) for  $\kappa$ , one could just as consistently approximate it by eq. (23.6.7), which would have given instead  $\kappa = 0.214 \text{ fm}^{-1}$  and thus  $E_{\text{bound}} = -1.91 \text{ MeV}$ .

<sup>‡</sup>The origin of the unit name refers to the fact that, by nuclear physics standards, this is not a small area, jokingly considered as easy to hit as “the broad side of a barn” in idiomatic North American English. Collider experiments at the present high-energy frontier often discuss cross-sections in units of picobarns or femtobarns.

sometimes called a **virtual bound state**. It can be interpreted as an indication that if the  $S = 0$  neutron-proton potential were just slightly more attractive, then the scattering length would be positive and a bound state would exist. Such virtual bound state poles should not be confused with the quasi-bound state resonances with positive energy discussed in section 23.6.

The initial neutron and proton spins are often random and unmeasured, so that the four spin states (1 for  $S = 0$  and 3 for  $S = 1$ ) are equally likely. Averaging over them, the prediction for the total neutron-proton cross-section in the  $E \rightarrow 0$  limit is

$$\sigma = \frac{1}{4}\sigma^{S=0} + \frac{3}{4}\sigma^{S=1} = 20.4 \text{ b}, \quad (23.8.10)$$

in good agreement with experimental observation.

The data eqs. (23.8.3) and (23.8.4) can also be used to make a crude model for the neutron-proton potential in the form of a spherical well as in (23.7.12), by solving eqs. (23.7.21)–(23.7.23) numerically for  $R$  and  $k_0 = \sqrt{-2\mu V_0}/\hbar$ . The results are

$$R^{S=1} = 2.07 \text{ fm}, \quad k_0^{S=1} = 0.91 \text{ fm}^{-1}, \quad V_0^{S=1} = -34 \text{ MeV}, \quad (23.8.11)$$

$$R^{S=0} = 2.59 \text{ fm}, \quad k_0^{S=0} = 0.58 \text{ fm}^{-1}, \quad V_0^{S=0} = -14 \text{ MeV}. \quad (23.8.12)$$

From the analysis of section 10.4, the condition for such a spherical-well potential to have  $n$  bound states with  $l = 0$  can be written as [see eq. (10.4.12), and recall that  $V_0$  and  $a$  there are  $-V_0$  and  $R$  here]

$$k_0 R / \pi > n - 1/2. \quad (23.8.13)$$

Putting in the numbers, the  $S = 1$  potential model predicts exactly one bound state, and the  $S = 0$  potential model barely misses having a bound state, in accord with the observed facts. The deuteron is surprisingly weakly bound, in the sense that the binding energy (2.2246 MeV) is more than an order of magnitude smaller than the depth of the potential (34 MeV).

## 23.9 Scattering of identical particles

So far, we have neglected the possibility that the scattering particles might be identical. To remedy this, first consider the scattering of two identical bosons with no spin (for example,  $\alpha$  particles, also known as  ${}^4\text{He}$  nuclei). The total wavefunction will then be of the form

$$\Psi(\vec{r}_1, \vec{r}_2) = e^{i\vec{p} \cdot (\vec{r}_1 + \vec{r}_2)/2\hbar} \psi(\vec{r}), \quad (23.9.1)$$

where, following eqs. (4.2.4) and (4.2.5),  $\vec{p}$  is the total momentum of the two particles,  $(\vec{r}_1 + \vec{r}_2)/2$  is the center-of-mass position, and

$$\vec{r} = \vec{r}_1 - \vec{r}_2 \quad (23.9.2)$$

is the relative position vector. By choosing the center-of-mass reference frame, we can take  $\vec{p} = 0$ . Because the particles are spinless bosons, this wavefunction must be symmetric under  $\vec{r}_1 \leftrightarrow \vec{r}_2$ , so the relative-position wavefunction must obey

$$\psi(\vec{r}) = \psi(-\vec{r}), \quad (23.9.3)$$

in other words, it must have even parity. Therefore, instead of a scattering wavefunction proportional to  $e^{ikz} + f_k(\theta) \frac{e^{ikr}}{r}$ , we must have (dropping the overall normalization in this section)

$$\psi(\vec{r}) = e^{ikz} + e^{-ikz} + [f_k(\theta) + f_k(\pi - \theta)] \frac{e^{ikr}}{r} \quad (23.9.4)$$

for the scattering wavefunction in the center-of-momentum frame. In this way, both incoming particles, and both outgoing particles, are described on an equal footing. The differential cross-section is therefore of the form

$$\frac{d\sigma}{d\Omega} = |f_k(\theta) + f_k(\pi - \theta)|^2 = |f_k(\theta)|^2 + |f_k(\pi - \theta)|^2 + 2\text{Re}[f_k^*(\theta)f_k(\pi - \theta)]. \quad (23.9.5)$$

This incorporates the fact that if one outgoing particle is detected at angle  $\theta$ , then by momentum conservation the other must be found at angle  $\pi - \theta$ . It is a necessary feature that the differential cross-section should be equal at those two angles, due to the intrinsic indistinguishability of the particles.

As a consequence, there must be constructive interference in the amplitudes for identical spinless bosons for right-angle scattering,

$$\left. \frac{d\sigma}{d\Omega} \right|_{\theta=\pi/2} = 4|f_k(\pi/2)|^2. \quad (23.9.6)$$

Also, only even- $l$  partial waves can contribute in the identical boson scattering case, because

$$f_k(\theta) + f_k(\pi - \theta) = \sum_{l=0}^{\infty} (2l+1) [P_l(\cos \theta) + P_l(\cos(\pi - \theta))] f_l(k) \quad (23.9.7)$$

$$= 2 \sum_{\text{even } l} (2l+1) P_l(\cos \theta) f_l(k), \quad (23.9.8)$$

where the terms with odd  $l$  have canceled because  $P_l(\cos(\pi - \theta)) = (-1)^l P_l(\cos \theta)$ .

Now consider the case of scattering of electrons, or other identical spin-1/2 fermions. This can be divided into two cases, depending on the total spin quantum number  $S$ .

First, suppose that  $S = 0$ . Since that is an antisymmetric spin state, the spatial wavefunction must be symmetric under exchange of the labels 1, 2, so  $\psi(\vec{r}) = \psi(-\vec{r})$ , and

$$\frac{d\sigma^{S=0}}{d\Omega} = |f_k(\theta) + f_k(\pi - \theta)|^2, \quad (23.9.9)$$

and so all of the remarks just made for scattering of identical bosons go through as before.

If instead the total spin state is the symmetric  $S = 1$  combination, then the spatial wavefunction must be antisymmetric under exchange of the labels 1, 2, so  $\psi(\vec{r}) = -\psi(-\vec{r})$  with odd parity. Therefore, the scattering wavefunction must be proportional to

$$\psi(\vec{r}) = e^{ikz} - e^{-ikz} + [f_k(\theta) - f_k(\pi - \theta)] \frac{e^{ikr}}{r}, \quad (23.9.10)$$

which implies

$$\frac{d\sigma}{d\Omega}^{S=1} = |f_k(\theta)|^2 + |f_k(\pi - \theta)|^2 - 2\text{Re}[f_k^*(\theta)f_k(\pi - \theta)]. \quad (23.9.11)$$

It follows that there is perfect destructive interference for right-angle scattering,

$$\left. \frac{d\sigma}{d\Omega}^{S=1} \right|_{\theta=\pi/2} = 0 \quad (23.9.12)$$

for identical fermions in the symmetric spin state. Furthermore, only partial waves with odd  $l$  contribute.

In many cases, the fermion spins are random and unmeasured. If so, then four spin states (1 for  $S = 0$  and 3 for  $S = 1$ ) are equally likely, so the observed unpolarized cross-section for identical fermions is the weighted average:

$$\left. \frac{d\sigma}{d\Omega} \right|_{\text{unpolarized}} = \frac{1}{4} \frac{d\sigma}{d\Omega}^{S=0} + \frac{3}{4} \frac{d\sigma}{d\Omega}^{S=1} \quad (23.9.13)$$

$$= |f_k(\theta)|^2 + |f_k(\pi - \theta)|^2 - \text{Re}[f_k^*(\theta)f_k(\pi - \theta)], \quad (23.9.14)$$

where the absence of a factor of 2 in front of the interference term is not a typographical error. For right angle scattering, the result is partial destructive interference.

In all of the cases of scattering of identical particles, it is important to note that when integrating the differential cross-section to get the total cross-section, one must include a factor of 1/2 to avoid double counting. This is because a state with one particle at  $(\theta, \phi)$  also has an identical particle at  $(\pi - \theta, \phi + \pi)$ . Therefore, the state specified by the presence of a particle at  $(\theta, \phi)$  and the state specified by  $(\pi - \theta, \phi + \pi)$  are actually the same state, and should not be counted twice. So, one has

$$\sigma = \frac{1}{2} \int d\Omega \frac{d\sigma}{d\Omega} \quad (\text{identical particles}) \quad (23.9.15)$$

for both the identical boson and identical fermion case. Note that this factor of 1/2 for identical particles comes in at the level of the total cross-section, not the differential cross-section.

## 23.10 Exercises

**Exercise 23.1.** Particles of mass  $m$  and incident momentum  $\vec{p} = \hbar k \hat{z}$  scatter from a potential

$$V(r) = \begin{cases} V_0 & (\text{for } r < a) \\ 0 & (\text{for } r > a), \end{cases} \quad (23.10.1)$$

where  $V_0$  and  $a$  are constants. Apply the first-order Born approximation to:

- (a) Find the differential scattering cross-section for small  $|V_0|$ . Write your answer in terms of  $q = 2k \sin(\theta/2)$ . Check that your answer has units of area.
- (b) Show that in the limit of small  $ka$ , the differential cross-section found in part (a) is constant with respect to the scattering angle, and show that the total cross-section is  $\sigma = nV_0^2 a^6$  where  $n$  is a quantity that you will find. (Check the units.)
- (c) Working now to the next-to-leading order in an expansion in small  $ka$ , show that the differential cross-section has the form  $d\sigma/d\Omega = b + c \cos(\theta)$ , and determine the quantities  $b$  and  $c$ .

**Exercise 23.2.** Particles of mass  $m$  and momentum  $\vec{p} = \hbar k \hat{z}$  scatter from a potential  $V = V_0 e^{-r/a}$ . Use the leading Born approximation to find the differential cross-section  $d\sigma/d\Omega$ , and integrate it to find the total cross-section.

**Exercise 23.3.** Consider the scattering of a spinless point particle with mass  $m$ , momentum  $\vec{p} = \hbar k \hat{z}$ , and charge  $-e$ , due to a point charge  $Ze$  fixed at the origin, which is screened at large distances by a uniform thin spherical shell of opposite charge at  $r = R$ , so that

$$V(r) = \begin{cases} -Ze^2/r & (\text{for } r < R), \\ 0 & (\text{for } r > R). \end{cases} \quad (23.10.2)$$

Use the first-order Born approximation for scattering to:

- (a) Find the differential cross section.
- (b) Find the lowest energy  $E$  for which  $d\sigma/d(\cos \theta)$  vanishes for backward scattering ( $\theta = \pi$ ).
- (c) Find the differential cross section and the total cross section in the low energy limit, keeping the leading and next-to-leading non-zero contributions.

**Exercise 23.4.** Suppose that the differential cross-section for particles of momentum  $\hbar \vec{k}$  to scatter from a potential  $V_0(\vec{r})$  centered at the origin is known to be  $d\sigma_0/d\Omega$  in the leading Born approximation. Now consider scattering from  $N$  such potentials centered at positions  $\vec{a}_n$ , so that the total potential is  $V(\vec{r}) = \sum_{n=1}^N V_0(\vec{r} - \vec{a}_n)$ . Show that in the leading Born approximation the differential cross-section is

$$\frac{d\sigma}{d\Omega} = \left| \sum_{n=1}^N \exp \left( i(\vec{k} - \vec{k}') \cdot \vec{a}_n \right) \right|^2 \frac{d\sigma_0}{d\Omega}. \quad (23.10.3)$$

**Exercise 23.5.** A beam of electrons with momentum  $\vec{p} = \hbar k \hat{z}$  scatters from the classical potential  $V(r)$  produced by a hydrogen atom in its ground state, which (from Gauss' law) obeys

$$\frac{dV}{dr} = \frac{q_e^2}{r^2} \left[ 1 - \int_0^r dr' 4\pi r'^2 |\psi_{1,0,0}(r')|^2 \right]. \quad (23.10.4)$$

Here, the electron charge is  $q_e$ , to distinguish it from the base of the natural logarithms.

(a) Obtain  $dV/dr$  from the above equation by doing the integral, and deduce that  $V(r) = -q_e^2(1/r + A/a_0)e^{-Br/a_0}$ , where  $A$  and  $B$  are certain constants that you will determine. [Hint: take the derivative of this form of  $V(r)$  and match it to the  $dV/dr$  that you found.]

(b) Find the differential cross section in the Born approximation, ignoring spin. Show that it can be written as  $d\sigma/d\Omega = n_1(N_2 + a_0^2 q^2)^2/(N_3 + a_0^2 q^2)^4$ , where  $n_1$  is a quantity that you will find, and  $N_2$  and  $N_3$  are certain positive integers, and  $q = 2k \sin(\theta/2)$ .

(c) In the low energy limit, what is  $d\sigma/d\Omega$ ? How does it depend on  $\theta$ ?

(d) In the high energy limit, show that  $d\sigma/d\Omega$  approaches the Rutherford differential cross section. (Can you explain why, even though the atom is electrically neutral?)

(e) Find the total cross section. Does it diverge, as for Rutherford scattering?

**Exercise 23.6.** For scattering of particles with mass  $m$  and momentum  $\vec{p} = \hbar k \hat{z}$  from a spherically symmetric potential  $V(r)$ , show that in the leading Born approximation the phases shifts in the partial wave expansion are

$$\delta_l = N \frac{mk}{\hbar^2} \int_0^\infty dr r^2 V(r) (j_l(kr))^2, \quad (23.10.5)$$

where  $N$  is a certain integer that you will discover.

**Exercise 23.7.** The differential cross-section of a beam of spinless particle of mass  $m$  with fixed low kinetic energy  $E$  on a target of spherically symmetric atoms is found to be

$$\frac{d\sigma}{d(\cos \theta)} = \frac{\pi \hbar^2}{mE} [c_0 + c_1 \cos(\theta) + c_2 \cos^2(\theta)]. \quad (23.10.6)$$

where  $c_0$ ,  $c_1$ , and  $c_2$  depend on  $E$  but not on  $\theta$ . Assume that the target is very small compared to the experimental apparatus, that multiple scatterings within the target are not an issue, and that the results should be interpreted as a combination of  $s$ -wave and  $p$ -wave scattering.

(a) Find real equations relating the phase shifts  $\delta_0$  and  $\delta_1$  to the measured quantities  $c_0$ ,  $c_1$ ,  $c_2$ .

(b) Use partial wave unitarity to find theoretical upper bounds on  $c_0$  and  $c_2$ .

(c) What are the theoretical upper and lower bounds on  $c_1$ ? [Hint: both upper and lower bounds are realized by phase shifts satisfying  $\delta_0 = \pi n_0/6$  and  $\delta_1 = \pi n_1/6$  where  $n_0$  and  $n_1$  are positive integers less than 6. You may use this fact without proving it.]

**Exercise 23.8.** The low-energy ( $s$ -wave) limit of scattering from a potential that vanishes outside of some range  $R$  is treated at the end of section 23.5. By expanding eq. (23.5.42) with  $l = 0$ , verify eq. (23.5.44) with the scattering length  $a$  given in eq. (23.5.45), and show that the effective range is  $r_0 = nR(1 - 1/\alpha_0 R + 1/(\alpha_0 R)^2)$  where  $n$  is a certain positive rational number that you will discover.

## 24 Subsystems: entanglement, evolution, and generalized measurements

### 24.1 Open and bipartite systems and entanglement

An **open system** is a quantum system that is part of a larger one. The rest of the larger system may be referred to as the **environment**, and we often want to study the dynamics of the open system in such a way that its interactions with the environment are to be simplified or averaged over in some way. Typically, the dynamics of the environment are not completely accessible to us, or are too complicated to understand in full detail, but will still have important impacts on the open system of primary interest.

More generally, we wish to consider **bipartite quantum systems**, defined to be those whose Hilbert space is the tensor product of two Hilbert spaces for subsystems  $a$  and  $b$ , so that  $\mathcal{H}_{ab} = \mathcal{H}_a \otimes \mathcal{H}_b$ . Often, subsystem  $a$  is the open system whose behavior we want to understand, and  $b$  is its environment (perhaps the whole rest of the universe). Another interpretation is that we might consider  $b$  to contain a quantum measurement apparatus that works through its correlations with the object of measurement, subsystem  $a$ . Yet another interpretation is that  $a$  and  $b$  might be associated with observers Alice and Bob who are in different locations, and therefore each have access to only part of a quantum state. Understanding the quantum correlations that arise between two subsystems gives insights into the dynamical origins of measurements, the emergence of macroscopic classical-like behavior in quantum mechanics, and the properties of information in quantum mechanics.

To describe the quantum physics of an open system, it is natural to use the density matrix operator description introduced in section 3.5. To understand why, take the simple example of the entangled state that we introduced in eq. (14.1.1) and used in our discussion of the EPR problem, which we re-write here in the notation

$$|\psi\rangle = \frac{1}{\sqrt{2}} \left( |\uparrow\rangle_a \otimes |\downarrow\rangle_b - |\downarrow\rangle_a \otimes |\uparrow\rangle_b \right). \quad (24.1.1)$$

Consider how Alice will describe the situation when she can observe only system  $a$ , assuming no communication with Bob. No matter what spin components of her particle Alice measures, if the experiments are repeated many times, the results are perfectly described as a random ensemble with equal probability  $1/2$  to be a spin-up or spin-down state. Considered as an open system, Alice can therefore describe her particle's spin  $a$  using a density operator

$$\rho_a = \frac{1}{2} \left( |\uparrow\rangle \langle\uparrow| + |\downarrow\rangle \langle\downarrow| \right). \quad (24.1.2)$$

Even if Bob is busily making measurements of spin  $b$ , as long as Alice does not (yet) know the results, she will still describe her measurements in terms of  $\rho_a$ . This density operator is that of



a completely random ensemble, and contains no information. In this way, the density operator formalizes the absence of instantaneous communication to Alice due to Bob's local interactions with spin  $b$ .

In contrast, suppose the combined state is instead a non-entangled ket, say

$$|\psi'\rangle = |\uparrow\rangle_a \otimes |\downarrow\rangle_b \quad (24.1.3)$$

for example. In this case, for the purposes of Alice's experiments, the subsystem  $a$  is always spin up, so the density operator is

$$\rho'_a = |\uparrow\rangle\langle\uparrow|, \quad (24.1.4)$$

corresponding to a pure ensemble.

As these examples suggest, if a combined system  $ab$  is in a product (non-entangled) pure state, then the subsystem  $a$  can always be described by a pure ensemble, but if the combined system is in an entangled state, then  $a$  is described by a mixed ensemble. We will now proceed to understand this in the general case.

Let us denote orthobases for the two subsystem Hilbert spaces as  $|\varphi_i\rangle_a$  with  $i = 1, \dots, d_a$  and  $|\chi_j\rangle_b$  with  $j = 1, \dots, d_b$ . Then the  $d_a d_b$  kets  $|\varphi_i\rangle \otimes |\chi_j\rangle$  form an orthobasis for  $\mathcal{H}_{ab}$ . Now, suppose Alice is measuring an observable  $A$  that operates only on  $\mathcal{H}_a$  and has eigenvalues (the possible measurement outcomes)  $\alpha$ . These outcomes are associated with projection operators

$$P_\alpha \otimes I, \quad (24.1.5)$$

where  $P_\alpha$  acts on subsystem  $a$ , and the identity operator acts on subsystem  $b$ . To be as general as possible, suppose that we have prepared a mixed ensemble of states of the system  $ab$ , which is therefore described by a density operator  $\rho_{ab}$ . Using eq. (3.5.11), the probability of getting the measurement result  $\alpha$  for  $A$  is

$$\mathcal{P}(\alpha) = \text{Tr}[\rho_{ab}(P_\alpha \otimes I)] = \sum_i \sum_j \left( \langle \varphi_i | \otimes \langle \chi_j | \right) \rho_{ab} (P_\alpha \otimes I) \left( | \varphi_i \rangle \otimes | \chi_j \rangle \right). \quad (24.1.6)$$

To put this into a nicer form, define the **reduced density operator** for subsystem  $a$ ,

$$\rho_a = \text{Tr}_b[\rho_{ab}] = \sum_j \langle \chi_j | \rho_{ab} | \chi_j \rangle. \quad (24.1.7)$$

Here we have taken the trace only over  $\mathcal{H}_b$ , called a **partial trace**. The notation is a bit tricky, since the objects  $\langle \chi_j | \rho_{ab} | \chi_j \rangle$  are matrix elements with respect to the Hilbert space  $\mathcal{H}_b$ , but are still operators in  $\mathcal{H}_a$ . Then eq. (24.1.6) becomes simply

$$\mathcal{P}(\alpha) = \text{Tr}_a[\rho_a P_\alpha] = \sum_i \langle \varphi_i | \rho_a P_\alpha | \varphi_i \rangle. \quad (24.1.8)$$

Thus,  $\rho_a$  as defined by eq. (24.1.7) contains all of the information that is available to Alice<sup>†</sup> by only observing subsystem  $a$ . It follows from eq. (24.1.8) that the average result of measurements of  $A$  is

$$\overline{A} = \text{Tr}_a[\rho_a A]. \quad (24.1.9)$$

These results are in accord with eqs. (3.5.11) and (3.5.12).

It is also a short exercise to check that  $\rho_a = \text{Tr}_b(\rho_{ab})$  obeys the three crucial properties required of any density operator. First, it is a Hermitian operator on  $\mathcal{H}_a$ . Second, it has unit trace,  $\text{Tr}_a(\rho_a) = 1$ . Third, it is a **positive operator**. In general, an operator  $A$  is said to be positive<sup>‡</sup> if  $\langle \psi | A | \psi \rangle \geq 0$  for any ket  $|\psi\rangle$ . More generally, if we wish to describe an open system  $a$  without any direct reference to a particular environment system, it is essential to require that it is described by a density operator  $\rho_a$  satisfying these three properties.

Consider the special case of a pure state for the combined system. The general form of such a state is

$$|\psi\rangle = \sum_i \sum_j c_{ij} |\varphi_i\rangle \otimes |\chi_j\rangle, \quad (24.1.10)$$

where  $c_{ij}$  are complex coefficients, which form the elements of a  $d_a \times d_b$  complex matrix  $C$ . Now, the **singular value decomposition** of linear algebra asserts that any complex matrix can be written in the form  $C = U^\dagger D V$ , where  $U$  and  $V$  are unitary  $d_a \times d_a$  and  $d_b \times d_b$  matrices, and  $D$  is a  $d_a \times d_b$  diagonal matrix, whose only non-zero entries are non-negative real numbers (called the **singular values** of  $C$ ) on the diagonal. Since the singular values appear only on the diagonal of  $D$ , the number of them that are non-zero is at most  $\min(d_a, d_b)$ . In our case,  $U$  simply acts as a change of orthobasis in the Hilbert space  $\mathcal{H}_a$ , and  $V$  similarly acts as a change of orthobasis in the Hilbert space  $\mathcal{H}_b$ . So, writing the singular values as  $\sqrt{p_k}$  with  $k = 1, \dots, \min(d_a, d_b)$ , we have just proved that the state can always be written in a particularly useful way:

**Theorem 24.1.1. (Singular value decomposition of a pure state in a bipartite system)**

*Consider a pure state  $|\psi\rangle$  of a bipartite system with subsystems  $a$  and  $b$  with Hilbert space dimensions  $d_a$  and  $d_b$ . Then one can always find a particular choice of orthobases  $|\varphi_i\rangle_a$  with  $i = 1, \dots, d_a$  and  $|\chi_j\rangle_b$  with  $j = 1, \dots, d_b$ , such that*

$$|\psi\rangle = \sum_{k=1}^{\min(d_a, d_b)} \sqrt{p_k} |\varphi_k\rangle \otimes |\chi_k\rangle, \quad (24.1.11)$$

*where  $p_k$  are non-negative real numbers, with  $\sum_k p_k = 1$  if the state is normalized to unity.*

---

<sup>†</sup>Conversely, Bob, who makes observations only on subsystem  $b$ , will describe his experiences using  $\rho_b = \text{Tr}_a[\rho_{ab}] = \sum_i \langle \varphi_i | \rho_{ab} | \varphi_i \rangle$ . Of course, the observables he can measure do not include  $A$ .

<sup>‡</sup>A better terminology would be “non-negative”, but it is hard to fight tradition.

The choice of bases needed to accomplish eq. (24.1.11) depends on the state  $|\psi\rangle$ . Equation (24.1.11) is also called the **Schmidt decomposition**, and the number of non-zero singular values is called the **Schmidt number**. A useful corollary relates the Schmidt number to the entanglement of the state:

**Theorem 24.1.2. (Schmidt number reveals entanglement)** *A pure state  $|\psi\rangle$  of a bipartite system is entangled if, and only if, the Schmidt number of non-zero coefficients  $\sqrt{p_k}$  in its singular value decomposition exceeds 1.*

Proof: the key point is that the list of singular values of a complex matrix  $C$  is uniquely determined. So, if there are at least 2 non-zero singular values, then there can be no way to choose a ket from each subsystem such that  $|\psi\rangle$  is their tensor product; therefore it is entangled by definition. Conversely, if only one of the  $p_k$  is non-zero, then  $|\psi\rangle$  is a product state, and so it is not entangled by definition. (If all of the  $p_k$  are zero, then  $|\psi\rangle$  is the null ket, of course.)  $\square$

In terms of the singular value decomposition coefficients and orthobases, the pure state  $|\psi\rangle$  of the combined system has a reduced density operator

$$\rho_a = \text{Tr}_b(|\psi\rangle\langle\psi|) = \sum_j \langle\chi_j|\psi\rangle\langle\psi|\chi_j\rangle = \sum_k p_k |\varphi_k\rangle\langle\varphi_k|. \quad (24.1.12)$$

So, the non-zero ensemble probabilities  $p_k$  of  $\rho_a$  in this basis are equal to the squares of the non-zero Schmidt coefficients of the combined system state. It follows that  $\rho_a$  describes a mixed state (more than one non-zero  $p_k$ ) if the combined system state  $|\psi\rangle$  is entangled, and is a pure state (only one non-zero  $p_k$ ) if  $|\psi\rangle$  is a product state.

The von Neumann entropy of the subsystem  $a$  density operator, following eq. (3.5.16), is

$$\sigma_a = - \sum_k p_k \ln p_k. \quad (24.1.13)$$

In the context of bipartite systems, this is known as the **entanglement entropy**. Of course, one can reverse the roles of subsystems  $a$  and  $b$ . Taking the trace over  $\mathcal{H}_a$  to find the density operator for subsystem  $b$ , one obtains  $\rho_b = \sum_k p_k |\chi_k\rangle\langle\chi_k|$ . In particular, this shows that if the combined system state is a pure one, then  $\rho_b$  has exactly the same list of non-zero ensemble probabilities, and therefore the same entanglement entropy,  $\sigma_b = \sigma_a$ . This may be somewhat surprising, since  $a$  and  $b$  can have very different properties considered as open systems by themselves, including hugely disparate degrees of freedom. For example,  $a$  might describe a single spin, while subsystem  $b$  might be the entire rest of the universe. The maximum entanglement entropy for a pure state of a bipartite system occurs when all of the non-zero  $p_k$  are equal to  $1/\min(d_a, d_b)$ . More generally, it is a single quantitative measure of the amount of entanglement of the pure combined state.

In the discussion leading to eq. (3.5.24), we found that the von Neumann entropy of a closed system does not change when it undergoes unitary time evolution. However, subsystems of the closed system often become entangled in this process. For example, an open system in a pure product state (with vanishing entanglement entropy) will often evolve into a state with non-zero entanglement entropy. Thus, the entanglement entropy of the subsystems can increase even when the closed system entropy is fixed. Although we will not prove it, the entropy of any combined system density operator  $\rho_{ab}$  and the reduced density operators  $\rho_a$  and  $\rho_b$  of its subsystems can be shown to always obey

$$|\sigma_a - \sigma_b| \leq \sigma_{ab} \leq \sigma_a + \sigma_b. \quad (24.1.14)$$

The second inequality becomes an equality in the special case of a product density matrix  $\rho_{ab} = \rho_a \otimes \rho_b$ , while  $\sigma_{ab} = 0$  and  $\sigma_a = \sigma_b$  in the special case of a pure combined state  $\rho_{ab} = |\psi\rangle\langle\psi|$ , as we have seen.

As examples of eq. (24.1.12) in action, it is straightforward to verify that the EPR state  $|\psi\rangle$  in eq. (24.1.1) leads to the completely random ensemble density operator  $\rho_a$  as claimed in eq. (24.1.2), and the unentangled state  $|\psi'\rangle$  in eq. (24.1.3) leads to the pure ensemble density operator  $\rho'_a$  of eq. (24.1.4). But let us generalize the first example slightly, by replacing the minus sign in eq. (24.1.1) by an arbitrary relative phase  $e^{i\theta}$ , so that

$$|\psi\rangle = \frac{1}{\sqrt{2}} \left( |\uparrow\rangle_a \otimes |\downarrow\rangle_b + e^{i\theta} |\downarrow\rangle_a \otimes |\uparrow\rangle_b \right). \quad (24.1.15)$$

To an observer who is capable of making measurements on the combined system  $ab$ , this relative phase is certainly physically relevant; for example, you can check that the observable  $\Omega = \sigma_x \otimes \sigma_x$  has expectation value  $\langle\psi|\Omega|\psi\rangle = \cos\theta$ . Now, the density operator for the combined system is

$$\begin{aligned} \rho_{ab} = |\psi\rangle\langle\psi| &= \frac{1}{2} \left[ (|\uparrow\rangle\langle\uparrow|)_a \otimes (|\downarrow\rangle\langle\downarrow|)_b + (|\downarrow\rangle\langle\downarrow|)_a \otimes (|\uparrow\rangle\langle\uparrow|)_b \right. \\ &\quad \left. + e^{i\theta} (|\downarrow\rangle\langle\uparrow|)_a \otimes (|\uparrow\rangle\langle\downarrow|)_b + e^{-i\theta} (|\uparrow\rangle\langle\downarrow|)_a \otimes (|\downarrow\rangle\langle\uparrow|)_b \right]. \end{aligned} \quad (24.1.16)$$

Taking the partial trace over the  $b$  spin using  $\text{Tr}_b(\rho_{ab}) = \langle\uparrow|_b \rho_{ab} |\uparrow\rangle_b + \langle\downarrow|_b \rho_{ab} |\downarrow\rangle_b$ , the last two terms in eq. (24.1.16) do not contribute, and we find

$$\rho_a = \frac{1}{2} \left( |\uparrow\rangle\langle\uparrow| + |\downarrow\rangle\langle\downarrow| \right). \quad (24.1.17)$$

The relative phase information present in the combined system state as the factor  $e^{i\theta}$  is therefore completely inaccessible to Alice alone.<sup>§</sup> Recall, from the discussion surrounding eqs. (3.5.5) and (3.5.6), that this loss of phase information is a characteristic feature of mixed ensembles.

---

<sup>§</sup>To see this without using the density operator language, but in a morally equivalent way, note that the most general combined Hilbert space Hermitian operator that acts non-trivially only on Alice's spin is  $A = (r_0 I + r_1 \sigma_x + r_2 \sigma_y + r_3 \sigma_z) \otimes I$ , where  $r_{0,1,2,3}$  are arbitrary real numbers. It is a short exercise to compute  $\langle\psi|A|\psi\rangle$  and check that it does not depend on  $\theta$ .

## 24.2 Dynamical maps and evolution of open systems

Consider an open system described at a given time by a density operator  $\rho$ . (Going forward, we refer to the subsystem  $a$  of a bipartite system  $ab$  as simply “the open system”, and often leave off the subscript  $a$  for its states and operators.) What is the density operator  $\rho'$  that describes the system at some unspecified later time? For the special case of a closed system, the answer was given in eq. (3.5.23), which says that  $\rho' = U\rho U^\dagger$  for some unitary operator  $U$ . However, this is not true for open systems. In general, we can write

$$\rho \rightarrow \rho' = \mathcal{E}(\rho), \quad (24.2.1)$$

where  $\mathcal{E}$  is a map taking operators to operators, sometimes called a **superoperator**. We will refer to  $\mathcal{E}$  as a **quantum dynamical map**, but it has many names; other sources refer to it as a **quantum channel**, or a **quantum operation**. Our goal is to understand the possible forms allowed by quantum mechanics for dynamical maps of open systems.

One way to investigate the issue is to introduce a specific environment subsystem with orthobasis  $|\chi_n\rangle_b$  where  $n = 1, \dots, d_b$ . Let us further assume that the initial condition for the combined system is a product density operator

$$\rho_{ab} = \rho \otimes (|0\rangle\langle 0|), \quad (24.2.2)$$

where  $|0\rangle$  is some reference state of subsystem  $b$ . We now let the combined system evolve in time, just as in eq. (3.5.23), which says that the final density operator is  $U\rho_{ab}U^\dagger$  for some unitary operator  $U$  that acts on the combined system. The reduced density operator for the open system after the evolution is obtained by taking the partial trace over the environment subsystem  $b$ ,

$$\mathcal{E}(\rho) = \text{Tr}_b(U\rho_{ab}U^\dagger) = \sum_n \langle \chi_n | U | 0 \rangle \rho \langle 0 | U^\dagger | \chi_n \rangle. \quad (24.2.3)$$

The objects  $\langle \chi_n | U | 0 \rangle$  and  $\langle 0 | U^\dagger | \chi_n \rangle$  are matrix elements of the environment subsystem  $b$ , but are still operators acting on the open system. They reflect the fact that the unitary evolution of the combined state causes the environment state to transition from  $|0\rangle$  to a superposition of basis states, correlated with the evolution of the open system. Giving them the names

$$K_n = \langle \chi_n | U | 0 \rangle, \quad K_n^\dagger = \langle 0 | U^\dagger | \chi_n \rangle \quad (n = 1, \dots, d_b), \quad (24.2.4)$$

we have the important result

$$\mathcal{E}(\rho) = \sum_n K_n \rho K_n^\dagger, \quad (24.2.5)$$

where it is easy to check from eq. (24.2.4), using completeness, that

$$\sum_n K_n^\dagger K_n = I. \quad (24.2.6)$$

In general, equations of the form (24.2.5) and (24.2.6) constitute an **operator-sum representation** for the dynamical map  $\mathcal{E}$ , and the  $K_n$  are called **Kraus operators**, after Karl Kraus.

If there is only a single non-zero Kraus operator, then eq. (24.2.6) tells us that it is a unitary operator. In that special case, we say that the dynamical map is unitary, as it has the same form as the unitary time evolution of a closed system in eq. (3.5.23), namely  $\rho' = U\rho U^\dagger$ .

Dynamical maps defined by operator-sum representations, as in eqs. (24.2.5) and (24.2.6), can be sequentially combined. This means that if  $\mathcal{E}_1$  and  $\mathcal{E}_2$  are dynamical maps with operator sum representations, then so is

$$\mathcal{E}_2 \circ \mathcal{E}_1(\rho) = \mathcal{E}_2(\mathcal{E}_1(\rho)). \quad (24.2.7)$$

A dynamical map  $\mathcal{E}$  has an inverse dynamical map  $\mathcal{E}^{-1}$ , so that  $\mathcal{E}^{-1}(\mathcal{E}(\rho)) = \rho$ , if and only if the map is unitary. The proofs of these properties are left as exercises. The most interesting dynamical maps for open systems are non-unitary and thus non-invertible.

As a simple example of a non-unitary dynamical map for an open system, consider a model for the decay of an atom in an excited state  $|1\rangle_a$  to its ground state  $|0\rangle_a$  by emitting a photon. These two states are the orthobasis for the open system. The environment orthobasis consists of states with and without a photon from the decay, denoted  $|1\rangle_b$  and  $|0\rangle_b$  respectively. The combined system thus has an orthobasis of four states  $|n\rangle_a \otimes |m\rangle_b$  with  $n, m = 0, 1$ . The atomic state can be either excited or not, but we assume that there is no photon present initially. This means that the combined system always starts in a superposition of the two orthobasis states

$$|0\rangle_a \otimes |0\rangle_b, \quad \text{and} \quad |1\rangle_a \otimes |0\rangle_b. \quad (24.2.8)$$

We now define a unitary map  $U$  to describe the evolution of the combined system,

$$U(|0\rangle \otimes |0\rangle) = |0\rangle \otimes |0\rangle, \quad (24.2.9)$$

$$U(|1\rangle \otimes |0\rangle) = \sqrt{1-p}|1\rangle \otimes |0\rangle + \sqrt{p}|0\rangle \otimes |1\rangle, \quad (24.2.10)$$

where  $p$  is a real number between 0 and 1. The interpretation of the first equation is that nothing happens if the atom starts in its ground state. The second equation says that if it starts in the excited state, then with probability  $p$  it will decay to the ground state while adding the photon to the environment state, and with probability  $1-p$  everything will remain as it is. The operator  $U$  preserves inner products, in the sense that the states in eqs. (24.2.9) and (24.2.10)

are orthogonal and each have norm 1. This is an important requirement for our assertion that  $U$  is a unitary operator on the complete Hilbert space.<sup>†</sup>

Following the construction of eq. (24.2.4), we see that there are two Kraus operators in this example, one for each of our two environment orthobasis states. We can use that equation as a recipe to construct their actions on the orthobasis of the open system,

$$K_0 |0\rangle_a = \langle 0|_b U (|0\rangle \otimes |0\rangle) = |0\rangle_a, \quad (24.2.11)$$

$$K_0 |1\rangle_a = \langle 0|_b U (|1\rangle \otimes |0\rangle) = \sqrt{1-p} |1\rangle_a \quad (24.2.12)$$

$$K_1 |0\rangle_a = \langle 1|_b U (|0\rangle \otimes |0\rangle) = 0, \quad (24.2.13)$$

$$K_1 |1\rangle_a = \langle 1|_b U (|1\rangle \otimes |0\rangle) = \sqrt{p} |0\rangle_a, \quad (24.2.14)$$

or, in matrix representation form,

$$K_0 = \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{1-p} \end{pmatrix}, \quad K_1 = \begin{pmatrix} 0 & \sqrt{p} \\ 0 & 0 \end{pmatrix}. \quad (24.2.15)$$

These indeed satisfy the completeness requirement  $\sum_n K_n^\dagger K_n = I$ . Now, given an initial density matrix of the open system (the atom),

$$\rho = \begin{pmatrix} \rho_{00} & \rho_{01} \\ \rho_{10} & \rho_{11} \end{pmatrix}, \quad (24.2.16)$$

the dynamical map resulting from the Kraus operator-sum representation of eq. (24.2.5) is

$$\mathcal{E}(\rho) = K_0 \rho K_0^\dagger + K_1 \rho K_1^\dagger = \begin{pmatrix} \rho_{00} + p\rho_{11} & \sqrt{1-p} \rho_{01} \\ \sqrt{1-p} \rho_{10} & (1-p)\rho_{11} \end{pmatrix}. \quad (24.2.17)$$

So far, we have not associated the dynamical map with a specific time interval. To remedy this, we can take eq. (24.2.17) to correspond to an infinitesimal time interval  $\Delta t$ , with an excited state atomic decay probability  $p = \Gamma \Delta t$ . We then apply the map  $N$  times. In doing so, we take the initial state of the environment to be the no-photon state  $|0\rangle_b$  each time, on the grounds that any photons from decays in earlier iterations of the dynamical map will have fled the scene anyway, and therefore can and should be ignored. The result is

$$\rho \rightarrow \begin{pmatrix} \rho_{00} + (1 - (1-p)^N) \rho_{11} & (\sqrt{1-p})^N \rho_{01} \\ (\sqrt{1-p})^N \rho_{10} & (1-p)^N \rho_{11} \end{pmatrix}. \quad (24.2.18)$$

---

<sup>†</sup>Actually, we have only bothered to specify how  $U$  acts on two of the four orthobasis states for the combined system, because that is all we will directly need, but you can easily complete it by constructing  $U(|0\rangle \otimes |1\rangle)$  and  $U(|1\rangle \otimes |1\rangle)$  such that  $U$  is unitary. More generally, it is true that if an operator preserves all inner products on a subspace, it can always be extended to a unitary operator on the full Hilbert space. The formal statement and a constructive proof of this assertion are postponed to Theorem 24.2.2 near the end of this section.

Now taking the limit  $N \rightarrow \infty$ , with  $t = N\Delta t$  held fixed, and using the defining property of the exponential function,

$$\lim_{N \rightarrow \infty} \left(1 - \frac{\Gamma t}{N}\right)^N = e^{-\Gamma t}, \quad (24.2.19)$$

we find, after making use of the trace condition  $\rho_{00} = 1 - \rho_{11}$ ,

$$\rho(t) = \begin{pmatrix} 1 - e^{-\Gamma t} \rho_{11} & e^{-\Gamma t/2} \rho_{01} \\ e^{-\Gamma t/2} \rho_{10} & e^{-\Gamma t} \rho_{11} \end{pmatrix}. \quad (24.2.20)$$

The interpretation of the diagonal entries is simply that a population of excited states decays to the ground state with a probability  $1 - e^{-\Gamma t}$ , which approaches 1 at late times. The off-diagonal entries of the density operator matrix also decay, with a lifetime twice as large. This dynamical map is non-invertible, and no matter what mixed state we started with, it always evolves into the pure ground state at late times. In particular, the entropy for this open system decreases to 0, unlike what we found for a unitary evolution of a closed system in eq. (3.5.24).

In the construction of eq. (24.2.4), the number of Kraus operators was  $d_b$ , the same as the dimension of the environment subsystem Hilbert space. This is not necessarily convenient, especially if the environment is a very large and complicated system. However, the operator-sum representation is not unique, and different versions for the same map  $\mathcal{E}(\rho)$  can have different numbers of Kraus operators. One can define new Kraus operators by

$$\tilde{K}_\mu = \sum_n v_{\mu n} K_n, \quad (24.2.21)$$

after which the same map can be rewritten as  $\mathcal{E}(\rho) = \sum_\mu \tilde{K}_\mu \rho \tilde{K}_\mu^\dagger$  with  $\sum_\mu \tilde{K}_\mu^\dagger \tilde{K}_\mu = I$ , provided that  $\sum_\mu v_{\mu n}^* v_{\mu n'} = \delta_{nn'}$ . This condition is equivalent to saying that the (non-square, in general) matrix of elements  $v_{\mu n}$  can be extended to a unitary matrix. Fortunately, for any given dynamical map  $\mathcal{E}$ , we will soon show that there is always an operator-sum representation with no more than  $d^2$  Kraus operators, where  $d$  is the dimension of the open system Hilbert space, regardless of the (possibly enormous) dimension  $d_b$  of the environment Hilbert space.

We can now inquire about more general dynamical maps, beyond the type we defined in eq. (24.2.3). One obvious possibility is to take the initial condition for the combined system to involve a general density operator for the environment, rather than a pure state  $|0\rangle$ , so that initially<sup>‡</sup>

$$\rho_{ab} = \rho \otimes \rho_b, \quad (24.2.22)$$

---

<sup>‡</sup>The initial condition of eq. (24.2.22) is still far from general, as it is a product of density operators for the subsystems, with no initial correlations between system and environment. It is an understatement to say that the general case of an entangled density operator for the initial condition is much more subtle. It will not be addressed here. See P. Pechukas, Phys. Rev. Lett. **73**, no.8, 1060 (1994); T. F. Jordan, A. Shaji, and E. C. G. Sudarshan, Phys. Rev. A **70**, 052110 (2004) [arXiv:quant-ph/0407083]; D. Schmid, K. Ried, and R.W. Spekkens, Phys. Rev. A **100**, 022112 (2019) [arXiv:1806.02381], and associated references.



and then  $\mathcal{E}(\rho) = \text{Tr}_b[U(\rho \otimes \rho_b)U^\dagger]$  for some unitary joint-state operator  $U$ . It is left to Exercise 24.3 to show that in this case the map  $\mathcal{E}(\rho)$  can be written in the same operator-sum form, but with  $d_b^2$  Kraus operators instead of only  $d_b$ , although as noted in the previous paragraph it is often possible to reduce this number.

Given this, it is a plausible guess that dynamical maps can always be written in the operator-sum form of eqs. (24.2.5) and (24.2.6). As we will now show, this is indeed true, provided we assume some reasonable conditions motivated by the requirement that  $\mathcal{E}(\rho)$  must be a legitimate density operator for the open system. These conditions are:

- (i)  $\mathcal{E}$  is a linear map, satisfying  $\mathcal{E}(c_1 A_1 + c_2 A_2) = c_1 \mathcal{E}(A_1) + c_2 \mathcal{E}(A_2)$  for complex numbers  $c_{1,2}$  and operators  $A_{1,2}$ . There are two motivating reasons to impose this condition. The first reason is that any collection of mixed ensembles can be viewed as a single mixed ensemble, so one must have for any density operators  $\rho_k$  with population ratios  $p_k$ ,

$$\mathcal{E}\left(\sum_k p_k \rho_k\right) = \sum_k p_k \mathcal{E}(\rho_k). \quad (24.2.23)$$

The second reason is that linearity implies that the map preserves Hermiticity; if  $\rho$  is Hermitian, then  $\mathcal{E}(\rho)$  will be also, as it must.

- (ii) The map  $\mathcal{E}$  preserves traces, so that  $\text{Tr}[\mathcal{E}(A)] = \text{Tr}(A)$ . When  $A = \rho$  and  $\text{Tr}(\rho) = 1$ , this is equivalent to the conservation of probability.

- (iii) The map  $\mathcal{E}$  is a completely positive map.

The last condition requires a terminology explanation. A superoperator  $\mathcal{E}$  is said to be a **positive map** if  $\mathcal{E}(A)$  is a positive operator whenever  $A$  is a positive operator. A positive map is said to be **completely positive** if every extension of it to include an environment system, on which it acts trivially, is also positive. In other words, a positive map  $\mathcal{E}(A)$  for operators  $A$  on  $\mathcal{H}_a$  is completely positive if, for any extended system  $\mathcal{H}_a \otimes \mathcal{H}_b$ , the map  $\mathcal{E}_{\text{ext}}$  defined by

$$\mathcal{E}_{\text{ext}}(A \otimes B) = \mathcal{E}(A) \otimes B \quad (24.2.24)$$

is also a positive map. A motivation for requiring dynamical maps to be completely positive is that if  $\mathcal{E}(\rho)$  is a valid quantum dynamics, then it ought to remain one if we add an arbitrary environment to which it is very weakly coupled. Naively, this might seem trivial, but it is not, because there do exist linear trace-preserving maps that are positive but not completely positive. A minimal example will be given at the end of this section.

It is a short exercise to confirm that a dynamical map with the operator-sum form of eqs. (24.2.5) and (24.2.6) satisfies the three conditions above. Conversely, we can now make, and prove, a strong statement about the general form of dynamical maps on open systems:

**Theorem 24.2.1. (Operator-sum representation of dynamical maps)** *If a map  $\rho \rightarrow \rho' = \mathcal{E}(\rho)$  for density operators on an open system Hilbert space of dimension  $d$  is linear, trace-preserving, and completely positive, then it has an operator-sum representation*

$$\mathcal{E}(\rho) = \sum_{n=1}^N K_n \rho K_n^\dagger, \quad \sum_{n=1}^N K_n^\dagger K_n = I, \quad (24.2.25)$$

where it is always possible to choose  $N$  to be  $d^2$  or less.

Proof: for the open system Hilbert space  $\mathcal{H}_a$ , choose an orthobasis  $|\varphi_i\rangle$  with  $i = 1, \dots, d$ . Now introduce<sup>§</sup> an auxiliary system Hilbert space  $\mathcal{H}_b$  with the same dimension  $d$ , so that we can choose an orthobasis  $|\chi_i\rangle$  for it, again with  $i = 1, \dots, d$ . This allows us to assign to each ket  $|\varphi_i\rangle$  a uniquely associated  $\langle\chi_i|$ , a fact that will be important soon. Now consider the following normalized state defined on the combined system with Hilbert space  $\mathcal{H}_a \otimes \mathcal{H}_b$ :

$$|\Psi\rangle = \frac{1}{\sqrt{d}} \sum_{i=1}^d |\varphi_i\rangle \otimes |\chi_i\rangle. \quad (24.2.26)$$

[As an aside, this is a maximally entangled ket, with Schmidt number  $d$  and entanglement entropy  $\ln(d)$ .] Consider the extension of  $\mathcal{E}$  to a map  $\mathcal{E}_{\text{ext}}$  for operators on the combined Hilbert space, defined in general as in eq. (24.2.24), so that

$$\mathcal{E}_{\text{ext}}(|\Psi\rangle\langle\Psi|) = \frac{1}{d} \sum_{i=1}^d \sum_{j=1}^d \mathcal{E}(|\varphi_i\rangle\langle\varphi_j|) \otimes (|\chi_i\rangle\langle\chi_j|). \quad (24.2.27)$$

Because  $\mathcal{E}$  is completely positive by assumption,  $\mathcal{E}_{\text{ext}}$  is a positive map, by definition. Therefore, since  $|\Psi\rangle\langle\Psi|$  is a valid density operator on  $\mathcal{H}_a \otimes \mathcal{H}_b$ , it follows that so is  $\mathcal{E}_{\text{ext}}(|\Psi\rangle\langle\Psi|)$ , since it satisfies the conditions of being a positive Hermitian operator with trace 1. As a density operator, it must be possible to express it in terms of an orthobasis of its eigenkets  $|\Phi_n\rangle$ , as in eq. (3.5.14),

$$\mathcal{E}_{\text{ext}}(|\Psi\rangle\langle\Psi|) = \sum_{n=1}^{d^2} p_n |\Phi_n\rangle\langle\Phi_n|. \quad (24.2.28)$$

Here we have used the fact that the number of orthobasis eigenkets  $|\Phi_n\rangle$  is  $d^2$ , because that is the dimension of  $\mathcal{H}_a \otimes \mathcal{H}_b$ . Equating (24.2.27) and (24.2.28), and then acting from the left with  $\langle\chi_i|$  and the right with  $|\chi_j\rangle$  and multiplying by  $d$ , we obtain

$$\mathcal{E}(|\varphi_i\rangle\langle\varphi_j|) = d \sum_{n=1}^{d^2} p_n \langle\chi_i|\Phi_n\rangle\langle\Phi_n|\chi_j\rangle. \quad (24.2.29)$$

---

<sup>§</sup>It is important that this auxiliary system  $b$  is introduced only as a calculation trick, and is not the physical environment system, which we will not need to make any reference to in this proof.

In the last expression,  $\langle \chi_i | \Phi_n \rangle$  is a matrix element in  $\mathcal{H}_b$ , but it is a ket in the open system Hilbert space  $\mathcal{H}_a$ . We now make the crucial observation that, because there is a unique association of  $|\varphi_i\rangle$  with  $\langle \chi_i |$ , we can define a linear operator  $K_n$  on  $\mathcal{H}_a$  by specifying its action on each of the orthobasis kets,

$$K_n |\varphi_i\rangle = \sqrt{dp_n} \langle \chi_i | \Phi_n \rangle. \quad (24.2.30)$$

This allows us to rewrite eq. (24.2.29) as

$$\mathcal{E}(|\varphi_i\rangle \langle \varphi_j|) = \sum_{n=1}^{d^2} K_n |\varphi_i\rangle \langle \varphi_j| K_n^\dagger. \quad (24.2.31)$$

Since  $\mathcal{E}$  is linear by assumption, and any density operator can be expressed as a linear combination  $\rho = \sum_{i,j} \rho_{ij} |\varphi_i\rangle \langle \varphi_j|$ , we obtain the result for  $\mathcal{E}(\rho)$  in the first part of eq. (24.2.25). Finally, we compute from eq. (24.2.31) that

$$\begin{aligned} \text{Tr}[\mathcal{E}(|\varphi_i\rangle \langle \varphi_j|)] &= \sum_k \sum_n \langle \varphi_k | K_n |\varphi_i\rangle \langle \varphi_j | K_n^\dagger | \varphi_k \rangle = \sum_n \langle \varphi_j | K_n^\dagger \left( \sum_k |\varphi_k\rangle \langle \varphi_k| \right) K_n | \varphi_i \rangle \\ &= \langle \varphi_j | \left( \sum_n K_n^\dagger K_n \right) | \varphi_i \rangle. \end{aligned} \quad (24.2.32)$$

Now, using the assumption that  $\mathcal{E}$  is trace-preserving, this must equal  $\text{Tr}(|\varphi_i\rangle \langle \varphi_j|) = \delta_{ij}$ , which shows that  $\sum_n K_n^\dagger K_n$  is the identity operator, the second part of eq. (24.2.25).  $\square$

We conclude this section by tying up two loose ends. First, in the footnote following eq. (24.2.10), it was claimed that any linear operator that preserves all inner products on a subspace can be extended to a unitary operator on the whole Hilbert space. More formally:

**Theorem 24.2.2. (Construction of a unitary operator from its action on a subspace)**

*Let  $\mathcal{H}$  be a Hilbert space with a subspace  $\mathcal{H}_s$ . Suppose that  $U_s$  is a linear operator that acts on kets in the subspace  $\mathcal{H}_s$ , mapping them to kets in the full space  $\mathcal{H}$  while preserving inner products. This means that for all kets  $|v\rangle$  and  $|w\rangle$  in  $\mathcal{H}_s$ , the kets  $U_s |v\rangle$  and  $U_s |w\rangle$  are elements of  $\mathcal{H}$ , and*

$$\langle w | U_s^\dagger U_s | v \rangle = \langle w | v \rangle. \quad (24.2.33)$$

*Then  $U_s$  can be extended to a unitary operator  $U$  defined on all of  $\mathcal{H}$ . In other words, one can find an operator  $U$  such that  $U |v\rangle = U_s |v\rangle$  for all  $|v\rangle$  in  $\mathcal{H}_s$ , and  $U^\dagger U = U U^\dagger = I$ .*

Proof: we will explicitly construct a unitary operator  $U$  that satisfies the theorem. Let the dimensions of  $\mathcal{H}$  and  $\mathcal{H}_s$  be  $d$  and  $d_s$ , respectively. Choose an orthobasis  $\{|\phi_j\rangle\}$  for  $\mathcal{H}_s$ , with  $j = 1, \dots, d_s$ . Now, let  $\mathcal{H}^\perp$  be the subspace of kets in  $\mathcal{H}$  that are perpendicular to all kets

in  $\mathcal{H}_s$ . Then  $\mathcal{H}^\perp$  is a Hilbert space, and one can use the Gram-Schmidt procedure to find an orthobasis  $\{|\bar{\phi}_k\rangle\}$  for it, with  $k = 1, \dots, d - d_s$ . Note that  $\{|\phi_j\rangle, |\bar{\phi}_k\rangle\}$  is an orthobasis for  $\mathcal{H}$ . Next, note that from eq. (24.2.33),  $U_s |\phi_j\rangle$  are a set of  $d_s$  orthonormal kets in  $\mathcal{H}$ . The set of kets perpendicular to all of them is a Hilbert space with dimension  $d - d_s$ , and one can therefore find an orthobasis  $\{|\tilde{\phi}_k\rangle\}$  for it, again with  $k = 1, \dots, d - d_s$ . Now define

$$U = \sum_{j=1}^{d_s} U_s |\phi_j\rangle \langle \phi_j| + \sum_{k=1}^{d-d_s} |\tilde{\phi}_k\rangle \langle \bar{\phi}_k|. \quad (24.2.34)$$

Since  $\langle \bar{\phi}_k | \phi_i \rangle = 0$  and  $\langle \phi_j | \phi_i \rangle = \delta_{ij}$ , it is clear that  $U |\phi_i\rangle = U_s |\phi_i\rangle$  for all orthobasis elements of  $\mathcal{H}_s$ , as required. Finally, we check that  $U$  is indeed unitary:

$$\begin{aligned} U^\dagger U &= \left( \sum_{j'} |\phi_{j'}\rangle \langle \phi_{j'}| U_s^\dagger + \sum_{k'} |\bar{\phi}_{k'}\rangle \langle \tilde{\phi}_{k'}| \right) \left( \sum_j U_s |\phi_j\rangle \langle \phi_j| + \sum_k |\tilde{\phi}_k\rangle \langle \bar{\phi}_k| \right) \\ &= \sum_{j,j'} |\phi_{j'}\rangle \langle \phi_{j'}| \phi_j\rangle \langle \phi_j| + \sum_{k,k'} |\bar{\phi}_{k'}\rangle \langle \tilde{\phi}_{k'}| \tilde{\phi}_k\rangle \langle \bar{\phi}_k| \\ &= \sum_j |\phi_j\rangle \langle \phi_j| + \sum_k |\bar{\phi}_k\rangle \langle \bar{\phi}_k| = I, \end{aligned} \quad (24.2.35)$$

where we have used eq. (24.2.33) and the orthonormality and completeness properties of  $|\phi_j\rangle$ ,  $|\bar{\phi}_k\rangle$ , and  $|\tilde{\phi}_k\rangle$  on their respective subspaces.  $\square$

The other loose end is that after eq. (24.2.24) we promised to exhibit a linear, trace-preserving, positive map that is not completely positive. A standard example is the transpose map,  $\mathcal{E}(\rho) = \rho^T$ , which is defined to act on subsystem  $a$  by taking the transpose of the density matrix in some orthobasis. This map is linear, and trace-preserving and positive because  $\rho$  and  $\rho^T$  have the same eigenvalues, and the trace is the sum of the eigenvalues while a positive operator is one with all non-negative eigenvalues. However, its extension  $\mathcal{E}_{\text{ext}}(\rho)$  as defined by eq. (24.2.24) is not a positive map. To see this in a minimal case, consider two 2-dimensional subspaces  $a$  and  $b$ , with orthobases  $|0\rangle_a, |1\rangle_a$  and  $|0\rangle_b, |1\rangle_b$  respectively. Let us use a notation  $|jk\rangle = |j\rangle_a \otimes |k\rangle_b$  for  $j, k = 0, 1$ . Then the map  $\mathcal{E}_{\text{ext}}(\rho)$  acts to exchange the subsystem  $a$  labels only, according to

$$\mathcal{E}_{\text{ext}}(|jk\rangle \langle nm|) = |nk\rangle \langle jm|. \quad (24.2.36)$$

Now consider the combined system density operator

$$\rho_{ab} = \frac{1}{2}(|00\rangle \langle 00| + |00\rangle \langle 11| + |11\rangle \langle 00| + |11\rangle \langle 11|). \quad (24.2.37)$$

This is the density operator for the pure state  $(|00\rangle + |11\rangle)/\sqrt{2}$ , so it is clearly a positive operator. Applying eq. (24.2.36) to it gives

$$\mathcal{E}_{\text{ext}}(\rho_{ab}) = \frac{1}{2}(|00\rangle \langle 00| + |10\rangle \langle 01| + |01\rangle \langle 10| + |11\rangle \langle 11|). \quad (24.2.38)$$

Now consider the state  $|\psi\rangle = (|01\rangle - |10\rangle)/\sqrt{2}$ . One finds

$$\langle\psi|\mathcal{E}_{\text{ext}}(\rho_{ab})|\psi\rangle = -1/4, \quad (24.2.39)$$

so  $\mathcal{E}_{\text{ext}}(\rho_{ab})$  is not a positive operator, and by definition  $\mathcal{E}(\rho)$  is not a completely positive map.

### 24.3 Generalized measurements

The postulates of quantum mechanics presented in section 3.1 give rules for observables and measurements in closed systems. The allowed results of a measurements are the eigenvalues  $\alpha$  of a Hermitian operator, and the probabilities and post-measurement state of the system are dictated by Hermitian projection operators  $P_\alpha$  constructed out of the corresponding eigenstates. This type of measurement, used up to now in this book, is called a **projective measurement**.

For open systems, a more general type of measurement process is available. As a model, the open system  $a$  is coupled to an environment subsystem  $b$  which we can call the “meter”. To conduct a measurement, the meter is initially prepared in a specific reference state. We then allow a unitary transformation<sup>†</sup> to take place, causing the combined system state to become entangled. In a sense, this is already a kind of measurement on the open system, since there are now correlations between it and the meter, and even if the open system was originally in a pure state, it is now described by a mixed ensemble. To expose this, a projective measurement is then performed on the meter subsystem. The combination of the unitary entanglement followed by the projective measurement on the meter is called a **generalized measurement** of the open system. It will give us probabilities associated with the possible measurement outcomes, and for each outcome will also tell us the post-measurement density operator of the open system.

To be specific, let the quantum meter subsystem have dimension  $d_b = N$ , with an orthobasis of states  $|\chi_n\rangle$  where  $n = 1, 2, \dots, N$ . To define the measurement process, we require that the meter system is always prepared in the same reference state  $|0\rangle$  initially. Then, for any open system state  $|\psi\rangle$ , define the entangling unitary transformation by

$$U(|\psi\rangle \otimes |0\rangle) = \sum_n (M_n |\psi\rangle \otimes |\chi_n\rangle), \quad (24.3.1)$$

where the operators  $M_n$  are called **measurement operators** on the open system. The  $M_n$  can be recognized as nothing other than the Kraus operators for a dynamical map. In fact, in the following we will essentially be revisiting the discussion surrounding eqs. (24.2.2)-(24.2.6), just with  $K_n$  replaced by  $M_n$  and in a cosmetically different way.

---

<sup>†</sup>The unitary transformation of the combined system is typically associated with some time evolution, but we need not be specific about the time interval or the Hamiltonian.

In order for  $U$  to be unitary, it must preserve inner products between any two states  $|\psi\rangle \otimes |0\rangle$  and  $|\phi\rangle \otimes |0\rangle$ . This implies  $\sum_k \sum_n \langle \phi | M_n^\dagger M_k | \psi \rangle \langle \chi_n | \chi_k \rangle = \langle \phi | \psi \rangle$ . Due to the orthonormality of the meter states  $\langle \chi_n | \chi_k \rangle = \delta_{nk}$ , this simplifies to  $\langle \phi | \sum_n M_n^\dagger M_n | \psi \rangle = \langle \phi | \psi \rangle$ . Since this is supposed to hold for any  $|\phi\rangle$  and  $|\psi\rangle$ , it is necessary that

$$\sum_n M_n^\dagger M_n = I, \quad (24.3.2)$$

as in eq. (24.2.6). Theorem 24.2.2 guarantees that eq. (24.3.2) is also sufficient; having defined  $U$  to preserve all inner products when acting on the subspace of states of the form  $|\psi\rangle \otimes |0\rangle$ , it can be extended to a unitary operator on the whole combined Hilbert space.

Now suppose, to be general, that we have a mixed ensemble of open system states, so that we start with a combined density operator of the form

$$\rho_{ab} = \rho \otimes |0\rangle \langle 0|. \quad (24.3.3)$$

Applying the unitary transformation  $U$ , the new density matrix of the combined system is

$$\rho'_{ab} = U \rho_{ab} U^\dagger = \sum_{k,m} M_k \rho M_m^\dagger \otimes |\chi_k\rangle \langle \chi_m|. \quad (24.3.4)$$

To complete the measurement process, we assume that an observer, with access to the meter only, makes a projective measurement to collapse the meter to one of the states  $|\chi_n\rangle$ , using mutually orthogonal projection operators that act nontrivially only on the meter subsystem,

$$P_n = I \otimes |\chi_n\rangle \langle \chi_n|. \quad (24.3.5)$$

Specifically, the Hermitian observable for this projective measurement is

$$Q = \sum_{n=1}^N n P_n, \quad (24.3.6)$$

which is a spectral decomposition in terms of the  $P_n$ . The outcome of the measurement is thus one of the  $Q$  eigenvalues,  $n = 1, \dots, N$ . Applying eq. (3.5.11), the probability of getting each outcome  $n$  is

$$\mathcal{P}(n) = \text{Tr}(\rho'_{ab} P_n) = \sum_{k,m} \text{Tr}(M_k \rho M_m^\dagger) \text{Tr}(|\chi_k\rangle \langle \chi_m| |\chi_n\rangle \langle \chi_n|), \quad (24.3.7)$$

which simplifies, using  $\text{Tr}(|\chi_k\rangle \langle \chi_m| |\chi_n\rangle \langle \chi_n|) = \delta_{mn} \langle \chi_n | \chi_k \rangle = \delta_{mn} \delta_{kn}$ , to

$$\mathcal{P}(n) = \text{Tr}(\rho M_n^\dagger M_n). \quad (24.3.8)$$

Also, if the measurement process outcome is  $n$ , then the post-measurement combined system density operator, obtained by applying the general formula eq. (3.5.29), will be

$$\rho_{ab}'' = \frac{P_n \rho_{ab}' P_n}{\mathcal{P}(n)} = \frac{M_n \rho M_n^\dagger \otimes |\chi_n\rangle \langle \chi_n|}{\text{Tr}(\rho M_n^\dagger M_n)}. \quad (24.3.9)$$

Taking the partial trace over the meter subsystem, using  $\text{Tr}(|\chi_n\rangle \langle \chi_n|) = \langle \chi_n | \chi_n \rangle = 1$ , gives

$$\rho_n = \frac{M_n \rho M_n^\dagger}{\mathcal{P}(n)} = \frac{M_n \rho M_n^\dagger}{\text{Tr}(\rho M_n^\dagger M_n)}. \quad (24.3.10)$$

for the density operator of the open subsystem after the outcome  $n$ .

To recapitulate and slightly augment the preceding discussion, a generalized measurement on an open system can be defined by choosing *any* set of measurement operators  $M_n$  that obey the completeness relation (24.3.2). The probability of getting the outcome  $n$  from the measurement is eq. (24.3.8), and the open system density operator after the measurement outcome  $n$  will be eq. (24.3.10). In the special case that the open system starts in a pure state  $|\psi\rangle$ , these imply

$$\mathcal{P}(n) = \langle \psi | M_n^\dagger M_n | \psi \rangle, \quad |\psi_n\rangle = \frac{M_n |\psi\rangle}{\sqrt{\mathcal{P}(n)}}. \quad (24.3.11)$$

Note that the post-measurement state  $|\psi_n\rangle$  is also a pure one, due to the collapse of the meter state. These rules can be taken as definitions of a generalized measurement that do not refer to a specific environment, although we had our meter subsystem in mind while deriving them. As consistency checks, note first that the sum of all of the measurement result probabilities is

$$\sum_n \mathcal{P}(n) = \sum_n \text{Tr}(\rho M_n^\dagger M_n) = \text{Tr}(\rho) = 1, \quad (24.3.12)$$

where the completeness relation,  $\sum_n M_n^\dagger M_n = I$ , plays the important role of ensuring the conservation of probability. Second, eq. (24.3.10) implies that the open system density operator after the measurement obeys  $\text{Tr}(\rho_n) = 1$  and  $\rho_n = \rho_n^\dagger$ , and that  $\rho_n$  is a positive operator (with positive eigenvalues), provided that the initial  $\rho$  satisfied these properties.

There is a close relationship between generalized measurements and dynamical maps on open systems. In this connection, the generalized measurement operators  $M_n$  can simply be identified with the Kraus operators  $K_n$  of a dynamical map. To see this more generally (without reference to the derivations above in terms of specific environment models), suppose we start with an open system density operator  $\rho$ , and make a generalized measurement with operators  $M_n$ , but do not know the result. Then the open system will be described by a mixed ensemble of states obtained by summing over the outcomes using eqs. (24.3.8) and (24.3.10),

$$\rho = \sum_n \mathcal{P}(n) \rho_n = \sum_n M_n \rho M_n^\dagger, \quad (24.3.13)$$

which is nothing but the dynamical map definition of eq. (24.2.25) with  $K_n = M_n$ . Thus, any dynamical evolution of an open system can be viewed as a generalized measurement of it by the environment, if the result of the generalized measurement is unknown or disregarded. Conversely, any generalized measurement can be viewed as a dynamical map followed by an ordinary projective measurement, but with the key feature that the collapse-of-the-state is banished from the open system being measured, and instead occurs in the meter system.

Projective measurements are special cases of generalized measurements, as the latter's name implies. They arise if one imposes the further conditions that the measurement operators  $M_n$  are Hermitian, and are orthogonal projection operators in the sense that for any  $n$  and  $m$ ,

$$M_n M_m = \delta_{nm} M_n \quad (\text{projective measurements}). \quad (24.3.14)$$

This implies that  $M_n^\dagger M_n = M_n$  can play the role of the orthogonal Hermitian projection operators  $P_\alpha$  in Postulates 4 and 5 of section 3.1, and their equivalents for density operators, eqs. (3.5.11) and (3.5.29).

Generalized measurements can do interesting things that projective measurements restricted to the open system cannot. First, recall that Postulate 3 tells us that the outcome of doing a projective measurement is always one of the eigenvalues  $\alpha$  of a Hermitian operator. In particular, the number of distinct possible outcomes cannot exceed the dimension of the state space. In contrast, the outcomes for generalized measurements are not necessarily associated with the eigenvalues of a Hermitian operator, and the number  $N$  of distinct possible outcomes can be arbitrarily large. Also, because the  $M_n$  are not necessarily orthogonal projection operators, if one does a generalized measurement on an open system and finds the result  $n$ , one is not guaranteed to get the same result if one immediately repeats the same measurement. For the same reason, the post-measurement states need not be orthogonal to each other for different outcomes of a generalized measurement, again unlike the case for projective measurements. Finally, while a projective measurement cannot decrease the entropy [as we showed in the discussion leading to eq. (3.5.35)], a generalized measurement can do so, as we will see by example soon.

In some applications, we may care only about the probabilities, and not what the density operator is after the measurement. In that case, we can define a measurement by specifying Hermitian operators<sup>‡</sup>  $E_n$  with the completeness requirement

$$\sum_n E_n = I, \quad (24.3.15)$$

and the positivity requirement  $\langle \psi | E_n | \psi \rangle \geq 0$  for each  $n$  and all  $|\psi\rangle$ . With these conditions, one can show that it is always possible to find operators  $M_n$  such that  $E_n = M_n^\dagger M_n$ . One can

---

<sup>‡</sup>The notation  $E_n$  for the operators  $M_n^\dagger M_n$  is a standard tradition, and they have nothing to do with energy.



therefore copy eq. (24.3.8) by assigning probability results

$$\mathcal{P}(n) = \text{Tr}(\rho E_n). \quad (24.3.16)$$

In this context, a choice of operators  $E_n$  is often called a **positive operator-valued measure**, or **POVM**. However, for any choice of POVM, there are actually many inequivalent generalized measurements, because there are many different operators  $M_n$  such that  $E_n = M_n^\dagger M_n$ . In particular, if a POVM  $\{E_n\}$  is obtained as  $E_n = M_n^\dagger M_n$  from a set  $\{M_n\}$ , then clearly it can also be obtained as  $E_n = M_n'^\dagger M_n'$  from the set  $\{M_n'\} = \{V_n M_n\}$ , where the  $V_n$  are arbitrary unitary operators chosen independently for each  $n$ . Choosing a POVM  $\{E_n\}$  fully determines the outcome probabilities, but it certainly does not determine the open system post-measurement density operator, because eq. (24.3.10) does not just involve the  $E_n$ . To completely define the generalized measurement, one must choose the specific operators  $M_n$ .

Suppose we want to measure a system to detect the presence of a photon. As a very simplified model, consider an open system with two orthobasis states  $|0\rangle$  (no photon) and  $|1\rangle$  (photon is present). One way to proceed is to define a projective measurement using

$$P_{\text{photon}} = |1\rangle\langle 1|, \quad (24.3.17)$$

which has eigenvalues  $\alpha = 0$  and  $1$ , with eigenstates  $|0\rangle$  and  $|1\rangle$ . For an input density matrix

$$|\psi\rangle = \begin{pmatrix} \rho_{00} & \rho_{01} \\ \rho_{10} & \rho_{11} \end{pmatrix}, \quad (24.3.18)$$

the probability of obtaining the result  $\alpha = 1$  (“yes, the photon is present”) will be  $\mathcal{P}(1) = \rho_{11}$ . After that result, the system will definitely be in the pure state  $|1\rangle$ ; according to the standard rules of projective measurements, detecting the photon ensures that any future measurement will also detect it. But this shows that our simple model is *too* simple. In the real world, the process of observing a photon typically destroys the photon by converting it and collecting its energy in a detector. A more realistic description of the measurement should have the property that if the photon was detected, the post-measurement state should be  $|0\rangle$ .

To do better while still maintaining simplicity, we can invent a generalized measurement, let us call it a **destructive measurement** of the photon, by defining two measurement operators

$$M_0 = |0\rangle\langle 0|, \quad M_1 = |0\rangle\langle 1|. \quad (24.3.19)$$

The corresponding POVM operators are then

$$E_0 = M_0^\dagger M_0 = |0\rangle\langle 0|, \quad E_1 = M_1^\dagger M_1 = |1\rangle\langle 1|, \quad (24.3.20)$$

so that the completeness condition  $\sum_n M_n^\dagger M_n = I$  is indeed satisfied. Because  $M_1^\dagger M_1$  is equal to  $P_{\text{photon}}$ , the probability for the outcome  $n = 1$  is exactly the same,  $\mathcal{P}(1) = \rho_{11}$ . The difference is entirely in the post-measurement state for this outcome. Applying eq. (24.3.10) gives

$$\rho_1 = \frac{M_1 \rho M_1^\dagger}{\mathcal{P}(1)} = \frac{1}{\rho_{11}} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \rho_{00} & \rho_{01} \\ \rho_{10} & \rho_{11} \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad (24.3.21)$$

which describes the pure state  $|0\rangle$  with no photon, as expected, since if it was there we destroyed it by measuring it. If we try to measure it again, we will not find it. Similarly, the results for the outcome  $n = 0$  are a probability  $\mathcal{P}(0) = \rho_{00}$  and a post-measurement density operator

$$\rho_0 = \frac{M_0 \rho M_0^\dagger}{\mathcal{P}(0)} = \frac{1}{\rho_{00}} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \rho_{00} & \rho_{01} \\ \rho_{10} & \rho_{11} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}. \quad (24.3.22)$$

Regardless of the outcome of the generalized measurement, the post-measurement result is the pure state with no photon, so the entropy has been reduced to 0.

As POVMs, both the projective measurement and the destructive measurement of the photon have the same operators  $E_0 = |0\rangle\langle 0|$  and  $E_1 = |1\rangle\langle 1|$ , which in this example happen to be projection operators. The only difference between them is that for the projective measurement we chose  $M_1 = |1\rangle\langle 1|$ , while for the destructive measurement we chose  $M_1 = |0\rangle\langle 1|$ , which is not a projection operator. Both choices gave the same probabilities, but the latter choice gave a better model for the impact of the environment (detector) on the photon. It is left to Exercise 24.4 to consider the case  $M_1 = \sqrt{p}|0\rangle\langle 1| + \sqrt{1-p}|1\rangle\langle 1|$ , which interpolates between the two choices for the detector impact on an initial-state photon while keeping the same POVM operator  $E_1$ .

We close this section with another example of the useful application of a generalized measurement: the problem of distinguishing two states. Suppose that Alice is presented with a spin state which is known to be one of two possibilities,  $|\psi_1\rangle$  or  $|\psi_2\rangle$ . Can she do a measurement to determine which it is?

If the two possible state are orthogonal, for example  $|\psi_1\rangle = |\uparrow\rangle$  and  $|\psi_2\rangle = |\downarrow\rangle$ , then the task is trivial. Alice can simply measure the projection operator observable  $P_\uparrow = |\uparrow\rangle\langle\uparrow|$ , which has eigenvalues 0 and 1. Depending on which result she gets, she knows which state she was given, with certainty.

However, if the two possible states are not orthogonal, then absolute certainty is not possible. Consider, for example, the problem of distinguishing

$$|\psi_1\rangle = |\uparrow\rangle, \quad |\psi_2\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle + |\downarrow\rangle). \quad (24.3.23)$$

Alice could again make a projective measurement with the observable  $P_\uparrow = |\uparrow\rangle\langle\uparrow|$ . If the state was  $|\psi_1\rangle$ , then she will certainly get the result 1, but then she cannot know which state it was,

because it could have been  $|\psi_2\rangle$ . On the other hand, if the state was  $|\psi_2\rangle$ , then she will have probability  $1/2$  to get each of the results 0 and 1 for  $P_{\uparrow}$ . If, and only if, the result was 0, then she will know with certainty that the state was not  $|\psi_1\rangle$ , so it must have been  $|\psi_2\rangle$ . Therefore, given an ensemble of experiments in which she is given the two states randomly with equal probability, Alice will know with certainty which state it was in only a fraction  $1/4$  of the trials.

Alice is clever and knows about generalized measurements, so she devises one with the following three POVM operators in matrix form in the  $|\uparrow\rangle, |\downarrow\rangle$  basis,

$$E_1 = \frac{a}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}, \quad E_2 = a \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad E_3 = \begin{pmatrix} 1 - a/2 & a/2 \\ a/2 & 1 - 3a/2 \end{pmatrix}, \quad (24.3.24)$$

where  $a$  is a real number to guarantee Hermiticity. Note that these operators are designed to satisfy  $\sum_n E_n = I$ . The positivity requirement limits the range of  $a$ , because requiring  $E_1$  and  $E_2$  to have non-negative eigenvalues and to be distinct operators implies  $a > 0$ , and requiring  $E_3$  to have non-negative eigenvalues implies  $a \leq 2 - \sqrt{2}$ . Now if Alice is given  $|\psi_1\rangle$ , you can use  $\mathcal{P}(n) = \langle \psi_1 | E_n | \psi_1 \rangle$  to calculate that the measurement outcome probabilities are

$$\mathcal{P}(1) = a/2, \quad \mathcal{P}(2) = 0, \quad \mathcal{P}(3) = 1 - a/2, \quad (24.3.25)$$

while if she is given  $|\psi_2\rangle$ , then the outcome probabilities are

$$\mathcal{P}(1) = 0, \quad \mathcal{P}(2) = a/2, \quad \mathcal{P}(3) = 1 - a/2. \quad (24.3.26)$$

So, if Alice measures the outcome  $n = 1$ , she will know with certainty that the state was  $|\psi_1\rangle$ , and if she measures  $n = 2$ , she will know that the state was definitely  $|\psi_2\rangle$ . Alice will never wrongly identify the state, but if instead the outcome is  $n = 3$ , then she gains no information. To maximize her discrimination ability, Alice should choose  $a$  to be the maximum allowed, which results in a probability  $a/2 = 1 - 1/\sqrt{2} \approx 0.293$  that she will be able to identify the state with absolute certainty, no matter which it is. In exchange for this opportunity to sometimes identify either state with certainty, the state will remain unknown if the outcome is  $n = 3$ , with probability  $1 - a/2 = 1/\sqrt{2} \approx 0.707$ .

## 24.4 Summary of rules of quantum mechanics for open systems

In the preceding sections of this chapter, we derived rules for open quantum systems based on models of interactions with an environment. An alternative is to treat these rules for open systems as fundamental principles, made without reference to an environment, and comparable to the postulates that we listed in section 3.1. For convenient reference and review, they are collected in this short section, for the case of a Hilbert space with finite dimension.

First, the state of an open system is specified, not by a ket, but by a density operator  $\rho$ , which must be Hermitian, unit-trace, and positive. In other words, it must satisfy  $\rho^\dagger = \rho$  and  $\text{Tr}(\rho) = 1$  and  $\langle \psi | \rho | \psi \rangle \geq 0$  for all kets  $|\psi\rangle$  in the Hilbert space on which it acts. Second, a generalized measurement corresponds to a collection of some number  $N$  (which can be arbitrarily large) of operators  $M_n$  that satisfy

$$\sum_{n=1}^N M_n^\dagger M_n = I. \quad (24.4.1)$$

In other words, the operators  $E_n = M_n^\dagger M_n$  form a positive operator-valued measure. Third, the possible outcomes of a generalized measurement are associated with the integers  $n = 1, \dots, N$  that label the measurement operators. Fourth, the probability of obtaining the result  $n$  for a generalized measurement is

$$\mathcal{P}(n) = \text{Tr}(\rho M_n^\dagger M_n) = \text{Tr}(\rho E_n). \quad (24.4.2)$$

Fifth, the state after the generalized measurement outcome  $n$  is given by the density operator

$$\rho_n = \frac{M_n \rho M_n^\dagger}{\mathcal{P}(n)}. \quad (24.4.3)$$

Sixth, the evolution of the open-system state is given by a trace-preserving completely positive dynamical map, which can be written in the form

$$\rho \rightarrow \rho' = \sum_n K_n \rho K_n^\dagger, \quad (24.4.4)$$

for some set of Kraus operators satisfying

$$\sum_n K_n^\dagger K_n = I. \quad (24.4.5)$$

This evolution is unitary only in the special case that there is only one Kraus operator.

Some sources give postulates allowing generalized measurements even for closed systems, which might be viewed as more permissive than the postulates we presented in section 3.1. However, since any system other than the whole universe is actually an open system, in practice this does not really allow anything new. This follows because we showed that generalized measurements can be derived from unitary time evolution and a projective measurement after adding an environment system to the open system being measured.

The sixth rule for open systems might be considered too vague or incomplete, since it contains no statement about how the Kraus operators are constructed or how they are related to the elapsed time of the evolution. This is because in general the time evolution of the density operator of an open system is only governed by a differential equation in an approximation, if at all. This is the subject of the next section.

## 24.5 Local time evolution approximation and the Lindblad equation

How do open systems evolve in time? For a closed system, we learned in eq. (3.5.21) that the density operator obeys a simple linear first-order differential equation,  $i\hbar\dot{\rho} = [H, \rho]$ , which was derived by applying the Schrödinger equation to the states in a mixed ensemble. It would be nice to have a similar differential equation for the density operator of an open system interacting with an environment, but it turns out that this is possible only in an approximation. Intuitively, the obstacle is that interactions transfer energy and information to the environment, from which their effects can return after a non-infinitesimal time delay. Even if the combined system has unitary time evolution, the effective dynamics of the open subsystem will not be local in time, and so cannot be expressed using a differential equation in  $t$  with a finite number of derivatives. Instead,  $\dot{\rho}(t)$  can depend not just on  $\rho(t)$ , but on  $\rho(t')$  for all earlier times  $t' < t$ .

Fortunately, in many cases the dynamics of an open system are still *approximately* local in time; this is sometimes called the **Markovian approximation**. This will occur if the environment tends to “forget” what has happened on time scales longer than some characteristic value  $\tau_{\text{env}}$ , for example by thermal averaging or dissipation processes. This is particularly plausible if the environment is very large, or if parts of it with direct coupling to the open system are a small subset of the whole. For example, the open system may emit light or vibrational quanta, which might be unlikely to return to it within any particular time interval. The fact that some of the energy flowing into the environment may be effectively lost forever also leads to damping of the open system on some time scale  $\tau_{\text{damping}}$ . Also relevant are time scales of order  $\hbar/\Delta E$ , where  $\Delta E$  are energy eigenstate differences for the open system treated in isolation, which would describe unitary time evolution of the open system if the environment were absent. Let us denote the smallest of those by  $\tau_{\text{unitary}}$ . If  $\tau_{\text{env}}$  is very short, then we can hope to describe the averaged dynamics, for  $dt$  much larger than  $\tau_{\text{env}}$  but much smaller than  $\tau_{\text{damping}}$  and  $\tau_{\text{unitary}}$ , by

$$\rho(t + dt) \approx \rho(t) + dt \mathcal{L}(\rho), \quad (24.5.1)$$

where

$$\mathcal{L}(\rho) = d\rho/dt \quad (24.5.2)$$

is called the **Liouvillian**. It is a superoperator map from the open system density operator to its approximate time derivative. Our next goal is to understand its form.

The density operator  $\rho(t + dt)$  can be obtained from  $\rho(t)$  by a dynamical map. As we learned in eq. (24.2.25), this can always be expressed in an operator-sum form with at most  $d^2$  Kraus operators,

$$\rho(t + dt) = \sum_{n=0}^{d^2-1} K_n \rho(t) K_n^\dagger, \quad (24.5.3)$$

where, for future convenience, we have chosen to start the index  $n$  from 0. To conserve probability, they must obey

$$\sum_{n=0}^{d^2-1} K_n^\dagger K_n = I \quad (24.5.4)$$

The assumed approximation of locality in time, eq. (24.5.1), implies that  $\rho(t + dt)$  must be linear in  $dt$ , and since it is quadratic in the Kraus operators, the latter must be quadratic in  $\sqrt{dt}$ . Therefore, we can write for each  $n$ ,

$$K_n = K_n^{(0)} + \sqrt{dt} K_n^{(1)} + dt K_n^{(2)}, \quad (24.5.5)$$

where we are consistently dropping contributions higher order in  $dt$ .

Now consider the limit  $dt = 0$ , which tells us that

$$\sum_{n=0}^{d^2-1} K_n^{(0)} \rho K_n^{(0)\dagger} = \rho \quad (24.5.6)$$

for all  $\rho$ . Choosing an orthobasis  $|\varphi_i\rangle$  for the open system, and looking at the specific case  $\rho = |\varphi_i\rangle \langle \varphi_i|$ , and applying  $\langle \varphi_j|$  from the left and  $|\varphi_j\rangle$  from the right, we learn that

$$\sum_{n=0}^{d^2-1} |\langle \varphi_j | K_n^{(0)} | \varphi_i \rangle|^2 = \delta_{ij} \quad (i, j = 1, \dots, d^2). \quad (24.5.7)$$

The left side is a sum of non-negative numbers, so this is only possible if  $\langle \varphi_j | K_n^{(0)} | \varphi_i \rangle = 0$  for each  $n$  when  $i \neq j$ , so the matrix representations of all  $K_n^{(0)}$  are diagonal. Since our orthobasis choice was arbitrary, and the only operators that are diagonal in every orthobasis are proportional to the identity, we must have  $K_n^{(0)} = k_n I$  for each  $n$ , for some complex numbers  $k_n$ , which must then satisfy  $\sum_n |k_n|^2 = 1$ . Now, by doing a unitary rotation among the Kraus operators of the form of eq. (24.2.21), we can arrange that one of the  $k_n$  (let us choose  $n = 0$ ) is equal to 1, and the others vanish.

Having selected  $K_0^{(0)} = I$ , and  $K_n^{(0)} = 0$  for all other  $n$ , next consider the parts of eqs. (24.5.3) and (24.5.4) proportional to  $\sqrt{dt}$ , which give, respectively,

$$K_0^{(1)} \rho + \rho K_0^{(1)\dagger} = 0, \quad K_0^{(1)} + K_0^{(1)\dagger} = 0. \quad (24.5.8)$$

It follows that  $K_0^{(1)}$  is anti-Hermitian and  $[K_0^{(1)}, \rho] = 0$  for every  $\rho$ , which together imply that  $K_0^{(1)} = 0$ . So, writing  $K_0^{(2)} = G - \frac{i}{\hbar} H$ , which defines Hermitian operators  $G$  and  $H$ , we have

$$K_0 = I + dt \left( G - \frac{i}{\hbar} H \right), \quad (24.5.9)$$

$$K_n = \sqrt{dt} L_n, \quad (n = 1, \dots, d^2 - 1). \quad (24.5.10)$$

In the last expression we have written  $K_n^{(1)} = L_n$  to agree with a traditional notation, and dropped  $K_n^{(2)}$  since it now only contributes to eq. (24.5.3) and (24.5.4) at order  $(dt)^{3/2}$ , due to the absence of  $K_n^{(0)}$ . Plugging eqs. (24.5.9) and (24.5.10) into eq. (24.5.4) gives

$$I = \sum_{n=0}^{d^2-1} K_n^\dagger K_n = I + dt \left( 2G + \sum_{n=1}^{d^2-1} L_n^\dagger L_n \right), \quad (24.5.11)$$

so that we can solve for

$$G = -\frac{1}{2} \sum_{n=1}^{d^2-1} L_n^\dagger L_n. \quad (24.5.12)$$

Therefore, using eqs. (24.5.9), (24.5.10), and (24.5.12) in eqs. (24.5.3) and then (24.5.1), we arrive at

$$\frac{d\rho}{dt} = \mathcal{L}(\rho) = -\frac{i}{\hbar} [H, \rho] + \sum_{n=1}^{d^2-1} \left( L_n \rho L_n^\dagger - \frac{1}{2} L_n^\dagger L_n \rho - \frac{1}{2} \rho L_n^\dagger L_n \right). \quad (24.5.13)$$

This is known as the **Lindblad equation** after Göran Lindblad; a slightly different form of it was obtained earlier by Vittorio Gorini, Andrzej Kossakowski, and E.C. George Sudarshan. It follows that the Liouvillian  $\mathcal{L}(\rho)$  is linear in  $\rho$ , and traceless, so that  $\text{Tr}[\rho(t)] = \text{Tr}[\rho(0)] = 1$ .

Equation (24.5.13) is the (Markovian approximation) generalization to open quantum systems of the unitary time evolution for closed systems, eq. (3.5.21). The operators  $L_n$  are known as **Lindblad operators**, or **jump operators** because the terms  $L_n \rho L_n^\dagger$  are responsible for transitions (“jumps”) in the open system that occur solely due to the influence of the environment. The Lindblad operator terms often produce damping due to a net energy flow from the open system to the environment, but on a much shorter time scale they can also cause an irreversible loss of coherence among the open system states. These effects will be explained and explored in more detail, with examples, in Chapter 25.

We can interpret  $H$  in eq. (24.5.13) as an effective Hamiltonian for the open system, although it need not be the same as the Hamiltonian for the limit in which interactions with the environment are removed. In fact,  $H$  is not even uniquely determined by the previous discussion. As an exercise, you can check that the Liouvillian  $\mathcal{L}(\rho)$  is unchanged under a transformation

$$L_n \rightarrow L_n + c_n I, \quad (24.5.14)$$

$$H \rightarrow H + \frac{i}{2} \hbar \sum_{n=1}^{d^2-1} (c_n L_n^\dagger - c_n^* L_n), \quad (24.5.15)$$

where the  $c_n$  are arbitrary complex numbers. This freedom can be used to choose a representation such that the Lindblad operators are each traceless,  $\text{Tr}(L_n) = 0$ . There is a further freedom to do unitary rotations of the Lindblad operators among each other without changing  $\mathcal{L}(\rho)$ , as  $L_n \rightarrow \sum_m v_{nm} L_m$ , where  $v_{nm}$  are the elements of a unitary  $(d^2 - 1) \times (d^2 - 1)$  matrix.

## 24.6 Exercises

**Exercise 24.1.** Consider the density matrix  $\rho_a = \text{Tr}_b(\rho_{ab})$  for subsystem  $a$  obtained by tracing over the complementary subsystem  $b$ , as in section 24.1.

- (a) Show that  $\rho_a$  is Hermitian, has trace 1, and is a positive operator.
- (b) Show that if there is a unitary time evolution for subsystem  $b$ , then  $\rho_a$  is unchanged.
- (c) Show that if an arbitrary projective measurement is made on subsystem  $b$ , but the results are unknown, then  $\rho_a$  is unchanged.

**Exercise 24.2.** Prove the following claims made about dynamical maps in section 24.2:

- (a) If  $\mathcal{E}_1$  and  $\mathcal{E}_2$  are dynamical maps with operator sum representations, then so is  $\mathcal{E}_2 \circ \mathcal{E}_1(\rho) = \mathcal{E}_2(\mathcal{E}_1(\rho))$ .
- (b) A dynamical map  $\mathcal{E}$  has an inverse  $\mathcal{E}^{-1}$  if and only if the map is unitary (can be defined with a single Kraus operator).

**Exercise 24.3.** For an arbitrary environment density operator  $\rho_b$  and unitary joint-state operator  $U$ , consider the dynamical map  $\mathcal{E}(\rho) = \text{Tr}_b[U(\rho \otimes \rho_b)U^\dagger]$ . Show that  $\mathcal{E}(\rho)$  can be written in operator-sum form with  $d_b^2$  Kraus operators.

**Exercise 24.4.** An open system has two orthobasis kets  $|0\rangle$  and  $|1\rangle$  representing the absence and presence of a photon respectively, as in the discussion involving eqs. (24.3.17)-(24.3.22). Consider a generalized measurement defined by operators

$$M_0 = |0\rangle\langle 0|, \quad M_1 = \sqrt{p}|0\rangle\langle 1| + \sqrt{1-p}|1\rangle\langle 1|, \quad (24.6.1)$$

which allows a photon some chance to survive the generalized measurement.

- (a) Check that the corresponding POVM operators are  $E_0 = |0\rangle\langle 0|$  and  $E_1 = |1\rangle\langle 1|$ .
- (b) For a general initial density matrix  $\begin{pmatrix} \rho_{00} & \rho_{01} \\ \rho_{10} & \rho_{11} \end{pmatrix}$ , find the probabilities of the outcomes  $n = 0$  and  $n = 1$ , and the post-measurement density operator and its entropy in each case.

**Exercise 24.5.** Show that the Liouvillian  $\mathcal{L}(\rho)$  defined by eq. (24.5.13) is not changed under the transformation of Lindblad operators in eqs. (24.5.14) and (24.5.15).

**Exercise 24.6.** Show that for time evolution governed by the Lindblad equation (24.5.13), the entropy cannot decrease if the Lindblad operators are all Hermitian.



## 25 Decoherence

### 25.1 Emergence of classical behavior

Quantum mechanics is most commonly applied to phenomena that, by the standards of everyday experience, involve very small amounts of time, distance, energy, and angular momentum. These processes include scattering and transitions involving electrons, photons, atoms, molecules, nuclei, and elementary particles. In contrast, our everyday experiences with larger objects are well-explained using classical physics. Some aspects of this are not too hard to understand. For example, the tiny value of  $\hbar$  when expressed in macroscopic measurement units immediately implies that in our daily lives we can usually ignore the position-momentum uncertainty principle, and the discrete quantization of angular momentum.

A more subtle feature is that superposition and interference never seem to play any role in our efforts to understand and describe our macroscopic experiences, despite being absolutely essential and ubiquitous features of quantum mechanics. When trying to hit, catch, or throw in baseball or cricket, it is not necessary to take into account the possibility that the ball might be in a superposition of states that suffer visible interference. While the smallness of  $\hbar$  again plays a role, there is still something to be explained: why is it so hard to prepare a baseball in a state where interference might become an obvious issue? Even the best pitchers in baseball and bowlers in cricket, whatever their other formidable skills in velocity, placement, and deception, never seem to achieve interference nodes or distinct wavepacket peaks separated on a scale of, say, centimeters. The explanation for this is **decoherence**, which can be defined as the quantum processes, due to interaction with the environment, by which certain non-classical-like coherent superposition states of macroscopic objects become impossible, for all practical purposes.

A famous example, known even in popular culture, is the thought experiment of Schrödinger's cat. A cat is confined to a box which also includes a radioactive nucleus which can decay in an infinitesimal time interval  $\Delta t$  with probability  $\Gamma \Delta t$ . If the decay occurs, an apparatus within the box kills the poor cat. As an idealization, the cat, the nucleus, and the lethal apparatus are all treated as a single quantum system undergoing unitary time evolution. After some time comparable to  $1/\Gamma$ , which could be hours or days for a suitably chosen nucleus, it might be supposed that the system will therefore be in a coherent superposition of states in which the cat has significant probability amplitudes to be both alive and dead, like

$$e^{-\Gamma t/2} |\text{alive}\rangle + \sqrt{1 - e^{-\Gamma t}} |\text{dead}\rangle. \quad (25.1.1)$$

This supposedly continues until we open the box and observe the contents, presumably only then collapsing the state to one in which the cat is definitely alive or dead.

Schrödinger presented this idealized picture as contrary to both common sense and experience. Real cats seem to always be in either a state  $|\text{alive}\rangle$  or  $|\text{dead}\rangle$ , and never in a superposition. However, general superpositions of states are perfectly valid and commonly encountered in quantum mechanics for electrons, and even for larger composite objects like nuclei and molecules. What makes the cat different? There are two obvious and related features to take into account. First, the cat is composed of a much larger number of constituent particles, something like  $10^{28}$  total electrons, protons, and neutrons, if it is well-fed. Surely this makes a difference. Second, contrary to the idealized assumption, the cat (and any apparatus potent enough to kill it) cannot be isolated in reality. Instead, the cat-apparatus-nucleus quantum system has an enormous number of interactions with the walls of the box, and even the photons in the cosmic microwave background radiation. These environmental interactions might be viewed as effectively conducting continuous measurements on the state of the cat's health, collapsing the state into one in which the cat is definitely dead or alive. While the influence of the environment is generally accepted to be crucial, the exact mechanism by which decoherence works is not obvious. Furthermore, we ought to be able to treat the walls of the box, and any photons within it, as also part of the quantum system undergoing unitary time evolution, although this might be prohibitively difficult to analyze.

Despite many remaining questions, a general picture of decoherence, pioneered by H. Dieter Zeh starting in the early 1970's and continuing with Wojciech Zurek a decade later, can be summarized qualitatively as follows. The dynamics of interacting subsystems entangles them, in such a way that the density operator of any particular open system evolves towards a mixed ensemble (not a superposition!) of special states, called **robust states** or **classical-like states** or **pointer states**. For macroscopic open systems, the entanglement involves an enormous number of degrees of freedom and is very rapid, even if the other effects of the environment are not so dramatic. The robust states are the ones that correspond to everyday experience. Other states, including almost all superpositions of the robust states, have such extremely low probabilities to endure after a very short time that they may as well not exist. The interactions of each open system with its environment selects the robust states for possible survival.

In terms of Schrödinger's thought experiment, the Hamiltonian coupling of the cat and its environment are supposed to pick out the states  $|\text{alive}\rangle$  and  $|\text{dead}\rangle$  as robust states. Instead of eq. (25.1.1), the density operator will evolve almost instantaneously towards the time-dependent diagonal form

$$\rho(t) = e^{-\Gamma t} |\text{alive}\rangle \langle \text{alive}| + (1 - e^{-\Gamma t}) |\text{dead}\rangle \langle \text{dead}|. \quad (25.1.2)$$

The missing off-diagonal terms in the density operator, proportional in this case to  $|\text{alive}\rangle \langle \text{dead}|$  and  $|\text{dead}\rangle \langle \text{alive}|$ , are called **coherences**, while the diagonal entries are called **populations**.

The key point is that the coherences are absent here because their coefficients are presumed to have a time dependence like  $e^{-\gamma_{\text{deco}} t}$  or perhaps  $e^{-\gamma_{\text{deco}}^2 t^2}$ , with an incredibly small decoherence time  $1/\gamma_{\text{deco}}$ , many orders of magnitude shorter than humans (or even cats) can resolve. In particular,  $1/\gamma_{\text{deco}}$  can easily be much shorter than the time scale  $1/\Gamma$  associated with the nuclear decay and the cat's demise. The absence of the coherences after a very short time asserts that, even before we open the box, for all practical purposes the cat is either dead or alive, rather than a coherent superposition. If we re-write this density operator in any other orthobasis, say,

$$\frac{1}{\sqrt{2}}(|\text{alive}\rangle + |\text{dead}\rangle), \quad \frac{1}{\sqrt{2}}(|\text{alive}\rangle - |\text{dead}\rangle), \quad (25.1.3)$$

it is not diagonal; evidently, these states are not robust because they do not decohere. Another way of saying the same thing is that for times much larger than  $1/\gamma_{\text{deco}}$ , the system is described almost perfectly as in eq. (25.1.2) by a mixed ensemble of only states in which the cat is either alive or dead, and not by the coherent superposition pure state in eq. (25.1.1),

Underpinning all of this is the idea that if we can calculate  $1/\gamma_{\text{deco}}$ , it will be shown to be tiny because  $\hbar$  is small in everyday units. However, calculating the details of this in terms of cats is too complicated (and cat-killing experiments are unlikely to be funded anyway), so we will resort to some much simpler models to illustrate how decoherence should work.

To explain in general terms how the coherences in a density operator could be reduced, consider an open system  $a$  coupled to an environment  $b$  in such a way that the unitary time evolution operators for the combined system have the special form

$$U = \sum_n (|\varphi_n\rangle \langle \varphi_n|) \otimes V_n, \quad (25.1.4)$$

where  $V_n$  are unitary operators acting on the environment, and the states  $|\varphi_n\rangle$  form an orthobasis, and will turn out to be the prototypical robust states of the open system in this example. Suppose that initially the density operator for the combined system has the form

$$\rho_{\text{com}} = \rho \otimes (|0\rangle \langle 0|), \quad (25.1.5)$$

where  $|0\rangle$  is some normalized reference environment state, perhaps its ground state. Writing  $\rho_{nm} = \langle \varphi_n | \rho | \varphi_m \rangle$  for the initial density operator matrix elements of the open system, the combined system density operator after the evolution is then

$$\rho'_{\text{com}} = U \rho_{\text{com}} U^\dagger = \sum_{n,m} \rho_{nm} (|\varphi_n\rangle \langle \varphi_m|) \otimes (V_n |0\rangle \langle 0| V_m^\dagger). \quad (25.1.6)$$

To find the density operator for the open system after the evolution, we trace  $\rho'_{\text{com}}$  over the environment subsystem using  $\text{Tr}_b(V_n |0\rangle \langle 0| V_m^\dagger) = \langle 0| V_m^\dagger V_n |0\rangle$ . Defining new unit-norm environment

states  $|\chi_n\rangle = V_n |0\rangle$ , we arrive at

$$\rho' = \sum_{n,m} \langle \chi_m | \chi_n \rangle \rho_{nm} |\varphi_n\rangle \langle \varphi_m|, \quad (25.1.7)$$

so that, for each  $n$  and  $m$ ,

$$\rho'_{nm} = \langle \chi_m | \chi_n \rangle \rho_{nm}. \quad (25.1.8)$$

Now, the unitarity of  $V_n$  ensures that  $\langle \chi_n | \chi_n \rangle = 1$ , so the diagonal elements  $\rho_{nn}$  of the open system density matrix are unaffected by the evolution. However, for the off-diagonal entries, the matrix elements  $\langle \chi_m | \chi_n \rangle$  have magnitude less than 1 for  $n \neq m$ , and so the coherences will be reduced in magnitude. In this example, the special set of states  $|\varphi_n\rangle$  that interact with the environment in the special form of eq. (25.1.4) are the ones whose populations remain intact, while their coherences are reduced.

The previous discussion leaves unanswered the question of why the coherences should decay exponentially (or faster) with time, rather than some more mild reduction. This is indeed the key question to explain decoherence for any particular system. For a toy model example to address this, consider an open system whose density operator evolves according to the Lindblad equation (24.5.13), but with the very simple assumption that there is only one non-zero Lindblad operator, which is proportional to the open system Hamiltonian itself,  $L = \sqrt{\alpha}H$ , where  $\alpha$  is a constant with units of  $1/(\text{energy})^2(\text{time})$ . We therefore have

$$\frac{d\rho}{dt} = -\frac{i}{\hbar}[H, \rho] + \alpha \left( H\rho H - \frac{1}{2}H^2\rho - \frac{1}{2}\rho H^2 \right). \quad (25.1.9)$$

This is straightforward to evaluate in an orthobasis of energy eigenstates satisfying  $H|\varphi_n\rangle = E_n|\varphi_n\rangle$ , with the result that the density operator matrix elements  $\rho_{nm} = \langle \varphi_n | \rho | \varphi_m \rangle$  satisfy

$$\frac{d}{dt}\rho_{nm} = \left[ -\frac{i}{\hbar}(E_n - E_m) - \frac{1}{2}\alpha(E_n - E_m)^2 \right] \rho_{nm}. \quad (25.1.10)$$

This is a particularly easy differential equation, with the solution

$$\rho_{nm}(t) = \exp \left( \left[ -\frac{i}{\hbar}(E_n - E_m) - \frac{1}{2}\alpha(E_n - E_m)^2 \right] t \right) \rho_{nm}(0). \quad (25.1.11)$$

The diagonal elements (populations) with  $n = m$  are constant in time. The off-diagonal elements (coherences) have a time-dependent phase, but more importantly their magnitudes suffer an exponential decay with mean lifetime  $2/\alpha(E_n - E_m)^2$ . In this particular simple example, the robust classical-like states that survive decoherence at late times are the energy eigenstates, but we will soon see that this is not a general feature.

## 25.2 Loss of coherence for a damped harmonic oscillator

In section 7.4, we observed that the energy eigenstates of the harmonic oscillator behave nothing like our macroscopic experience, as their expectation values for position and momentum do not change in time. This motivated us to introduce the quasi-classical, or coherent, states, which behave as much as possible like classical oscillators. These are indeed the robust states called for by the decoherence idea. Note that in this case, the robust states are not energy eigenstates. They also do not form an orthobasis, since no two coherent states are orthogonal.

However, noting the mere existence of such states is only solving half of the problem of classical behavior; the other half is to understand why they are inevitable for macroscopic masses and displacements. What would happen if we were somehow able to prepare a superposition of such states with a macroscopic position gap? How quickly does decoherence occur? Why are non-classical-like states essentially impossible to obtain? In this section we will show firstly that coherent states robustly survive the introduction of damping by an environment, and secondly that these states will decohere from each other at an incredibly fast rate.

Let us take a simple model for the damped harmonic oscillator as an open system, based on the Lindblad equation (24.5.13). To incorporate damping, we choose a single non-zero Lindblad operator

$$L = \sqrt{\gamma}a, \quad (25.2.1)$$

where  $\gamma$  is a positive real parameter with units of 1/(time), and  $a$  is the lowering operator of the oscillator. As we will confirm soon, this Lindblad operator evolves the open system by continuously lowering the energy, which is transferred to the environment. The environment therefore is effectively providing a frictional force. For the open system Hamiltonian in the Lindblad equation, we take  $H = \hbar\omega(a^\dagger a + 1/2)$ . So, the equation governing the time evolution of the density operator of the oscillator is

$$\frac{d\rho}{dt} = -i\omega[a^\dagger a, \rho] + \gamma\left(a\rho a^\dagger - \frac{1}{2}a^\dagger a\rho - \frac{1}{2}\rho a^\dagger a\right). \quad (25.2.2)$$

Our first goal is to show that there are pure coherent state solutions  $|\alpha(t)\rangle$  to this equation. Referring to the formula for coherent states, eq. (7.4.8), we therefore try

$$\rho = |\alpha\rangle\langle\alpha| = e^{-|\alpha|^2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} \frac{\alpha^{*m}}{\sqrt{m!}} |n\rangle\langle m|, \quad (25.2.3)$$

where  $\alpha$  is a function of time to be determined. It is convenient to proceed by defining the matrix elements

$$\rho_{nm} = \langle n|\rho|m\rangle = e^{-|\alpha|^2} \frac{\alpha^n}{\sqrt{n!}} \frac{\alpha^{*m}}{\sqrt{m!}}, \quad (25.2.4)$$

so that

$$\rho = \sum_{n,m} \rho_{nm} |n\rangle \langle m|. \quad (25.2.5)$$

One can now compute the various pieces needed in eq. (25.2.2),

$$\frac{d\rho}{dt} = \sum_{n,m} \left( \frac{n}{\alpha} \frac{d\alpha}{dt} + \frac{m}{\alpha^*} \frac{d\alpha^*}{dt} - \frac{d}{dt} |\alpha|^2 \right) \rho_{nm} |n\rangle \langle m|, \quad (25.2.6)$$

$$a^\dagger a \rho = \sum_{n,m} n \rho_{nm} |n\rangle \langle m|, \quad (25.2.7)$$

$$\rho a^\dagger a = \sum_{n,m} m \rho_{nm} |n\rangle \langle m|, \quad (25.2.8)$$

$$a \rho a^\dagger = \sum_{n,m} \sqrt{n} \sqrt{m} \rho_{nm} |n-1\rangle \langle m-1| = |\alpha|^2 \sum_{n,m} \rho_{nm} |n\rangle \langle m|, \quad (25.2.9)$$

where the last equality was obtained by using  $\sqrt{n} \sqrt{m} \rho_{nm} = |\alpha|^2 \rho_{n-1,m-1}$  and then relabeling the summation indices according to  $n-1 \rightarrow n$  and  $m-1 \rightarrow m$ . Plugging eqs. (25.2.6)-(25.2.9) into eq. (25.2.2), and collecting like terms, one finds for each  $n$  and  $m$

$$\left[ n \left( \frac{d}{dt} \ln \alpha + i\omega + \gamma/2 \right) + m \left( \frac{d}{dt} \ln \alpha^* - i\omega + \gamma/2 \right) - \frac{d}{dt} |\alpha|^2 - \gamma |\alpha|^2 \right] \rho_{nm} = 0. \quad (25.2.10)$$

These equations have the neat common solution

$$\alpha(t) = \alpha_0 \exp[-(i\omega + \gamma/2)t], \quad (25.2.11)$$

where  $\alpha_0$  is an arbitrary complex number that serves as the initial condition. Thus, a pure coherent state remains a pure coherent state, with a simple time dependence including an exponential damping. Note that with  $\gamma = 0$ , we recover the time evolution of the coherent state found in eq. (7.4.33). Recalling from eq. (7.4.37) that  $|\alpha|$  is proportional to the position amplitude of the coherent state oscillation, we see that the parameter  $\gamma$  is associated with a damping factor  $e^{-\gamma t/2}$  in the amplitude, and therefore  $e^{-\gamma t}$  in the energy.

To address the question of decoherence, suppose we somehow manage to prepare a pure state that is a superposition of two coherent states,

$$|\psi\rangle = \frac{1}{\sqrt{N}} (|\alpha_0\rangle + |\beta_0\rangle). \quad (25.2.12)$$

(Recall that coherent states are never orthogonal, so  $N = 2 + 2\text{Re}(\langle\beta_0|\alpha_0\rangle)$  is not equal to 2.)

The initial condition for the density operator is then

$$\rho(0) = |\psi\rangle \langle\psi| = \frac{1}{N} (|\alpha_0\rangle \langle\alpha_0| + |\beta_0\rangle \langle\beta_0| + |\alpha_0\rangle \langle\beta_0| + |\beta_0\rangle \langle\alpha_0|). \quad (25.2.13)$$

We have already seen how the first two terms in this density operator evolve, so we can make an educated guess

$$\rho(t) = \frac{1}{N}(|\alpha\rangle\langle\alpha| + |\beta\rangle\langle\beta| + F(t)|\alpha\rangle\langle\beta| + F(t)^*|\beta\rangle\langle\alpha|), \quad (25.2.14)$$

where, as in eq. (25.2.11),

$$\alpha = \alpha_0 \exp[-(i\omega + \gamma/2)t], \quad \beta = \beta_0 \exp[-(i\omega + \gamma/2)t], \quad (25.2.15)$$

and the other time dependence is in the decoherence factor  $F(t)$ , which remains to be found.

Plugging the guess of eq. (25.2.14) into the Lindblad equation (25.2.2), and using the same sort of strategy as in eqs. (25.2.4)-(25.2.10), one finds success provided that

$$\frac{d}{dt}F(t) = \gamma\left(-\frac{1}{2}|\alpha|^2 - \frac{1}{2}|\beta|^2 + \alpha\beta^*\right)F(t), \quad (25.2.16)$$

which can be rewritten as

$$\frac{d}{dt}\ln(F) = \left(-\frac{1}{2}|\alpha_0|^2 - \frac{1}{2}|\beta_0|^2 + \alpha_0\beta_0^*\right)\gamma e^{-\gamma t}. \quad (25.2.17)$$

Integrating both sides with respect to  $t$ , then exponentiating, and imposing the initial condition  $F(0) = 1$ , we find after rewriting the  $\alpha_0, \beta_0$  part slightly,

$$F(t) = \exp\left[\left(-\frac{1}{2}|\alpha_0 - \beta_0|^2 + i\text{Im}[\alpha_0\beta_0^*]\right)(1 - e^{-\gamma t})\right]. \quad (25.2.18)$$

When  $\alpha_0$  and  $\beta_0$  are macroscopic and different, this decoherence factor quickly approaches 0, eliminating the off-diagonal terms in  $\rho(t)$ . Expanding in the short time limit  $\gamma t \ll 1$  gives

$$F(t) \approx e^{i\Omega t} e^{-\gamma_{\text{deco}} t}, \quad (25.2.19)$$

where  $\Omega = \gamma\text{Im}(\alpha_0\beta_0^*)$  is a mostly uninteresting phase, and

$$\gamma_{\text{deco}} = \frac{1}{2}\gamma|\alpha_0 - \beta_0|^2 \quad (25.2.20)$$

is the much more interesting decoherence rate.

Equation (25.2.20) shows that the environmental energy damping  $\gamma$  is accompanied by a faster exponential damping of the coherences between classical-like states. Recall, from eq. (7.4.37), that  $\alpha_0$  is related to the position amplitude by  $\alpha_0 = \sqrt{\frac{m\omega}{2\hbar}}x_0$ . So, writing

$$|\alpha_0 - \beta_0|^2 = \frac{m\omega}{2\hbar}(\Delta x)^2, \quad (25.2.21)$$

we have

$$\frac{\gamma_{\text{deco}}}{\gamma} = \frac{m\omega(\Delta x)^2}{4\hbar}. \quad (25.2.22)$$

This shows that the very short decoherence time  $1/\gamma_{\text{deco}}$  for macroscopic states is indeed due to the smallness of  $\hbar$ . Let us put in some numbers to get an idea. Take the same undergraduate lab oscillator discussed in section 7.4, with  $m = 0.2$  kg and  $\omega = 10$  radians/second, and consider the decoherence of two coherent states with peaks  $\Delta x = 1$  centimeter apart. If the energy damping rate is a generous  $1/\gamma = 1000$  seconds, corresponding to a quite low-friction environment, then the time scale needed for a loss of coherence is

$$1/\gamma_{\text{deco}} \approx 2 \times 10^{-27} \text{ seconds}, \quad (25.2.23)$$

which is a tiny fraction of the time needed for light to traverse a proton's diameter.

One should certainly question whether the above estimate is accurate for a realistic case, since the physical processes leading to damping operate on much larger time scales than the  $1/\gamma_{\text{deco}}$  that we found. Real-world macroscopic oscillators have internal structures with all kinds of vibrational modes, and there are numerous environmental factors that contribute to damping in different ways not captured in our simple model. However, the key conceptual result stands: decoherence implies a ridiculously short time for a macroscopic system to become almost perfectly described by a mixed ensemble of robust coherent states, and never a superposition of such states subject to interference.

### 25.3 Measurement and decoherence: phase damping of a spin

The process of **phase damping** of a spin gives another instructive simple example of decoherence, without appealing to the Lindblad equation approximation.

Consider an open system  $a$  consisting of a single spin with orthobasis  $|\uparrow\rangle$  and  $|\downarrow\rangle$ , coupled to an environment  $b$  with three orthobasis states labeled  $|0\rangle$ ,  $|1\rangle$ , and  $|2\rangle$ . The environment starts in the reference state  $|0\rangle$ , and it subsequently evolves according to a dynamical map defined by a unitary operator on the combined system, with<sup>†</sup>

$$U(|\uparrow\rangle \otimes |0\rangle) = \sqrt{1-p} |\uparrow\rangle \otimes |0\rangle + \sqrt{p} |\uparrow\rangle \otimes |1\rangle, \quad (25.3.1)$$

$$U(|\downarrow\rangle \otimes |0\rangle) = \sqrt{1-p} |\downarrow\rangle \otimes |0\rangle + \sqrt{p} |\downarrow\rangle \otimes |2\rangle, \quad (25.3.2)$$

where  $0 < p < 1$ . This map has the property that it does not change the spin at all, but instead has some probability amplitude to move the environment to a different state that is correlated with the spin. If the spin is up, then the environment may be changed to the state  $|1\rangle$ , but if the spin is down, then it may be changed to the state  $|2\rangle$  instead. In this way, the dynamical map can

---

<sup>†</sup>Equations (25.3.1)-(25.3.2) only explicitly define  $U$  acting on 2 of the 6 orthobasis states for the combined system, but Theorem 24.2.2 assures us that, since it preserves inner products, it can be completed to a unitary operator on the whole combined Hilbert space. We will do this soon.



be thought of as making a sort of imperfect measurement on the open system, by entangling it with the environment. An observer who projectively measures the post-map environment state will have completed a generalized measurement, as discussed in section 24.3. If the observer finds it to be  $|1\rangle$ , the open system must be in the spin-up state, and if it is  $|2\rangle$ , it is spin-down. If found to be  $|0\rangle$ , the spin will remain unknown.

The dynamical map on the open system induced by  $U$  can be written in operator-sum form in terms of three Kraus operators  $K_0$ ,  $K_1$ , and  $K_2$ , one for each of the environment orthobasis states  $|0\rangle$ ,  $|1\rangle$ , and  $|2\rangle$ . To construct them, we compute their actions on the two open system orthobasis kets using eq. (24.2.4),

$$K_0 |\uparrow\rangle = \langle 0| U (|\uparrow\rangle \otimes |0\rangle) = \sqrt{1-p} |\uparrow\rangle, \quad (25.3.3)$$

$$K_0 |\downarrow\rangle = \langle 0| U (|\downarrow\rangle \otimes |0\rangle) = \sqrt{1-p} |\downarrow\rangle, \quad (25.3.4)$$

$$K_1 |\uparrow\rangle = \langle 1| U (|\uparrow\rangle \otimes |0\rangle) = \sqrt{p} |\uparrow\rangle, \quad (25.3.5)$$

$$K_1 |\downarrow\rangle = \langle 1| U (|\downarrow\rangle \otimes |0\rangle) = 0, \quad (25.3.6)$$

$$K_2 |\uparrow\rangle = \langle 2| U (|\uparrow\rangle \otimes |0\rangle) = 0, \quad (25.3.7)$$

$$K_2 |\downarrow\rangle = \langle 2| U (|\downarrow\rangle \otimes |0\rangle) = \sqrt{p} |\downarrow\rangle, \quad (25.3.8)$$

from which we learn

$$K_0 = \sqrt{1-p} I, \quad K_1 = \sqrt{p} |\uparrow\rangle \langle \uparrow|, \quad K_2 = \sqrt{p} |\downarrow\rangle \langle \downarrow|. \quad (25.3.9)$$

These satisfy  $K_0^\dagger K_0 + K_1^\dagger K_1 + K_2^\dagger K_2 = I$  as required, and the dynamical map is  $\mathcal{E}(\rho) = K_0 \rho K_0^\dagger + K_1 \rho K_1^\dagger + K_2 \rho K_2^\dagger$ . Given an initial density matrix for the spin,

$$\rho = \begin{pmatrix} \rho_{\uparrow\uparrow} & \rho_{\uparrow\downarrow} \\ \rho_{\downarrow\uparrow} & \rho_{\downarrow\downarrow} \end{pmatrix}, \quad (25.3.10)$$

the result of the dynamical map is found to be

$$\mathcal{E}(\rho) = \begin{pmatrix} \rho_{\uparrow\uparrow} & (1-p)\rho_{\uparrow\downarrow} \\ (1-p)\rho_{\downarrow\uparrow} & \rho_{\downarrow\downarrow} \end{pmatrix}. \quad (25.3.11)$$

In words, the spin populations (the diagonal entries) are unchanged, while the spin coherences in this basis are reduced by a factor  $1-p$ .

So far we have not associated the dynamical map with a specific elapsed time interval. This can be remedied by treating the dynamical map as corresponding to an infinitesimal time  $\Delta t$ , and then repeating the process a large number of times  $N$  to build up the result for a finite time  $t = N\Delta t$ . However, there are two ways to do this, which lead to quite different outcomes. For the first way, which we will call Version 1, we can suppose that each iteration starts from

the combined system state left over from the previous iteration. Since the environment is left in a different state than it started, this means that each dynamical map is actually a slightly different one than its predecessor. The second way, Version 2, proceeds by assuming that, after each iteration, the environment state is restored to  $|0\rangle$ , perhaps by some relaxation process not captured in our simple model. For example, the states  $|1\rangle$  and  $|2\rangle$  might quickly turn back into  $|0\rangle$  by emitting a photon that always leaves the scene without further effect. In that case, we just use the same map eq. (25.3.11) over and over.<sup>‡</sup> Let us see what these two versions predict.

To implement Version 1, the simplest way is to go back to the unitary map  $U$  defined in eq. (25.3.1)-(25.3.2), and complete it to the whole combined Hilbert space. Working in a matrix representation with the orthobasis  $|\uparrow\rangle \otimes |0\rangle, |\uparrow\rangle \otimes |1\rangle, |\downarrow\rangle \otimes |0\rangle, |\downarrow\rangle \otimes |2\rangle, |\uparrow\rangle \otimes |2\rangle, |\downarrow\rangle \otimes |1\rangle$ , in that order, you can check that the map is implemented by the unitary matrix

$$U = \begin{pmatrix} \sqrt{1-p} & -\sqrt{p} & 0 & 0 & 0 & 0 \\ \sqrt{p} & \sqrt{1-p} & 0 & 0 & 0 & 0 \\ 0 & 0 & \sqrt{1-p} & -\sqrt{p} & 0 & 0 \\ 0 & 0 & \sqrt{p} & \sqrt{1-p} & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}. \quad (25.3.12)$$

To realize this as an exponential of infinitesimal time maps, we invent a Hermitian Hamiltonian

$$H = \hbar\omega \begin{pmatrix} 0 & -i & 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -i & 0 & 0 \\ 0 & 0 & i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad (25.3.13)$$

so that  $U = \exp(-iHt/\hbar)$ , where we can now identify

$$\sqrt{p} = \sin(\omega t), \quad \sqrt{1-p} = \cos(\omega t). \quad (25.3.14)$$

In this interpretation, eq. (25.3.11) becomes

$$\mathcal{E}(\rho) = \begin{pmatrix} \rho_{\uparrow\uparrow} & \cos^2(\omega t) \rho_{\uparrow\downarrow} \\ \cos^2(\omega t) \rho_{\downarrow\uparrow} & \rho_{\downarrow\downarrow} \end{pmatrix}, \quad (\text{Version 1}), \quad (25.3.15)$$

where  $\cos^2(\omega t)$  has replaced  $1-p$  as the parameterization of the dynamical map. This is the result if the environment evolves continuously without being reset.

---

<sup>‡</sup>Note that the treatment of the decaying atomic state example surrounding eqs. (24.2.18)-(24.2.20) corresponds to Version 2 here, for the good reason explained there that the outgoing photon should not be relevant to the atom. This would be revealed in a more detailed treatment that took into account the photon kinematics.

To implement Version 2, we take  $p = \Gamma\Delta t = \Gamma t/N$  in the infinitesimal map, and then use  $\lim_{N \rightarrow \infty} (1 - \Gamma t/N)^N = e^{-\Gamma t}$  to write the  $N$ -fold iteration of eq. (25.3.11) as

$$\mathcal{E}(\rho) = \begin{pmatrix} \rho_{\uparrow\uparrow} & e^{-\Gamma t} \rho_{\uparrow\downarrow} \\ e^{-\Gamma t} \rho_{\downarrow\uparrow} & \rho_{\downarrow\downarrow} \end{pmatrix}, \quad (\text{Version 2}). \quad (25.3.16)$$

This is the result if the environment is continually being restored to the reference state  $|0\rangle$ .

Both versions lead to decoherence in the preferred basis of robust states  $|\uparrow\rangle$  and  $|\downarrow\rangle$ , but in different ways. In Version 1, the decoherence is quadratic in  $t$  for short times, with  $1 - p \approx 1 - \omega^2 t^2/2$ , and becomes complete at time  $t = \pi/2\omega$ . However, the density operator is periodic, and the coherences are restored at regular time intervals. In Version 2, the decoherence is linear in  $t$  for short times, but is monotonic, irreversible, and complete for long times. This irreversibility can be viewed as due to the fact that the information flowing to the environment is eliminated by the resetting of the environment state, so that any coherence present in the initial spin state will be lost forever. In both versions, after the dynamical map has taken place, the completion of a generalized measurement by projectively measuring the environment may provide some information about the open system state. The probability that the spin state will be successfully measured is equal to  $p$ , and so is maximized if  $p = 1$ , corresponding to complete decoherence. Both versions are only simplified models for how an environment will cause decoherence of a spin state, but in a realistic scenario there will almost always be some irreversible component to the loss of coherence. In the rapidly growing enterprise of quantum information, one often seeks practical ways to minimize this.

## 25.4 Decoherence for a spin in an exactly solvable model

In the example of decoherence of a damped harmonic oscillator in section 25.2, we used the Lindblad equation to model the interaction with the environment. However, the Lindblad equation is an approximation, so one might have residual doubts about the results obtained. In the case of phase damping of a spin in section 25.3, we studied two versions. In the first version, an exact unitary time evolution of the combined system resulted in complete decoherence, but this was temporary and the coherences were restored at regular intervals. The second version resulted in permanent decoherence at late times, but at the cost of appealing to a non-unitary reset of the environment state.

In this section we will study a slightly more complicated model<sup>†</sup> for decoherence of a spin, which can be solved exactly and leads to effectively complete decoherence at late times in a preferred basis of robust states, despite having quasi-periodic time-dependence. Since we can

---

<sup>†</sup>W.H. Zurek, Phys. Rev. D 26, 1862, (1982).

solve it exactly, we will not have to use the Markovian approximation of local time dependence for the open system, or appeal to non-unitary time evolution in the environment.

The open system of interest will consist of a single spin with orthobasis  $|\uparrow\rangle, |\downarrow\rangle$ , which will turn out to be the robust states. It interacts with an environment consisting of a large number  $N$  of spins, each with orthobasis  $|\uparrow\rangle_k, |\downarrow\rangle_k$  for  $k = 1, \dots, N$ . The Hamiltonian is taken to be

$$H = \sigma_z \otimes H_{\text{env}}, \quad (25.4.1)$$

where  $\sigma_z$  acts on the open-system spin, and the environment part of the Hamiltonian is

$$H_{\text{env}} = \frac{\hbar}{2} \sum_{k=1}^N \omega_k \sigma_z^{(k)}, \quad (25.4.2)$$

where  $\sigma_z^{(k)}$  acts on the environment spin  $k$ , with coupling constants  $\omega_k$  that have units of angular frequency. Since  $H$  is diagonal acting on each spin in our chosen orthobasis, it will be simple to exponentiate.

We start at time  $t = 0$  in a product density operator for the combined system,

$$\rho_{\text{com}}(0) = \rho(0) \otimes |0\rangle \langle 0|, \quad (25.4.3)$$

where the initial environment state is itself a product state,

$$|0\rangle = \bigotimes_{k=1}^N (\alpha_k |\uparrow\rangle_k + \beta_k |\downarrow\rangle_k), \quad (25.4.4)$$

with  $|\alpha_k|^2 + |\beta_k|^2 = 1$  for each  $k$ , and we write the initial open system density operator as

$$\rho(0) = \rho_{\uparrow\uparrow} |\uparrow\rangle \langle \uparrow| + \rho_{\uparrow\downarrow} |\uparrow\rangle \langle \downarrow| + \rho_{\downarrow\uparrow} |\downarrow\rangle \langle \uparrow| + \rho_{\downarrow\downarrow} |\downarrow\rangle \langle \downarrow|. \quad (25.4.5)$$

Now, using  $U = e^{-iHt/\hbar}$ , we obtain the time evolutions

$$U(|\uparrow\rangle \otimes |0\rangle) = |\uparrow\rangle \otimes |\chi_+\rangle, \quad U(|\downarrow\rangle \otimes |0\rangle) = |\downarrow\rangle \otimes |\chi_-\rangle, \quad (25.4.6)$$

where the unit-norm environment states correlated with the open system spins are

$$|\chi_{\pm}\rangle = \bigotimes_{k=1}^n (\alpha_k e^{\pm i\omega_k t/2} |\uparrow\rangle_k + \beta_k e^{\mp i\omega_k t/2} |\downarrow\rangle_k). \quad (25.4.7)$$

The combined system density operator at time  $t$  is therefore

$$\begin{aligned} \rho_{\text{com}}(t) = U \rho_{\text{com}}(0) U^\dagger &= \rho_{\uparrow\uparrow} |\uparrow\rangle \langle \uparrow| \otimes |\chi_+\rangle \langle \chi_+| + \rho_{\uparrow\downarrow} |\uparrow\rangle \langle \downarrow| \otimes |\chi_+\rangle \langle \chi_-| + \\ &\quad \rho_{\downarrow\uparrow} |\downarrow\rangle \langle \uparrow| \otimes |\chi_-\rangle \langle \chi_+| + \rho_{\downarrow\downarrow} |\downarrow\rangle \langle \downarrow| \otimes |\chi_-\rangle \langle \chi_-|. \end{aligned} \quad (25.4.8)$$

To obtain the reduced density operator for the open system, we need to trace  $\rho_{\text{com}}$  over all of the environment spins. This can be done easily by noticing that

$$\text{Tr}(|\chi_+\rangle \langle \chi_+|) = \langle \chi_+ | \chi_+ \rangle = 1, \quad (25.4.9)$$

$$\text{Tr}(|\chi_-\rangle \langle \chi_-|) = \langle \chi_- | \chi_- \rangle = 1, \quad (25.4.10)$$

$$\text{Tr}(|\chi_+\rangle \langle \chi_-|) = \langle \chi_- | \chi_+ \rangle = F(t), \quad (25.4.11)$$

where

$$F(t) = \prod_{k=1}^N (|\alpha_k|^2 e^{i\omega_k t} + |\beta_k|^2 e^{-i\omega_k t}), \quad (25.4.12)$$

which has initial value  $F(0) = 1$ . The resulting density operator, in the matrix representation in the  $|\uparrow\rangle, |\downarrow\rangle$  basis, is

$$\rho(t) = \begin{pmatrix} \rho_{\uparrow\uparrow} & F(t)\rho_{\uparrow\downarrow} \\ F(t)^*\rho_{\downarrow\uparrow} & \rho_{\downarrow\downarrow} \end{pmatrix}. \quad (25.4.13)$$

We therefore recognize that  $F(t)$  is the decoherence factor responsible for suppressing the coherences of the robust states  $|\uparrow\rangle$  and  $|\downarrow\rangle$ . Using the normalization condition  $|\alpha_k|^2 + |\beta_k|^2 = 1$ , you can show from eq. (25.4.12) that

$$|F(t)| = \prod_{k=1}^N \sqrt{1 - 4|\alpha_k|^2 |\beta_k|^2 \sin^2(\omega_k t)}, \quad (25.4.14)$$

which is a product of  $N$  factors, each between 0 and 1. If the frequencies  $\omega_k$  are not all commensurate, then  $|F(t)| < 1$  for all  $t > 0$ . However, because it is a product of periodic functions, it is quasi-periodic for any finite  $N$ , which means that it will return arbitrarily close to 1 for some sufficiently large  $t$ . This property is called<sup>‡</sup> **Poincaré recurrence**.

However, the Poincaré recurrence time  $t$  to have  $|F(t)| > 1 - \epsilon$  for any fixed  $\epsilon$  grows extremely quickly with  $N$ . So, for a sufficiently large environment with slightly different  $\omega_k$  values, it will take many orders of magnitude longer than the age of the universe, and therefore has no practical significance. For large  $N$ , we expect that the suppression of coherences is otherwise extremely efficient. An analysis by W.H. Zurek, F.M. Cucchietti, and J.P. Paz, Phys. Rev. A 72, 052113,

---

<sup>‡</sup>The recurrence property was originally pointed out as a general feature of Hamiltonian classical physics by Henri Poincaré in 1890. Any dynamical system that preserves phase-space volumes and has only bounded orbits will return arbitrarily close to its initial conditions, with a recurrence time that is finite, but quite possibly irrelevantly large, especially if the number of degrees of freedom is more than a few. For example, if you opened a bottle of classical gas molecules inside a sealed room, the molecules will in principle eventually all return to the open bottle, but obviously it is pointless to try to observe this phenomenon. For every practical purpose, it cannot happen, because the recurrence time scale is much too large.

arXiv:quant-ph/0312207, argues that for large  $N$  and  $\omega_k$  distributed randomly and with a finite standard deviation, for almost all  $t$  one will have Gaussian decoherence,

$$|F(t)| \approx e^{-\gamma_{\text{deco}}^2 t^2}, \quad \gamma_{\text{deco}}^2 = 2 \sum_{k=1}^N |\alpha_k|^2 |\beta_k|^2 \omega_k^2. \quad (25.4.15)$$

Indeed, this has the same expansion for small  $t$  as eq. (25.4.14). We conclude that the form of the environmental interactions selects  $|\uparrow\rangle$  and  $|\downarrow\rangle$  as robust states, and for large  $N$  ensures that at late times the open system will be described by a mixed ensemble of them, rather than a coherent superposition.

## 25.5 Exercises

**Exercise 25.1.** A spin-1/2 system obeys the Lindblad equation (24.5.13) with  $H = \omega S_z$  and a single jump operator  $L = \sqrt{\gamma} S_z / \hbar$ , where  $\omega$  and  $\gamma$  are constants with units 1/(time). This could be interpreted as a coupling of the spin to a magnetic field pointing in the  $\hat{z}$  direction, which also acts as an environment that monitors the spin.

(a) Find the differential equations for the four elements of the  $2 \times 2$  density matrix, and solve for them in terms of their initial conditions. What are the robust states that always survive decoherence at late times?

(b) Suppose that the system starts in the pure state  $(4|\uparrow\rangle + |\downarrow\rangle)/\sqrt{5}$ . Write the density matrix in the  $|\uparrow\rangle, |\downarrow\rangle$  basis as a function of  $t$ . Use it to find  $\langle S_z \rangle$  and  $\langle S_x \rangle$  as functions of time.

(c) Express the density matrix solution from the previous part in the diagonal form  $\rho(t) = p_1 |\psi_1\rangle \langle \psi_1| + p_2 |\psi_2\rangle \langle \psi_2|$ , identifying the eigenvalues  $p_1, p_2$ , and eigenvectors  $|\psi_1\rangle, |\psi_2\rangle$ .

**Exercise 25.2.** A (very) simplified model for quantum Brownian motion of a particle with mass  $m$  moving in one dimension uses a single Lindblad operator  $L = \sqrt{2\gamma} X / \lambda_{\text{th}}$ , where  $X$  is the position operator,  $\gamma$  is the dissipation rate characterizing energy exchange with the environment at temperature  $T$ , and  $\lambda_{\text{th}} = \hbar / \sqrt{2mk_B T}$  is the **thermal de Broglie wavelength**. Solve the Lindblad equation (24.5.13), ignoring the  $H$  term for extra simplicity, in position space in terms of  $\rho(x, x', t) = \langle x | \rho(t) | x' \rangle$ , to show that the exponential decay of spatial coherences is

$$\rho(x, x', t) = e^{-\gamma_{\text{deco}} t} \rho(x, x', 0), \quad (25.5.1)$$

where the ratio of the decoherence rate for wavefunction separations  $|x - x'|$  to the dissipation rate is

$$\frac{\gamma_{\text{deco}}}{\gamma} = \frac{(x - x')^2}{\lambda_{\text{th}}^2} = \frac{2mk_B T (x - x')^2}{\hbar^2}. \quad (25.5.2)$$

The diagonal part of the density matrix with  $x = x'$  is unaffected. What is  $\gamma_{\text{deco}}/\gamma$  for  $x - x' = 10^{-9}$  meters, for a nanoparticle of mass  $m = 10^{-18}$  grams at room temperature?

## 26 Invitation to quantum information

### 26.1 Qubits and the Bloch ball

In this chapter, we will briefly introduce some of the basic concepts of quantum information theory. This is an exciting and rapidly developing field, with the potential to revolutionize how computations are done and what they can accomplish. However, the practical realization of quantum computations faces significant challenges. Although some simple quantum computers have been built, it is not clear at present which of many types of physical implementations will scale in a useful way. Accordingly, our scope will be limited to theoretical considerations, which represent an idealized limit for what might be attainable in the real world. We will avoid some of the specialized terminology and tools that are very useful in quantum information theory, including quantum circuit diagrams. There are many references that treat the subject in much more depth and detail, including the free lecture notes of John Preskill, and the book *Quantum Computation and Quantum Information* by Michael A. Nielsen and Isaac L. Chuang.

In classical information theory, the basic unit of information is a bit, which can be either 0 or 1, and can be thought of as the answer to a single yes/no question. To encode or process information, one can use a register, which is an ordered collection of  $N$  bits. Since each bit has two possible values, there are  $2^N$  possible classical states for the register. The binary representation of the register is an integer  $n$ , in the range from 0 to  $2^N - 1$ . Thus, we can write

$$n = (01011010 \cdots 1) \quad (26.1.1)$$

to express the fact that the classical bit register can be used to store or transmit the information corresponding to  $n$ . Classical computations consist of operations on the register to change its value in some desired way.

The analogous basic unit of quantum information is a quantum bit, or **qubit**, a system with two orthobasis states. Physically, a qubit could be the spin state of an electron or nucleus, the polarization state of a photon or electromagnetic field mode in a cavity, a pair of energy levels for a trapped ion or a neutral atom, a spin or charge state of a semiconductor quantum dot, or the state of a superconducting circuit. In the quantum information context it is traditional to label the orthobasis states as  $|0\rangle$  and  $|1\rangle$ , rather than, say,  $|\uparrow\rangle$  and  $|\downarrow\rangle$  in the case of a spin-1/2 system. The most general pure state of a qubit is

$$|\psi\rangle = a|0\rangle + b|1\rangle, \quad (26.1.2)$$

where  $a$  and  $b$  are two complex numbers. Because the norm of the state has no physical significance, we can impose the restriction  $|a|^2 + |b|^2 = 1$ , eliminating one real degree of freedom. The

global phase is also not physical, eliminating a second real degree of freedom. Therefore, the physical information in a qubit amounts to one complex number, or two real numbers.

However, measurements on a quantum state can only extract partial information about it. For a single qubit, the most general observable can only have two possible results, since that is the dimension of the state space. For example, if the observable is  $\sigma_z$ , the measurement will project onto the basis vectors  $|0\rangle$  and  $|1\rangle$ . For a general qubit state  $a|0\rangle + b|1\rangle$ , the outcome of this measurement is not deterministic, with probabilities  $|a|^2$  and  $|b|^2$  for the two possible outcomes. Furthermore, the act of measurement collapses the qubit to one of  $|0\rangle$  or  $|1\rangle$ , permanently erasing some of the information it contained, as more measurements will reveal nothing about what it was initially. To do better, it is necessary to devise measurements that act on multiple qubits in a carefully chosen way, but even this can only determine some of the information contained in the quantum state. Much of quantum information theory amounts to extracting a desired subset of information from multiple entangled qubits.

A qubit will certainly interact with its environment. Sometimes this is what we want, as in planned operations such as quantum computations, or measurements to reveal partial information about its state. However, interactions with the environment can also be a curse that we want to suppress as much as possible. The environment can induce decoherence of the qubit, corrupting it in a way analogous to noisy errors in classical computations. In both cases, treating the qubit as an open system by tracing over its environment means that it can be described by a density matrix, as explained in section 24.1.

The density matrix for a qubit in the  $|0\rangle, |1\rangle$  basis is a  $2 \times 2$  Hermitian matrix with trace 1, for which the most general form is

$$\rho = \frac{1}{2} \begin{pmatrix} 1 + b_3 & b_1 - ib_2 \\ b_1 + ib_2 & 1 - b_3 \end{pmatrix}, \quad (26.1.3)$$

for some real numbers  $b_1, b_2$ , and  $b_3$ . By constructing the vector  $\vec{b} = b_1\hat{x} + b_2\hat{y} + b_3\hat{z}$ , we can rewrite the density matrix as

$$\rho = \frac{1}{2} \left( I + \vec{b} \cdot \vec{\sigma} \right), \quad (26.1.4)$$

where  $\vec{\sigma} = \sigma_x\hat{x} + \sigma_y\hat{y} + \sigma_z\hat{z}$  is the vector of Pauli matrices. The eigenvalues of  $\rho$  are

$$p_{\pm} = \frac{1}{2} \left( 1 \pm \sqrt{b_1^2 + b_2^2 + b_3^2} \right) = \frac{1}{2} \left( 1 \pm |\vec{b}| \right). \quad (26.1.5)$$

As we learned in section 3.5,  $p_+$  and  $p_-$  can be interpreted as the mixed ensemble probabilities for the orthobasis states consisting of the eigenvectors of  $\rho$ . As probabilities, they must be in the range from 0 to 1 inclusive, so  $\vec{b}$  must be a vector with length  $|\vec{b}| \leq 1$ .






Figure 26.1.1: The entanglement entropy  $\sigma = -p_+ \log(p_+) - p_- \log(p_-)$  for the density matrix of a qubit, as a function of the larger eigenvalue  $p_+$ . The normalization of  $\sigma$  is fixed here by using logarithms base 2. The entropy is maximum for the completely random ensemble with  $p_+ = 1/2$  and minimum for pure states with  $p_+ = 1$ , decreasing monotonically for larger information content.

The vector  $\vec{b}$  that specifies the density matrix of a qubit is called its **Bloch vector**, and it lies in the **Bloch ball** of radius 1, both named for Felix Bloch. In the case that  $\vec{b}$  is a unit vector, lying anywhere on the spherical surface of the Bloch ball, the qubit is in a pure state. A complete specification of such a pure state with  $p_+ = 1$  and  $p_- = 0$  requires the two spherical coordinate direction angles  $(\theta_b, \phi_b)$  of the Bloch vector, corresponding to the two real degrees of freedom mentioned above. The pure state is the eigenstate of  $\vec{b} \cdot \vec{\sigma}$  with eigenvalue 1,

$$|\psi\rangle = \cos(\theta_b/2) |0\rangle + e^{i\phi_b} \sin(\theta_b/2) |1\rangle. \quad (26.1.6)$$

To specify a mixed state of the qubit requires not just the direction but the length  $|\vec{b}|$ . The unique density matrix for the completely random ensemble has  $\vec{b} = 0$ , the point at the center of the Bloch ball, with  $p_+ = p_- = 1/2$ .

The information content in a qubit can be measured by the von Neumann entropy

$$\sigma = -p_+ \log(p_+) - p_- \log(p_-), \quad (26.1.7)$$

as defined in eq. (3.5.16), which we revisited as entanglement entropy in the context of bipartite systems in eq. (24.1.13). In eq. (26.1.7), we have normalized  $\sigma$  by using logarithms base 2. If the qubit's density matrix is that of a completely random ensemble, so that we have no information about its state, then the entropy is  $-2(1/2) \log(1/2) = 1$ . At the other extreme, if the qubit's state is perfectly known, so that the information is maximized, then  $p_+ = 1$  and  $p_- = 0$ , and the entropy vanishes. Figure 26.1.1 shows how  $\sigma$  depends on  $p_+$ , illustrating that the entropy decreases monotonically with more information content.

## 26.2 Quantum registers and parallelism

A **register** of  $N$  qubits is a tensor product state of the form

$$|\Psi\rangle = |\psi_1\rangle \otimes |\psi_2\rangle \otimes \cdots \otimes |\psi_N\rangle = |\psi_1 \psi_2 \cdots \psi_N\rangle. \quad (26.2.1)$$

The register has a canonical orthobasis of unentangled states of the form, for example,

$$|n\rangle = |01011010\cdots 1\rangle, \quad (26.2.2)$$

where  $n$  is a convenient integer label for the binary representation. This means that the complete information about the state of the register can be specified in terms of  $2^N$  complex numbers coefficients  $c_n$ , as

$$|\Psi\rangle = \sum_{n=0}^{2^N-1} c_n |n\rangle. \quad (26.2.3)$$

The norm of  $|\Psi\rangle$  has no physical significance, and can be set to 1 by requiring  $\sum_n |c_n|^2 = 1$ , eliminating one real degree of freedom. The global phase of  $|\Psi\rangle$  is also not observable, eliminating another real degree of freedom (say, the phase of one of the non-zero  $c_n$ ). This leaves  $2^N - 1$  complex numbers to completely specify the state of the register.

There are two general types of things that we can do with a register of qubits. First, we can apply a unitary transformation to it, which is equivalent to evolving it in time with some appropriate Hamiltonian, as we learned in section 3.4. However, in theoretical quantum computation studies, one often does not bother to specify either the elapsed time or the Hamiltonian. Instead, one simply assumes that *any* linear unitary operation on the qubit register is allowed, and can be realized somehow. The corresponding transformation is called a **quantum gate**, by analogy with logic gates in ordinary computation. The simplest gates operate on only one or two qubits at a time, and more complicated gates can be constructed out of simpler ones. Since unitary transformations are invertible, the information contained in the register can be recovered and is not changed. The second type of thing that one can do to a register is to conduct measurements on it to find out partial information about its state, for example to read a message or to learn something about the result of a computation. Measurements collapse the state, so some of the information that had been present in the register will necessarily be destroyed.

A first clue as to the power of quantum information over classical information is the enormous difference in information content. As we have just seen, a quantum register of  $N$  qubits stores  $2^N - 1$  continuous complex numbers worth of information, while the classical  $N$  bit register only stores an integer in the range from 0 to  $2^N - 1$ , equivalent to the responses to  $N$  yes/no questions. The possibility of entanglement of the qubits in the register is a big part of the quantum advantage, as most of the states with more than one non-zero coefficient  $c_n$  are entangled states. Entanglement is not just a puzzling phenomenon to be understood, but the key resource behind much of quantum information and computing capabilities.

To see the benefit of entanglement, consider a function  $f$ , which maps an integer  $n$  in the range from 0 to  $2^N - 1$  to another integer  $f(n)$  in the range from 0 to  $2^M - 1$ . The numbers  $n$

and  $f(n)$  can each be given a binary representation in an  $N$ -bit and  $M$ -bit register, respectively. If we want to evaluate all of the values of the function  $f$ , we would need to do the classical computation  $2^N$  times, one for each possible value of  $n$ . Quantum computation provides a powerful alternative. Consider a tensor product of an  $N$ -qubit register with an  $M$ -qubit register, and the map  $U_f$  defined by

$$U_f(|n\rangle \otimes |m\rangle) = |n\rangle \otimes |m + f(n)\rangle, \quad (26.2.4)$$

where the sum  $m + f(n)$  is taken modulo  $2^M$ . Since the states labeled  $|n\rangle \otimes |m\rangle$  form an orthobasis, by linearity this completely specifies the map for any state of the combined system. It is also a very short exercise to show that this map is unitary.

Now we apply  $U_f$  to a particular useful choice. For each qubit, define the orthobasis<sup>†</sup> of eigenstates of  $\sigma_x$ ,

$$|+\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle), \quad |-\rangle = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle). \quad (26.2.5)$$

and construct from  $|+\rangle$  the  $N$ -qubit register state  $|++\cdots+\rangle = |+\rangle \otimes |+\rangle \otimes \cdots \otimes |+\rangle$ . By expanding the tensor product, we find

$$|++\cdots+\rangle = \frac{1}{\sqrt{2^N}} \sum_{n=0}^{2^N-1} |n\rangle. \quad (26.2.6)$$

This is a very useful state, as it is an equal-weight superposition of all  $2^N$  canonical basis states of the  $N$ -qubit register. By linearity, the map defined by eq.(26.2.4) acts according to

$$U_f(|++\cdots+\rangle \otimes |m\rangle) = \frac{1}{\sqrt{2^N}} \sum_{n=0}^{2^N-1} |n\rangle \otimes |m + f(n)\rangle. \quad (26.2.7)$$

For each  $m$ , this single state is a superposition containing information about all of the results  $f(n)$  for every  $n$ . It does this by entangling the information for  $f(n)$  with the information for  $n$ . The point is that, in principle, we can use a single operation to perform the computational task simultaneously for all of the different  $n$ , which is  $2^N$  computations at once. This is not possible in classical computations, except by duplicating the hardware or by extending the computation time, which quickly becomes impractical for increasing  $N$ . This beautiful aspect of quantum information is called **parallelism**.

However, for parallelism to be useful it is not enough to perform the calculation; it is also necessary to find a way to extract information from the state. Although quantum registers

---

<sup>†</sup>Some jargon: in quantum information and computing, the unitary transformation between the  $\{|0\rangle, |1\rangle\}$  and  $\{|+\rangle, |-\rangle\}$  orthobases is called the **Hadamard transformation** or **Hadamard gate**, and is usually denoted  $H$  (not to be confused with a Hamiltonian). It obeys  $H = H^{-1} = H^\dagger = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$  in either basis.

can store and process vastly more information than classical registers, only a subset of the information can be extracted through measurement. For example, one might gain information about  $f(n)$  for some particular  $n$  by conducting a measurement that projects onto the input register state  $|n\rangle$ . But this collapses the state, permanently erasing all of the information about  $f(n)$  for other values of  $n$ . Fortunately, some global information about the function (that is, information that depends on the values of  $f(n)$  for multiple different inputs  $n$ ) can be extracted by making further use of superposition and devising measurements that are cleverer than just projecting onto  $|n\rangle$ .

To illustrate how this might work in a simple example, consider the following problem in quantum computation, due to Ethan Bernstein and Umesh Vazirani as a variation on earlier problems proposed and solved by David Deutsch and Richard Jozsa. To make things simple, suppose that the result for the function  $f(n)$  is either 0 or 1, so that the second register in the discussion above has  $M = 1$ . Furthermore, suppose the function  $f(n)$  is not known, but is coded into the unitary operation  $U_f$ , and is asserted to be of the form

$$f(n) = a \cdot n \pmod{2}, \quad (26.2.8)$$

for some unknown integer  $a$  in the same domain as  $n$ , from 0 to  $2^N - 1$ . The dot notation means

$$a \cdot n \equiv \sum_{k=1}^N a_k n_k, \quad (26.2.9)$$

where  $(a_1 a_2 \dots a_N)$  and  $(n_1 n_2 \dots n_N)$  are the binary representations of the integers  $a$  and  $n$ , with each  $a_k$  and  $n_k$  equal to 0 or 1. We would like to determine what  $a$  is. In classical information theory, we would need  $N$  evaluations of  $f(n)$  in order to determine the  $N$  bits  $a_k$ . However, using quantum mechanics, the problem can be solved using only one operation of  $U_f$  on a cleverly chosen input state, followed by one measurement.

To see how it works, let us first remark on a useful change of basis. The original canonical basis of  $N$ -qubit register states, based on  $|0\rangle$  and  $|1\rangle$  for the individual qubits, is

$$|n\rangle = |n_1 n_2 \dots n_N\rangle, \quad (\text{each } n_k = 0 \text{ or } 1). \quad (26.2.10)$$

Now, using the orthobasis  $|+\rangle$ ,  $|-\rangle$  for each qubit, we can construct another canonical basis for the  $N$ -qubit states,

$$|\tilde{s}\rangle = |\tilde{s}_1 \tilde{s}_2 \dots \tilde{s}_N\rangle, \quad (\text{each } \tilde{s}_k = + \text{ or } -). \quad (26.2.11)$$

Expanding the implicit tensor product gives the relation between the two orthobases,

$$|\tilde{s}\rangle = |\tilde{s}_1 \tilde{s}_2 \dots \tilde{s}_N\rangle = \frac{1}{\sqrt{2^N}} \sum_{n=0}^{2^N-1} (-1)^{s \cdot n} |n\rangle, \quad (26.2.12)$$

where the dot notation means

$$s \cdot n \equiv \sum_{k=1}^N s_k n_k, \quad \text{where } s_k = \begin{cases} 0 & \text{if } \tilde{s}_k = +, \\ 1 & \text{if } \tilde{s}_k = -. \end{cases} \quad (26.2.13)$$

Note that eq. (26.2.6) was a special case of this, in which all  $\tilde{s}_k = +$ , so that  $s \cdot n = 0$  for all  $n$ . Conversely, by taking the real inner product  $\langle n | \tilde{s} \rangle = \langle \tilde{s} | n \rangle$ , we find

$$|n\rangle = \frac{1}{\sqrt{2^N}} \sum_{\tilde{s}_k = \pm} (-1)^{s \cdot n} |\tilde{s}\rangle. \quad (26.2.14)$$

To solve the Bernstein–Vazirani problem, we start with the state  $|++ \cdots +\rangle \otimes |-\rangle$  and act on it with the unitary operator  $U_f$  defined as in eq. (26.2.4). The result, using eq. (26.2.7), is

$$U_f(|++ \cdots +\rangle \otimes |-\rangle) = \frac{1}{\sqrt{2^{N+1}}} \sum_{k=0}^{2^N-1} |n\rangle \otimes (|f(n)\rangle - |f(n)+1\rangle). \quad (26.2.15)$$

Recall that the addition in the last ket label is taken mod 2. It follows that  $|f(n)\rangle - |f(n)+1\rangle = (-1)^{f(n)}(|0\rangle - |1\rangle) = (-1)^{a \cdot n}(|0\rangle - |1\rangle)$ , and therefore

$$U_f(|++ \cdots +\rangle \otimes |-\rangle) = \frac{1}{\sqrt{2^N}} \sum_{k=0}^{2^N-1} (-1)^{a \cdot n} |n\rangle \otimes |-\rangle. \quad (26.2.16)$$

Using eq. (26.2.12), this can be rewritten as simply

$$U_f(|++ \cdots +\rangle \otimes |-\rangle) = |\tilde{a}_1 \tilde{a}_2 \cdots \tilde{a}_N\rangle \otimes |-\rangle, \quad (26.2.17)$$

with

$$\tilde{a}_k = \begin{cases} + & \text{if } a_k = 0, \\ - & \text{if } a_k = 1. \end{cases} \quad (26.2.18)$$

Now we conduct a measurement by projecting the state onto the  $|\tilde{s}\rangle$  basis, using the observable

$$Q = \left( \sum_{\tilde{s}_k = \pm} s |\tilde{s}\rangle \langle \tilde{s}| \right) \otimes I. \quad (26.2.19)$$

This solves the problem, since exactly one of the  $|\tilde{s}\rangle$  is  $|\tilde{a}_1 \tilde{a}_2 \cdots \tilde{a}_N\rangle$ . With probability 1, the result of the measurement of  $Q$  on the state  $U_f(|++ \cdots +\rangle \otimes |-\rangle)$  will be the number  $a$ .

As promised, quantum information methods solve the Bernstein–Vazirani problem with only one unitary transformation and one measurement, a task that would take  $N$  evaluations of  $f(n)$  with classical computation methods. This is not necessarily very impressive, as doing 1 computation with  $N+1$  qubits is not obviously easier (with present technology) than doing  $N$  classical computations on  $N$  bits. However, it shows the general principle that quantum

parallelism provides advantages. Although it is beyond our scope to show here, there are some computation problems that are believed to be not feasible in practical terms using classical methods, but become easy (in principle) by exploiting quantum parallelism in more sophisticated ways that exploit entanglement and superposition.

The most famous example is Peter Shor's 1994 breakthrough quantum computing algorithm for factoring an integer  $n$  as a product of primes. The reason for its fame is that it cracks the most popular modern public-key encryption schemes, with profound implications for human events far beyond mere scientific curiosity. In these schemes, anyone can encrypt a message using a public key that depends on some very large integer  $n$ , but to decrypt and read the message one must know the prime factorization of  $n$ , which is not publicly provided. Thus, the security of the code relies on the widely held, but unproved, belief that if  $n$  has more than a few hundred digits, prime factorization is a hard problem for classical computing,<sup>‡</sup> with a computation time that grows faster than any polynomial in the number of bits of  $n$ . The running time for the best known algorithm (the general number field sieve) scales for large  $n$  approximately like  $\exp([(64/9)(\log n)(\log \log n)^2]^{1/3})$ . In contrast, Shor's factoring algorithm on a quantum computer should have a running time that scales like only  $(\log n)^2(\log \log n)(\log \log \log n)$ . The quantum algorithm is not deterministic, and gives the answer only with high probability rather than unit probability, but checking a candidate factorization is trivial in terms of computation time, as it requires only one multiplication. As of this writing, the immediate implications of Shor's factoring algorithm are only academic, since implementation of it for large  $n$  on a quantum computer is not feasible yet,<sup>§</sup> for mundane practical reasons. However, those with important secrets to keep should not bet on this remaining true forever.

Shor's factoring algorithm, and algorithms for many other problems that are classically hard, makes use of the **quantum Fourier transform**, a unitary map defined by

$$U_{\text{FT}} |n\rangle = \frac{1}{\sqrt{2^N}} \sum_{k=0}^{2^N-1} \exp(2\pi i n k / 2^N) |k\rangle \quad (26.2.20)$$

This provides a change of basis that is useful for parallelizing calculations and extracting relevant information through measurement. It is crucial that there exists an efficient way of implementing the quantum Fourier transform as a subroutine acting on qubit registers. The factoring problem is a special case of a more general type of problem called the hidden subgroup problem, which contains many other interesting and useful examples that can be attacked in similar ways by quantum computing. An example of a completely different type of problem is Grover's search algorithm, discovered by Lov Grover in 1996, which uses quantum parallelism to give a quadratic

---

<sup>‡</sup>At least, this is true for *publicly known* classical algorithms.

<sup>§</sup>At least, this is true for *publicly known* quantum computing hardware.

speed improvement over classical search methods. It is also worth mentioning that quantum computation may be the natural way to attack difficult quantum mechanics problems.

## 26.3 Sending information using entanglement: teleportation and dense coding

Transmission of information using classical bits is a well-exploited feature of modern life, underlying much familiar technology. An electromagnetic wave signal can be modulated in various patterns to represent the 0's and 1's in a register. In quantum information, the problem is more subtle, since even a single qubit contains a continuum of information.

Let us establish some notation before continuing. First, the Pauli matrices acting on qubits will occur often, and position operators will not appear at all, so throughout the rest of this chapter we will follow a common practice in the quantum information literature by writing

$$X \equiv \sigma_x, \quad Y \equiv \sigma_y, \quad Z \equiv \sigma_z, \quad (26.3.1)$$

so that  $Z|0\rangle = |0\rangle$  and  $Z|1\rangle = -|1\rangle$ , and  $X|0\rangle = |1\rangle$  and  $X|1\rangle = |0\rangle$ , and  $Y|0\rangle = i|1\rangle$  and  $Y|1\rangle = -i|0\rangle$ . When there is more than one qubit, a subscript on  $X$ ,  $Y$ , or  $Z$  may be used to indicate which qubit is being acted on. The simplest cases will involve entanglement of two qubits, like the one familiar to Alice and Bob from the Einstein–Podolsky–Rosen and Bohm problem in section 14.1. Let us define four maximally entangled states of a 2-qubit system,

$$|\phi^\pm\rangle = \frac{1}{\sqrt{2}}(|00\rangle \pm |11\rangle), \quad (26.3.2)$$

$$|\psi^\pm\rangle = \frac{1}{\sqrt{2}}(|01\rangle \pm |10\rangle). \quad (26.3.3)$$

These are called **Bell states** or **EPRB states**. In many, but not all, cases of interest, the two qubits are spatially separated, after which they cannot be directly accessed simultaneously by a single observer. If we identify  $|0\rangle$  and  $|1\rangle$  with spin-up and spin-down respectively, then the state featured in the EPRB discussion in section 14.1 was  $|\psi^-\rangle$  in the present notation, but the problem could have been equally well formulated in terms of any one of the other Bell states instead. Note that  $|\phi^+\rangle$ ,  $|\phi^-\rangle$ ,  $|\psi^+\rangle$ ,  $|\psi^-\rangle$  form a complete orthobasis for the two-qubit system, and the four unentangled states  $|00\rangle$ ,  $|01\rangle$ ,  $|10\rangle$ , and  $|11\rangle$  form another orthobasis.

There is a very useful measurement that projects any 2-qubit state onto the four Bell states. The corresponding observable may be defined explicitly as

$$Q = |\phi^+\rangle\langle\phi^+| + 2|\phi^-\rangle\langle\phi^-| + 3|\psi^+\rangle\langle\psi^+| + 4|\psi^-\rangle\langle\psi^-|, \quad (26.3.4)$$

with possible outcomes equal to the four eigenvalues 1, 2, 3, and 4. The result of the measurement of  $Q$  can therefore be expressed using 2 classical bits. This is an example of a **collective measurement**, as it acts nontrivially on more than one qubit.

Now consider the following problem: Alice has a qubit state

$$|\gamma\rangle_c = c_0 |0\rangle + c_1 |1\rangle, \quad (26.3.5)$$

which she would like to transmit to Bob in a distant location, without sending the physical qubit. Can this be accomplished by transmitting only classical bits? Naively, it might seem impossible, since the qubit contains two continuous real numbers worth of information, while the classical bits to be transmitted are only a discrete set. Indeed, it will not do for Alice to make any kind of measurement on the qubit  $|\gamma\rangle_c$  alone. For example, measuring  $Z$  by projecting onto the basis  $|0\rangle$  and  $|1\rangle$  will not allow reconstruction of  $|\gamma\rangle_c$ , and even worse, it will collapse the state, making further measurements useless. However, it can be done, provided that Alice and Bob already share a known Bell state, which was prepared for this occasion, and upon which they have not yet conducted any measurements. It is not important which entangled Bell state they share, but to be concrete let us suppose it is  $|\phi^+\rangle_{ab}$ . Alice can act on the first qubit of this entangled pair either by measuring it or by unitarily evolving it, but she is not able to do anything with the second qubit. The converse is true for Bob in the obvious way.

To accomplish the task, Alice makes a collective measurement on the two qubits that she has access to, labeled  $a$  (the one entangled with Bob's qubit  $b$ ) and  $c$  (the one to be sent). Specifically, Alice measures the observable  $Q_{ac}$  from eq. (26.3.4), projecting onto the four Bell orthobasis states  $|\phi^+\rangle_{ac}$ ,  $|\phi^-\rangle_{ac}$ ,  $|\psi^+\rangle_{ac}$ , and  $|\psi^-\rangle_{ac}$ . (This is equivalent to the simultaneous measurement of the two commuting observables  $Z_a Z_c$  and  $X_a X_c$ , as you will show in Exercise 26.3.) The result of Alice's measurement will be one of four possibilities, corresponding to these four orthogonal states. She then sends this information, equivalent to only two classical bits, to Bob by some ordinary communication channel. Bob then applies a unitary operator  $U_b$  to his qubit of the shared Bell state, following this rule:

$$U_b = I, \quad \text{if Alice's measurement result corresponded to } |\phi^+\rangle_{ac}, \quad (26.3.6)$$

$$U_b = Z, \quad \text{if Alice's measurement result corresponded to } |\phi^-\rangle_{ac}, \quad (26.3.7)$$

$$U_b = X, \quad \text{if Alice's measurement result corresponded to } |\psi^+\rangle_{ac}, \quad (26.3.8)$$

$$U_b = -iY, \quad \text{if Alice's measurement result corresponded to } |\psi^-\rangle_{ac}. \quad (26.3.9)$$

As we are about to show, this will put Bob's qubit into the state  $(c_0 |0\rangle + c_1 |1\rangle)_b$ , an exact copy of the state that Alice wanted to transfer. This is called **quantum teleportation** of the qubit from Alice to Bob.

To see why the procedure works, first note that we can write the initial joint state of all



three qubits as

$$|\phi^+\rangle_{ab} \otimes |\gamma\rangle_c = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)_{ab} \otimes (c_0|0\rangle + c_1|1\rangle)_c \quad (26.3.10)$$

$$= \frac{1}{\sqrt{2}}(c_0|000\rangle + c_0|110\rangle + c_1|001\rangle + c_1|111\rangle)_{abc}. \quad (26.3.11)$$

Now the key trick: you can check by a little routine expansion that this is also equal to

$$\frac{1}{2}|\gamma\rangle_b \otimes |\phi^+\rangle_{ac} + \frac{1}{2}(Z|\gamma\rangle)_b \otimes |\phi^-\rangle_{ac} + \frac{1}{2}(X|\gamma\rangle)_b \otimes |\psi^+\rangle_{ac} + \frac{1}{2}(iY|\gamma\rangle)_b \otimes |\psi^-\rangle_{ac}. \quad (26.3.12)$$

When Alice makes her measurement of  $Q_{ac}$ , the state is projected onto one of these four terms, and her ordinary 2-bit message to Bob tells him which one. Bob's application of the appropriate unitary operator selected from the list in eqs. (26.3.6)-(26.3.9) then turns qubit  $b$  into the state  $|\gamma\rangle_b$ , due to the Pauli-matrix relations  $X^2 = Y^2 = Z^2 = I$ .

The teleportation of the qubit has several interesting features to keep in mind. First, it is only possible because of entanglement; using an unentangled qubit pair in place of  $|\phi^+\rangle_{ab}$  would certainly not work. Second, Alice's original qubit  $|\gamma\rangle_c$  is necessarily destroyed by the process of teleporting it to Bob, in the sense that its state is collapsed by Alice's measurement on it. This means that the qubit has been transferred, but not cloned, which turns out to be impossible on general grounds as we will show in section 26.4. Third, neither Alice nor Bob has learned anything about the teleported state during the process. No matter what the teleported qubit was, Alice's measurement has equal probabilities 1/4 for each of the four outcomes, and the measurement result only gives information about the joint state of the qubits  $a$  and  $c$ , not  $c$  alone. For his part, Bob has only applied a unitary transformation, not measured anything. Finally, the teleportation of the state is definitely not instantaneous communication, as the transfer speed is limited by that of the two classical bits sent from Alice to Bob.

Teleporting a whole register of  $N$  qubits (including their possible entanglements with each other) is a straightforward generalization, as can be understood using the same ideas, with some proliferation of indices. Alice and Bob will need to start with  $N$  prepared Bell pairs,

$$|\Psi\rangle_{ab} = |\phi^+\rangle_{a_1b_1} \otimes \cdots \otimes |\phi^+\rangle_{a_Nb_N}, \quad (26.3.13)$$

which can be thought of as an entangled state living in the tensor product of an  $N$ -qubit register controlled by Alice and an  $N$ -qubit register controlled by Bob. Let us write the register to be teleported from Alice to Bob as

$$|\Gamma\rangle_c = \sum_{\{\gamma_k\}} c_{\gamma_1 \cdots \gamma_N} |\gamma_1 \cdots \gamma_N\rangle, \quad (26.3.14)$$

where each  $\gamma_k$  is summed over 0 and 1. The initial joint state is therefore a  $3N$ -qubit register,

$$|\Psi\rangle_{ab} \otimes |\Gamma\rangle_c = \sum_{\{\gamma_k\}} c_{\gamma_1, \dots, \gamma_N} \left( |\phi^+\rangle_{a_1b_1} \otimes |\gamma_1\rangle_{c_1} \right) \otimes \cdots \otimes \left( |\phi^+\rangle_{a_Nb_N} \otimes |\gamma_N\rangle_{c_N} \right). \quad (26.3.15)$$

Now each of the  $N$  factors in parentheses can be re-expressed, using exactly the same trick as for a single qubit, as

$$\begin{aligned} |\phi^+\rangle_{a_k b_k} \otimes |\gamma_k\rangle_{c_k} &= \frac{1}{2} |\gamma_k\rangle_{b_k} \otimes |\phi^+\rangle_{a_k c_k} + \frac{1}{2} (Z |\gamma_k\rangle)_{b_k} \otimes |\phi^-\rangle_{a_k c_k} \\ &\quad + \frac{1}{2} (X |\gamma_k\rangle)_{b_k} \otimes |\psi^+\rangle_{a_k c_k} + \frac{1}{2} (iY |\gamma_k\rangle)_{b_k} \otimes |\psi^-\rangle_{a_k c_k}. \end{aligned} \quad (26.3.16)$$

For each  $k = 1, \dots, N$ , Alice makes a projective measurement  $Q_{a_k c_k}$  on the four Bell states for the two qubits  $a_k, c_k$ , and sends the results to Bob as  $2N$  classical bits of information. Bob then applies an appropriate unitary transformation

$$U_b = U_{b_1} \otimes U_{b_2} \otimes \dots \otimes U_{b_N} \quad (26.3.17)$$

on his register, with each  $U_{b_k}$  chosen just as in eqs. (26.3.6)-(26.3.9). The result will be that the  $N$ -qubit register Bob controls will be left in the state  $|\Gamma\rangle_b$ .

Now we turn to a question that may at first seem unrelated, but is actually just the inverse of quantum teleportation. Suppose that Alice needs to send a message composed of classical bits to Bob. She decides to encode the information using qubits,<sup>†</sup> since she knows that they are denser in information than classical bits. How much classical bit information can Alice send to Bob, encoded in a register of  $N$  qubits?

First consider what happens if Alice only sends a single unentangled qubit. Clearly, she can use it to encode a single classical bit, for example by preparing the qubit in one of the two states  $|0\rangle$  and  $|1\rangle$ , which will then be measured by Bob using the observable  $Z$ . A little thought reveals that this is also the best that can be done. No matter how Alice prepares the qubit, any measurement that Bob can do will have one of at most 2 results, since that is the dimension of the 1-qubit state space. Importantly, Bob's measurement will also collapse the qubit's state into a completely known one, making all further measurements useless. So only one classical bit can be sent with a single unentangled qubit.

But now suppose that Alice and Bob share a known Bell state, prepared earlier, say  $|\phi^+\rangle$ , with each having excess to one qubit of the pair. To send her message, Alice does one of the 4 unitary transformations  $U = I, Z, X$ , or  $iY$  on the qubit  $a$  that she controls, causing the Bell state to change in one of the following ways:

$$I_a |\phi^+\rangle = |\phi^+\rangle, \quad (26.3.18)$$

$$Z_a |\phi^+\rangle = |\phi^-\rangle, \quad (26.3.19)$$

$$X_a |\phi^+\rangle = |\psi^+\rangle, \quad (26.3.20)$$

$$iY_a |\phi^+\rangle = |\psi^-\rangle. \quad (26.3.21)$$

---

<sup>†</sup>The physical delivery of a qubit over a distance is a challenge, not the least because one must avoid decoherence due to environmental interactions. Photonic polarization qubits can be transmitted with lasers or over optical fibers, but direct transportation of particle spins is much more difficult. We consider the idealized case.

Alice sends the qubit  $a$  to Bob, who now controls both qubits. Bob can measure the observable  $Q$  from eq. (26.3.4), projecting onto the Bell state basis and giving the result 1, 2, 3, or 4 depending on the choice Alice made. Thus, Alice has successfully transmitted two classical bits to Bob, by sending only one qubit. The transfer of two classical bits by sending one entangled qubit is known as **superdense coding**, or slightly more modestly as **dense coding**. Entanglement effectively doubles the information capacity of the communication.

One might object that dense coding requires the involvement of two qubits to encode and send two classical bits. However, it is also true that Alice never had direct access to Bob's qubit. In fact, the Bell pair could have been created and separated to the two parties long before Alice decided what message to send. Bob could have even prepared the Bell state himself, then sent Alice's share to her. Then, by modifying the one qubit and sending it back, Alice accomplished the mission (perhaps without even knowing for sure that Bob's share of the Bell pair existed).

Another benefit to the dense coding protocol, likely more important than information density, is security. Alice and Bob have an evil nemesis, Eve, who wants to eavesdrop on their conversations. If Alice sent classical bits, then Eve might be able to read them by listening in. If a single unentangled qubit is sent instead, then Eve could still read it by measuring the observable  $Z$ , provided that she can intercept the qubit and knows that it is encoded in the  $|0\rangle$ ,  $|1\rangle$  basis. Eve could then send the qubit on its way to Bob, without Alice or Bob ever knowing. But, in the dense coding method, any measurement done by Eve provides her no information, because Eve only has access to the half of the entangled pair that was sent by Alice. A formal way of expressing this is that the density matrix for the qubit sent by Alice, obtained by tracing over the qubit held by Bob, is  $\rho_a = I/2$ , corresponding to a completely random ensemble.

A new pristine entangled qubit is needed for each pair of classical bits transmitted, as they are "used up" by the measurement collapse needed by Bob to extract the message. Therefore, by sending a register of  $N$  qubits, Alice can transmit  $2N$  classical bits of information extractable by Bob. This is the converse of the result found above that Alice can send  $2N$  classical bits to teleport a general  $N$ -qubit register.

We have seen that in teleportation, sending two classical bits is enough to transmit one qubit, while in dense coding sending one qubit is enough to transmit two classical bits. In that sense, they are opposites or inverses. However, one should be careful not to claim that a qubit is equivalent to two classical bits; the statements refer to the optimal extraction of information aided by entanglement, rather than equivalence of information contained.

## 26.4 Copying quantum information: the no-cloning theorem and redundant encoding

Suppose we have a system in a state  $|\psi\rangle$ , which we would like to copy without changing the original. Let us try to do this by producing the copy on another system of the same type, but starting in some state  $|\beta\rangle$  that is not of interest and will be overwritten. In other words, we might like to devise a method of starting with the tensor product state  $|\psi\rangle \otimes |\beta\rangle$  and ending up with  $|\psi\rangle \otimes |\psi\rangle$ , no matter what  $|\psi\rangle$  is. If this were possible, it would be very useful for quantum information applications. For example, we could make many exact copies of a quantum state, and then measure it in many different ways, including with non-commuting operators, to determine its exact form. However, it cannot be done.

To see why, it is easiest to take the simple case that  $|\psi\rangle$  is a single qubit. Suppose that we have a device or method, realized as a linear unitary operator, whose effect is the transformation

$$|0\rangle \otimes |\beta\rangle \rightarrow |0\rangle \otimes |0\rangle, \quad (26.4.1)$$

$$|1\rangle \otimes |\beta\rangle \rightarrow |1\rangle \otimes |1\rangle. \quad (26.4.2)$$

Then, acting on a general qubit  $|\psi\rangle = a|0\rangle + b|1\rangle$ , by linearity we must have

$$(a|0\rangle + b|1\rangle) \otimes |\beta\rangle \rightarrow a|0\rangle \otimes |0\rangle + b|1\rangle \otimes |1\rangle. \quad (26.4.3)$$

The point is simply that this is not the same as

$$|\psi\rangle \otimes |\psi\rangle = a^2|0\rangle \otimes |0\rangle + ab|0\rangle \otimes |1\rangle + ab|1\rangle \otimes |0\rangle + b^2|1\rangle \otimes |1\rangle, \quad (26.4.4)$$

except in the very special case that  $ab = 0$  and  $a^2 = a$  and  $b^2 = b$ , which implies  $a = 1, b = 0$  or  $a = 0, b = 1$ . A straightforward generalization of this proof gives the following no-go result, obtained in 1982 by William Wootters, Wojciech Zurek, and Dennis Dieks:

**Theorem 26.4.1. (No cloning)** *It is not possible to produce an identical copy of a general state  $|\psi\rangle$  without modifying the original. In other words, given a system consisting of a tensor product of two subsystems of the same type, there does not exist a linear operator that takes  $|\psi\rangle \otimes |\beta\rangle \rightarrow |\psi\rangle \otimes |\psi\rangle$  for every state  $|\psi\rangle$ .*

This is consistent with the example of teleportation in section 26.3, in which Alice had to destroy her qubit by measurement collapse in the process of sending it to Bob. More generally, the impossibility of cloning a state avoids paradoxes involving instantaneous communication. Consider an entangled Bell state  $|\phi^+\rangle_{ab}$  shared by Alice and Bob. Alice wants to send a message to Bob, either yes or no. To say “yes”, she measures  $Z_a$ , and to say “no”, she instead measures  $X_a$ . (The actual result of the measurement is not relevant to the argument.) Now, suppose Bob

could clone his qubit. He makes a large number of copies, and then measures  $Z_b$  on half of them and  $X_b$  on the other half. If Alice's message was "yes", Bob's measurements of  $Z_b$  will all have the same result, and his measurements of  $X_b$  will appear random. Conversely, if Alice's message was "no", then Bob's measurements of  $Z_b$  will appear random and his measurements of  $X_b$  will all have the same result. By cloning his qubit, Bob could (almost always) learn Alice's message immediately. But this is just fiction; in reality Bob has access to only one copy of his qubit, and no instantaneous communication can occur.

Although cloning of states is prohibited, it is possible to do something slightly different, and still very useful, called **redundant encoding**. Consider the linear unitary transformation of eqs. (26.4.1)-(26.4.3), which we can express in a more compact way by leaving out the irrelevant qubit  $|\beta\rangle$ , and using a double arrow to denote an encoding, as

$$a|0\rangle + b|1\rangle \Rightarrow a|00\rangle + b|11\rangle \quad (26.4.5)$$

Although this perfectly valid operation does not clone the original qubit, what we have managed to do is to encode its information in a 2-qubit register, in which the second qubit is redundant. By iterating this procedure, it is possible to encode a qubit  $|\psi\rangle = a|0\rangle + b|1\rangle$  in a register with as many redundant qubits as we wish,

$$a|0\rangle + b|1\rangle \Rightarrow a|000\cdots 0\rangle + b|111\cdots 1\rangle. \quad (26.4.6)$$

The right-hand side is sometimes known as a **logical qubit** or a **synthetic qubit**, to distinguish it from the physical qubits that it is made out of. Its information content is exactly the same as in the original qubit, one complex degree of freedom. Redundant encoding does not violate the no-cloning theorem, because the redundant qubits that form the synthetic qubit are fully entangled with each other. The key difference is that a cloned copy of the original qubit would have been an independent copy, not entangled with the original.

Redundant encoding can be done in usefully different ways, depending on the choice of basis. For example, we might choose to use for the redundant physical qubits the basis

$$|+\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle), \quad |-\rangle = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle), \quad (26.4.7)$$

which are eigenstates of  $X$  with eigenvalues 1 and  $-1$ , respectively. Then one can construct a redundant encoding of a logical qubit that takes the form

$$a|0\rangle + b|1\rangle \Rightarrow a|++++\cdots+\rangle + b|----\cdots-\rangle. \quad (26.4.8)$$

One can also nest this encoding procedure, by bundling synthetic qubits into bigger synthetic qubits. As we will see in section 26.6, these synthetic qubit tricks are very useful for quantum error correction.

## 26.5 Keeping secrets with quantum private key generation

As noted at the end of section 26.2, the security of many public-key encryption schemes is threatened by the prospect of quantum computers solving problems that were previously believed to be too hard, based on classical computation algorithms. Shor's quantum factorization algorithm is the most famous example of this, but there are others. One way to improve public-key codes is to devise decryption problems for which the solution would take too long even for the best possible quantum computers, and some proposals have been made along these lines. However, a different approach is to use private keys. In this section, we will show how quantum information enables secure generation of private keys and their distribution over large distances.

The one-time private key encryption method is unbreakable, even classically. Suppose that Alice and Bob each have a copy of a prepared and secret register of random bits,  $\kappa = (\kappa_1 \kappa_2 \cdots \kappa_N)$ . This is the private key. Now Alice has a message that she wants to send to Bob, consisting of bits  $\alpha = (\alpha_1 \alpha_2 \cdots \alpha_N)$ . To encrypt the message, she adds the two classical registers bitwise to get  $\epsilon = \alpha + \kappa = (\epsilon_1 \epsilon_2 \cdots \epsilon_N)$ , where  $\epsilon_k = \alpha_k + \kappa_k \pmod{2}$  for each  $k = 1, \dots, N$ . Alice then sends the encrypted message  $\epsilon$  to Bob, who can read the message by bitwise subtracting (or adding, which is the same thing) the key,  $\alpha = \epsilon - \kappa$ . An eavesdropper Eve who intercepts the message will gain absolutely no information without knowing the private key, because if the key  $\kappa$  consists of completely random bits, so does the encrypted message  $\epsilon$ .

However, this absolute security requires that one cannot reuse the key or any part of it. Otherwise, it is conceivable that a clever Eve might be able to discern patterns that allow the key to be discovered. If Alice and Bob are at different locations, they will need to have a new private key with a new random bit for each bit of message to be sent. What if they run out of their supply of private key bits? Any classical communication of a new private key provides an opportunity for Eve to intercept it. Fortunately for Alice and Bob, there are quantum methods for private key generation and distribution, which can work even when they are spatially separated, and allow for any eavesdropping by Eve to be detected.

It should be emphasized that the overriding concern here is security, not speed or efficiency of communication. Also, only the private key generation and distribution, not the encryption or decryption, relies on the principles of quantum mechanics.

One way of generating and distributing private keys is the **EPR protocol**. Alice and Bob are distant from each other, but share a large collection of Bell pairs, all in the same state

$$|\phi^+\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle) = \frac{1}{\sqrt{2}}(|++\rangle + |--\rangle). \quad (26.5.1)$$

When they need to generate a private key, Alice and Bob each measure  $X$  or  $Z$ , chosen randomly and independently, sequentially for each qubit that they control, getting one of two results in

each case. They then communicate to each other the sequence of measurements that they made, but not the results of those measurements. There is no need to hide this communication, which is therefore considered public. In about half of the cases, they will have chosen to make different measurements, and their results will be uncorrelated; those are discarded. In the other cases, their measurements were either both  $X$ , or both  $Z$ . In those cases, Alice and Bob will always get the same result, and those will define the bits of their private key. (Alice and Bob agree that  $X = 1$  or  $Z = 1$  stand for a classical bit 0, and  $X = -1$  or  $Z = -1$  for a classical bit 1.) For each sequence of  $N$  Bell states that they process in this way, they can therefore expect to generate a private key of length approximately  $N/2$ , with random bits known only to them. This private key can then be used to encrypt and decrypt a message composed of ordinary classical bits.

Alice and Bob should worry that Eve might be trying to learn their private key. Eve could do this by intercepting and modifying the Bell pairs before they are used. For example, perhaps Eve is actually the vendor that sold the supposed Bell pairs to Alice and Bob, and she substituted states of her own choosing in place of  $|\phi^+\rangle$ . Fortunately for Alice and Bob, there is an easy way to detect this. After generating their private key, they tell each other publicly what their results are for a random selection of the private key bits, which of course will not be used for actual secret messages. If those bits fail to match, they will know that the 2-qubit state provided to them was not actually  $|\phi^+\rangle$ , exposing Eve's fraud. By sacrificing a sufficient fraction of their private key bits, they can be reasonably sure that the states were not tampered with, and feel safe in using the other bits. This relies on the fact that  $|\phi^+\rangle$  is the unique 2-qubit state that will always give the same result to Alice and Bob regardless of whether it was  $X$  or  $Z$  that they both measured. This also explains why they had to go to the trouble of randomly measuring  $X$  and  $Z$ , and then discarding the results when their choices did not match. If Eve knew in advance that they were both going to measure  $Z$ , then she could substitute the state  $|00\rangle$  or  $|11\rangle$  in place of the state  $|\phi^+\rangle$ , ensuring that the private key bit for that case would be 0 or 1, at Eve's discretion, and that Alice and Bob would not see anything amiss.

The use of Bell pairs in the preceding might give the impression that entanglement is a necessary ingredient for quantum private key distribution, but this is not true. To see this, consider the following variation on the scheme. Alice creates the Bell pairs  $|\phi^+\rangle_{ab}$  herself, measures  $X$  or  $Z$  randomly on qubit  $a$ , and then sends qubit  $b$  to Bob. Then Bob measures either  $X$  or  $Z$  on qubit  $b$ , and the rest of the protocol is the same. This is formally equivalent to a protocol in which qubit  $a$  never existed at all, and Alice simply prepares qubit  $b$  randomly in one of the four states  $|0\rangle$  or  $|1\rangle$  (the  $Z$  basis) or  $|+\rangle$  or  $|-\rangle$  (the  $X$  basis). She then sends this qubit to Bob, who measures either  $X$  or  $Z$ . Alice and Bob announce to each other over a public channel their choices of the  $X$  or  $Z$  basis, and the rest of the protocol is the same. This



version, which uses only a one-way sending of qubits and no entanglement, is called the **BB84 protocol**, after its 1984 proposal by Charles Bennett and Gilles Brassard. This has the clear advantage that Alice and Bob do not need a large supply of nonlocally entangled Bell states.

In the BB84 scheme, it is again crucial that Alice and Bob use both the  $X$  and  $Z$  bases randomly. If Eve can intercept the qubits that Alice sends, and knew in advance that a qubit was definitely an eigenstate of  $X$  or of  $Z$ , she would be able to measure it in the corresponding eigenstate basis and send on an exact copy to Bob. Without this knowledge, the no-cloning theorem prevents Eve from making a duplicate and keeping one copy for herself, to be measured after Alice and Bob have announced whether they both chose  $X$  or  $Z$ . Any measurements that Eve makes will collapse the qubits unless Eve happens to guess whether they are eigenstates of  $X$  and  $Z$ . So, Alice and Bob will again discover any eavesdropping by sacrificing a portion of their private key bits and noting that sometimes they do not match.

In the real world, the qubits are typically encoded in photon polarizations, and sent either through optical fibers or through empty space. There are of course practical limitations, including noise, decoherence, and photon loss, which become more acute for long-distance communications. Also, the detection of Eve's possible shenanigans is only probabilistic, and one must account for the fact that in some cases the check bits will fail to match because of non-malicious faults. The need for tradeoffs and optimizations has given rise to more sophisticated protocols. The relative simplicity of one-way non-entangled photonic qubit transmission, and the market demand for privacy in communication, has already lead to the availability of commercial implementations of quantum key generation and distribution.

## 26.6 Fighting decoherence and other faults with error correction

An important issue in both classical and quantum information is dealing with errors that may change a register in undesirable ways. In the case of classical bits, errors can be introduced by stray electromagnetic fields, mechanical defects, thermal fluctuations, radioactivity, or even cosmic rays. There are well-developed methods for preventing such errors, and detecting and correcting them when they occur, so that the engineering problem for classical bits is essentially a solved one. For qubits, the same sort of errors can occur, but there is the additional and especially pernicious problem of decoherence due to interactions with the environment. As we have seen in Chapter 25, decoherence is very hard to avoid completely, especially in systems with even small couplings to macroscopic degrees of freedom. Correction of decoherence errors is therefore a necessity in order to build practical large-scale quantum computers. In this section, we will give a brief account of error-correction strategies for qubits.

For the case of a single classical bit, a simple strategy for error correction is redundant



encoding followed by majority-rule correction. For example, one might redundantly encode each logical bit as three physical bits, according to the rule

$$0 \Rightarrow (000), \quad 1 \Rightarrow (111). \quad (26.6.1)$$

Then, if only one physical bit is modified by an error, the logical bit 0 might change to (100) or (010) or (001), while the logical bit 1 might change to (011) or (101) or (110). Such an error is easily diagnosed by the fact that the three physical bits are not all the same. The error is then corrected by changing the different bit to agree with the majority. This strategy fails if an error affects two or three physical bits at the same time, but this is hopefully rare, especially if the error correction is applied repeatedly after short time intervals. An obvious way to improve the reliability is to use more than three physical bits for each logical bit, but there are more sophisticated and efficient schemes for classical error-correcting codes.

Error correction of qubits is more difficult for several reasons. First, the no-cloning theorem declares that it is not possible to just copy the qubit to three independent identical copies as in the classical case; therefore we will have to use redundant encoding as described in section 26.4. Second, a measurement on a single physical qubit will inevitably collapse the state, erasing the information it contains. So, we will have to use cleverly chosen collective measurements, which act on more than one physical qubit at a time in such a way as to not destroy the information in the logical qubit. Third, there is only one type of error for a classical bit; it can flip, changing from 0 to 1 or 1 to 0. In contrast, qubits have multiple different error syndromes, because the general qubit state is an arbitrary linear combination of the basis states, including relative phases that have physical meaning.

To expand on the last point, let us consider several different error types. The most obvious counterpart to the classical bit error for a qubit is the basis bit-flip error, which changes the qubit basis state  $|0\rangle$  to  $|1\rangle$  and vice versa, so that the general physical qubit state changes according to

$$a|0\rangle + b|1\rangle \rightarrow a|1\rangle + b|0\rangle \quad (\text{bit-flip error}). \quad (26.6.2)$$

Another type of error is a phase flip, which changes the relative phase between the kets  $|0\rangle$  and  $|1\rangle$  by a factor of  $-1$ , so that

$$a|0\rangle + b|1\rangle \rightarrow a|0\rangle - b|1\rangle \quad (\text{phase-flip error}). \quad (26.6.3)$$

The error could also be a combination of a phase flip and a bit flip,

$$a|0\rangle + b|1\rangle \rightarrow a|1\rangle - b|0\rangle \quad (\text{phase-and-bit-flip error}). \quad (26.6.4)$$

These three error types are realized by unitary operators, respectively,

$$U_{\text{err}} = X, \quad (\text{bit-flip error}), \quad (26.6.5)$$

$$U_{\text{err}} = Z, \quad (\text{phase-flip error}), \quad (26.6.6)$$

$$U_{\text{err}} = -iY, \quad (\text{phase-and-bit-flip error}). \quad (26.6.7)$$

These are “large” errors, as they square to the identity (up to a global phase). More generally, a qubit error could be an arbitrary unitary transformation. Since the identity matrix and the Pauli matrices together form a basis for addition of all  $2 \times 2$  matrices, one can always write (again up to an irrelevant global phase factor),

$$U_{\text{err}} = \frac{I + i(\epsilon_X X + \epsilon_Y Y + \epsilon_Z Z)}{\sqrt{1 + \epsilon_X^2 + \epsilon_Y^2 + \epsilon_Z^2}}, \quad (26.6.8)$$

where the unitarity condition  $U_{\text{err}}^\dagger U_{\text{err}} = I$  requires that  $\epsilon_X$ ,  $\epsilon_Y$ , and  $\epsilon_Z$  are real numbers, which may or may not be small. The three special cases of the bit flip, the phase flip, and the combined phase-and-bit flip are recovered in the limits of large  $\epsilon_X$ , large  $\epsilon_Z$ , and large  $\epsilon_Y$ , respectively (once again, up to a global phase). Also included in this general parameterization are “small” errors that are continuously connected to the no-error case, but can build up over time. Ideally, a quantum error-correction method should be able to fix any error of this type, even if we have no way of knowing which error occurred. We will see that this is indeed true.

The strategy for quantum error correction can be summarized as follows. First, each logical qubit is redundantly encoded as a register of physical qubits. After some time, an error may have been introduced in one of the physical qubits, but we don’t know which error or which qubit, if any. Therefore, one makes measurements on the register to diagnose which error syndrome has occurred, but this must be done in a careful way so that the state is unaffected if there was actually no error. The joint result of the measurements will be one of a discrete finite set. If an error is detected, the measurements also project the register state onto a corresponding discrete set of known error-syndrome states. Finally, one applies an appropriate unitary operator, determined by the result of the diagnostic measurements, to correct the register back to its pre-error state. This error-correction process should be repeated at regular time intervals, short enough so that the probability of multiple errors within each interval is small.

For simplicity, the following discussion is framed in terms of pure-state qubits affected by an error that takes the form of a unitary transformation  $U_{\text{err}}$ , as above. However, it is very important that this class of algorithms also corrects decoherence errors that evolve the pure state of the qubit to a mixed state, through entanglement with the environment. The reason is that the diagnostic measurement step automatically projects the density matrix of the register

onto a pure error-syndrome state, unentangled with the environment. The ensuing correction is a unitary operator on the register alone, so that the final restored state is always a pure one.

As a warmup example, let us first consider the simplified case that errors are restricted to the bit-flip type on a single physical qubit. Inspired by the majority-rule scheme for classical bits, we use the triply redundant encoding rule

$$a|0\rangle + b|1\rangle \Rightarrow a|000\rangle + b|111\rangle. \quad (26.6.9)$$

A bit-flip error on one of the three physical qubits could change this state to any one of the following error-syndrome states:

$$a|100\rangle + b|011\rangle \quad (\text{bit-flip error on qubit 1}), \quad (26.6.10)$$

$$a|010\rangle + b|101\rangle \quad (\text{bit-flip error on qubit 2}), \quad (26.6.11)$$

$$a|001\rangle + b|110\rangle \quad (\text{bit-flip error on qubit 3}). \quad (26.6.12)$$

These states are illegal, according to the original encoding, and so must be corrected. Now, to diagnose the error, it would be bad to simply measure the three qubits one at a time, the strategy that worked in the classical case. If we measure  $Z_1$  for the first qubit, that would give results 1 and  $-1$  with probabilities  $|a|^2$  and  $|b|^2$  or vice versa, but this does not distinguish with certainty which qubit may have suffered an error. Even worse, the information stored will have been destroyed by the measurement collapse, which projects onto only states with definite  $Z_1 = 1$  or  $-1$ , rather than the superposition in the original state. Measurement of a single physical qubit would irrevocably destroy the logical qubit we are trying to correct.

Instead, the correct strategy is to use only collective measurements, which act on more than one physical qubit. Specifically, we measure the compatible operators<sup>†</sup>

$$A_{12} = Z_1 Z_2, \quad \text{and} \quad A_{13} = Z_1 Z_3, \quad (26.6.13)$$

which compare pairs of qubits without measuring them individually. Each of the four states appearing in eqs. (26.6.9)-(26.6.12) is always an eigenstate of both  $A_{12}$  and  $A_{13}$ , no matter what  $a$  and  $b$  are. This means that measuring one or both of them does not change the state, and there is no loss of information from collapse. Furthermore, the four possible results for the measurement of  $(A_{12}, A_{13})$  correspond to the four possible error syndromes (including the trivial syndrome of no error), telling us which error occurred and how to correct it. If we measure  $(A_{12}, A_{13}) = (-1, -1)$ , the error was a bit-flip of qubit 1, as in eq.(26.6.10). We can therefore correct the error by applying the unitary operator  $X_1$ . If instead  $(A_{12}, A_{13}) = (-1, 1)$ , the error was a bit-flip of qubit 2, and we should correct the error by applying  $X_2$ . If  $(A_{12}, A_{13}) = (1, -1)$ ,

---

<sup>†</sup>The notation means the same thing as  $A_{12} = Z \otimes Z \otimes I$  and  $A_{13} = Z \otimes I \otimes Z$ .

Error syndrome	$A_{12}$	$A_{13}$	$U_{\text{recovery}}$
No error	1	1	$I$
Bit flip on qubit 1	-1	-1	$X_1$
Bit flip on qubit 2	-1	1	$X_2$
Bit flip on qubit 3	1	-1	$X_3$

Table 26.6.5: Error correction for single-qubit bit-flips, using the 3-physical-qubit encoding of one logical qubit. Simultaneous measurement of the compatible observables  $A_{12}$  and  $A_{13}$  diagnoses the error syndrome and projects onto the corresponding error syndrome state. The unitary operator  $U_{\text{recovery}}$  then corrects the error.

then the error was a bit-flip of qubit 3, and we should apply the unitary operator  $X_3$ . Finally, the case  $(A_{12}, A_{13}) = (1, 1)$  corresponds to no error, so no unitary correction is applied. This error-correction algorithm is summarized in Table 26.6.5.

The algorithm actually corrects a more general class of errors on a single qubit, namely those obtained with  $\epsilon_Y = \epsilon_Z = 0$  and arbitrary  $\epsilon_X$  in eq. (26.6.8). For example, suppose that the error hits the first physical qubit, according to the unitary transformation

$$a|000\rangle + b|111\rangle \rightarrow \frac{1}{\sqrt{1+\epsilon^2}} [a|000\rangle + b|111\rangle + i\epsilon(a|100\rangle + b|011\rangle)], \quad (26.6.14)$$

where  $\epsilon$  is a real number whose magnitude may or may not be small. (Note that  $\epsilon = 0$  corresponds to no error, while  $\epsilon \rightarrow \infty$  is the large bit-flip error case.) Now when the measurement of  $(A_{12}, A_{13})$  takes place, there are two possible results. With probability  $1/(1+\epsilon^2)$ , the measurement will give the result  $(1, 1)$  and collapse the state to  $a|000\rangle + b|111\rangle$ . In this case, no unitary recovery transformation is applied. With probability  $\epsilon^2/(1+\epsilon^2)$ , the measurement will give the result  $(-1, -1)$ , diagnosing the error syndrome as a bit-flip on qubit 1, and collapsing the state to  $a|100\rangle + b|011\rangle$ . In that case, the algorithm calls for applying the unitary operator  $U_{\text{recovery}} = X_1$ , changing the state back to  $a|000\rangle + b|111\rangle$ . So, in both cases, the final result for the state is the correct pre-error state. Even though there is non-trivial measurement collapse in this case, it is actually beneficial, because it forces the state into one of the discrete error syndromes that we know how to fix. The measurement allows us to correctly repair the state using one of three operations, every time, no matter what the continuous parameter  $\epsilon$  was.

However, the preceding example does not protect at all against phase-flip errors. As a second warmup, let us consider an algorithm that corrects all single-qubit phase-flip errors (but does not handle bit-flip errors). The strategy is actually very similar to the previous case, based on the observation that a phase-flip in the basis  $|0\rangle, |1\rangle$  is equivalent to a bit-flip in the basis  $|+\rangle, |-\rangle$ , and vice versa. We use the following 3-qubit redundant encoding:

$$a|0\rangle + b|1\rangle \Rightarrow a|+++ \rangle + b|--- \rangle, \quad (26.6.15)$$

where  $|\pm\rangle = (|0\rangle \pm |1\rangle)/\sqrt{2}$  are the eigenstates of  $X$  with eigenvalues  $\pm 1$ . A phase-flip error amounts to changing  $|+\rangle \leftrightarrow |-\rangle$  on one of the qubits, so the possible error-syndrome states are

$$a| -++\rangle + b| +--\rangle \quad (\text{phase-flip error on qubit 1}), \quad (26.6.16)$$

$$a| +-+\rangle + b| -+-\rangle \quad (\text{phase-flip error on qubit 2}), \quad (26.6.17)$$

$$a| ++-\rangle + b| --+\rangle \quad (\text{phase-flip error on qubit 3}). \quad (26.6.18)$$

It is straightforward to check that measurement of the compatible operators

$$B_{12} = X_1 X_2, \quad \text{and} \quad B_{13} = X_1 X_3 \quad (26.6.19)$$

will uniquely diagnose the phase-flip error syndrome, with  $(B_{12}, B_{13}) = (-1, -1)$ ,  $(-1, 1)$ ,  $(1, -1)$ , or  $(1, 1)$  for, respectively, a phase-flip of qubit 1, 2, 3, or no error. This also works for the case of any  $U_{\text{err}}$  in eq. (26.6.8) with arbitrary  $\epsilon_Z$ , as long as  $\epsilon_X = \epsilon_Y = 0$ . In general, the measurement projects onto one of the phase-flip error-syndrome states. The error is then corrected by applying the unitary operator  $Z_1$ ,  $Z_2$ ,  $Z_3$ , or  $I$ , respectively.

In 1995, Shor showed how to neatly generalize these ideas into an algorithm that corrects any single-qubit error. One starts with the following 9-qubit redundant encoding:

$$|0\rangle \Rightarrow \frac{1}{\sqrt{2^3}}(|000\rangle + |111\rangle) \otimes (|000\rangle + |111\rangle) \otimes (|000\rangle + |111\rangle), \quad (26.6.20)$$

$$|1\rangle \Rightarrow \frac{1}{\sqrt{2^3}}(|000\rangle - |111\rangle) \otimes (|000\rangle - |111\rangle) \otimes (|000\rangle - |111\rangle). \quad (26.6.21)$$

This encodes each logical qubit as three bundles of three physical qubits each. It can be viewed as a nested encoding, first triply redundantly in the  $|+\rangle$ ,  $|-\rangle$  basis, and then triply redundantly in the  $|0\rangle$ ,  $|1\rangle$  basis.

Now, if a phase-flip error occurs in any single physical qubit, it will change the relative sign of the two terms within one of the three bundles. The resulting state does not depend on which of the three qubits in the bundle suffered the error. Therefore, phase-flip errors can be completely diagnosed by simultaneously measuring the two compatible operators

$$B_{123456} = X_1 X_2 X_3 X_4 X_5 X_6, \quad (26.6.22)$$

$$B_{123789} = X_1 X_2 X_3 X_7 X_8 X_9, \quad (26.6.23)$$

each with eigenvalues  $\pm 1$ . If, and only if,  $B_{123456}$  is found to be  $-1$ , then the phase-flip error must have been in either the first or second bundle, and similarly if  $B_{123789}$  is found to be  $-1$ , then the error was in either the first or third bundle. There are thus four possible error syndromes for the combined measurement of the two operators. After the measurement projects the state onto one of them, the phase-flip error can be corrected by applying a unitary transformation. For

example, if it is determined that the phase-flip error occurred on the second bundle, one can undo the error by applying  $U_{\text{recovery}}^{\text{phase}} = Z_4$  (although using  $Z_5$  or  $Z_6$  or the symmetric combination  $Z_4Z_5Z_6$  instead would work just as well). The same encoding also allows one to detect and correct any bit-flip error. By measuring each of the pairs

$$A_{12} = Z_1Z_2, \quad A_{13} = Z_1Z_3, \quad (26.6.24)$$

$$A_{45} = Z_4Z_5, \quad A_{46} = Z_4Z_6, \quad (26.6.25)$$

$$A_{78} = Z_7Z_8, \quad A_{79} = Z_7Z_9, \quad (26.6.26)$$

one can detect which of the 9 qubits was bit-flipped, and apply the appropriate unitary recovery operator. All eight of these observables commute, and measurement collapse has no effect on the state in the special case that there was no error. The error correction algorithm is summarized in Table 26.6.6. As you will check explicitly by doing Exercise 26.5, the algorithm corrects every possible single-qubit error of the general form given in eq. (26.6.8). The essential reason for this is that the general error transformation in eq. (26.6.8) is always a linear combination (superposition) of  $I$ ,  $X$ ,  $Z$ , and  $Y$ , which means no change, a bit flip, a phase flip, or a combination of phase and bit flips.

This error correction strategy relies on the assumption that only one physical qubit was affected. This may be a reasonable (but not guaranteed) assumption if errors are rare on the time scale between applications of the algorithm. If two qubits both had bit-flip errors, the algorithm would wrongly flip the qubit that did not change, instead of the two that did. Similarly, if two bundles had phase-flip errors, the wrong bundle would have its phase flipped.

Although Shor's encoding and error-correction algorithm is relatively simple to understand, it is no longer the most efficient. It turns out to be possible to do the same thing with synthetic qubits containing only 5 (instead of 9) physical qubits. There are also more sophisticated methods than can, with high probability, correct errors that occur in more than one physical qubit simultaneously.

Error syndrome	$B_{123456}$	$B_{123789}$	$U_{\text{recovery}}^{\text{phase}}$
No phase-flip error	1	1	$I$
Phase flip on qubit 1, 2, or 3	-1	-1	$Z_1$
Phase flip on qubit 4, 5, or 6	-1	1	$Z_4$
Phase flip on qubit 7, 8, or 9	1	-1	$Z_7$
Error syndrome	$A_{12}$	$A_{13}$	$U_{\text{recovery}}^{123}$
No bit-flip error on qubit 1, 2, or 3	1	1	$I$
Bit flip on qubit 1	-1	-1	$X_1$
Bit flip on qubit 2	-1	1	$X_2$
Bit flip on qubit 3	1	-1	$X_3$
Error syndrome	$A_{45}$	$A_{46}$	$U_{\text{recovery}}^{456}$
No bit-flip error on qubit 4, 5, or 6	1	1	$I$
Bit flip on qubit 4	-1	-1	$X_4$
Bit flip on qubit 5	-1	1	$X_5$
Bit flip on qubit 6	1	-1	$X_6$
Error syndrome	$A_{78}$	$A_{79}$	$U_{\text{recovery}}^{789}$
No bit-flip error on qubit 7, 8, or 9	1	1	$I$
Bit flip on qubit 7	-1	-1	$X_7$
Bit flip on qubit 8	-1	1	$X_8$
Bit flip on qubit 9	1	-1	$X_9$

Table 26.6.6: Error correction for the Shor 9-physical-qubit encoding of one logical qubit based on measurement of the 8 compatible observables  $B_{123456}$ ,  $B_{123789}$ ,  $A_{12}$ ,  $A_{13}$ ,  $A_{45}$ ,  $A_{46}$ ,  $A_{78}$ , and  $A_{79}$ . Measurement of  $B_{123456}$  and  $B_{123789}$  diagnoses any single-qubit phase-flip error, and projects onto the corresponding error-syndrome state, which can then be corrected by applying the corresponding unitary operator  $U_{\text{recovery}}^{\text{phase}}$ . Measurement of  $A_{12}$  and  $A_{13}$  diagnoses any bit-flip error of qubit 1, 2, or 3, which is then corrected by the corresponding  $U_{\text{recovery}}^{123}$ , and similarly for bit-flip errors on qubits 4, 5, 6, 7, 8, or 9. This corrects any error on any single physical qubit, restoring the 9-qubit register to its pre-error state.

## 26.7 Exercises

**Exercise 26.1.** The decoherence of a pure state qubit to a mixed state can be described using a dynamical map using Kraus operators  $K_n$ , according to  $\rho \rightarrow \sum_n K_n \rho K_n^\dagger$ , as discussed in section 24.2. Consider the set of points on the unit-sphere surface of the Bloch ball, corresponding to all pure states of the qubit. What surface results from applying each of the following maps?

- (a) the **depolarizing channel** map defined by  $K_0 = \sqrt{1-3p/4}I$  and  $K_1 = \sqrt{p/4}\sigma_x$  and  $K_2 = \sqrt{p/4}\sigma_y$  and  $K_3 = \sqrt{p/4}\sigma_z$ . [Answer: you should find that the effect of the map is to shrink the Bloch vector according to  $\vec{b} \rightarrow (1-p)\vec{b}$ . Thus, pure states on the unit-sphere surface of the Bloch ball are mapped to mixed states on a sphere of radius  $1-p$ .]
- (b) the **bit-flip decoherence** channel defined by  $K_0 = \sqrt{1-p}I$  and  $K_1 = \sqrt{p}\sigma_x$ .
- (c) the **phase-flip decoherence** channel defined by  $K_0 = \sqrt{1-p}I$  and  $K_1 = \sqrt{p}\sigma_z$ .
- (d) the **bit-and-phase-flip decoherence** channel defined by  $K_0 = \sqrt{1-p}I$  and  $K_1 = \sqrt{p}\sigma_y$ .

**Exercise 26.2.** Show that the map  $U_f$  defined by eq.(26.2.4) is unitary.

**Exercise 26.3.** Consider the two operators  $A = Z_1 Z_2$  and  $B = X_1 X_2$  acting on a system of two qubits labeled 1 and 2. Show that  $[A, B] = 0$ , and that their common eigenstates are the maximally entangled states  $|\phi^\pm\rangle$  and  $|\psi^\pm\rangle$  given in eq. (26.3.2) and (26.3.3). Find the corresponding eigenvalues. Relate the eigenvalues of  $(A, B)$  to those of the operator  $Q$  in eq. (26.3.4).

**Exercise 26.4.** Consider the teleportation of a qubit  $|\gamma\rangle_c$  from Alice to Bob as discussed in section 26.3, but suppose that the initial shared Bell pair is in the state  $|\psi^-\rangle_{ab}$ , instead of  $|\phi^+\rangle_{ab}$ . What rules will Bob follow to receive the teleported state, replacing eqs. (26.3.6)-(26.3.9)?

**Exercise 26.5.** For Shor's 9-qubit encoding of a logical qubit  $a|0\rangle + b|1\rangle$  using eqs. (26.6.20) and (26.6.21), consider the most general unitary error transformation of the form in eq. (26.6.8) applied to the first physical qubit. For the simultaneous measurement of the operators  $B_{123456}$ ,  $B_{123789}$ ,  $A_{12}$ , and  $A_{13}$ , give the possible outcomes and their probabilities, in terms of  $\epsilon_X$ ,  $\epsilon_Y$ , and  $\epsilon_Z$ . Give the corresponding post-measurement states, and check that the error-correction algorithm summarized in Table 26.6.6 always restores the qubit to the pre-error state. (Since there is nothing special about the first physical qubit, this generalizes to a general unitary error transformation on any single qubit.)



## 27 Relativistic quantum mechanics

### 27.1 Special relativity, four-vectors, and Lorentz transformations

Einstein's special relativity places the four dimensions of spacetime on an equal footing. This presents a challenge for the quantum mechanics of a single particle as formulated above, where the three spatial coordinates are operators, but time is treated completely differently, as a mere parameter. In this chapter we will discuss how quantum mechanical wave equations can be made consistent with the symmetries of special relativity, leading to the Klein-Gordon equation for spin-0 particles and the Dirac equation for spin-1/2 particles.

Classically, a four-vector coordinate can be assigned to each event,

$$(ct, x, y, z) = (x^0, x^1, x^2, x^3) = x^\mu. \quad (27.1.1)$$

Greek indices  $\mu, \nu, \rho, \dots$  run over the values 0, 1, 2, 3, and  $c$  is the speed of light in vacuum. It is a guiding principle of special relativity that the laws of physics should take the same form in any **inertial reference frame**, which means that the coordinates of a free classical particle do not accelerate. It is often useful to change our coordinate system from one inertial reference frame to another, according to

$$x^\mu \rightarrow x'^\mu = L^\mu{}_\nu x^\nu. \quad (27.1.2)$$

Here  $L^\mu{}_\nu$  is a constant  $4 \times 4$  real matrix satisfying an orthogonality constraint to be found soon, and the repeated (and therefore implicitly summed) index  $\nu$  is said to be “contracted”. Such a change of coordinates specified by  $L^\mu{}_\nu$  is called a **Lorentz transformation**. By convention, contracted indices must have opposite heights.

As a matter of terminology,  $x^\mu$  (with a raised index) is an example of a **contravariant** four-vector. By definition, all contravariant four-vectors transform in the same way,

$$a^\mu \rightarrow a'^\mu = L^\mu{}_\nu a^\nu. \quad (27.1.3)$$

Another contravariant four-vector is the **four-momentum** consisting of the energy  $E$  and spatial momentum  $\vec{p}$  of a particle,

$$p^\mu = (E/c, \vec{p}). \quad (27.1.4)$$

In the rest frame of a particle of mass  $m$ , its four-momentum is  $p^\mu = (mc, 0, 0, 0)$ .

A simple example of a Lorentz transformation is the rotation about the  $z$ -axis by a fixed angle  $\alpha$ . In the new coordinate system,  $ct' = ct$ ,  $x' = x \cos \alpha + y \sin \alpha$ ,  $y' = -x \sin \alpha + y \cos \alpha$ , and  $z' = z$ . As another example, called a **boost**, we could go to a frame moving with respect

to the original frame with velocity  $v$  along the  $\hat{z}$  direction, with the origins of the two frames coinciding at time  $t = t' = 0$ . Then,

$$ct' = \gamma(ct - vz/c), \quad x' = x, \quad y' = y, \quad z' = \gamma(z - vt), \quad (27.1.5)$$

where

$$\gamma = 1/\sqrt{1 - v^2/c^2}. \quad (27.1.6)$$

It follows that the Lorentz transformation for a boost  $v$  along the  $\hat{z}$  direction is

$$L^\mu{}_\nu = \begin{pmatrix} \gamma & 0 & 0 & -\gamma v/c \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -\gamma v/c & 0 & 0 & \gamma \end{pmatrix}, \quad (27.1.7)$$

and the inverse Lorentz transformation is

$$a^\mu = (L^{-1})^\mu{}_\nu a'^\nu, \quad (L^{-1})^\mu{}_\nu = \begin{pmatrix} \gamma & 0 & 0 & \gamma v/c \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \gamma v/c & 0 & 0 & \gamma \end{pmatrix}. \quad (27.1.8)$$

As one might expect, the inverse of a boost simply replaces the boost velocity  $v\hat{z}$  by  $-v\hat{z}$ .

For any two events one can define the **proper distance**, which is independent of the choice of Lorentz frame and therefore tells us how far apart the two events are in a coordinate-independent sense. Consider two events occurring at  $x^\mu$  and  $x^\mu + d^\mu$ , where  $d^\mu$  is some four-vector displacement. The proper distance between the events is

$$(\Delta x)^2 = -(d^0)^2 + (d^1)^2 + (d^2)^2 + (d^3)^2 = g_{\mu\nu} d^\mu d^\nu, \quad (27.1.9)$$

where the **metric tensor** is

$$g_{\mu\nu} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (27.1.10)$$

Furthermore, one can define an inverse metric  $g^{\mu\nu}$  so that

$$g^{\mu\nu} g_{\nu\rho} = \delta^\mu_\rho, \quad (27.1.11)$$

where  $\delta^\mu_\nu = 1$  if  $\mu = \nu$ , and otherwise  $= 0$ . In special relativity, the matrix forms of  $g^{\mu\nu}$  and  $g_{\mu\nu}$  have the same numerical entries, and do not depend on the inertial reference frame.

The metric tensor allows us to define **covariant** four-vectors by lowering an index,

$$x_\mu = g_{\mu\nu}x^\nu = (-ct, x, y, z), \quad (27.1.12)$$

$$p_\mu = g_{\mu\nu}p^\nu = (-E/c, p_x, p_y, p_z), \quad (27.1.13)$$

which always just flips the sign of the time-like component. More generally,

$$a_\mu = g_{\mu\nu}a^\nu, \quad a^\mu = g^{\mu\nu}a_\nu. \quad (27.1.14)$$

It follows that covariant four-vectors transform as

$$a'_\mu = L_\mu{}^\nu a_\nu, \quad (27.1.15)$$

where we define (note the positions of the indices)

$$L_\mu{}^\nu = g_{\mu\rho}g^{\nu\sigma}L^\rho{}_\sigma. \quad (27.1.16)$$

If  $a^\mu$  and  $b^\mu$  are contravariant four-vectors, then

$$g_{\mu\nu}a^\mu b^\nu = g^{\mu\nu}a_\mu b_\nu = a_\mu b^\mu = a^\mu b_\mu. \quad (27.1.17)$$

Since the four-vector indices have been contracted, this quantity should be Lorentz-invariant, meaning that it is independent of the choice of inertial frame coordinates. This implies  $g_{\mu\nu}a'^\mu b'^\nu = g_{\mu\nu}a^\mu b^\nu$ , so that

$$g_{\mu\nu}L^\mu{}_\rho L^\nu{}_\sigma a^\rho b^\sigma = g_{\rho\sigma}a^\rho b^\sigma. \quad (27.1.18)$$

Since  $a^\rho$  and  $b^\sigma$  are arbitrary, it must be true that

$$g_{\mu\nu}L^\mu{}_\rho L^\nu{}_\sigma = g_{\rho\sigma}. \quad (27.1.19)$$

This is the orthogonality constraint that a Lorentz transformation matrix must satisfy. It has the equivalent form

$$L_\nu{}^\kappa L^\nu{}_\sigma = \delta_\sigma^\kappa, \quad (27.1.20)$$

obtained by contracting eq. (27.1.19) with  $g^{\rho\kappa}$  and using eq. (27.1.16).

Let us now consider some particular Lorentz transformations. To begin, we note that as a matrix, eq. (27.1.20) reads  $L^T L = I$ , which in turn implies  $\det(L) = \pm 1$  because of the general facts  $\det(AB) = \det A \det B$  and  $\det A^T = \det A$ . An **improper Lorentz transformation** is one with  $\det(L) = -1$ . An example is

$$L^\mu{}_\nu = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (27.1.21)$$

This just flips the sign of the time coordinate, and is therefore known as time reversal,

$$x'^0 = -x^0 \quad x'^1 = x^1 \quad x'^2 = x^2 \quad x'^3 = x^3. \quad (27.1.22)$$

Another improper Lorentz transformation is parity or space inversion,

$$L^\mu{}_\nu = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad (27.1.23)$$

so that

$$x'^0 = x^0, \quad x'^1 = -x^1, \quad x'^2 = -x^2, \quad x'^3 = -x^3. \quad (27.1.24)$$

It was once thought that the laws of physics must be invariant under these operations. Although this might seem intuitively reasonable, it was shown experimentally in the 1950s that parity is violated in the weak interactions, specifically in the weak decays of the  $^{60}\text{Co}$  nucleus and the  $K^\pm$  mesons. Likewise, experiments in the 1960s on the decays of  $K^0$  mesons showed that time-reversal invariance is not a symmetry of the Hamiltonian governing the weak interactions.

However, all experiments up to now are consistent with invariance of the laws of physics under the subset of Lorentz transformations that are continuously connected to the identity; these are known as **proper Lorentz transformations** and have  $\det(L) = +1$ . They can all be built up out of infinitesimal Lorentz transformations

$$L^\mu{}_\nu = \delta^\mu{}_\nu + \omega^\mu{}_\nu + \mathcal{O}(\omega^2). \quad (27.1.25)$$

For these, eq. (27.1.19) gives  $g_{\mu\nu}(\delta^\mu{}_\rho + \omega^\mu{}_\rho + \cdots)(\delta^\nu{}_\sigma + \omega^\nu{}_\sigma + \cdots) = g_{\rho\sigma}$ , or after simplification,  $g_{\rho\sigma} + \omega_{\sigma\rho} + \omega_{\rho\sigma} + \cdots = g_{\rho\sigma}$ . Therefore,

$$\omega_{\sigma\rho} = -\omega_{\rho\sigma} \quad (27.1.26)$$

is an antisymmetric  $4 \times 4$  matrix, with 6 independent entries. These nonzero elements correspond to 3 rotations ( $\rho, \sigma = 1, 2$  or  $1, 3$  or  $2, 3$ ) and 3 boosts ( $\rho, \sigma = 0, 1$  or  $0, 2$  or  $0, 3$ ). It is a mathematical fact that any Lorentz transformation can be built up out of repeated infinitesimal boosts and rotations, perhaps combined with time-reversal and/or space inversion. Lorentz transformations obey the mathematical properties of a group, known as the **Lorentz group**. The subset of Lorentz transformations that can be built out of repeated infinitesimal boosts and rotations form a smaller group, called the proper Lorentz group.

A **Lorentz tensor** is a multi-component object that can carry an arbitrary number of spacetime vector indices, and transforms appropriately when one goes to a new reference frame.

The objects  $g^{\mu\nu}$  and  $g_{\mu\nu}$  and  $\delta^\mu_\nu$  are examples of constant tensors. Contravariant and covariant four-vectors are also tensors. In general, the defining characteristic of a tensor function  $T^{\mu_1\mu_2\cdots}_{\nu_1\nu_2\cdots}(x)$  is that under a change of reference frame, it transforms so that in the primed coordinate system, the corresponding tensor  $T'$  is:

$$T'^{\mu_1\mu_2\cdots}_{\nu_1\nu_2\cdots}(x') = L^{\mu_1}_{\rho_1} L^{\mu_2}_{\rho_2} \cdots L^{\sigma_1}_{\nu_1} L^{\sigma_2}_{\nu_2} \cdots T^{\rho_1\rho_2\cdots}_{\sigma_1\sigma_2\cdots}(x). \quad (27.1.27)$$

A Lorentz scalar function is the special case of a Lorentz tensor that carries no vector indices, and is therefore independent of the choice of inertial reference frame.

For example, if  $p^\mu$  and  $q^\mu$  are the contravariant four-momenta of any particles, then  $p^\mu q_\mu$  is a scalar, and must not depend on which inertial reference frame it is computed or measured in. In particular, this implies that a particle with mass  $m$  satisfies the **on-shell condition**

$$p^\mu p_\mu = -E^2/c^2 + \vec{p}^2 = -m^2 c^2, \quad (27.1.28)$$

where the right side is obtained by evaluation in the rest frame with  $p^\mu = (mc, 0, 0, 0)$ .

Spacetime derivatives of functions are four-vectors. Suppose that  $f(x^\mu) = f(t, \vec{r})$  is a scalar function of  $x^\mu$ . Then

$$\partial_\mu f \equiv \frac{\partial f}{\partial x^\mu} = \left( \frac{1}{c} \frac{\partial f}{\partial t}, \vec{\nabla} f \right) \quad (27.1.29)$$

satisfies the Lorentz transformation rule of a covariant four-vector, eq. (27.1.15). By raising the index, one obtains a contravariant four-vector function

$$\partial^\mu f = g^{\mu\nu} \partial_\nu f = \left( -\frac{1}{c} \frac{\partial f}{\partial t}, \vec{\nabla} f \right). \quad (27.1.30)$$

One can obtain another scalar function by acting twice with the 4-dimensional derivative operator on  $f$ , contracting the indices,

$$\partial^\mu \partial_\mu f = -\frac{1}{c^2} \frac{\partial^2 f}{\partial t^2} + \nabla^2 f. \quad (27.1.31)$$

The differential operator  $\partial^\mu \partial_\mu$  is thus a four-dimensional generalization of the Laplacian.

## 27.2 Klein-Gordon wave equation

As a first attempt at relativistic quantum mechanics, let us consider the simple case of a free spinless particle of mass  $m$ , whose wavefunction  $\Phi(x^\mu)$  satisfies the Schrödinger equation

$$H\Phi = i\hbar \frac{\partial \Phi}{\partial t}. \quad (27.2.1)$$

To construct a Hamiltonian consistent with the symmetry of invariance under Lorentz transformations, we note that for a free particle, there should be an orthobasis of states  $\Phi$  that are also

eigenstates of  $\vec{P} = -i\hbar\vec{\nabla}$  with eigenvalues  $\vec{p}$ , the three-momentum of the particle. Now, using the special relativistic form  $E = \sqrt{m^2c^4 + c^2\vec{p}^2}$ , one can try

$$H = \sqrt{m^2c^4 + c^2\vec{P}^2}, \quad (27.2.2)$$

where  $\vec{P}^2 = -\hbar^2\nabla^2$  acting on wavefunctions. This has a series expansion,

$$H = mc^2 - \frac{\hbar^2}{2m}\nabla^2 - \frac{\hbar^4}{8m^3c^2}(\nabla^2)^2 + \dots \quad (27.2.3)$$

If we keep only the first two terms, then we recover the standard nonrelativistic quantum mechanics of a free particle, since the constant rest-energy contribution  $mc^2$  contributes only an unobservable phase to the time evolution. However, in the fully relativistic case in which the series is not truncated, the presence of an infinite number of terms with arbitrarily large numbers of derivatives is quite problematic, including apparently nonlocal effects. Furthermore, the Schrödinger equation in this case is not manifestly relativistic, since it involves only a single time derivative but an infinite number of spatial derivatives.

Instead, one can consider  $H^2\Phi$ , avoiding the square root. Setting this equal to  $(i\hbar\partial/\partial t)^2$  acting on  $\Phi$ , in order to be consistent with the Schrödinger equation, we find

$$\left(-\hbar^2\frac{\partial^2}{\partial t^2} + c^2\hbar^2\nabla^2 - m^2c^4\right)\Phi = 0. \quad (27.2.4)$$

Using eq. (27.1.31), this can be written in the manifestly Lorentz-invariant form

$$(\hbar^2\partial^\mu\partial_\mu - m^2c^2)\Phi = 0. \quad (27.2.5)$$

This wave equation is called the **Klein-Gordon** equation after the 1926 proposals of Oskar Klein and Walter Gordon.

It is easy to guess the solutions of the Klein-Gordon equation. If we try

$$\Phi(x^\mu) = \Phi_0 e^{ip_\mu x^\mu/\hbar}, \quad (27.2.6)$$

where  $\Phi_0$  is a normalization constant and  $p^\mu = (p^0, \vec{p})$  is a constant four-vector, then  $\partial_\mu\Phi = i(p_\mu/\hbar)\Phi$ , and so

$$\partial^\mu\partial_\mu\Phi = -(p^\mu p_\mu/\hbar^2)\Phi. \quad (27.2.7)$$

Therefore, we obtain a basis of solutions by imposing  $p^\mu p_\mu = -m^2c^2$ . Any  $\Phi$  of this form is an eigenstate of  $H$  and  $\vec{P}$ , with energy eigenvalue  $E = cp^0$  and three-momentum eigenvalue  $\vec{p}$ , provided that they satisfy the on-shell condition  $E^2 = m^2c^4 + \vec{p}^2c^2$ .

However, there is a big problem with this. For every positive-energy solution  $p^\mu = (E/c, \vec{p})$  with  $E = \sqrt{m^2c^4 + \vec{p}^2c^2} > 0$ , there is a corresponding negative-energy solution  $p^\mu = (-E/c, \vec{p})$ . By increasing  $|\vec{p}|$ , one can have arbitrarily large  $|E|$ , so the negative energies are not bounded from below. This would lead to the release of an infinite amount of energy as negative-energy particles interact, acquiring larger and larger three-momenta without bound.

### 27.3 Dirac equation

In 1928, Dirac suggested an alternative, based on the observation that the problem with the Klein-Gordon equation seemed to be that it is quadratic in  $H$ ; this leads to the sign ambiguity for  $E$ . Dirac could also have been motivated by the fact<sup>†</sup> that the electron has spin; since it has more than one intrinsic degree of freedom, trying to describe it with a single wavefunction  $\Phi(x^\mu)$  is doomed to failure. Instead, Dirac proposed to write a relativistic Schrödinger equation, linear in  $H$ , for a multi-component wavefunction  $\Psi_i(x^\mu)$ , where the index  $i = 1, 2, \dots, n$  runs over the components. These indices will be suppressed from here on, as we write  $\Psi$  as a column vector when its explicit form is needed.

Dirac reasoned that since the Schrödinger wave equation is linear in  $\partial/\partial t$ , and since relativity places  $t$  on the same footing as  $x, y, z$ , it should also be linear in derivatives of the spatial coordinates. Therefore, the equation ought to take the form

$$i\hbar \frac{\partial \Psi}{\partial t} = H\Psi = (c\vec{\alpha} \cdot \vec{P} + mc^2\beta)\Psi, \quad (27.3.1)$$

where  $\alpha_x, \alpha_y, \alpha_z$ , and  $\beta$  are dimensionless  $n \times n$  matrices. Equation (27.3.1) is known as the Dirac equation, and  $\Psi$  is called a **Dirac spinor**.

To determine what  $\vec{\alpha}$  and  $\beta$  should be, consider  $\partial^2\Psi/\partial t^2$ , which can be evaluated in two ways. First, the on-shell condition for  $E^2$  gives, just as in the Klein-Gordon case,

$$-\hbar^2 \frac{\partial^2 \Psi}{\partial t^2} = (-c^2\hbar^2\nabla^2 + m^2c^4)\Psi. \quad (27.3.2)$$

On the other hand, expressing  $H^2$  in terms of the right-hand side of eq. (27.3.1), we find

$$-\hbar^2 \frac{\partial^2 \Psi}{\partial t^2} = \left[ -c^2\hbar^2\alpha_a\alpha_b \frac{\partial^2}{\partial x^a\partial x^b} - i\hbar mc^3(\alpha_a\beta + \beta\alpha_a) \frac{\partial}{\partial x^a} + m^2c^4\beta^2 \right] \Psi, \quad (27.3.3)$$

where  $a$  and  $b$  are summed over  $x, y, z$ . Since partial derivatives commute,

$$\alpha_a\alpha_b \frac{\partial^2}{\partial x^a\partial x^b} = \frac{1}{2}(\alpha_a\alpha_b + \alpha_b\alpha_a) \frac{\partial^2}{\partial x^a\partial x^b}. \quad (27.3.4)$$

So, in order for eqs. (27.3.2) and (27.3.3) to agree, we must have

$$\beta^2 = I, \quad (27.3.5)$$

$$\alpha_a\beta + \beta\alpha_a = 0, \quad (27.3.6)$$

$$\alpha_a\alpha_b + \alpha_b\alpha_a = 2\delta_{ab}I. \quad (27.3.7)$$

It is also necessary that  $\vec{\alpha}$  and  $\beta$  are Hermitian matrices, in order for  $H$  to be Hermitian.

---

<sup>†</sup>He evidently realized this quickly, but only in hindsight.

For particles with non-zero mass, a solution for  $\vec{\alpha}$  and  $\beta$  turns out to require at least  $n = 4$  spinor indices.<sup>‡</sup> This may be somewhat surprising, since naively one only needs  $n = 2$  to describe a spin-1/2 particle like the electron. As we will see, the Dirac equation necessarily describes negative energy electrons in a way similar to the Klein-Gordon equation, accounting for the doubling. It is convenient to write the  $4 \times 4$  matrices in  $2 \times 2$  blocks. The necessary conditions are satisfied by

$$\vec{\alpha} = \begin{pmatrix} 0 & \vec{\sigma} \\ \vec{\sigma} & 0 \end{pmatrix}, \quad \beta = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad (27.3.8)$$

where  $\vec{\sigma}$  are the Pauli matrices,  $I$  is the  $2 \times 2$  identity, and “0” means a  $2 \times 2$  block of 0’s.

The Dirac equation automatically incorporates intrinsic angular momentum, or spin. To see this, we note that the Hamiltonian for a free particle should have the symmetry of rotations generated by a total angular momentum operator  $\vec{J}$ , so that  $[\vec{J}, H] = 0$ . Let us first consider how orbital angular momentum fits in. Since  $\vec{L} = \vec{R} \times \vec{P}$ , its components can be written as  $L_a = \epsilon_{abc} R_b P_c$ . Now we find the commutator of  $L_a$  with the Dirac Hamiltonian  $H$ . The  $mc^2\beta$  term has no coordinate dependence, so it does not contribute to the commutator. Therefore,  $[L_a, H] = c\alpha_b[L_a, P_b] = c\alpha_b\epsilon_{adc}[R_d, P_b]P_c = i\hbar c\alpha_b\epsilon_{adc}\delta_{bd}P_c = i\hbar c\epsilon_{abc}\alpha_bP_c$ , so that

$$[\vec{L}, H] = i\hbar c\vec{\alpha} \times \vec{P}. \quad (27.3.9)$$

This does not vanish, so there must be another contribution to  $\vec{J}$ . By inspired guesswork, we try defining the spin operator

$$\vec{S} = \frac{\hbar}{2} \begin{pmatrix} \vec{\sigma} & 0 \\ 0 & \vec{\sigma} \end{pmatrix}. \quad (27.3.10)$$

To check this guess, note that the non-zero commutator of  $\vec{S}$  with  $H$  is due entirely to the matrix nature of both,

$$[S_a, H] = cP_b[S_a, \alpha_b] = \frac{\hbar}{2}cP_b \begin{pmatrix} 0 & [\sigma_a, \sigma_b] \\ [\sigma_a, \sigma_b] & 0 \end{pmatrix} = i\hbar c\epsilon_{abc}P_b\alpha_c, \quad (27.3.11)$$

where we have made use of  $[\sigma_a, \sigma_b] = 2i\epsilon_{abc}\sigma_c$ . This gives

$$[\vec{S}, H] = -i\hbar c\vec{\alpha} \times \vec{P} = -[\vec{L}, H]. \quad (27.3.12)$$

Thus  $[\vec{J}, H] = 0$ , so that the Dirac Hamiltonian is indeed invariant under rotations generated by the total angular momentum operator  $\vec{J} = \vec{L} + \vec{S}$ .

---

<sup>‡</sup>The fact that Dirac spinors and Lorentz vectors both have 4 components is coincidental, and one must be careful not to confuse the two types of 4-dimensional spaces. If we lived in  $D$ -dimensional spacetime, Dirac spinors would have  $n = 2^{D/2}$  components.



The Dirac Hamiltonian also respects parity, but the way this works is slightly tricky. We can start by defining a “coordinate parity” operator  $\Pi_c$  that is proportional to the identity matrix in spinor space, and obeys

$$\Pi_c \vec{R} \Pi_c = -\vec{R}, \quad (27.3.13)$$

from which it follows that

$$\Pi_c \vec{P} \Pi_c = -\vec{P} \quad (27.3.14)$$

as well. However, it is then easy to see that  $\Pi_c H \Pi_c \neq H$ , because the  $\vec{\alpha} \cdot \vec{P}$  term flips sign, but the  $\beta$  term does not. Therefore, we define a matrix factor of the parity operator,

$$\Pi_m = \beta, \quad (27.3.15)$$

so that  $\Pi_m \vec{\alpha} \Pi_m = -\vec{\alpha}$  and  $\Pi_m \beta \Pi_m = \beta$ . Then the complete parity operator defined by

$$\Pi = \Pi_c \Pi_m = \begin{pmatrix} \Pi_c & 0 \\ 0 & -\Pi_c \end{pmatrix}, \quad (27.3.16)$$

satisfies  $\Pi H \Pi = H$ , and the Dirac Hamiltonian commutes with parity.

The Dirac equation (27.3.1) can be rewritten in a nicer way by multiplying it on the left by the matrix  $\beta$ , and defining gamma matrices  $\gamma^\mu$ , carrying a raised Lorentz index, by

$$\gamma^0 = \beta, \quad \gamma^a = \beta \alpha_a, \quad (a = 1, 2, 3). \quad (27.3.17)$$

Since  $\beta^2 = I$ , the Dirac equation can be expressed as

$$\left[ i\hbar \left( \gamma^0 \frac{\partial}{\partial x^0} + \gamma^1 \frac{\partial}{\partial x^1} + \gamma^2 \frac{\partial}{\partial x^2} + \gamma^3 \frac{\partial}{\partial x^3} \right) - mc \right] \Psi = 0, \quad (27.3.18)$$

or, even more nicely, as

$$(i\hbar \gamma^\mu \partial_\mu - mc) \Psi = 0. \quad (27.3.19)$$

The  $\gamma^\mu$  matrices defined above are explicitly given by

$$\gamma^0 = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad \gamma^a = \begin{pmatrix} 0 & \sigma^a \\ -\sigma^a & 0 \end{pmatrix}, \quad (a = 1, 2, 3). \quad (27.3.20)$$

However, this representation is not unique. To see this, suppose  $U$  is any constant unitary  $4 \times 4$  matrix satisfying  $U^\dagger U = U U^\dagger = I$ . Then the Dirac equation implies

$$U(i\hbar \gamma^\mu \partial_\mu - mc) U^\dagger U \Psi = 0, \quad (27.3.21)$$

from which it follows that, writing  $\gamma'^\mu = U\gamma^\mu U^\dagger$ , and  $\Psi' = U\Psi$ ,

$$(i\hbar\gamma'^\mu\partial_\mu - mc)\Psi' = 0. \quad (27.3.22)$$

So, the new  $\gamma'^\mu$  matrices together with the new spinor  $\Psi'$  are just as good as the old pair  $\gamma^\mu, \Psi$ ; there are an infinite number of different, equally valid choices. The choice in eqs. (27.3.8) and (27.3.20) is called the Dirac representation. Another choice, popular in treatments of high-energy particle physics and quantum field theory, is the chiral representation (see Exercise 27.1).

Some useful properties of the  $\gamma^\mu$  matrices, independent of the representation choice, are

$$\gamma^{0\dagger} = \gamma^0, \quad \gamma^{a\dagger} = -\gamma^a, \quad (a = 1, 2, 3), \quad (27.3.23)$$

$$\gamma^0\gamma^{\mu\dagger}\gamma^0 = \gamma^\mu, \quad (27.3.24)$$

$$\text{Tr}(\gamma^\mu\gamma^\nu) = -4g^{\mu\nu}, \quad (27.3.25)$$

$$\gamma^\mu\gamma_\mu = -4, \quad (27.3.26)$$

$$\gamma^\mu\gamma^\nu + \gamma^\nu\gamma^\mu = \{\gamma^\mu, \gamma^\nu\} = -2g^{\mu\nu}. \quad (27.3.27)$$

On the right sides of each of eqs. (27.3.26) and (27.3.27), there is an implicit  $4 \times 4$  unit matrix.

How does a Dirac spinor  $\Psi(x)$  transform under a Lorentz transformation? It carries no vector index, so it is not a tensor. On the other hand, the fact that the Hamiltonian “mixes up” the four components of  $\Psi(x)$  is a clue that it does not transform like an ordinary scalar function either. Instead, we might expect that the primed frame spinor is given by

$$\Psi'(x') = \Lambda\Psi(x), \quad (27.3.28)$$

where  $\Lambda$  is a  $4 \times 4$  matrix that depends on the Lorentz transformation matrix  $L^\mu{}_\nu$ . It is left to Exercise 27.2 to show that Lorentz invariance of the Dirac equation implies that, for an infinitesimal Lorentz transformation  $L^\mu{}_\nu = \delta^\mu{}_\nu + \omega^\mu{}_\nu$ ,

$$\Lambda = I - \omega_{\mu\nu}[\gamma^\mu, \gamma^\nu]/8 + \dots. \quad (27.3.29)$$

To obtain the result for a non-infinitesimal proper Lorentz transformation, we apply the same infinitesimal transformation  $N$  times, and take  $N \rightarrow \infty$ . Letting  $\Omega_{\mu\nu} = N\omega_{\mu\nu}$ , we obtain

$$L^\mu{}_\nu = \lim_{N \rightarrow \infty} (\delta^\mu{}_\nu + \Omega^\mu{}_\nu/N)^N = [\exp(\Omega)]^\mu{}_\nu \quad (27.3.30)$$

by using  $\lim_{N \rightarrow \infty} (1 + x/N)^N = e^x$ . For the Dirac spinor, one finds similarly

$$\Psi'(x') = \lim_{N \rightarrow \infty} (1 - \Omega_{\mu\nu}[\gamma^\mu, \gamma^\nu]/8N)^N \Psi(x) = \exp(-\Omega_{\mu\nu}[\gamma^\mu, \gamma^\nu]/8) \Psi(x). \quad (27.3.31)$$

So, the Lorentz transformation matrix for Dirac spinors as defined in eq. (27.3.28) is

$$\Lambda = \exp(-\Omega_{\mu\nu}[\gamma^\mu, \gamma^\nu]/8), \quad (27.3.32)$$

corresponding to the  $L^\mu_\nu$  that appears in eq. (27.3.30):

We conclude this section by constructing plane-wave solutions to the free-particle Dirac equation. Consider first the special case of the rest frame of the particle, so that the  $\vec{\alpha} \cdot \vec{P}$  term can be neglected. The Dirac equation then has the diagonal form

$$i\hbar \frac{\partial \Psi}{\partial t} = mc^2 \beta \Psi. \quad (27.3.33)$$

Defining 2-component spinors

$$\chi_\uparrow = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \chi_\downarrow = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad (27.3.34)$$

it is easy to check that a basis of four solutions is

$$\Psi_s^+ = e^{-imc^2 t/\hbar} \begin{pmatrix} \chi_s \\ 0 \end{pmatrix}, \quad \Psi_s^- = e^{imc^2 t/\hbar} \begin{pmatrix} 0 \\ \chi_s \end{pmatrix}, \quad (27.3.35)$$

with positive and negative energy eigenvalues  $E = mc^2$  and  $E = -mc^2$  for the superscript labels  $+$  and  $-$ , and  $S_z$  eigenvalues  $\hbar/2$  and  $-\hbar/2$  for the subscript labels  $s = \uparrow$  and  $\downarrow$ . These spin assignments follow immediately from the spin operator that we found in eq. (27.3.10).

Dirac spinor wavefunctions for arbitrary non-zero three-momenta can now be constructed at will from the rest-frame solutions by applying the Lorentz transformation for boosts and/or rotations using eq. (27.3.32). Alternatively, one can find the eigenspinors of the Dirac Hamiltonian directly. Let us do this for the special case of three-momentum  $p$  along the  $\hat{z}$  direction. Factoring out the plane-wave time and position dependence, we try

$$\Psi(\vec{r}, t) = e^{i(pz - Et)/\hbar} u, \quad (27.3.36)$$

where  $u$  is a constant spinor. Plugging this into the Dirac equation [either eq. (27.3.1), or its equivalent eq. (27.3.19)] gives

$$\begin{pmatrix} (mc^2 - E)I & pc\sigma_z \\ pc\sigma_z & (-mc^2 - E)I \end{pmatrix} u = 0, \quad (27.3.37)$$

which has two energy eigenvalue solutions  $E = \pm \sqrt{m^2 c^4 + p^2 c^2}$ , each with degeneracy 2. Up to normalization, the corresponding orthogonal basis spinors are, for  $E = +\sqrt{m^2 c^4 + p^2 c^2}$ ,

$$u_\uparrow^+ = \begin{pmatrix} \sqrt{m^2 c^4 + p^2 c^2} + mc^2 \\ 0 \\ pc \\ 0 \end{pmatrix}, \quad u_\downarrow^+ = \begin{pmatrix} 0 \\ \sqrt{m^2 c^4 + p^2 c^2} + mc^2 \\ 0 \\ -pc \end{pmatrix}, \quad (27.3.38)$$

and for  $E = -\sqrt{m^2 c^4 + p^2 c^2}$ ,

$$u_\uparrow^- = \begin{pmatrix} -pc \\ 0 \\ \sqrt{m^2 c^4 + p^2 c^2} + mc^2 \\ 0 \end{pmatrix}, \quad u_\downarrow^- = \begin{pmatrix} 0 \\ pc \\ 0 \\ \sqrt{m^2 c^4 + p^2 c^2} + mc^2 \end{pmatrix}. \quad (27.3.39)$$

These are again eigenstates of  $S_z$ , with eigenvalues  $\hbar/2$  for  $\uparrow$  and  $-\hbar/2$  for  $\downarrow$ .

Like the Klein-Gordon equation, the Dirac equation has the embarrassment of solutions with negative energies with arbitrarily large magnitudes, despite the fact that it is linear in  $H$  and  $\partial/\partial t$ . Undaunted, Dirac proposed to get around the problem of negative energy states by using the fact that spin-1/2 particles are fermions. Recall that the Pauli exclusion principle dictates that two fermions cannot occupy the same quantum state. This motivated Dirac to suggest that all of the negative energy states are normally occupied. This prevents electrons with positive energy from making disastrous transitions to the  $E < 0$  states. The infinite number of filled  $E < 0$  states is called the **Dirac sea**.

If one of the states in the Dirac sea becomes unoccupied, it leaves behind a “hole”. Since a hole is the absence of an  $E < 0$  state, it effectively has energy  $-E > 0$ . An electron has charge  $-e$ , so the hole corresponding to its absence effectively has the opposite charge,  $+e$ . Since both electrons and holes obey  $p^\mu p_\mu = m^2$ , they have the same mass. Dirac’s proposal therefore predicts the existence of “anti-electrons” or positrons, with positive energy and positive charge. Because no other candidate was known at the time, he briefly entertained the incorrect suggestion that the anti-electron was the proton, until the positron was discovered in 1932 in cosmic ray experiments.

Despite this impressive and revolutionary success, the modern solution of the problem of negative energy states for both the Klein-Gordon and Dirac cases is to abandon the whole approach of using operators for positions and momenta. Instead, relativistic particles are described by quantum field theories, in which the position coordinates  $\vec{r}$  join the time  $t$  as parameters, not operators. In quantum field theories, there are still objects  $\Phi(x^\mu)$  for spinless particles and  $\Psi(x^\mu)$  for spin-1/2 particles, but they are operators (“quantum fields”) rather than wavefunctions. This automatically avoids negative energy states and introduces antiparticles without appealing to Dirac’s rather *ad hoc* sea.<sup>§</sup> However, the Dirac equation formalism and Dirac spinors are still essential in setting up the kinematics of quantum field theories for spin-1/2 particles. There is much more that can be said about the Dirac wave equation and its solutions, and but we mostly leave this to texts that specialize in relativistic quantum field theory. In the remainder of this chapter, we will only discuss two important successes of the Dirac equation having to do with the coupling of the electron to electromagnetic fields.

---

<sup>§</sup>In any case, Dirac’s sea trick cannot work for the Klein-Gordon equation, or any other relativistic theory of bosons, since they do not have a Pauli exclusion principle. Unlike Dirac, we are privileged to know of a scalar particle that is quite possibly fundamental, the Higgs boson. There are also spin-1 bosons that are plausibly fundamental based on all presently available evidence (the photon, the  $W$  and  $Z$  particles, and the gluon of the Standard Model), and many composite integer-spin mesons.

## 27.4 Electromagnetic fields and the Dirac equation

In this section, we will work out how the Dirac equation incorporates the effects of background electromagnetic fields, and then recover the nonrelativistic limit.

As Einstein famously showed in 1905, Maxwell's electromagnetism is already a relativistic theory. To illustrate this, we start by revisiting some of the key equations of section 4.3, putting them into a manifestly Lorentz-covariant form. The classical electromagnetic potentials are unified into a contravariant four-vector,

$$A^\mu = (\Phi, \vec{A}). \quad (27.4.1)$$

The gauge transformation  $\vec{A} \rightarrow \vec{\nabla}\Lambda$  with  $\Phi \rightarrow \Phi - \frac{1}{c}\frac{\partial\Lambda}{\partial t}$  is then realized as

$$A_\mu \rightarrow A_\mu + \partial_\mu\Lambda \quad (27.4.2)$$

for an arbitrary scalar function  $\Lambda$ . The relations between the potentials and the fields,  $\vec{E} = -\vec{\nabla}\Phi - \frac{1}{c}\frac{\partial\vec{A}}{\partial t}$  and  $\vec{B} = \vec{\nabla} \times \vec{A}$ , can be expressed as

$$\begin{pmatrix} 0 & -E_x & -E_y & -E_z \\ E_x & 0 & B_z & -B_y \\ E_y & -B_z & 0 & B_x \\ E_z & B_y & -B_x & 0 \end{pmatrix} = F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu, \quad (27.4.3)$$

which defines the antisymmetric field strength tensor  $F_{\mu\nu}$ . The classical densities of charge and current can also be unified into a contravariant four-vector, as

$$j^\mu = (c\rho, \vec{j}). \quad (27.4.4)$$

The sourced Maxwell equations (Gauss' Law  $\vec{\nabla} \cdot \vec{E} = 4\pi\rho$  and Ampère's Law  $\vec{\nabla} \times \vec{B} = \frac{1}{c}\frac{\partial\vec{E}}{\partial t} + \frac{4\pi}{c}\vec{j}$ ) are then given by

$$\partial_\nu F^{\mu\nu} = \frac{4\pi}{c}j^\mu, \quad (27.4.5)$$

or equivalently,

$$\partial_\nu \partial^\mu A^\nu - \partial_\nu \partial^\nu A^\mu = \frac{4\pi}{c}j^\mu. \quad (27.4.6)$$

The remaining Maxwell equations ( $\vec{\nabla} \cdot \vec{B} = 0$  and Faraday's Law  $\vec{\nabla} \times \vec{E} = -\frac{1}{c}\frac{\partial\vec{B}}{\partial t}$ ) are equivalent to the identity

$$\partial_\rho F_{\mu\nu} + \partial_\mu F_{\nu\rho} + \partial_\nu F_{\rho\mu} = 0, \quad (27.4.7)$$

for  $\mu, \nu, \rho = 0, 1, 2, 3$ . Note that this equation is automatically true by construction in terms of  $A_\mu$ , as can be seen using eq. (27.4.3).

Also, because  $F^{\mu\nu}$  is antisymmetric and partial derivatives commute, we have  $\partial_\mu \partial_\nu F^{\mu\nu} = 0$ . It follows, by comparing with  $\partial_\mu$  acting on eq. (27.4.5), that

$$\partial_\mu j^\mu = 0, \quad (27.4.8)$$

which expresses the local conservation of charge, as it can be rewritten  $\vec{\nabla} \cdot \vec{j} + \partial\rho/\partial t = 0$ .

To see how to incorporate electromagnetic effects into the Dirac equation, we recall from eq. (4.3.16) that the space and time derivatives in the Schrödinger equation are modified, in the presence of  $\vec{A}$  and  $\Phi$ , according to

$$\vec{P} \rightarrow \vec{P} - \frac{q}{c}\vec{A}, \quad i\hbar\frac{\partial}{\partial t} \rightarrow i\hbar\frac{\partial}{\partial t} - q\Phi. \quad (27.4.9)$$

Applying this rule to eq. (27.3.1), we arrive at the version of the Dirac Hamiltonian with electromagnetic potentials included,

$$H = c\vec{\alpha} \cdot \left(\vec{P} - \frac{q}{c}\vec{A}\right) + mc^2\beta + q\Phi. \quad (27.4.10)$$

Also, the two parts of eq. (27.4.9) can be unified neatly into a Lorentz-covariant rule

$$i\hbar\partial_\mu \rightarrow i\hbar\partial_\mu + \frac{q}{c}A_\mu. \quad (27.4.11)$$

Applying this to eq. (27.3.19), we arrive at the covariant form of the Dirac wave equation,

$$\left[\gamma^\mu \left(i\hbar\partial_\mu + \frac{q}{c}A_\mu\right) - mc\right] \Psi = 0, \quad (27.4.12)$$

for a particle of mass  $m$  and charge  $q$  in the presence of electromagnetic potentials.

We now wish to recover the nonrelativistic limit of the Dirac equation; this is a useful thing to do because it fixes the  $g$ -factor for the electron to be  $g_e = 2$  (before corrections coming from virtual photons of the quantized electromagnetic field), and provides the fine-structure terms in the Hamiltonian for the hydrogen atom. To accomplish this, it is useful to write the Dirac spinor as two 2-component spinors  $\varphi$  and  $\chi$ , and factor out the largest part of the time dependence for positive energy states,

$$\Psi = e^{-imc^2t/\hbar} \begin{pmatrix} \varphi \\ \chi \end{pmatrix}. \quad (27.4.13)$$

The Schrödinger equation for the Hamiltonian in eq. (27.4.10) then becomes

$$i\hbar\frac{\partial\varphi}{\partial t} = c\vec{\Pi} \cdot \vec{\sigma}\chi + q\Phi\varphi, \quad (27.4.14)$$

$$i\hbar\frac{\partial\chi}{\partial t} = c\vec{\Pi} \cdot \vec{\sigma}\varphi + (q\Phi - 2mc^2)\chi, \quad (27.4.15)$$

where  $\vec{\Pi} = \vec{P} - (q/c)\vec{A}$ . Now, in the nonrelativistic limit (small  $p$ ) limit, we see from eq. (27.3.38) that the positive-energy solutions are mostly  $\varphi$ , and from eq. (27.3.38) that the negative-energy solutions are mostly  $\chi$ . Since our goal is to take the nonrelativistic limit for the positive-energy solutions, our strategy will be to eliminate the small components  $\chi$  in favor of the large components  $\varphi$ . Accordingly, we rewrite eq. (27.4.15) as

$$\chi = \frac{1}{2mc}\Omega\varphi + \frac{1}{2mc^2}\left(-i\hbar\frac{\partial}{\partial t} + q\Phi\right)\chi, \quad (27.4.16)$$

where we have defined, for convenience, the Hermitian operator

$$\Omega = \vec{\Pi} \cdot \vec{\sigma}. \quad (27.4.17)$$

By iterative elimination of  $\chi$  in the limit of large  $c$ , eq. (27.4.16) gives

$$\chi = \frac{1}{2mc}\left[I + \frac{1}{2mc^2}\left(-i\hbar\frac{\partial}{\partial t} + q\Phi\right) + \cdots\right]\Omega\varphi. \quad (27.4.18)$$

Putting this into eq. (27.4.14) gives an uncoupled differential equation for  $\varphi$ ,

$$i\hbar\frac{\partial\varphi}{\partial t} = \left[\frac{1}{2m}\Omega^2 + q\Phi + \frac{1}{4m^2c^2}\Omega\left(-i\hbar\frac{\partial}{\partial t} + q\Phi\right)\Omega + \cdots\right]\varphi, \quad (27.4.19)$$

where terms with 4 or more powers of  $c$  in the denominator have been dropped.

It might be tempting to interpret eq. (27.4.19) as the wave equation for the positive energy solutions, but this is wrong for a subtle reason: unitary time evolution does not preserve the norm of  $\varphi$ . To see this, note that the squared norm of the full Dirac spinor obeys

$$1 = \int d^3\vec{r} \Psi^\dagger \Psi = \int d^3\vec{r} (\varphi^\dagger \varphi + \chi^\dagger \chi) = \int d^3\vec{r} \left( \varphi^\dagger \varphi + \varphi^\dagger \frac{\Omega^2}{4m^2c^2} \varphi + \cdots \right), \quad (27.4.20)$$

where in the last expression we have used the leading approximation for  $\chi$  from eq. (27.4.18).

This shows that the two-component spinor  $\psi$  defined by

$$\varphi = \left(I - \frac{\Omega^2}{8m^2c^2} + \cdots\right)\psi \quad (27.4.21)$$

will have constant norm, and therefore unitary time evolution. Thus,  $\psi$  defined in this way (and not  $\varphi$ ) is the correctly normalized wavefunction describing the nonrelativistic limit of a spin-1/2 particle with leading relativistic corrections. Plugging eq. (27.4.21) into eq. (27.4.19), and consistently dropping terms of order  $1/c^4$ , after a little algebra the result can be written

$$i\hbar\frac{\partial\psi}{\partial t} = \left(\frac{1}{2m}\Omega^2 + q\Phi + \Delta H + \cdots\right)\psi, \quad (27.4.22)$$

where the leading-order relativistic correction is

$$\Delta H = \frac{1}{4m^2c^2} \left( \Omega^2 \left( -i\hbar \frac{\partial}{\partial t} + q\Phi \right) - \frac{1}{2} \left[ \Omega, \left[ \Omega, -i\hbar \frac{\partial}{\partial t} + q\Phi \right] \right] \right). \quad (27.4.23)$$

To put the nonrelativistic part into a more familiar form, we evaluate

$$\Omega^2 = (\vec{\Pi} \cdot \vec{\sigma})^2 = \vec{\Pi}^2 + i\vec{\sigma} \cdot (\vec{\Pi} \times \vec{\Pi}) = \left( \vec{P} - \frac{q}{c} \vec{A} \right)^2 - \frac{\hbar q}{c} \vec{\sigma} \cdot (\vec{\nabla} \times \vec{A}), \quad (27.4.24)$$

where the second equality uses the Pauli matrix product identity  $(\vec{a} \cdot \vec{\sigma})(\vec{b} \cdot \vec{\sigma}) = \vec{a} \cdot \vec{b} + i(\vec{a} \times \vec{b}) \cdot \vec{\sigma}$ , and the last term uses the position-representation differential form  $\vec{P} = -i\hbar \vec{\nabla}$ . Now, using  $\vec{\nabla} \times \vec{A} = \vec{B}$ , and  $\vec{\sigma} = 2\vec{S}/\hbar$ , we arrive at

$$i\hbar \frac{\partial \psi}{\partial t} = \left( \frac{1}{2m} \left( \vec{P} - \frac{q}{c} \vec{A} \right)^2 + q\Phi - \frac{q}{mc} \vec{S} \cdot \vec{B} + \Delta H + \dots \right) \psi. \quad (27.4.25)$$

This agrees with the form of eq. (4.3.28), provided that the gyromagnetic ratio is  $\gamma = q/mc$ . Indeed, for the electron with  $q = -e$ , we have learned that the gyromagnetic ratio is successfully predicted by the Dirac equation. Comparing with eq. (4.3.19),

$$\gamma_e = -\frac{g_e e}{2mc} = -\frac{e}{mc}, \quad (27.4.26)$$

so that  $g_e = 2$ . The same holds for the muon and the tau lepton (just with different masses), as they are also fundamental fermions. However, the Dirac equation cannot give the correct gyromagnetic ratios for composite fermions like the proton and the neutron, because of large contributions from their internal structure.

Returning to the leading-order relativistic correction in eq. (27.4.23), the first term simplifies because, from eq. (27.4.19),  $(-i\hbar \frac{\partial}{\partial t} + q\Phi)\psi = -\frac{\Omega^2}{2m}\psi$ , up to higher order terms in  $1/c$ . So

$$\Delta H = -\frac{1}{8m^3c^2} \Omega^4 - \frac{1}{8m^2c^2} \left[ \Omega, \left[ \Omega, -i\hbar \frac{\partial}{\partial t} + q\Phi \right] \right]. \quad (27.4.27)$$

From here on, we consistently drop terms in  $\Delta H$  with  $-(q/c)\vec{A}$ . So, at leading order in  $1/c$ , we can write  $\Omega^4 = (P^2)^2$ . Also, we can evaluate

$$\left[ \Omega, -i\hbar \frac{\partial}{\partial t} + q\Phi \right] = -i\hbar q \vec{\sigma} \cdot \vec{\nabla} \Phi = i\hbar q \vec{\sigma} \cdot \vec{E}, \quad (27.4.28)$$

and then,

$$\left[ \Omega, \left[ \Omega, -i\hbar \frac{\partial}{\partial t} + q\Phi \right] \right] = i\hbar q \left( \vec{\sigma} \cdot \vec{P} \vec{\sigma} \cdot \vec{E} - \vec{\sigma} \cdot \vec{E} \vec{\sigma} \cdot \vec{P} \right) \quad (27.4.29)$$

$$= \hbar^2 q \vec{\nabla} \cdot \vec{E} + 2\hbar q \vec{\sigma} \cdot (\vec{E} \times \vec{P}) + i\hbar^2 q \vec{\sigma} \cdot (\vec{\nabla} \times \vec{E}), \quad (27.4.30)$$



after using the Pauli matrix product identity again. Dropping the last term because  $\vec{\nabla} \times \vec{E} = -\frac{1}{c} \frac{\partial \vec{E}}{\partial t}$  has an extra power of  $1/c$ , we finally get

$$\Delta H = -\frac{1}{8m^3c^2}(P^2)^2 - \frac{q}{2m^2c^2}\vec{S} \cdot (\vec{E} \times \vec{P}) - \frac{\hbar^2 q}{8m^2c^2}\vec{\nabla} \cdot \vec{E}. \quad (27.4.31)$$

These three terms are, respectively, the relativistic kinematic, spin-orbit, and Darwin terms that appeared in the fine-structure Hamiltonian for an electron in a central electrostatic potential, eqs. (17.1.3)-(17.1.5). In making this comparison, one can use  $q = -e$ ,  $m = m_e$ ,  $\vec{\nabla} \cdot \vec{E} = -\nabla^2 \Phi$ , and  $\vec{E} = -\frac{1}{R} \frac{d\Phi}{dR} \vec{R}$  for a spherically symmetric potential, followed by  $\vec{R} \times \vec{P} = \vec{L}$ . In particular, the Dirac equation automatically gives the Thomas precession factor in the spin-orbit Hamiltonian, and the correct normalization of the Darwin term.

## 27.5 Dirac equation solutions for the hydrogen atom

In the previous section, we obtained the relativistic corrections to the Hamiltonian for a spin-1/2 charged particle, at leading order in  $1/c^2$ . These can in turn be used to obtain the fine-structure energy corrections for the hydrogen atom, as we had already done in section 17.1 using first-order perturbation theory. We will now show that these effects can also be obtained directly from an exact<sup>†</sup> solution of the Dirac equation, without relying on perturbation theory and to all orders in  $1/c$ . As a bonus, we will find the relativistic wavefunction.

Our task is to solve the equation

$$H\Psi(\vec{r}) = E\Psi(\vec{r}) \quad (27.5.1)$$

with the Dirac Hamiltonian in the position representation,

$$H = -i\hbar c \vec{\alpha} \cdot \vec{\nabla} + mc^2 \beta - Ze^2/r, \quad (27.5.2)$$

which arises from  $q = -e$  and the four-vector potential  $A^\mu = (Ze^2/r, 0, 0, 0)$  for a nucleus of charge  $Ze$ . Now, eq. (27.3.9) implies that  $L_z$  and  $L^2$  do not commute with  $H$ , so we cannot hope to find energy eigenstates that are also eigenstates of orbital angular momentum with fixed  $l$ . However, it follows from eq. (27.3.12) that  $\vec{J} = \vec{L} + \vec{S}$  does commute with  $H$ , and the parity operator  $\Pi$  defined in eq. (27.3.16) does as well. Therefore, an appropriate CSCO for our problem is  $H$ ,  $J^2$ ,  $J_z$ , and  $\Pi$ .

We will use spherical coordinates  $(r, \theta, \phi)$ , and begin by constructing two-component spinor eigenstates of  $J^2$  and  $J_z$  with fixed  $L^2$  eigenvalue  $\hbar^2 l(l+1)$ , called **spinor harmonics**. Then we

---

<sup>†</sup>However, the results we will obtain below are not fully exact because nuclear spin, nuclear structure, and virtual photon effects from quantization of the electromagnetic field are not included in the Dirac equation. Thus, the hyperfine splitting and Lamb shift are neglected in this section.

will form the four-component Dirac spinor energy eigenstates by combining two spinor harmonics with different  $l$ , but the same  $J^2$  and  $J_z$  eigenvalues  $\hbar^2 j(j+1)$  and  $\hbar m$ .

Fortunately, the problem of adding orbital and spin-1/2 angular momentum was already done in section 12.3. For a given  $j$  and  $m$ , we can construct two distinct spinor harmonics, one with  $l = j - 1/2$  and another with  $l = j + 1/2$ . From eqs. (12.3.11) and (12.3.14), we see that they are respectively

$$\phi_{jm}^+ = \frac{1}{\sqrt{2l+1}} \begin{pmatrix} \sqrt{l+1/2+m} Y_l^{m-1/2} \\ \sqrt{l+1/2-m} Y_l^{m+1/2} \end{pmatrix}, \quad (l = j - 1/2), \quad (27.5.3)$$

$$\phi_{jm}^- = \frac{1}{\sqrt{2l+1}} \begin{pmatrix} \sqrt{l+1/2-m} Y_l^{m-1/2} \\ -\sqrt{l+1/2+m} Y_l^{m+1/2} \end{pmatrix}, \quad (l = j + 1/2), \quad (27.5.4)$$

where in each case the upper and lower components have  $S_z$  eigenvalues  $\hbar/2$  and  $-\hbar/2$ , and we have chosen their overall phases for future convenience. Note that  $m$  is always half-integer, and  $Y_l^{m+1/2}$  and  $Y_l^{m-1/2}$  are the usual spherical harmonic functions. For a given  $j$ , the spherical harmonics  $\phi_{jm}^+$  and  $\phi_{jm}^-$  have  $l$  differing by one unit, so they have opposite coordinate parity  $\Pi_c$  eigenvalues  $(-1)^l$ .

Because the potential has spherical symmetry and we will be using spherical coordinates, it is useful to consider the operator

$$\vec{\sigma} \cdot \hat{r} = \begin{pmatrix} \cos \theta & e^{-i\phi} \sin \theta \\ e^{i\phi} \sin \theta & -\cos \theta \end{pmatrix}. \quad (27.5.5)$$

Remarkably, this acts on the spinor harmonics by turning them into each other,

$$\phi_{jm}^+ = \vec{\sigma} \cdot \hat{r} \phi_{jm}^-, \quad \phi_{jm}^- = \vec{\sigma} \cdot \hat{r} \phi_{jm}^+. \quad (27.5.6)$$

To prove it, note that as an operator,  $\vec{\sigma} \cdot \hat{r}$  commutes with  $J^2$  and  $J_z$  and has odd parity with respect to the operator  $\Pi_c$  defined in eq. (27.3.13). This implies that  $\vec{\sigma} \cdot \hat{r} \phi_{jm}^-$  must have the same  $j$  and  $m$  and opposite parity as  $\phi_{jm}^-$ . The unique spinor harmonic with this property is  $\phi_{jm}^+$ , so it must be true that  $\vec{\sigma} \cdot \hat{r} \phi_{jm}^- = C \phi_{jm}^+$  for some constant of proportionality  $C$ . To evaluate  $C$ , we can consider the special case  $\theta = 0$ , so that  $\hat{r} = \hat{z}$  and  $\vec{\sigma} \cdot \hat{z} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ . Now, using  $Y_l^{m'}(0, \phi) = \delta_{m'0} \sqrt{\frac{2l+1}{4\pi}}$  as found in eq. (8.6.53), we have

$$\phi_{jm}^+ = \sqrt{\frac{2j+1}{8\pi}} \begin{pmatrix} \delta_{m, \frac{1}{2}} \\ \delta_{m, -\frac{1}{2}} \end{pmatrix}, \quad \phi_{jm}^- = \sqrt{\frac{2j+1}{8\pi}} \begin{pmatrix} \delta_{m, \frac{1}{2}} \\ -\delta_{m, -\frac{1}{2}} \end{pmatrix} \quad (\text{for } \theta = 0). \quad (27.5.7)$$

This shows that we must have  $C = 1$ , establishing the first part of eq. (27.5.6). The identity

$$(\vec{\sigma} \cdot \hat{r})^2 = I \quad (27.5.8)$$

then immediately implies the second part of eq. (27.5.6) from the first.

We seek eigenstates of  $H$  that are also eigenstates of  $J^2$ ,  $J_z$ , and  $\Pi$ . Given the spinor harmonic definitions of eqs. (27.5.3) and (27.5.4), and the form of the parity operator  $\Pi = \Pi_c \beta$  defined in eq. (27.3.16), we see that the four-component Dirac spinors

$$\Psi_{jm}^{\pm}(r, \theta, \phi) = \frac{1}{r} \begin{pmatrix} G_{jm}^{\pm}(r) \phi_{jm}^{\pm} \\ -i F_{jm}^{\pm}(r) \phi_{jm}^{\mp} \end{pmatrix} \quad (27.5.9)$$

have  $J^2$ ,  $J_z$ , and  $\Pi$  eigenvalues  $\hbar^2 j(j+1)$ ,  $\hbar m$ , and  $(-1)^{j \mp 1/2}$ , respectively. The radial wavefunctions  $G_{jm}^{\pm}(r)$  and  $F_{jm}^{\pm}(r)$  are to be determined by solving  $H\Psi = E\Psi$ , and the factors of  $1/r$  and  $-i$  are introduced for later convenience.

To evaluate the first term in  $H\Psi$ , we need to find  $\vec{\sigma} \cdot \vec{\nabla}$  acting on our spherical harmonics multiplied by radial functions  $f(r)$ . We therefore calculate:

$$\vec{\sigma} \cdot \vec{\nabla} (f(r) \phi_{jm}^{\pm}) = \vec{\sigma} \cdot \hat{r} \vec{\sigma} \cdot \hat{r} \vec{\sigma} \cdot \vec{\nabla} f(r) \phi_{jm}^{\pm} = \vec{\sigma} \cdot \hat{r} \left( \hat{r} \cdot \vec{\nabla} + \frac{i}{r} \vec{\sigma} \cdot (\vec{r} \times \vec{\nabla}) \right) f(r) \phi_{jm}^{\pm} \quad (27.5.10)$$

$$= \vec{\sigma} \cdot \hat{r} \left( \frac{\partial}{\partial r} - \frac{1}{r} \frac{2}{\hbar^2} \vec{S} \cdot \vec{L} \right) f(r) \phi_{jm}^{\pm}. \quad (27.5.11)$$

The first equality used  $(\vec{\sigma} \cdot \hat{r})^2 = I$ , the second employed the Pauli matrix product identity  $(\vec{a} \cdot \vec{\sigma})(\vec{b} \cdot \vec{\sigma}) = \vec{a} \cdot \vec{b} + i(\vec{a} \times \vec{b}) \cdot \vec{\sigma}$ , and the third used  $\vec{S} = \hbar \vec{\sigma}/2$  and the position representation identity  $\vec{L} = \vec{R} \times \vec{P} = \vec{r} \times (-i\hbar \vec{\nabla})$ . Now we use  $2\vec{S} \cdot \vec{L}/\hbar^2 = j(j+1) - l(l+1) - 3/4$ , which can in turn be simplified using  $l = j \mp 1/2$ . The result can be written as

$$\vec{\sigma} \cdot \vec{\nabla} (f(r) \phi_{jm}^{\pm}) = \left( \frac{df}{dr} + (1 + \kappa_{\pm}) \frac{f}{r} \right) \phi_{jm}^{\mp}, \quad (27.5.12)$$

where, for convenience, we have defined

$$\kappa_{\pm} = \mp(j + 1/2). \quad (27.5.13)$$

Now plugging eq. (27.5.9) into the Dirac equation (27.5.1) with the Hamiltonian eq. (27.5.2), and using eqs. (27.5.12) with (27.5.13), gives

$$\left( E - mc^2 + \frac{Ze^2}{r} \right) G + \hbar c \left( \frac{d}{dr} - \frac{\kappa}{r} \right) F = 0, \quad (27.5.14)$$

$$\left( E + mc^2 + \frac{Ze^2}{r} \right) F - \hbar c \left( \frac{d}{dr} + \frac{\kappa}{r} \right) G = 0, \quad (27.5.15)$$

where each of  $G$ ,  $F$ , and  $\kappa$  has an implicit  $\pm$ , suppressed along with the subscripts  $jm$  from here on. For further simplicity, define dimensionless quantities

$$\mathcal{E} = E/mc^2, \quad s = mcr/\hbar, \quad (27.5.16)$$

so that our first-order coupled differential equations become

$$\left(\frac{d}{ds} - \frac{\kappa}{s}\right) F - \left(1 - \mathcal{E} - \frac{Z\alpha}{s}\right) G = 0, \quad (27.5.17)$$

$$\left(\frac{d}{ds} + \frac{\kappa}{s}\right) G - \left(1 + \mathcal{E} + \frac{Z\alpha}{s}\right) F = 0. \quad (27.5.18)$$

Note that for a bound state,  $E < mc^2$ , so  $\mathcal{E} < 1$ .

We now examine the large distance behaviors of  $G$  and  $F$ . For large  $s$ , the terms proportional to  $\kappa$  and  $Z\alpha$  can be neglected, so we get  $F' = (1 - \mathcal{E})G$  and  $G' = (1 + \mathcal{E})F$ , which imply  $G'' = (1 - \mathcal{E}^2)G$  and  $F'' = (1 - \mathcal{E}^2)F$ . The solutions at large  $s$  therefore behave like

$$G, F \sim e^{-\sqrt{1-\mathcal{E}^2}s} \quad (\text{for large } s). \quad (27.5.19)$$

Conversely, for small  $s$ , one finds  $F' = (\kappa F - Z\alpha G)/s$  and  $G' = (-\kappa G + Z\alpha F)/s$ , which can be disentangled to give

$$\left(s\frac{d}{ds}\right)^2 G = \gamma^2 G, \quad \left(s\frac{d}{ds}\right)^2 F = \gamma^2 F, \quad (27.5.20)$$

with

$$\gamma = \sqrt{\kappa^2 - Z^2\alpha^2}. \quad (27.5.21)$$

This is solved by a simple power-law behavior

$$G, F \sim s^\gamma, \quad (\text{for small } s), \quad (27.5.22)$$

where the possibility  $s^{-\gamma}$  is rejected so that the wavefunction has normalizable behavior near  $s = 0$ . We now factor out these large- and small-distance behaviors by writing

$$G(s) = s^\gamma e^{-\sqrt{1-\mathcal{E}^2}s} g(s), \quad F(s) = s^\gamma e^{-\sqrt{1-\mathcal{E}^2}s} f(s). \quad (27.5.23)$$

This defines new functions  $g(s)$  and  $f(s)$ , which should be finite and non-zero at  $s = 0$ , and should not grow exponentially for large  $s$ . Plugging these into eqs. (27.5.17) and (27.5.18) gives

$$\left(\frac{d}{ds} - \sqrt{1-\mathcal{E}^2} + \frac{\gamma - \kappa}{s}\right) f - \left(1 - \mathcal{E} - \frac{Z\alpha}{s}\right) g = 0, \quad (27.5.24)$$

$$\left(\frac{d}{ds} - \sqrt{1-\mathcal{E}^2} + \frac{\gamma + \kappa}{s}\right) g - \left(1 + \mathcal{E} + \frac{Z\alpha}{s}\right) f = 0. \quad (27.5.25)$$

Since  $g(s)$  and  $f(s)$  are finite and non-zero at  $s = 0$ , we try series solutions

$$g(s) = \sum_{q=0}^{\infty} a_q s^q, \quad f(s) = \sum_{q=0}^{\infty} b_q s^q. \quad (27.5.26)$$

Putting this into eqs. (27.5.24) and (27.5.25) gives the coupled recurrence relations

$$(1 - \mathcal{E})a_{q-1} + \sqrt{1 - \mathcal{E}^2} b_{q-1} - Z\alpha a_q - (q + \gamma - \kappa)b_q = 0, \quad (27.5.27)$$

$$(1 + \mathcal{E})b_{q-1} + \sqrt{1 - \mathcal{E}^2} a_{q-1} + Z\alpha b_q - (q + \gamma + \kappa)a_q = 0. \quad (27.5.28)$$

Consider first the case  $q = 0$ . Since we are assuming  $a_{-1} = b_{-1} = 0$ , we learn that

$$\frac{b_0}{a_0} = \frac{Z\alpha}{\kappa - \gamma} = \frac{\kappa + \gamma}{Z\alpha}. \quad (27.5.29)$$

Note that the last equality is a check on  $\gamma$  in eq. (27.5.21).

For general  $q$ , multiply eq. (27.5.27) by  $\sqrt{1 + \mathcal{E}}$  and eq. (27.5.28) by  $\sqrt{1 - \mathcal{E}}$ , and then take the difference. The coefficients of  $a_{q-1}$  and  $b_{q-1}$  both cancel, and we find that the coefficients of the two series obey the proportionality

$$\frac{b_q}{a_q} = \frac{(q + \gamma + \kappa)\sqrt{1 - \mathcal{E}} - Z\alpha\sqrt{1 + \mathcal{E}}}{(q + \gamma - \kappa)\sqrt{1 + \mathcal{E}} + Z\alpha\sqrt{1 - \mathcal{E}}}. \quad (27.5.30)$$

Using these in eqs. (27.5.27) and (27.5.28), it follows that for large  $q$ ,

$$\frac{a_q}{a_{q-1}} \approx \frac{b_q}{b_{q-1}} \approx \frac{2\sqrt{1 - \mathcal{E}^2}}{q}. \quad (27.5.31)$$

We can therefore see that both series must terminate. Otherwise, we would find  $g, f \sim e^{2\sqrt{1 - \mathcal{E}^2}s}$ , and thus  $G, F \sim e^{\sqrt{1 - \mathcal{E}^2}s}$ , and the wavefunctions would grow exponentially at large distances, failing to be normalizable.

Therefore, we must have  $a_{n'+1} = 0$  for some non-negative integer  $n'$ , and then eq. (27.5.30) implies that  $b_{n'+1} = 0$  as well. Then eqs. (27.5.27) and (27.5.28) with  $q = n' + 1$  tell us

$$\frac{b_{n'}}{a_{n'}} = -\sqrt{\frac{1 - \mathcal{E}}{1 + \mathcal{E}}}. \quad (27.5.32)$$

Requiring that this is equal to eq. (27.5.30) with  $q = n'$ , we find the condition for a normalizable bound state,

$$Z\alpha\mathcal{E} = \sqrt{1 - \mathcal{E}^2}(n' + \gamma). \quad (27.5.33)$$

This has the positive-energy solution

$$\mathcal{E} = \left[ 1 + \left( \frac{Z\alpha}{n' + \gamma} \right)^2 \right]^{-1/2}, \quad (27.5.34)$$

or, restoring  $E$ ,  $j$ , and  $Z\alpha$  using eqs. (27.5.16), (27.5.21), and then (27.5.13),

$$E = mc^2 \left[ 1 + \left( \frac{Z\alpha}{n' + \sqrt{(j + 1/2)^2 - Z^2\alpha^2}} \right)^2 \right]^{-1/2}. \quad (27.5.35)$$

Now let  $n = n' + j + 1/2$ , which is a positive integer, since  $n'$  is non-negative and  $j + 1/2$  is a positive integer. Then we can rewrite the bound-state energy eigenvalues as

$$E = mc^2 \left[ 1 + \frac{Z^2 \alpha^2}{(n + \delta_j)^2} \right]^{-1/2}, \quad (27.5.36)$$

where

$$\delta_j = \sqrt{(j + 1/2)^2 - Z^2 \alpha^2} - (j + 1/2). \quad (27.5.37)$$

Expanding in small  $Z\alpha$ ,

$$E = mc^2 \left[ 1 - \frac{Z^2 \alpha^2}{2n^2} + \frac{Z^4 \alpha^4}{2n^3} \left( \frac{3}{4n} - \frac{1}{j + 1/2} \right) + \dots \right]. \quad (27.5.38)$$

The first term is the electron's rest energy, and the second term is the usual nonrelativistic energy eigenvalue  $-Z^2 e^2 / 2a_0 n^2$ , so we recognize  $n$  as the principal quantum number. The third term is the total fine-structure correction to the energy that we found (with  $Z = 1$ ) in eq. (17.1.26).

Finally, we should carefully enumerate the valid solutions that we have found. For each  $j$  and  $m$ , there are two solutions with the same  $E$ , with wavefunctions  $\Psi_{jm}^\pm$  in eq. (27.5.9), but with one exception. The exception is that if  $n' = 0$ , then we have from eqs. (27.5.29) and (27.5.32) that  $b_0/a_0 = -\sqrt{(1 - \mathcal{E})/(1 + \mathcal{E})} = (\kappa + \gamma)/Z\alpha$ , but this is only possible if  $\kappa = \mp(j + 1/2)$  is negative, because  $\gamma$  and  $Z\alpha$  are positive. This means that in the special case  $n' = 0$  the would-be solution  $\Psi_{jm}^-$  in eq. (27.5.9) is not valid. Therefore, for each principal quantum number  $n = 1, 2, 3, \dots$ , there are two bound state solutions  $\Psi_{jm}^\pm$  with  $j = 1/2, 3/2, \dots, n - 3/2$ , and only one bound state solution  $\Psi_{jm}^+$  with  $j = n - 1/2$ . Each of these has further degeneracy  $2j + 1$  from the  $m$  quantum number. This is in accord with the results found at the end of section 17.1 from the perturbative approach to the fine structure of the hydrogen atom.

## 27.6 Exercises

**Exercise 27.1.** The chiral representation of gamma matrices for the Dirac equation is

$$\gamma^0 = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, \quad \gamma^a = \begin{pmatrix} 0 & \sigma^a \\ -\sigma^a & 0 \end{pmatrix}, \quad (a = 1, 2, 3).. \quad (27.6.1)$$

Find a unitary matrix  $U$  such that  $\gamma_{\text{chiral}}^\mu = U \gamma_{\text{Dirac}}^\mu U^\dagger$ , where  $\gamma_{\text{Dirac}}^\mu$  are the matrices in eq. (27.3.20).

**Exercise 27.2.** In this problem, we will check the Lorentz invariance of the Dirac equation, and in the process determine the Lorentz transformation rule for Dirac spinors. Suppose that two coordinate systems are related by a Lorentz transformation  $x'^\mu = L^\mu_\nu x^\nu$ . The Dirac wavefunction  $\Psi'(x')$  as reported by an observer in the primed frame should be related to that in the

unprimed frame by  $\Psi'(x') = \Lambda\Psi(x)$  where  $\Lambda$  is a  $4 \times 4$  matrix. Now, the Dirac equation in the unprimed frame is  $(i\gamma^\mu \frac{\partial}{\partial x^\mu} - m)\Psi(x) = 0$ , and in the primed frame it is  $(i\gamma^\mu \frac{\partial}{\partial x'^\mu} - m)\Psi'(x') = 0$ .

(a) Show that these equations are consistent provided that  $\Lambda^{-1}\gamma^\rho L_\rho{}^\mu \Lambda = \gamma^\mu$ .

(b) Now suppose that  $L^\mu{}_\nu = \delta^\mu{}_\nu + \omega^\mu{}_\nu$  with  $\omega^\mu{}_\nu$  infinitesimal. Prove that the equation found in part (a) is satisfied if  $\Lambda = 1 + \omega^{\mu\nu}[\gamma_\mu, \gamma_\nu]/8$ .

**Exercise 27.3.** Consider a Dirac spinor wavefunction  $\Psi$  satisfying the Dirac equation (27.4.12) with electromagnetic potentials. Derive the local conservation of probability,

$$\frac{\partial \rho}{\partial t} = -\vec{\nabla} \cdot \vec{J}, \quad (27.6.2)$$

where  $\rho = \Psi^\dagger \Psi$  is the probability density and you will need to identify the probability current  $\vec{J}$ . [Hint: you may want to review the strategy used in the nonrelativistic case in section 5.6.]

**Exercise 27.4.** What is the next order ( $\alpha^6$ ) contribution to the hydrogen atom stationary state energies in eq. (27.5.38)? Assess its numerical impact for the  $n = 1$  and  $n = 2$  levels with  $Z = 1$ .

## 28 Feynman path integral approach

### 28.1 Propagators

Consider a quantum system with a complete set of commuting observables  $Q_a$ , where  $a$  is an index  $1, \dots, n$ . For example, for a single spinless particle moving in three-dimensional space, we have  $n = 3$ , and the  $Q_a$  with  $a = 1, 2, 3$  could be chosen to be the components of the position coordinate vector  $\vec{R}$ . Alternatively, we could choose the  $Q_a$  to be the components of the momentum vector  $\vec{P}$ . For simplicity, we will often suppress the index  $a$ , and let the whole set of CSCO eigenvalues be represented simply as the generic symbol  $q$ . We do this to make room for other subscript labels that will distinguish the values of the  $q$ 's at various times. Given the state ket  $|\psi\rangle$  of the system, we can then write a wavefunction

$$\psi(q, t) = \langle q | \psi(t) \rangle, \quad (28.1.1)$$

where  $|q\rangle$  are the CSCO eigenstates in which each of the observables  $Q_a$  is known to be equal to the corresponding eigenvalue  $q_a$ .

Recall, from section 3.4, that if the state of the system at time  $t_0$  was  $|\psi(t_0)\rangle$ , then the state at time  $t$  is

$$|\psi(t)\rangle = U(t, t_0) |\psi(t_0)\rangle, \quad (28.1.2)$$

where  $U(t, t_0)$  is the unitary time evolution operator. Using completeness of the  $Q$  eigenstates at some initial time  $t_0$ , the wavefunction at any later time  $t$  is

$$\psi(q, t) = \int dq_0 \langle q | U(t, t_0) | q_0 \rangle \langle q_0 | \psi(t_0) \rangle. \quad (28.1.3)$$

It is now useful to define the **propagator** (or **kernel**, or **transition amplitude**) as the function obtained as the relevant matrix element of the unitary time-evolution operator,

$$U(q, t; q_0, t_0) \equiv \langle q | U(t, t_0) | q_0 \rangle, \quad (28.1.4)$$

so that

$$\psi(q, t) = \int dq_0 U(q, t; q_0, t_0) \psi(q_0, t_0). \quad (28.1.5)$$

The utility of the propagator is that, given the wavefunction  $\psi(q_0, t_0)$  at an initial time  $t_0$ , the wavefunction at any other time  $t$  can be obtained by doing this integration over  $q_0$ .

The propagator is the solution to a differential equation, which follows from eq. (3.4.2),

$$i\hbar \frac{\partial}{\partial t} U(q, t; q_0, t_0) = i\hbar \langle q | \frac{\partial}{\partial t} U(t, t_0) | q_0 \rangle = \langle q | H U(t, t_0) | q_0 \rangle. \quad (28.1.6)$$



Consider the special case of a spinless particle moving in three dimensions in a potential  $V$ , so that the Hamiltonian is  $H = P^2/2m + V(\vec{R})$ . Interpreting  $q$  as the position  $\vec{r}$ , eq. (28.1.5) becomes

$$\psi(\vec{r}, t) = \int d^3\vec{r}_0 U(\vec{r}, t; \vec{r}_0, t_0) \psi(\vec{r}_0, t_0). \quad (28.1.7)$$

where eq. (28.1.6) tells us that

$$\left[ i\hbar \frac{\partial}{\partial t} + \frac{\hbar^2 \nabla^2}{2m} - V(\vec{r}) \right] U(\vec{r}, t; \vec{r}_0, t_0) = 0. \quad (28.1.8)$$

Since this is a first-order differential equation in  $t$ , in principle it can be solved given the boundary condition at  $t = t_0$ , which is  $U(\vec{r}, t_0; \vec{r}_0, t_0) = \delta^{(3)}(\vec{r} - \vec{r}_0)$ . In practice, this might be quite difficult, depending on the choice of potential.

Returning to the general case, suppose that the Hamiltonian does not depend explicitly on time, so that  $U(t, t_0) = e^{-i(t-t_0)H/\hbar}$ , from eq. (3.4.3). If we know all of the  $H$  eigenstates  $|n\rangle$ , with energies  $E_n$ , then using completeness we get the spectral decomposition

$$U(q, t; q_0, t_0) = \sum_n e^{-i(t-t_0)E_n/\hbar} \langle q|n\rangle \langle n|q_0\rangle = \sum_n e^{-i(t-t_0)E_n/\hbar} \psi_n(q) \psi_n(q_0)^*, \quad (28.1.9)$$

where  $\psi_n(q) = \langle q|n\rangle$  are the wavefunctions of the orthonormal Hamiltonian eigenstates with energies  $E_n$ . Thus, the propagator can be evaluated as a sum over products of energy eigenstate wavefunctions, weighted by phases that vary linearly with time.

As a simple example, consider the propagator for a free particle of mass  $m$ , moving in one dimension, with the position  $x$  playing the role of  $q$  in the preceding. We have already done the work for this case, in section 6.2, and the result of eq. (6.2.14) can be rewritten as

$$U(x, t; x_0, t_0) = \left[ \frac{m}{2\pi i\hbar(t-t_0)} \right]^{1/2} \exp \left[ \frac{im(x-x_0)^2}{2\hbar(t-t_0)} \right]. \quad (28.1.10)$$

Repeating the procedure for a spinless free particle moving in three dimensions gives

$$U(\vec{r}, t; \vec{r}_0, t_0) = \left[ \frac{m}{2\pi i\hbar(t-t_0)} \right]^{3/2} \exp \left[ \frac{im|\vec{r} - \vec{r}_0|^2}{2\hbar(t-t_0)} \right] \quad (28.1.11)$$

for the position-representation propagator.

For a somewhat more involved example, consider the one-dimensional harmonic oscillator, with the familiar Hamiltonian  $H = P^2/2m + m\omega^2 X^2/2$ . One way of writing the propagator is to use the spectral decomposition as in eq. (28.1.9), which says

$$U(x, t; x_0, t_0) = \sum_{n=0}^{\infty} \psi_n(x) \psi_n(x_0)^* e^{-i\omega(t-t_0)(n+1/2)}, \quad (28.1.12)$$

where  $\psi_n(x)$  are the stationary state wavefunctions of eq. (7.2.31), involving Hermite polynomials. It may therefore come as a surprise that the propagator can be written in a nice form that does not involve Hermite polynomials or an infinite sum,

$$U(x, t; x_0, t_0) = \left[ \frac{m\omega}{2\pi i \hbar \sin(\omega(t - t_0))} \right]^{1/2} \exp(iS_{\text{cl}}[x, t; x_0, t_0]/\hbar), \quad (28.1.13)$$

where, for general initial and final positions and times,

$$S_{\text{cl}}[x_f, t_f; x_i, t_i] = \frac{m\omega}{2 \sin(\omega(t_f - t_i))} \{ (x_f^2 + x_i^2) \cos(\omega(t_f - t_i)) - 2x_i x_f \}. \quad (28.1.14)$$

Remarkably, this is the action for the classical trajectory that starts at  $x_i$  at time  $t_i$ , and ends at  $x_f$  at time  $t_f$ . Recall that in general the **action** is defined by

$$S = \int_{t_i}^{t_f} dt L(x, \dot{x}), \quad (28.1.15)$$

where  $L(x, \dot{x})$  is the Lagrangian. In the present case,  $L = \frac{1}{2}m\dot{x}^2 - \frac{1}{2}m\omega^2 x^2$ , and you can check that eq. (28.1.14) follows from

$$x_{\text{cl}}(t) = \frac{x_f \sin(\omega(t - t_i)) - x_i \sin(\omega(t - t_f))}{\sin(\omega(t_f - t_i))}, \quad (28.1.16)$$

which is the classical trajectory that satisfies the equation of motion  $\ddot{x} = -\omega^2 x$  and the initial and final boundary conditions.

The quantum harmonic oscillator propagator claimed in eq. (28.1.13) is somewhat non-trivial to derive from scratch, but once it has been written down, it is not too hard to verify. This consists of checking that it obeys the first-order (in time) differential equation

$$\left( i\hbar \frac{\partial}{\partial t} + \frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} - \frac{1}{2}m\omega^2 x^2 \right) U(x, t; x_0, t_0) = 0, \quad (28.1.17)$$

as in eq. (28.1.8), and that it satisfies the correct boundary condition

$$U(x, t_0; x_0, t_0) = \delta(x - x_0), \quad (28.1.18)$$

which follows from taking the  $t \rightarrow t_0$  limit of eq. (28.1.13), with the help of eq. (2.2.21).

The propagator for the oscillator in eq. (28.1.13) has several features worthy of note. First, it is manifestly periodic in time, with the same period  $2\pi/\omega$  as the classical oscillator. Second, one recovers the free particle propagator of eq. (28.1.10) by taking the limit  $\omega \rightarrow 0$ . Finally, it is intriguing that the only dependence of the quantum propagator on the starting and ending positions  $x$  and  $x_0$  comes through the  $e^{iS/\hbar}$  factor for the classical trajectory that connects them. This is a suggestive clue for the developments in the next section.

## 28.2 Summing over paths

An alternative way of computing the propagator for a quantum system uses the **Feynman sum over paths** or **sum over histories**, developed by Richard P. Feynman in his 1942 PhD thesis. Let us write, very schematically for now,

$$U(q_f, t_f; q_i, t_i) \sim \sum_{\text{paths}} (\text{something}). \quad (28.2.1)$$

In this section, we will denote the initial and final CSCO eigenvalue variables and time by  $(q_i, t_i)$  and  $(q_f, t_f)$  respectively, in order to distinguish them from intermediate times  $t$  and positions  $q$  in the following discussion. The “paths” here are all trajectories  $q(t)$  starting from  $q_i$  at time  $t_i$  and ending with  $q_f$  at time  $t_f$ . In the case of a spinless particle in three dimensions, a path could be a trajectory given by a function  $\vec{r}(t)$ , but by a different choice of CSCO it could just as easily be instead a function  $\vec{p}(t)$  for the momentum vector of the particle. After choosing the CSCO eigenvalue variables, these paths are arbitrary, in the sense that *all* trajectories (not just those satisfying the classical equations of motion!) are included as long as they obey the initial and final boundary conditions.

The task before us now is to make eq. (28.2.1), including its mysterious summand, more precise. To motivate this form, first note that

$$U(q_f, t_f; q_i, t_i) = \langle q_f | U(t_f, t_1) U(t_1, t_i) | q_i \rangle = \int dq_1 \langle q_f | U(t_f, t_1) | q_1 \rangle \langle q_1 | U(t_1, t_i) | q_i \rangle, \quad (28.2.2)$$

where we have chosen an arbitrary time  $t_1$  between  $t_i$  and  $t_f$ , divided the unitary time evolution operator accordingly using eq. (3.4.30), and then used completeness of the eigenstates  $|q_1\rangle$ . This can be rewritten as a composition rule for propagators,

$$U(q_f, t_f; q_i, t_i) = \int dq_1 U(q_f, t_f; q_1, t_1) U(q_1, t_1; q_i, t_i). \quad (28.2.3)$$

In words, to propagate from time  $t_i$  to time  $t_f$ , we can integrate over all of the ways to propagate from  $t_i$  to  $t_1$ , and then from  $t_1$  to  $t_f$ , as indicated in Figure 28.2.1.

There is an intuitive analogy, **Feynman’s disappearing screen**, for the preceding mathematical derivation of eq. (28.2.3) and Figure 28.2.1. Consider particles whose vertical positions are described by  $q$ . The particles are emitted at a source point  $q_i$ , and detected at a point  $q_f$ , which can be varied by moving the detector. The probability of detection is proportional to the square of the complex propagator amplitude. Now if we place a large impenetrable screen between the source and the detector, then the amplitude will certainly vanish. But suppose that we then drill a small hole in the screen at a single point  $q_1$ . In that case, the only non-zero contribution corresponds to the particle first moving from  $q_i$  to  $q_1$ , and then from  $q_1$  to  $q_f$ , so




Figure 28.2.1: Diagrammatic representation of the composition law eq. (28.2.3). Each line between dots represents a propagator. The propagator  $U(q_f, t_f; q_i, t_i)$  is equal to the integral (or sum, if the allowed values are discrete) over all possible  $q_1$  of the product  $U(q_f, t_f; q_1, t_1)U(q_1, t_1; q_i, t_i)$ , for any choice of intermediate time  $t_1$ .

the propagator amplitude is the product of two separate amplitudes. If we drill a second hole at a different  $q_1$ , then the total amplitude will be the sum of two such product amplitudes, which interfere to give a diffraction pattern of maxima and minima as we vary  $q_f$ . Drilling more holes in the screen will just add more product amplitudes to the superposition, corresponding to more intermediate points  $q_1$ . In the limit that we have drilled through every point on the screen, it will have completely disappeared. So the amplitude for no screen at all must be proportional to the superposition of all amplitudes for all possible points  $q_1$ , as on the right side of eq. (28.2.3) and its pictorial version in Figure 28.2.1.<sup>†</sup>

We now extend the same idea to subdivide the time interval from  $t_i$  to  $t_f$  into  $N$  time steps instead of just two, with intermediate times  $t_1, t_2, \dots, t_{N-1}$ , as shown in Figure 28.2.2. At each time step, we integrate (or sum, if the allowed values are discrete) over all possible  $q_j$  for  $j = 1, 2, \dots, N-1$ , using completeness of the states  $|q_j\rangle$  at each step. The result is

$$U(q_f, t_f; q_i, t_i) = \int dq_1 \int dq_2 \cdots \int dq_{N-1} U(q_f, t_f; q_{N-1}, t_{N-1}) U(q_{N-1}, t_{N-1}; q_{N-2}, t_{N-2}) \cdots U(q_2, t_2; q_1, t_1) U(q_1, t_1; q_i, t_i). \quad (28.2.4)$$

Although this is not the ultimate expression of the sum over paths idea, it already illustrates the basic concept that we can find the propagator by adding up all possible ways to get from  $q_i$  at time  $t_i$  to  $q_f$  at time  $t_f$ .

Feynman's sum over paths can be thought of as the continuum limit of the discretized picture in Figure 28.2.2. He proposed that the propagator  $U(q_f, t_f; q_i, t_i)$  can be computed as a sum over all possible space-time paths of a phase determined by the classical action for that path, fulfilling an inspiring but less concrete suggestion made earlier by Dirac. The paths to be summed over are functions  $q(t)$  defined for  $t_i \leq t \leq t_f$  and constrained to obey the boundary conditions

$$q(t_i) = q_i, \quad q(t_f) = q_f. \quad (28.2.5)$$

<sup>†</sup>A slight imperfection in the simple pictorial version of the analogy is that the line containing  $q_1$  in Figure 28.2.1 represents a fixed time, while the vanishing screen instead occupies a fixed horizontal position.




Figure 28.2.2: One possible trajectory, or path,  $q(t)$  with initial boundary condition  $q_i$  at time  $t_i$  and final boundary condition  $q_f$  at time  $t_f$ . Time increases moving to the right in the figure, and is divided into equal slices, with variables  $q_1, q_2, \dots, q_{N-1}$  at intermediate times  $t_1, t_2, \dots, t_{N-1}$ . Each short line between dots represents a propagator. The path integral is obtained by taking the limit of  $N \rightarrow \infty$  time steps with infinitesimal time intervals  $\epsilon = t_j - t_{j-1} = (t_f - t_i)/N$ , and integrating over all  $q_1, q_2, \dots, q_{N-1}$ .

The action for a given path is

$$S[q(t)] = \int_{t_i}^{t_f} dt L(q, \dot{q}, t), \quad (28.2.6)$$

where  $L(q, \dot{q}, t)$  is the classical Lagrangian for the system, often the kinetic energy minus the potential energy. For example, for a spinless particle moving in three dimensions, the position vector  $\vec{r}(t)$  plays the role of  $q(t)$ , and the Lagrangian is

$$L(\vec{r}, \dot{\vec{r}}, t) = \frac{1}{2} m \dot{\vec{r}}^2 - V(\vec{r}). \quad (28.2.7)$$

The action is an example of a **functional**, an object that takes as its input a function [here the path  $q(t)$ ] and returns a number. Feynman therefore proposed that eq. (28.2.1) should be

$$U(q_f, t_f; q_i, t_i) \propto \sum_{\text{paths } q(t)} e^{iS[q(t)]/\hbar}, \quad (28.2.8)$$

following a suggestion by Dirac.

Because the number of paths is continuously infinite, it makes more sense to think of integrating over all paths rather than summing them. This suggests the arguably better names **Feynman path integral** or **functional integral**, and the notational change

$$\sum_{\text{paths } q(t)} e^{iS[q(t)]/\hbar} \rightarrow \int d[q(t)] e^{iS[q(t)]/\hbar}. \quad (28.2.9)$$

Just as ordinary definite integration takes a function and returns a number, a functional integral takes a functional (in this case, the action) and returns a number. The constant of proportionality implied in eq. (28.2.8) can then be absorbed into the normalization for the functional differential, denoted by  $d[q(t)]$ .

In summary, the following can now be taken as an alternative to the Schrödinger equation Postulate 6 of section 3.1:

**Postulate 6': Time evolution as a sum over paths.** For a system with eigenvalues  $\vec{q}$  for a complete set of commuting observables, the propagator is

$$U(\vec{q}_f, t_f; \vec{q}_i, t_i) = \int d[\vec{q}(t)] \exp(iS[\vec{q}(t)]/\hbar). \quad (28.2.10)$$

where  $S[\vec{q}(t)] = \int_{t_i}^{t_f} dt L(\vec{q}, \dot{\vec{q}})$  is the action functional, and the functional integral is over all paths satisfying  $\vec{q}(t_i) = \vec{q}_i$  and  $\vec{q}(t) = \vec{q}_f$ . Equivalently, the wavefunction satisfies

$$\psi(\vec{q}_f, t_f) = \int d[\vec{q}(t)] \exp(iS[\vec{q}(t)]/\hbar) \psi(\vec{q}_i, t_i), \quad (28.2.11)$$

but now the functional integral is over all paths satisfying  $\vec{q}(t_f) = \vec{q}_f$ , but with no constraint on  $\vec{q}(t_i)$ , since those initial values are being integrated over, as in eq. (28.1.5).

Note that while the Schrödinger version of Postulate 6 requires us to specify the Hamiltonian, the Feynman version Postulate 6' instead requires that the Lagrangian is the object to be specified to determine the dynamics of the system.

Although path integration has proved to be a very useful tool, especially in quantum field theory, it is not as straightforward to define as the Schrödinger differential equation. Physicists often simply ignore the associated problems of mathematical precision and rigor. One way to define the functional integral, inspired by the composition rule for the propagator as in eq. (28.2.4) and Figure 28.2.2, is to discretize the path into steps, integrate over the values of  $q$  on each step, and then take the limit in which the number of steps is infinitely large.

As a specific example, consider the case of a spinless particle moving in a potential  $V(\vec{r})$  in three dimensions, with the coordinates  $\vec{r}$  playing the role of  $\vec{q}$ . To compute the path integral, we partition the time interval  $(t_i, t_f)$  into  $N$  equal steps of duration

$$\epsilon = (t_f - t_i)/N, \quad (28.2.12)$$

so that

$$t_j = t_i + j\epsilon, \quad (j = 0, 1, 2, \dots, N), \quad (28.2.13)$$

with  $t_i = t_0$  and  $t_f = t_N$ . We then define the discretized version of a path from  $\vec{r}_i$  to  $\vec{r}_f$  by positions  $\vec{r}_j$  for each  $t_j$  for  $j = 1, \dots, N-1$ , with  $\vec{r}_0 = \vec{r}_i$  and  $\vec{r}_N = \vec{r}_f$ . The velocity of the particle in the interval between times  $t_{j-1}$  and  $t_j$  is  $\vec{v} = (\vec{r}_j - \vec{r}_{j-1})/\epsilon$ , so the kinetic energy is  $\frac{1}{2}mv^2 = \frac{1}{2}m(|\vec{r}_j - \vec{r}_{j-1}|/\epsilon)^2$ . Therefore, we can write the discretized version of the action for the path as

$$S[\text{path}] = S(\vec{r}_i, \vec{r}_1, \vec{r}_2, \dots, \vec{r}_{N-1}, \vec{r}_f) = \epsilon \sum_{j=1}^N \left[ \frac{m|\vec{r}_j - \vec{r}_{j-1}|^2}{2\epsilon^2} - V(\vec{r}_j) \right]. \quad (28.2.14)$$

In the limit  $\epsilon \rightarrow 0$  with  $N = (t_f - t_i)/\epsilon \rightarrow \infty$ , this will give the action for the continuous path. We now write eq. (28.2.10) as

$$\psi(\vec{r}_f, t_f) = \lim_{N \rightarrow \infty} \int \frac{d^3 \vec{r}_0}{a^3} \int \frac{d^3 \vec{r}_1}{a^3} \int \frac{d^3 \vec{r}_2}{a^3} \cdots \int \frac{d^3 \vec{r}_{N-1}}{a^3} \exp[iS[\text{path}]/\hbar] \psi(\vec{r}_0, t_0), \quad (28.2.15)$$

where  $a$  is a normalization factor, with units of [length], to be found soon. So, removing the integration over  $\vec{r}_0$  and the initial wavefunction  $\psi(\vec{r}_0, t_0)$ ,

$$U(\vec{r}_f, t_f; \vec{r}_i, t_i) = \lim_{N \rightarrow \infty} \frac{1}{a^3} \int \frac{d^3 \vec{r}_1}{a^3} \int \frac{d^3 \vec{r}_2}{a^3} \cdots \int \frac{d^3 \vec{r}_{N-1}}{a^3} \exp[iS[\text{path}]/\hbar], \quad (28.2.16)$$

in which it is important that there is one more factor of  $1/a^3$  than the remaining number of integrations, in accord with the fact that the propagator has dimension  $1/[\text{length}]^3$ . Equation (28.2.16) is one possible meaning of the notation

$$U(\vec{r}_f, t_f; \vec{r}_i, t_i) = \int d[\vec{r}(t)] \exp(iS[\vec{r}(t)]/\hbar) \quad (28.2.17)$$

for the path integral.

Let us now show that the preceding proposal is indeed equivalent to the Schrödinger equation. It is sufficient to consider only time evolution over a single infinitesimal time interval, since longer time intervals can be obtained using the composition law eq. (28.2.3). For simplicity, we write  $t_i = t_0 = t$ , and after one infinitesimal time step,  $t_f = t + \epsilon$ . Then, using the  $N = 1$  version of eq. (28.2.15), the wavefunction at the final time in terms of the initial wavefunction is

$$\psi(\vec{r}, t + \epsilon) = \int \frac{d^3 \vec{r}_0}{a^3} e^{iS(\vec{r}, \vec{r}_0)/\hbar} \psi(\vec{r}_0, t). \quad (28.2.18)$$

Expanding the left side in small  $\epsilon$  gives

$$\text{LS of (28.2.18)} = \psi(\vec{r}, t + \epsilon) = \psi(\vec{r}, t) + \epsilon \frac{\partial}{\partial t} \psi(\vec{r}, t) + \cdots. \quad (28.2.19)$$

The right side of eq. (28.2.18) is

$$\text{RS of (28.2.18)} = \int \frac{d^3 \vec{r}_0}{a^3} \exp\left(\frac{i}{\hbar} \left[ \frac{m|\vec{r} - \vec{r}_0|^2}{2\epsilon} - \epsilon V(\vec{r}) \right]\right) \psi(\vec{r}_0, t), \quad (28.2.20)$$

or, expanding the exponential part involving  $V(\vec{r})$  to order  $\epsilon$ , and then shifting the integration variable according to  $\vec{r}_0 \rightarrow \vec{r}_0 + \vec{r}$ ,

$$\text{RS of (28.2.18)} = \left[1 - i\frac{\epsilon}{\hbar} V(\vec{r})\right] \int \frac{d^3 \vec{r}_0}{a^3} e^{imr_0^2/2\hbar\epsilon} \psi(\vec{r} + \vec{r}_0, t). \quad (28.2.21)$$

Now, in the limit of small  $\epsilon$ , there will be almost complete cancellation from the rapidly varying phase inside the integral. The only region that contributes as  $\epsilon \rightarrow 0$  is where  $r_0^2 \approx 0$ , so that the phase factor  $e^{imr_0^2/2\hbar\epsilon}$  is close to 1. Therefore, we can use the expansion for small  $r_0$ ,

$$\psi(\vec{r} + \vec{r}_0, t) = \psi(\vec{r}, t) + \vec{r}_0 \cdot \vec{\nabla} \psi(\vec{r}, t) + \frac{1}{2} (\vec{r}_0 \cdot \vec{\nabla})^2 \psi(\vec{r}, t) + \cdots. \quad (28.2.22)$$

Inside the integration over  $\vec{r}_0$ , this can be simplified because the part odd under  $(x_0, y_0, z_0) \rightarrow -(x_0, y_0, z_0)$  will cancel. Thus, the linear term does not contribute at all, and using rectangular coordinates and the spherical symmetry of the integrand you can check that inside the integral the last term can be replaced according to  $(\vec{r}_0 \cdot \vec{\nabla})^2 \rightarrow \frac{1}{3}r_0^2 \nabla^2$ . So, we have

$$\int d^3\vec{r}_0 e^{imr_0^2/2\hbar\epsilon} \psi(\vec{r} + \vec{r}_0, t) = \psi(\vec{r}, t) \int d^3\vec{r}_0 e^{imr_0^2/2\hbar\epsilon} + \frac{1}{6} \nabla^2 \psi(\vec{r}, t) \int d^3\vec{r}_0 r_0^2 e^{imr_0^2/2\hbar\epsilon} + \dots, \quad (28.2.23)$$

in which the neglected terms have higher powers of  $r_0^2$ , and so can be checked to contribute only at higher order in  $\epsilon$ . Using

$$\int d^3\vec{r}_0 e^{imr_0^2/2\hbar\epsilon} = 4\pi \int_0^\infty dr_0 r_0^2 e^{imr_0^2/2\hbar\epsilon} = \left( \frac{2\pi i\hbar\epsilon}{m} \right)^{3/2}, \quad (28.2.24)$$

$$\int d^3\vec{r}_0 r_0^2 e^{imr_0^2/2\hbar\epsilon} = 4\pi \int_0^\infty dr_0 r_0^4 e^{imr_0^2/2\hbar\epsilon} = i \frac{3\hbar\epsilon}{m} \left( \frac{2\pi i\hbar\epsilon}{m} \right)^{3/2}, \quad (28.2.25)$$

and plugging into eq. (28.2.21) gives

$$\text{RS of (28.2.18)} = \frac{1}{a^3} \left( \frac{2\pi i\hbar\epsilon}{m} \right)^{3/2} \left[ 1 + i \frac{\epsilon}{\hbar} \left( \frac{\hbar^2 \nabla^2}{2m} - V(\vec{r}) \right) + \mathcal{O}(\epsilon^2) \right] \psi(\vec{r}, t). \quad (28.2.26)$$

Now compare eqs. (28.2.19) and (28.2.26). Matching at leading order in  $\epsilon$  just informs us that the normalization factor is

$$a = \left( \frac{2\pi i\hbar\epsilon}{m} \right)^{1/2}. \quad (28.2.27)$$

Then, matching the terms at first order in  $\epsilon$  gives

$$i\hbar \frac{\partial}{\partial t} \psi(\vec{r}, t) = \left[ -\frac{\hbar^2 \nabla^2}{2m} + V(\vec{r}) \right] \psi(\vec{r}, t), \quad (28.2.28)$$

which is the Schrödinger equation in the position wavefunction representation. This completes the demonstration that the Feynman sum over paths formulation is equivalent to the Schrödinger equation time evolution.

### 28.3 Evaluation of the path integral for the harmonic oscillator

In the previous section we demonstrated the equivalence of the path integral formulation and the Schrödinger equation, but this is not the same as actually computing a useful result directly in terms of the path integral. We will now show how to arrive at the propagator for the one-dimensional harmonic oscillator, already given in eq. (28.1.13), in the path integral approach.



Consider the propagator as a path integral, starting at time  $t = 0$  and ending at time  $t = T$ ,

$$U(x_f, T; x_i, 0) = \int d[x(t)] e^{iS[x(t)]/\hbar}. \quad (28.3.1)$$

In the case of the harmonic oscillator, we have a great advantage, that there is a unique classical trajectory, given by eq. (28.1.16) with  $t_f = T$  and  $t_i = 0$ . Accordingly, it will help us greatly to do a change of variables

$$x(t) = x_{\text{cl}}(t) + y(t), \quad (28.3.2)$$

where the new coordinate  $y(t)$  can be interpreted as the quantum fluctuation. It satisfies the boundary conditions

$$y(0) = y(T) = 0. \quad (28.3.3)$$

Now we can evaluate the action functional, with the result

$$S[x(t)] = S[x_{\text{cl}}(t)] + \frac{m}{2} \int_0^T dt (y^2 - \omega^2 y^2), \quad (28.3.4)$$

where the cross-terms involving both  $x_{\text{cl}}$  and  $y$  have conveniently canceled. To see this cancellation, note that those cross-terms are

$$m \int_0^T dt (\dot{x}_{\text{cl}} \dot{y} - \omega^2 x_{\text{cl}} y) = -m \int_0^T dt (\ddot{x}_{\text{cl}} + \omega^2 x_{\text{cl}}) y, \quad (28.3.5)$$

where we have integrated by parts, making use of the vanishing of the boundary terms from eq. (28.3.3). The last expression vanishes due to the equation of motion satisfied by  $x_{\text{cl}}$ .

Now, since  $S[x_{\text{cl}}(t)]$  is a constant with respect to the new functional integration variable  $y(t)$ , we can pull it out of the path integral, to get

$$U(x_f, T; x_i, 0) = I e^{iS[x_{\text{cl}}(t)]/\hbar}, \quad (28.3.6)$$

where it remains to evaluate the function integral over  $y(t)$ ,

$$I \equiv \int d[y(t)] \exp \left[ \frac{im}{2\hbar} \int_0^T dt (\dot{y}^2 - \omega^2 y^2) \right]. \quad (28.3.7)$$

We have already made good progress, since we have successfully obtained the correct  $e^{iS[x_{\text{cl}}(t)]/\hbar}$  factor in eq. (28.1.13). Indeed, from its definition, the remaining factor  $I$  clearly has no dependence on  $x_f$  or  $x_i$ . Our remaining task is to show that, in accord with eq. (28.1.13),

$$I = \left( \frac{m\omega}{2\pi i \hbar \sin(\omega T)} \right)^{1/2}. \quad (28.3.8)$$

We will do the path integral to get  $I$  in two different ways.

First, we use the discretization of the same type used in eq. (28.2.16),

$$I = \lim_{N \rightarrow \infty} \frac{1}{a} \left( \prod_{j=1}^{N-1} \int_{-\infty}^{\infty} \frac{dy_j}{a} \right) \exp \left( \frac{im}{2\hbar\epsilon} \sum_{k=1}^N [(y_k - y_{k-1})^2 - \epsilon^2 \omega^2 y_k^2] \right), \quad (28.3.9)$$

where  $\epsilon = T/N$ , and we have already obtained the normalization  $a$  in eq. (28.2.27). Then, letting  $z_j = y_j/a$  and assembling  $z_1, z_2, \dots, z_{N-1}$  into an  $(N-1)$ -dimensional vector  $\vec{z}$ , we have

$$I = \lim_{N \rightarrow \infty} \frac{1}{a} \int d^{N-1} \vec{z} \exp(-\pi \vec{z}^T B_N \vec{z}) \quad (28.3.10)$$

where  $B_N$  is an  $(N-1) \times (N-1)$  matrix, which takes the form

$$B_N = \begin{pmatrix} 2-\delta & -1 & 0 & 0 & \cdots & 0 \\ -1 & 2-\delta & -1 & 0 & \cdots & 0 \\ 0 & -1 & 2-\delta & -1 & \cdots & 0 \\ 0 & 0 & -1 & 2-\delta & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & -1 \\ 0 & 0 & 0 & 0 & -1 & 2-\delta \end{pmatrix}, \quad (28.3.11)$$

where for convenience we have defined

$$\delta \equiv \epsilon^2 \omega^2 = \omega^2 T^2 / N^2. \quad (28.3.12)$$

Now, since  $B$  is a real symmetric matrix, we can do a rotation in the  $(N-1)$ -dimensional space to new coordinates  $\vec{u} = O\vec{z}$ , where  $O$  is an orthogonal matrix, in such a way that the rotated matrix  $\tilde{B}_N = O B_N O^T$  is diagonal. Since a real symmetric matrix is just a special kind of Hermitian matrix, and an orthogonal matrix is just a special kind of unitary matrix, Theorems 2.6.7 and 2.6.9 apply to tell us that the diagonal entries of  $\tilde{B}_N$  are the same as the eigenvalues of  $B_N$ . Calling those eigenvalues  $\lambda_j$ ,

$$I = \lim_{N \rightarrow \infty} \frac{1}{a} \prod_{j=1}^{N-1} \int_{-\infty}^{\infty} du_j e^{-\pi \lambda_j u_j^2}. \quad (28.3.13)$$

Now each of the  $u_j$  integrals can be done separately, using  $\int_{-\infty}^{\infty} du e^{-\pi \lambda u^2} = \lambda^{-1/2}$ , giving

$$I = \lim_{N \rightarrow \infty} \frac{1}{a} \prod_{j=1}^{N-1} (\lambda_j)^{-1/2} = \lim_{N \rightarrow \infty} \frac{1}{a} (\det B_N)^{-1/2} \quad (28.3.14)$$

where we have used the fact from matrix algebra that the determinant of  $B_N$  is equal to the product of its eigenvalues.

Let us now find the determinant of the  $(N-1) \times (N-1)$  matrix  $B_N$ , temporarily taking  $\delta$  to be general even though in our case it depends on  $N$  through eq. (28.3.12). We have

$$\det B_2 = 2 - \delta, \quad (28.3.15)$$

$$\det B_3 = (2 - \delta)^2 - 1, \quad (28.3.16)$$

and for larger  $N$  we can evaluate  $\det B_N$  in terms of its minors for the two non-zero entries in the first row. Conveniently, these involve determinants of versions of the same matrix with dimension smaller by 1 and 2. This gives the recurrence relation

$$\det B_N = (2 - \delta)\det B_{N-1} - \det B_{N-2}, \quad (N = 4, 5, 6, \dots). \quad (28.3.17)$$

It is possible to solve this recurrence relation systematically, but we will take the shortcut of writing down the answer and checking that it works. To do so, we define numbers  $\beta_{1,2}$  by

$$\beta_1 + \beta_2 = 2 - \delta, \quad \beta_1 \beta_2 = 1. \quad (28.3.18)$$

Then we claim that

$$\det B_N = \frac{\beta_1^N - \beta_2^N}{\beta_1 - \beta_2}. \quad (28.3.19)$$

This is easily checked to work for  $N = 2$  and 3, and then for the recurrence relation eq. (28.3.17), by direct substitution.

We are now ready to take the large  $N$  limit. First, solving eq. (28.3.18), we find that

$$\beta_{1,2} = 1 - \frac{\delta}{2} \pm i\sqrt{\delta - \delta^2/4} = 1 \pm i\omega T/N + \dots, \quad (28.3.20)$$

where the ellipses involves terms of higher order in  $1/N$ . Therefore, for large  $N$ ,

$$\det B_N = \frac{(1 + i\omega T/N)^N - (1 - i\omega T/N)^N}{2i\omega T/N} = \frac{e^{i\omega T} - e^{-i\omega T}}{2i\omega T/N} = \frac{N \sin(\omega T)}{\omega T}, \quad (28.3.21)$$

where the definition of the exponential function,  $e^x = \lim_{N \rightarrow \infty} (1 + x/N)^N$ , was used to get the second equality. Using this in eq. (28.3.14) along with  $a = (2\pi i \hbar T/mN)^{1/2}$ , we find

$$I = \lim_{N \rightarrow \infty} \left( \frac{mN}{2\pi i \hbar T} \right)^{1/2} \left( \frac{\omega T}{N \sin(\omega T)} \right)^{1/2} = \left( \frac{m\omega}{2\pi i \hbar \sin(\omega T)} \right)^{1/2}, \quad (28.3.22)$$

which agrees with the result found before.

Greedy for more insight, let us now consider a second way to calculate  $I$ , using a different realization of the path integral. Instead of enumerating the possible paths by discretizing in

time, we write the quantum fluctuation as a general Fourier series expansion of terms that satisfy the boundary conditions  $y(0) = y(T) = 0$ ,

$$y(t) = \sum_{n=1}^{\infty} y_n \sin(n\pi t/T). \quad (28.3.23)$$

We then interpret the path integral in eq. (28.3.7) as an integration over all of the Fourier coefficients  $y_n$ ,

$$I = \frac{1}{c} \left( \prod_{n=1}^{\infty} \int_{-\infty}^{\infty} \frac{dy_n}{b_n} \right) \exp \left[ \frac{im}{2\hbar} \int_0^T dt ( \dot{y}^2 - \omega^2 y^2 ) \right] \quad (28.3.24)$$

where we have introduced normalization factors  $c$  and  $b_n$  with dimensions of [length]. Now the action integral over  $t$  in the exponent can be easily computed, yielding

$$I = \frac{1}{c} \left( \prod_{n=1}^{\infty} \int_{-\infty}^{\infty} \frac{dy_n}{b_n} \right) \exp \left[ - \sum_{n=1}^{\infty} \frac{imT}{4\hbar} (\omega^2 - n^2\pi^2/T^2) y_n^2 \right]. \quad (28.3.25)$$

This nicely separates into one-dimensional integrals over the  $y_n$ , as

$$I = \frac{1}{c} \prod_{n=1}^{\infty} \left( \int_{-\infty}^{\infty} \frac{dy_n}{b_n} \exp \left[ - \frac{imT}{4\hbar} (\omega^2 - n^2\pi^2/T^2) y_n^2 \right] \right). \quad (28.3.26)$$

Doing the individual Gaussian integrals,

$$I = \frac{1}{c} \prod_{n=1}^{\infty} \frac{1}{b_n} \left( \frac{4\pi i\hbar}{mT(n^2\pi^2/T^2 - \omega^2)} \right)^{1/2}. \quad (28.3.27)$$

To make further progress, we make use of a famous formula due to Euler,

$$\prod_{n=1}^{\infty} \left( 1 - \frac{x^2}{n^2\pi^2} \right) = \frac{\sin(x)}{x}, \quad (28.3.28)$$

with  $x = \omega T$  in our present case, to obtain

$$I = \frac{1}{c} \left( \frac{\omega T}{\sin(\omega T)} \right)^{1/2} \prod_{n=1}^{\infty} \frac{1}{b_n} \left( \frac{4i\hbar T}{n^2\pi m} \right)^{1/2} \quad (28.3.29)$$

Comparing with eq. (28.3.22), we see that we have correctly obtained the  $\omega$  dependence, and that the whole formula for  $I$  is correct provided that the normalization factors are

$$b_n = \left( \frac{4i\hbar T}{n^2\pi m} \right)^{1/2}, \quad c = \left( \frac{2\pi i\hbar T}{m} \right)^{1/2}. \quad (28.3.30)$$

The advantage of this Fourier expansion method for the harmonic oscillator path integral is that the separation into individual integrals was automatic. The disadvantage is that the  $\omega$ -independent normalization factors  $1/c$  and  $1/b_n$  associated with the path integral had to be

obtained by comparison to the known result for  $I$ . In the discretized version of the path integral, the normalization factor  $1/a$  was known already from the case of infinitesimal  $T$  in section 28.2.

The case of a free particle can be obtained from the  $\omega \rightarrow 0$  limit. In the preceding, we were able to make use of a very special property of the free particle and the harmonic oscillator, namely that the action is quadratic in the configuration variable(s) and its time derivatives, leading to Gaussian integrations in either eq. (28.3.13) or eq. (28.3.26), which we were therefore able to do analytically. Another simple application is given in the next section. For other problems in nonrelativistic quantum mechanics, the path integral method is only rarely efficient compared to methods based on the Schrödinger equation. However, the path integral approach is still extremely valuable, for in addition to the beautiful conceptual understanding it brings, it turns out to provide a practically useful way of systematically organizing certain approximation methods. In quantum field theories, this includes both Feynman rules for perturbative calculations, and stochastic methods for non-perturbative calculations, topics beyond the scope of this book.

## 28.4 The Ehrenberg–Siday and Aharonov–Bohm effect

The classical trajectory of a charged particle in an electromagnetic field is determined uniquely by its initial conditions and the electromagnetic fields  $\vec{E}$  and  $\vec{B}$  that it encounters locally. In quantum mechanics, things are different. The Hamiltonian in the Schrödinger equation, and the Lagrangian in the path integral approach, depend on the gauge-dependent potentials  $\Phi$  and  $\vec{A}$ . These potentials contain global information that is gauge invariant, but not accessible to an observer on a classical trajectory making only local measurements of  $\vec{E}$  and  $\vec{B}$ . As a result, there can be observable consequences of electromagnetic fields even if the particle is absolutely forbidden to enter the region where  $\vec{E}$  and  $\vec{B}$  are nonzero!

The simplest example of this purely quantum phenomenon is most commonly known as the Aharonov–Bohm effect, after a 1959 paper by Yakir Aharonov and David Bohm, although it was actually proposed a decade earlier by Werner Ehrenberg and Raymond Siday. Consider a spinless charged particle with coordinates  $\vec{r}(t)$  subject to a static but non-uniform magnetic field  $\vec{B} = \vec{\nabla} \times \vec{A}$ . By specializing eq. (4.3.11), the Lagrangian is

$$L = \frac{1}{2}m \left( \frac{d\vec{r}}{dt} \right)^2 + \frac{q}{c} \frac{d\vec{r}}{dt} \cdot \vec{A}(\vec{r}). \quad (28.4.1)$$

The path integral expression for the propagator,

$$U(\vec{r}_f, t_f; \vec{r}_i, t_i) = \int d[\vec{r}(t)] \exp \left( \frac{i}{\hbar} S_{\text{path}}[\vec{r}(t)] \right), \quad (28.4.2)$$




Figure 28.4.1: The dependence of the difference in actions for two alternative paths for a charged spinless particle on a static magnetic field is proportional to the magnetic flux  $\Phi_m$  through an area subtended by the two paths.

therefore naturally divides into two parts, according to

$$S_{\text{path}} = S_{\text{path}}^{\vec{A}=0} + \Delta S_{\text{path}} \quad (28.4.3)$$

where the first term is the free-particle action, and the remaining part can be rewritten as a line integral along the path specified from  $\vec{r}_i$  to  $\vec{r}_f$  by  $\vec{r}(t)$ ,

$$\Delta S_{\text{path}} = \frac{q}{c} \int_{t_i}^{t_f} dt \frac{d\vec{r}}{dt} \cdot \vec{A}(\vec{r}) = \frac{q}{c} \int_{\vec{r}_i}^{\vec{r}_f} d\vec{r} \cdot \vec{A}(\vec{r}). \quad (28.4.4)$$

Thus each path contributes with a phase whose dependence on the magnetic field is proportional to the line integral of  $\vec{A}$  along the path.

When evaluating the probability of detecting the particle at  $\vec{r}_f$  at time  $t_f$ , we need the relative phases between different paths. Consider any two paths contributing to the propagator, call them “path 1” and “path 2”. (See Figure 28.4.1.) Then the dependence on  $\vec{A}$  of the relative phase between their contributions to the propagator is

$$\Delta\varphi \equiv (\Delta S_{\text{path 1}} - \Delta S_{\text{path 2}})/\hbar = \frac{q}{\hbar c} \left( \int_{\vec{r}_i}^{\vec{r}_f} d\vec{r} \cdot \vec{A}(\vec{r}) - \int_{\vec{r}_i}^{\vec{r}_f} d\vec{r} \cdot \vec{A}(\vec{r}) \right). \quad (28.4.5)$$

Using the minus sign to reverse the direction of the line integral over path 2, we can combine the two paths into a single closed path  $P$ . Then applying Stokes’ Theorem gives the key result

$$\Delta\varphi = \frac{q}{\hbar c} \oint_P d\vec{r} \cdot \vec{A} = \frac{q}{\hbar c} \int_{\text{area}} d\vec{a} \cdot (\vec{\nabla} \times \vec{A}) = \frac{q}{\hbar c} \int_{\text{area}} d\vec{a} \cdot \vec{B} = \frac{q\Phi_m}{\hbar c}. \quad (28.4.6)$$

Here  $\Phi_m$  is the magnetic field flux through a surface area subtending the two paths, obtained following the right-hand rule. As usual when applying Stokes’ Theorem, the surface is not unique; any such area subtending the two paths must<sup>†</sup> give the same result.

A two-slit interference version of the **Ehrenberg–Siday–Aharonov–Bohm effect** is depicted in Figure 28.4.2. A source of electrons is separated from a flat detection screen by a

<sup>†</sup>This relies on the standard assumption that magnetic monopoles do not exist, so that the magnetic flux through any closed surface is 0. For the possibility that magnetic monopoles do exist, see section 28.5.




Figure 28.4.2: The Ehrenberg–Siday–Aharonov–Bohm effect for a two-slit interference experiment for electrons. The shaded circle represents the cross-section of a solenoid, carrying a magnetic field out of the plane. The presence of the magnetic field in the solenoid affects the interference pattern visible on the detection screen, even though the electrons are absolutely forbidden to enter the region with non-zero  $\vec{B}$ .

diffraction screen with two small holes. Behind the diffraction screen is a solenoid (or perhaps a permanent magnet), which carries in its interior a magnetic field  $\vec{B}$  coming out of the plane of the page. The solenoid is tightly wound and long enough so that the magnetic field  $\vec{B}$  outside of it can be neglected, and is shielded so that electrons cannot enter it. Therefore, classically the presence of the magnetic field inside the solenoid can have no effect on the electrons.

In quantum mechanical reality, it is important that the vector potential outside of the solenoid is necessarily non-zero. For example, in terms of cylindrical coordinates  $(r, \phi, z)$  coaxial with a solenoid of radius  $R$ , the magnetic field has the form

$$\vec{B} = \begin{cases} B\hat{z} & (\text{for } r < R), \\ 0 & (\text{for } r > R), \end{cases} \quad (28.4.7)$$

and the associated vector potential can be chosen to be

$$\vec{A} = \begin{cases} (Br/2)\hat{\phi} & (\text{for } r \leq R), \\ (BR^2/2r)\hat{\phi} & (\text{for } r \geq R). \end{cases} \quad (28.4.8)$$

The resulting magnetic flux for any closed path encircling the solenoid once is  $\Phi_m = \pi R^2 B$ . Accordingly, the phase difference, for electrons going on any path below the solenoid compared to those going on any path above the solenoid, due to the existence of the magnetic field, is, from eq. (28.4.6) with  $q = -e$ ,

$$\Delta\varphi = -2\pi\Phi_m/\Phi_0, \quad (28.4.9)$$

where we have defined the relevant quantum unit of magnetic flux,<sup>‡</sup>

$$\Phi_0 = 2\pi\hbar c/e = 4.13567 \times 10^{-7} \text{ gauss} \cdot \text{cm}^2. \quad (28.4.10)$$

To observe the effect, start with the solenoid turned off, and consider a reference point at the center of the detection screen where the interference is maximally constructive, with paths from

---

<sup>‡</sup>This same quantity also appeared in eq. (9.3.45), in the discussion of Landau levels.

both holes contributing equally to the peak intensity  $I_0$ . Now when the magnetic field is turned on, the intensity at that point will be changed to

$$\left| \frac{1}{2} + \frac{1}{2} e^{i\Delta\varphi} \right|^2 I_0 = \frac{1}{4} |1 + e^{-2\pi i \Phi_m / \Phi_0}|^2 I_0, \quad (28.4.11)$$

where the two terms in the first expression correspond to the contributions from all paths through the top hole and all paths through the bottom hole, respectively. So, if the magnetic field strength  $B$  is adjusted so that  $\Phi_m = (n+1/2)\Phi_0$  where  $n$  is any integer, the interference will become completely destructive, and the intensity will vanish at the reference point. If  $\Phi_m = n\Phi_0$ , the interference will be maximally constructive again. This effect has been confirmed in many increasingly sophisticated experiments.

One can interpret the effect as illustrating that the vector potential  $\vec{A}$  contains genuine physical information that is not apparent through local interaction of the charge with the field  $\vec{B}$ . This is the meaning of the first term in  $\Delta\varphi$  in eq. (28.4.6). However, the effect can also be described entirely in terms of the non-trivial magnetic flux,  $\Phi_m$ , in the last expression in eq. (28.4.6), which depends only on the gauge-invariant magnetic field  $\vec{B}$ . This dependence of interference on the flux is a topological one, in the sense that it occurs because the allowed paths of the charged particle cannot be deformed into each other without crossing into the region with nonzero  $\vec{B}$ . In the path integral approach, we understand this in terms of the particle's ability (in fact, obligation) to coherently sample phases from all available paths, even though none of those paths actually encounters the non-zero  $\vec{B}$  field.

## 28.5 Dirac quantization condition for magnetic monopoles

The Ehrenberg–Siday–Aharonov–Bohm effect is an experimentally well-established phenomenon. The key result leading to it, eq. (28.4.6), can also be applied to learn something quite non-trivial about a more speculative hypothesis, the possibility of magnetic monopoles. Dirac showed that the extension of Maxwell's electrodynamics to include magnetic monopoles is only consistent with quantum mechanics if both magnetic and electric charges are quantized. There are several different ways to derive Dirac's result. In this section we will do it using eq. (28.4.6).

The standard assumption that magnetic monopoles do not exist is one of the four Maxwell equations,  $\vec{\nabla} \cdot \vec{B} = 0$ . To include magnetic monopoles in the theory, one generalizes this to

$$\vec{\nabla} \cdot \vec{B} = 4\pi\rho_m, \quad (28.5.1)$$

where  $\rho_m$  is the density of magnetic charge. A point monopole with magnetic charge  $q_m$ , fixed at the origin, would then give rise to a magnetic field

$$\vec{B} = q_m \hat{r} / r^2 \quad (28.5.2)$$






Figure 28.5.1: A monopole of magnetic charge  $q_m$  is fixed at the origin, and a particle with electric charge  $q$  is restricted to a circle of fixed radius around it. As shown in the left panel, there are contributions to the propagator path integral from trajectories with net counterclockwise direction (paths 1) and net clockwise direction (paths 2). The presence of the monopole affects the relative phase between these classes of paths through  $\Delta\varphi$ , which can be obtained from the first expression in eq. (28.4.6) as a line integral of the vector potential around the closed circle. Alternatively, it can be obtained from the magnetic flux through either the upper hemisphere or the lower hemisphere, as depicted in the right panel.

in spherical coordinates. Equations (28.5.1) and (28.5.2) are the hypothetical analogs of, respectively, Gauss' Law  $\vec{\nabla} \cdot \vec{E} = 4\pi\rho$  and  $\vec{E} = q\hat{r}/r^2$ , for ordinary electric charge density  $\rho$  and a point electric charge  $q$ . This similarity between electric and magnetic charges is one reason to take magnetic monopoles seriously as a hypothesis. Another is that they can be shown to be necessarily present in some extensions of the Standard Model of particle physics that may be motivated for other reasons, notably Grand Unified Theories. Although magnetic monopoles have not been observed despite extensive searches, it is possible that this is because they are extremely rare in the present universe. For example, they could be too heavy to produce with collision energies available today, and the cosmological inflation that is believed (for other reasons) to have occurred in the early universe would have diluted the density of them left over from the Big Bang to a level that makes it extremely unlikely for us to encounter one.

Suppose that there is a magnetic monopole  $q_m$  at rest at the origin, with  $\vec{B}$  as given in eq. (28.5.2). Consider the quantum mechanics of an ordinary particle with charge  $q$ , which is assumed to be restricted to a circle of radius  $R$  in the  $xy$  plane with its center at the origin, as shown in the left panel of Figure 28.5.1. The propagator for the particle to start at some point  $\vec{r}_i$  at time  $t_i$  and end up at  $\vec{r}_f$  at time  $t_f$  can be written as a sum over contributions from paths that wind around the circle in different ways. Thus we can write

$$U(\vec{r}_i, t_i; \vec{r}_f, t_f) = \int_{\text{paths 1}} d[\vec{r}(t)] \exp\left(\frac{i}{\hbar} S[\vec{r}(t)]\right) + \int_{\text{paths 2}} d[\vec{r}(t)] \exp\left(\frac{i}{\hbar} S[\vec{r}(t)]\right). \quad (28.5.3)$$

The “paths 1” contribution is from paths that have a net counterclockwise direction in the figure,

while “paths 2” have a net clockwise direction; of course, these paths  $\vec{r}(t)$  need not be monotonic in direction around the circle. (We neglect contributions from paths that wind around the circle more than once. This is not essential to the following argument.) The propagator can now be written in the form

$$U(\vec{r}_i, t_i; \vec{r}_f, t_f) = e^{i\alpha} (e^{i\Delta\varphi} c_1 + c_2), \quad (28.5.4)$$

where  $c_1$  and  $c_2$  are the contributions from paths 1 and paths 2 that would be present if the magnetic monopole were absent,  $e^{i\alpha}$  is a common phase factor due to the magnetic monopole, and  $e^{i\Delta\varphi}$  is the relative phase between the two classes of paths due to the magnetic monopole, as given in eq. (28.4.6). If the particle was known to be at  $\vec{r}_i$  at time  $t_i$ , then the probability that it will be found at  $\vec{r}_f$  at time  $t_f$  is

$$\mathcal{P} = |e^{i\Delta\varphi} c_1 + c_2|^2. \quad (28.5.5)$$

Since the initial and final boundary conditions are arbitrary, and other potentials felt by the charge  $q$  are not specified,  $c_1$  and  $c_2$  can be considered arbitrary complex numbers. Now  $\Delta\varphi$  can be evaluated in two distinct ways, either of which leads to the Dirac quantization condition.

First, one can evaluate  $\Delta\varphi$  using the first equality in eq. (28.4.6), the line integral of  $\vec{A}$  around the full circle in the  $xy$  plane at radius  $R$  from the origin. However, there is a problem associated with the vector potential, because the equation  $\vec{\nabla} \times \vec{A} = \vec{B}$  assumes no magnetic monopoles, since the vector calculus identity  $\vec{\nabla} \cdot (\vec{\nabla} \times \vec{A}) = 0$  implies  $\vec{\nabla} \cdot \vec{B} = 0$ . This shows that it is impossible to write down a consistent  $\vec{A}$  in any volume that includes the origin. Fortunately, this difficulty can be evaded by recognizing that to compute  $\Delta\varphi$ , we only need  $\vec{A}$  on the path arcs at  $r = R$  with  $\theta = \pi/2$ . Accordingly, working in spherical coordinate, we can take either

$$\vec{A}_I = \frac{q_m}{r} \left( \frac{\sin \theta}{1 + \cos \theta} \right) \hat{\phi}, \quad \text{or} \quad \vec{A}_{II} = -\frac{q_m}{r} \left( \frac{\sin \theta}{1 - \cos \theta} \right) \hat{\phi}. \quad (28.5.6)$$

These both give  $\vec{\nabla} \times \vec{A} = q_m \hat{r}/r^2$ , except at the origin, and on the negative  $z$  axis for  $\vec{A}_I$  and on the positive  $z$  axis for  $\vec{A}_{II}$ , where the respective denominators vanish. Since we only need  $r = R$  and  $\theta = \pi/2$  to evaluate the line integral,  $\vec{A}_I$  and  $\vec{A}_{II}$  are both equally valid for our purposes. However, they give two different results for  $\Delta\varphi$ ,

$$\Delta\varphi_I = -\Delta\varphi_{II} = \frac{q}{\hbar c} \int_0^{2\pi} R d\phi \frac{q_m}{R} = \frac{2\pi q q_m}{\hbar c}. \quad (28.5.7)$$

To avoid an unacceptable inconsistency in the physical prediction for the probability  $|e^{i\Delta\varphi} c_1 + c_2|^2$  following from the two equally valid choices for  $\vec{A}$ , we must have

$$e^{i\Delta\varphi_I} = e^{i\Delta\varphi_{II}}, \quad (28.5.8)$$

since  $c_1$  and  $c_2$  are arbitrary complex numbers. This is equivalently  $4\pi qq_m/\hbar c = 2\pi n$  for some integer  $n$ , or simply

$$\frac{2qq_m}{\hbar c} = n, \quad (n = \text{integer}). \quad (28.5.9)$$

This is the **Dirac quantization condition for magnetic monopole charges**.

The second (alternative) way to evaluate  $\Delta\varphi$  avoids any mention of the vector potential by using the last equality of eq. (28.4.6). So we have

$$\Delta\varphi = \frac{qq_m}{\hbar c} \int_{\text{area}} d\vec{a} \cdot \frac{\hat{r}}{r^2} = \frac{q\Phi_m}{\hbar c}. \quad (28.5.10)$$

Here we must confront the issue that there are actually two equally good choices of area subtended by the circle in the  $xy$  plane, one above the origin and one below. For simplicity of computation, it is convenient to take these to be the upper and lower hemispheres of radius  $R$ , labeled I and II in the second panel of Figure 28.5.1. The directions of their area elements point outward and inward, respectively, as dictated by the orientation of the closed circular path and the right-hand rule in Stokes' Theorem. Therefore, the magnetic fluxes through them are opposite in sign, and we have the two equally valid choices

$$\Phi_m^{(\text{I})} = -\Phi_m^{(\text{II})} = (2\pi R^2)(q_m/R^2) = 2\pi q_m. \quad (28.5.11)$$

Using these two choices in eq. (28.5.10) immediately gives exactly the same ambiguity as in eq. (28.5.7), with the same resolution: the Dirac quantization condition eq. (28.5.9) must hold.

Since we can pick  $q$  to be the proton charge  $e$  or the electron charge  $-e$ , we learn from Dirac's result that magnetic charges  $q_m$  must be quantized in integer multiples of a fundamental unit

$$q_m^{\text{Dirac}} = \frac{\hbar c}{2e} = 3.29106 \times 10^{-16} \text{ Tesla} \cdot \text{meters}^2. \quad (28.5.12)$$

Conversely, if even one magnetic monopole with charge  $q_m^{\text{Dirac}} = \hbar c/2e$  exists, anywhere in the universe, then electric charges must be integer multiples of  $e$ . Thus magnetic monopoles, if they exist, would give a compelling reason for the observed quantization of electric charge.

There is a subtlety here, because quarks and antiquarks exist with charges that are integer multiples of  $e/3$ . Therefore, one might naively think that the fundamental quantum of magnetic charge should actually be 3 times  $q_m^{\text{Dirac}}$ . However, quarks are confined inside hadrons by quantum chromodynamics (QCD), also known as the strong force. The QCD gauge potentials also contribute to the path integral, and it turns out (although showing it in detail is well beyond our scope here) that one can still have magnetic monopoles with charge  $q_m^{\text{Dirac}}$ , provided that they also carry magnetic-QCD charges, and are likewise confined. Thus Dirac's  $q_m^{\text{Dirac}}$  is the true quantum of magnetic charge, even though he did not know about quarks or QCD.




Figure 28.6.1: A path  $q(t)$  (solid line), and a nearby path  $q(t) + \delta q(t)$  (dashed line), subject to the boundary conditions  $q(t_i) = q_i$  and  $q(t_f) = q_f$  and  $\delta q(t_i) = \delta q(t_f) = 0$ . The path is a stationary path if the change in the action  $\delta S$  vanishes for every infinitesimal change in path  $\delta q(t)$ .

## 28.6 Classical limit from the sum over paths

In this section, we will show how the equations of motion for classical physics can be derived as an appropriate limit of quantum mechanics in the path integral formulation.

Consider a path  $q(t)$  subject to the boundary conditions  $q(t_i) = q_i$  and  $q(t_f) = q_f$ , and suppose we make a small change  $\delta q(t)$  to it,

$$q(t) \rightarrow q(t) + \delta q(t), \quad (28.6.1)$$

as illustrated in Figure 28.6.1. This will result in a change in the action, which we write as

$$S[q(t)] \rightarrow S[q(t)] + \delta S. \quad (28.6.2)$$

Because we must maintain the boundary conditions at  $t = t_i$  and  $t = t_f$  when computing the propagator, we must also require

$$\delta q(t_i) = \delta q(t_f) = 0. \quad (28.6.3)$$

In general, if  $\delta S$  is non-zero, then the integrand of the path integral will acquire an additional phase  $e^{i\delta S/\hbar}$  compared to the result from the original path. If we take  $\hbar \rightarrow 0$ , which we associate with the classical limit, this additional phase varies rapidly as  $\delta S$  changes, leading to very efficient cancellation between the contributions for nearby paths.

Therefore, in the classical limit, the propagator is dominated by **stationary paths**, defined to be those for which  $\delta S = 0$  for every  $\delta q(t)$  that satisfies the boundary conditions eq. (28.6.3). The point is that those are the only paths that do not suffer efficient cancellations from the neighboring paths, with cancellation becoming perfectly efficient in the  $\hbar \rightarrow 0$  limit. So, we can write the condition for a stationary path  $q(t)$  as the vanishing of the **functional derivative**,

$$\frac{\delta S[q(t)]}{\delta q(t)} \equiv \lim_{\delta q(t) \rightarrow 0} \frac{S[q(t) + \delta q(t)] - S[q(t)]}{\delta q(t)} = 0. \quad (28.6.4)$$

In quantum mechanics, the path integral samples all paths, but in the classical limit  $\hbar \rightarrow 0$ , only stationary paths contribute non-negligibly.

Suppose we are given a Lagrangian  $L(q, \dot{q}, t)$  for a system, so that

$$S[q(t)] = \int_{t_i}^{t_f} dt L(q, \dot{q}, t). \quad (28.6.5)$$

The change in the action due to the change in path  $\delta q(t)$  can be written as

$$\delta S = \int_{t_i}^{t_f} dt \left[ \delta q(t) \frac{\partial L}{\partial q} + \delta \dot{q}(t) \frac{\partial L}{\partial \dot{q}} \right], \quad (28.6.6)$$

or, using  $\delta \dot{q}(t) = \frac{d}{dt} \delta q(t)$ ,

$$\delta S = \int_{t_i}^{t_f} dt \left[ \delta q(t) \frac{\partial L}{\partial q} + \frac{d}{dt} \delta q(t) \frac{\partial L}{\partial \dot{q}} \right]. \quad (28.6.7)$$

Integrating the second term by parts, we have

$$\delta S = \int_{t_i}^{t_f} dt \delta q(t) \left[ \frac{\partial L}{\partial q} - \frac{d}{dt} \left( \frac{\partial L}{\partial \dot{q}} \right) \right] + \left( \delta q(t) \frac{\partial L}{\partial \dot{q}} \right) \Big|_{t_i}^{t_f}. \quad (28.6.8)$$

The last surface term is 0 because of the boundary conditions eq. (28.6.3). In order to have  $\delta S = 0$  for all small variations  $\delta q(t)$ , we conclude that the integrand in eq. (28.6.8) must vanish,

$$\frac{\partial L}{\partial q} - \frac{d}{dt} \left( \frac{\partial L}{\partial \dot{q}} \right) = 0. \quad (28.6.9)$$

This is the Lagrangian equation of motion familiar from classical mechanics. More generally, if there are multiple variables  $q_a$  with  $a = 1, \dots, N$ , with a Lagrangian  $L(q_a, \dot{q}_a, t)$ , then one can show by the same argument that the classical limit implies the equations of motion

$$\frac{\partial L}{\partial q_a} - \frac{d}{dt} \left( \frac{\partial L}{\partial \dot{q}_a} \right) = 0 \quad (28.6.10)$$

for each  $a$ . Thus we have derived classical mechanics from the  $\hbar \rightarrow 0$  limit of Feynman's path integral formulation of quantum mechanics.

In the case of a single spinless particle moving in three dimensions, we could of course take our CSCO eigenvalues to be  $(q_1, q_2, q_3) \rightarrow \vec{r}$ . However, in more general situations, we often need to describe processes that change the number of particles. Such processes include decays of one particle into two or more particles, like the decay of a Higgs boson into a quark-antiquark pair, or two gluons, or two photons. They also include the joining of two particles to make particles of a different type, for example the annihilation of an electron and positron pair to make a  $Z$  boson, which then turns into a quark-antiquark pair. In such cases, the way we have

been treating quantum mechanics is hopelessly clumsy, because even the number of coordinates needed to describe the positions or momenta of the particles changes depending on the state; in other words, they are not a good CSCO.

Furthermore, special relativity teaches us that time and space are on the same footing, while we have been treating spatial coordinates as operators, but using the time coordinate as a parameter, not an operator. This is particularly embarrassing when we realize that the symmetries of special relativity mix space and time coordinates.

The simultaneous resolution of these issues is that instead of associating the position coordinates  $\vec{r}$  with operators, they are taken to be labels, just like the time  $t$ . In other words, there are different operators

$$\Phi(\vec{r}, t) \tag{28.6.11}$$

for each label  $(\vec{r}, t)$ . The operators  $\Phi(\vec{r}, t)$ , called **quantum fields**. They are associated with the different types of particles in the theory, and can also carry indices associated with their spin and other quantum numbers. The field operators can act on a vacuum state  $|0\rangle$  (which describes empty space with no particles, and is the lowest energy state) to give other states that have any number of particles and antiparticles. In the path integral formulation, the classical action functional looks schematically like

$$S[\phi(\vec{r}, t)] = \int_{t_i}^{t_f} dt \int d^3\vec{r} \mathcal{L}(\phi, \dot{\phi}, \vec{\nabla}\phi), \tag{28.6.12}$$

where  $\mathcal{L}$  is called the classical Lagrangian density, and  $\phi$  is the classical field corresponding to the quantum field  $\Phi$ .

The resulting type of theory, which also obeys the other postulates of section 3.1, is called a **quantum field theory**. Relativistic quantum field theories have the important property that energies can be bounded from below, which is not the case in relativistic one-particle quantum mechanics, as we saw in Chapter 27. In quantum field theory, the Schrödinger equation formulation is valid, but often less useful than the equivalent path integral formulation, which provides elegant methods for both conceptual understanding and practical calculations, using both perturbative and non-perturbative methods. But that's another story...

## 28.7 Exercises

**Exercise 28.1.** Consider the general spectral decomposition of the propagator in eq. (28.1.9). Compare it to the specific result for the harmonic oscillator, in eqs. (28.1.13) and (28.1.14) with  $T = t - t_0$ , to deduce the known result  $E_n = \hbar\omega(n + 1/2)$  for  $n = 0, 1, 2, \dots$ , and check the known wavefunctions for  $n = 0$  and  $n = 1$ . [Hint: write  $\cos(\omega T) = (1/z + z)/2$  and  $\sin(\omega T) = (1/z - z)/2i$ , where  $z = e^{-i\omega T}$ . Then do a formal expansion in small  $z$ .]

# Index

- 21 centimeter line (from hyperfine splitting of hydrogen), 271, 372, 478
- A and B coefficients, Einstein, 469–471
- $\alpha$  (fine structure constant), 259–260
- $\alpha$  particles, 8
- absorption (harmonic time-dependent perturbation), 458–461
- absorption of light, 464–468
  - electric dipole approximation, 472–473
- absorption spectrum, hydrogen atom, 261
- action, 98, 622
- addition of angular momenta, 266–285
  - formula for spherical harmonics, 206–208
  - general case, 275–282
  - more than two, 283–285
  - orbital and spin, 272–275
  - two spins, 268–270
- adjoint, 32, 36–37
- Aharonov–Bohm (Ehrenberg–Siday) effect, 633–636
- Airy function and equation, 154–156
- alkali metals, 403
- allowed (electric dipole) transitions, 475
- allowed results of measurements, 71
- ambiguity
  - operator ordering, 74
  - phase, 28, 45, 111
- angular frequency, Planck–Einstein relation to photon energy, 13
- angular momentum
  - addition of, 266–285
  - and spherical harmonics, 201–205
  - and the Laplacian, 199–200, 206
  - as generator of rotations, 115–120
  - commutation relations, 115, 119
  - conservation from rotation symmetry, 120–121
  - dot product trick, 267–268, 270–271, 367, 377
  - eigenvalue problem, general, 186–190
  - eigenvalue problem, orbital, 200–202
  - general, 119
  - intrinsic, 16–17, 119, 191
  - multiplet, 190
  - orbital, 115
  - relative and center-of-mass, 120
  - spin, 16–17, 119, 191
  - squared, 119
  - total, 120
- anharmonicity parameter, 426
- annihilation operator (harmonic oscillator), 169
- anomalous magnetic moment of electron, 105
- anti-Hermitian operator, 37
- anticommutator, 34
- antisymmetric (singlet) spin combination, 269–271
- antisymmetric Levi-Civita symbol  $\epsilon_{abc}$ , 116
- antisymmetrization for fermions, 271–272, 385–388
- Argonne, interference with Batavia, 393–394
- Aspect’s experiments, 315–319
- associated Laguerre polynomials, 244–245
  - differential equation, 245
  - explicit form, 245
  - for 3-d harmonic oscillator, 244–245
  - for hydrogen atom, 252–253
  - orthogonality
    - for 3-d harmonic oscillator, 245
    - for hydrogen atom, 253
- associated Legendre functions, 203
- associativity for operators, 34
- associativity for vector addition, 24
- asymmetric top, 428
  - water molecule, 436–437
- atom
  - classical instability, 8–9
  - hydrogen-like, 248–261
  - multi-electron, 398–412
- azimuthal symmetry, 486
- Baker–Campbell–Hausdorff formula, 39
- Balmer series, 261
- barn (unit of area), 525
- basis, 25
  - orthonormal, 28–29



- Batavia, interference with Argonne, 393–394
- Bayes' Theorem, 91
- BB84 private key protocol, 587–588
- Bell inequality, 311–316
- Bell states, 579
- Bernstein–Vazirani problem, 576–578
- Bessel functions, spherical, 231–232
  - modified, 233
- beta decay
  - inverse, 131
  - of tritium, 448
- binomial coefficient, 408
- bipartite system, 532
- black-body radiation, 10–12
- Bloch ball (or sphere) and vector, 572–573
- Bohm's version of the EPR problem, 304–309
- Bohr magneton ( $\mu_B$ ), 106
- Bohr radius of hydrogen atom, 251, 259
- Boltzmann entropy, 90
- Boltzmann factor, 11, 94, 470
  - derivation of, 93–94
- Boltzmann's constant, 11, 90
- boost (relativistic), 597
- Born approximation, 502–507
  - beyond first-order, 505–507
  - diagrammatic representation, 506–507
  - first-order, 502–504
  - validity condition, 503
- Born rule (probabilistic interpretation of the wavefunction), 58, 65, 72
- Born–Oppenheimer approximation, 415–418
- Bose–Einstein condensate, 390–391
- Bose–Einstein statistics, 385
- boson, 272
- bound state, 60, 138, 139
  - energy quantization, 9
  - in 1 dimension, 139–140
- boundary condition
  - bound state wavefunction at infinite distance, 138, 139
  - continuity of wavefunction, 229
  - derivative of wavefunction, 139
  - finiteness of radial wavefunction at  $r = 0$ , 228–229
  - impenetrable walls, 59
  - periodic, 10, 60, 226
  - radial wavefunction, 229
  - wavefunction, 138–139
- box
  - particle in a 1-d, 140–142
  - very large, 60
- bra vector, 32
- bra-ket notation, 23, 32
- Brackett series, 261
- Breit–Wigner (Lorentzian) lineshape, 468, 517
- Brillouin–Wigner perturbation theory, 334–335
- $c$  (speed of light in vacuum), 9
- canonical commutation relations, 99
- canonical ensemble, 93–95
- canonical momentum, 98–99
  - and translations, 114
  - vs. kinetic momentum, 104
- Cartan subalgebra, 109
- Cartesian tensor operator, 294
- Cauchy–Schwarz inequality, 27
- center-of-mass coordinates and momenta, 101
- central-field method for multi-electron atoms, 399–400
- centrifugal term, 229, 242, 518
- change of orthobasis, unitary, 42–43
- channel (dynamical map), 537
- characteristic equation for eigenvalues, 44
- charge of electron, 8, 103
- charge radius of proton, 330–331
- classical instability of charged matter, 8–9
- classical limit, 74, 252, 259, 640–641
  - coherent states, 174–180
- classical-like states, 558
- Clebsch–Gordan coefficients, 276–283
  - computation algorithm, 279–282
  - phase convention, 276–277
- closure (completeness) relation, 35
- closure property
  - for Lie groups, 108–109
  - for vector spaces, 23
- cofactor matrix, 41
- coherences, 87, 558
- coherent states of harmonic oscillator, 174–180, 561–564
  - time evolution, 178–179
- coherent superposition, 87



- collapse of state, 72–73
  - controversy and unease, 84–85
  - for generalized measurement, 547
  - for mixed ensembles, 91–92
- collective measurement, 579
- combination formula, 407–408
- commutativity property of vector space, 24
- commutator, 34
  - canonical, 99
  - position and momentum, 63, 64
- compatible (commuting) operators, 54–56
- complete orthobasis and observables, 53
- complete set of commuting observables (CSCO), 55–56
- completely positive map, 541–545
- completeness relation, 35, 71
  - angular momentum, 186, 190, 201
  - Kraus operators, 538, 542, 546
  - momentum (1-dimensional), 60
  - momentum (3-dimensional), 64
  - position (1-dimensional), 57
  - position (3-dimensional), 64
  - spherical harmonics, 201
- complex linear vector space, 23–24
  - basis and dimension, 25
  - examples, 24
  - subspace, 26
- components of a vector, 25, 26
  - for an orthobasis, 32
- composition rule for propagators, 623–624
- Compton wavelength, 259
- computation, 571–596
- Condon–Shortley phase convention, 202
- confluent hypergeometric function, 249–250
- conservation laws
  - and symmetries, 108–111
  - angular momentum, 120–121
  - charge, 103, 610
  - energy, 111
  - general theorem (Noether’s principle), 110
  - momentum, 113–115
  - probability, 128–131
- constant of motion, 110, 439
- continuity of wavefunctions, 58, 138–139, 229
- contravariant four-vector, 597
- conventions for units, 8, 11, 13, 102–103
- Copenhagen interpretation, 84
- correlation for entangled states, 306–307, 311–319
- correspondence principle, 90, 99–100
- Coulomb gauge, 107, 463
- Coulomb potential, 248
  - unbound states, 261–264
- covariant four-vector, 598–599
- creation operator (harmonic oscillator), 169
- cross-section
  - absorption of light, 465, 468
  - electric dipole approximation, 472–473
  - differential, 486
  - photo-electric effect, 482–484
  - relation to scattering amplitude, 490
  - scattering, 486
- CSCO (Complete Set of Commuting Observables), 55–56
- current density
  - electric, 103, 129
  - probability, 128–130
- cycloid motion, 227
- cyclotron frequency, 218–219
- Dalgarno–Lewis method, 335–337
  - applied to quadratic Stark effect, 345–346
- Darwin term, 365–366, 368
  - from Dirac equation, 613
- Davisson–Germer experiments, 15–16
- de Broglie wavelength, 15
  - thermal, 570
- decay of unstable states, 130–131
- decoherence, 557–570
  - and error correction, 588–594
  - Schrödinger’s cat, 557–559
  - damped harmonic oscillator, 561–564
  - exactly solvable spin model, 567–570
  - of diffraction pattern, 87–88
  - phase damping of a spin, 564–567
  - qubit, 596
- degeneracy label, 48, 71
  - and CSCO, 56
  - continuous, 53
- degeneracy of eigenvalue, 44, 48
- degeneracy of energy levels
  - 3-dimensional harmonic oscillator, 181–182

- absence in 1-dimensional bound states, 139–140
- charged particle in magnetic field, 221–227
- diatomic molecular states, 430
- hydrogen atom bound states, 251
- particle confined to a sphere, 238
- rigid body rotations, 428–429
- delta function, 28–30
  - 3-dimensional, 64
  - and Fourier transforms, 30
  - integral representation, 29–30
- delta, Kronecker, 28
- dense coding, 582–583
- density (matrix) operator, 88–95
  - reduced, 533
- density of modes
  - electromagnetic radiation, 10–11
  - electron states in cubic box, 481–482
  - energy eigenstates, 455
- depolarization of qubits, 596
- derivative operator, 58–59
- destruction operator (harmonic oscillator), 169
- detailed balance, 459, 471
- determinant of matrix, 41
- deuteron, 523–526
- Deutsch–Jozsa problems, 576
- diagonalization
  - of Hermitian matrix by a unitary matrix, 49, 55
  - of real symmetric matrix by orthogonal matrix, 160
- diatomic molecules
  - $\text{H}_2^+$  ion, 419–422
  - identical nuclei, 431–433
  - rotational excitations, 429–434
  - sample empirical energy parameters, 430
  - vibrational excitations, 422–426, 429–431
- differential cross-section, 486
  - relation to scattering amplitude, 490
- diffraction, 14
  - and decoherence, 87–88
  - for electrons, 15–16
- diffuse ( $l = 2$  in spectroscopic notation), 252
- dimension of vector space, 25
- dipole selection rules, 298–299
- Dirac bra-ket notation, 23, 32
- Dirac delta function, 28–30
  - 3-dimensional, 64
  - and Fourier transforms, 30
  - integral representation, 29–30
- Dirac equation, 366, 368, 603–618
  - hydrogen atom solutions, 613–618
  - plane-wave solutions, 607–608
- Dirac magnetic monopoles, 636–639
- Dirac orthonormality, 28–29, 33, 53, 71
  - momentum eigenstates, 60, 64
  - position eigenstates, 57, 64
  - spherical coordinate eigenstates, 200
- Dirac picture (interaction picture), 441–445
  - Dyson series solution, 443–444
- Dirac sea, 608
- distribution, 29
- distributive property of vector space, 24
- dot product of angular momenta trick, 267–268, 270–271, 367, 377
- dots and lines trick, 182
- double factorial notation, 231
- double-counting avoidance in identical particle scattering, 528
- dual vector space, 32–33
- duality, Gaussian position and momentum wavefunctions, 133
- duality, wave-particle, 13–15
- dynamical map, 537
  - amplitude damping example, 538–540
  - conditions for, 541–543
  - operator-sum form, 537–538, 541–543
  - phase damping of spin example, 564–567
- Dyson series, 83
  - interaction picture, 443–444
- $e$  (proton charge in Gaussian cgs units), 8, 103
- effective potential, 229, 242, 518
- effective range, 513–514
  - for neutron-proton scattering, 524–525
  - for spherical well scattering, 523
- Ehrenberg–Siday (Aharonov–Bohm) effect, 633–636
- Ehrenfest’s Theorem, 81–82, 99
- eigenstate (eigenket), 44
- eigenvalues, 44
  - and trace of operator, 52

- characteristic equation, 44
- continuous, 56–57
- degenerate, 44, 48
- Hermitian operator, 47
- identity operator, 45
- multi-component, 64, 73–74
- parity, 121
- projection operator, 45
- unitary operator, 52
- eigenvector, 44
- Einstein A and B coefficients, 469–471
- Einstein’s energy quantization for photons, 13
- Einstein–Podolsky–Rosen (EPR) problem, 304–309
- electric dipole (long wavelength) approximation, 471–475
  - selection rules, 474–475
- electric quadrupole (E2) transitions, 476–479
  - selection rules, 478–479
- electromagnetic energy density, 103
- electromagnetic potentials, 103
  - four-vector, 609
- electromagnetic wave, 463–464
  - absorption, 464–468
    - electric dipole approximation, 472–473
  - emission, 468–471
    - electric dipole approximation, 473
- electron
  - charge, 8, 103
  - $g$ -factor, 105, 612
  - mass, 9
  - spin, 16
- electron configuration for multi-electron atom, 400–404
- elliptical coordinates, 420–421
- emission (general harmonic time-dependent perturbation), 458–461
- emission of light, 468–471
  - electric dipole approximation, 473
  - induced (stimulated), 468–469
  - spontaneous, 469–471
- encoding, redundant, 585
- energy conservation from time translation symmetry, 111
- energy density, electromagnetic, 103
- ensemble
  - average over, 86
  - completely random, 89
  - mixed, 85–95
  - probabilities, 86, 89
  - pure, 76
- entanglement, 67
  - and subsystems, 532–536
  - Bell states, 579
  - dense coding, 582–583
  - entropy, 535
  - nonlocal, 305
  - Schmidt (singular value) decomposition, 534–535
  - teleportation, 579–582, 596
- entropy, 89–90
  - entanglement, 535
  - for a qubit, 573
- environment, 532
- EPR problem, 304–309
  - reduced density operator, 532–533
- EPR protocol for private key distribution, 586–587
- EPRB (Einstein–Podolsky–Rosen–Bohm) states, 579
- epsilon symbol  $\epsilon_{abc}$ , 116
- equal-time commutation relations, 440
- equations of motion
  - Hamiltonian, 99
  - Lagrangian, 98
  - Lagrangian, derived, 641
- error correction, 588–594
  - Shor 9-qubit encoding, 593–594, 596
- Euler angles, 198
- Everett, Hugh, 84
- exchange degeneracy, 385–386
- exchange density, 393
- exclusion principle for identical fermions, 272, 386–388, 391, 401
- expectation value, 76, 86
- explicit time dependence, 80
- exponentiation of operator, 37–39, 43
- Fermi’s golden rule, 456–457
  - harmonic perturbations, 458–459
- Fermi–Dirac statistics, 285, 385–386, 393, 396, 401

- fermion, 272
- Feynman sum over paths (path integral), 623–642
  - classical limit, 640–641
  - discretized version, 626–627
  - equivalence to Schrödinger equation, 627–628
- Feynman's disappearing screen, 623–624
- fiction, convenient, 25
- fine structure
  - from Dirac equation, 613, 617–618
  - of hydrogen atom, 365–369, 375
  - of multi-electron atoms, 404–406
- fine structure constant  $\alpha$ , 259–260
- finite range scattering potential, 489, 511–514
- flux of particles
  - in 1-dimensional scattering, 148
  - in 3-dimensional scattering, 486
- forbidden transitions involving light, 475
- four-momentum, 597
- four-vector, 597–599
- Fourier transform, 30–31
  - 3-dimensional, 65
  - momentum and position wavefunctions, 62
- Fourier transform (quantum gate), 578
- free energy, Helmholtz, 95
- free particle
  - in 1 dimension, 60, 134–137
    - wavefunction, 61–62
  - in 3 dimensions, 64
  - in spherical coordinates, 229–236
  - time evolution, 136, 621
- frequency, Planck–Einstein relation to photon energy, 11, 13
- full width at half maximum
  - Breit–Wigner (Lorentzian) lineshape, 468, 517
  - Gaussian, 30, 132
- functional, 625
- functional derivative, 640
- functional integral, 625–628
  - for harmonic oscillator, 628–633
- functions of operators, 37–39
  - spectral decomposition, 54
- fundamental ( $l = 3$  in spectroscopic notation), 252
- fundamental theorem of algebra, 44, 48
- fundamental theorem of calculus, 139, 148
- $g$ -factor
  - electron, 105
    - from Dirac equation, 612
  - Landé
    - hydrogen atom, 377
    - multi-electron atom, 410
  - neutron, 105
  - proton, 105, 369
- $\Gamma(z)$  (Gamma function), 244–245
- gamma matrices (for Dirac equation), 605–606
- $\Gamma$  (for unstable states), 130
- $\Gamma$  (width in resonant scattering), 516–517
- gate, 574
- gauge transformation, 103
  - in quantum mechanics, 124–125
  - Lorentz covariant, 609
- gauge-invariant observable, 125–127
- Gaussian wavefunction, 132–137
  - full width at half maximum, 132
  - momentum, 133
  - time evolution for free particle, 136, 137
  - uncertainty relation for, 134
- generalized (non-normalizable) kets, 29, 33, 57
- generalized (Robertson–Schrödinger) uncertainty relation, 77–78
- generalized measurement, 545–551
  - photon detection example, 549
  - positive operator-valued measure (POVM), 548–549
  - state distinguishing example, 550–551
- generator
  - of symmetry, 109–111
  - of transformation, 108–109
  - rotations (with spin), 120
  - rotations (without spin), 116–117
  - translations, 113
- Gibbs entropy, 90
- giraffes hiding in kitchens, 252
- global phase, 80
- gluon, 8, 105, 210, 386, 608, 641
- golden rule, 456–457
  - harmonic perturbations, 458–459
- good quantum number, 56

- Gorini, Vittorio, 555
- Gram–Schmidt process, 31–32
- Grand Unified Theories, 103
- graphical solution
  - 1-dimensional square well, 144
  - particle confined to a sphere, 237–238
- Green function for 3-d scattering, 497–499
- Greenberger–Horne–Zeilinger–Mermin state, 313–314
- grotesque (silly name for  $l = 4$  in spectroscopic notation), 252
- ground state, 24
- group velocity
  - of Gaussian wavepacket, 137
  - vs. phase velocity, 134–135
- Grover search algorithm, 579
- gyromagnetic ratio, 105
  - and spin precession, 212–215
- $\hbar$ , 13
- H<sub>2</sub>O molecule, 435–437
- H<sub>2</sub><sup>+</sup> ion molecule, 419–422
- Hadamard transformation (gate), 575
- half-life, 474
- Hall effect, 225
- Hamiltonian, 73
  - classical, 98
  - Dirac equation, 603, 610
  - equations of motion, 99
  - free-particle, 134
  - harmonic oscillator, 1-d, 162
  - harmonic oscillator, 3-d isotropic, 240
  - hydrogen-like atoms, 248
  - non-Hermitian, 130–131
  - particle in electromagnetic field, 106–107
  - time-dependent, 82–84
- Hankel functions, spherical, 232–233, 512
- hard-sphere scattering
  - classical, 487–488
  - quantum, 519–521
- harmonic oscillator, 159–182, 240–245
  - algebraic method, 169–174
  - anisotropic 3-d, 180–181
  - coherent states, 174–180, 561–564
  - differential equation method, 162–169
  - energy representation, 169–174
  - ground-state wavefunction, 166–167
  - Hamiltonian, 162
  - isotropic 3-d, 181–182, 240–245
  - momentum representation, 168–169
  - number operator, 171
  - path integral approach, 628–633
  - position representation, 162–168
  - positivity of energy eigenvalues, 162
  - propagator, 621–622
  - stationary state wavefunctions, 166–168
- Heisenberg equation of motion, 440
- Heisenberg picture, 438–441
- Heisenberg uncertainty relation, 78
- helium atom
  - excited states, 396–398
  - ground state
    - and spin, 395–396
    - first-order perturbation theory, 332–334
    - variational method, 361–363
  - Hamiltonian, 332
  - hyperfine structure, 397
  - para (total spin 0) and ortho (total spin 1), 396–398
- Helmholtz free energy, 95
- Hermite polynomials, 165–166
  - algebraic derivation, 173
- Hermitian adjoint, 32, 36–37
- Hermitian operator, 37
  - constraint for infinite-dimensional Hilbert space, 59
  - corresponding orthobasis, 47–48
- heteronuclear diatomic molecules, 430–431
- hidden variables, 21, 309–319
  - demise of, 315–319
- Hilbert space, 26, 71
- hole (absence of electron), 409, 608
- homonuclear diatomic molecules, 431–434
- horrendous (silly name for  $l = 5$  in spectroscopic notation), 252
- Humphreys series, 261
- Hund’s rules, 406–409
- hydrogen atom, 248–261
  - absorption spectrum, 261
  - classical instability, 8–9
  - Dirac equation, 613–618
  - energy spectrum and degeneracy, 250–251



- expectation values of powers of  $R$ , 256–258, 341–342
- fine structure, 365–369, 375, 613–618
- hyperfine structure, 271, 369–375
- impact of proton radius, 330–331
- in external magnetic field, 375–381
- Paschen–Back effect, 378–381
- polarizability, 346
- relativistic corrections, 260, 365–369
- Stark effect, 343–348
- stationary state wavefunctions, 252–256
- unbound states, 261–264
- wavefunction at the origin, 254, 331
- Zeeman effect, 375–381
- hyperfine structure
  - helium atom, 397
  - hydrogen atom, 271, 369–375
- identical particles, 271–272, 383–412
  - and spin, 391–395
  - constraint on Hamiltonian, 385
  - constraint on observables, 384
  - constraint on states, 385–386
  - factor of 1/2 for cross-section, 528
  - nuclei in diatomic molecules, 431–433
  - pair-exchange operators, 384–385
  - permutation operators, 386–387
  - scattering, 526–528
- identity operator, 34–35
- impact parameter, 486–487
- improper Lorentz transformation, 599
- incoherent superposition of harmonic perturbations, 459–460
- incompatible operators, 56, 73–74
- index relabeling trick, 164, 242, 286, 300
- index summation convention, 116, 160
- induced (stimulated) emission, 468–469
  - electric dipole approximation, 473
- inelastic photon scattering (Raman effect), 433–434
- inert (noble) gases, 402, 407
  - electron scattering, 152–153, 523
- inertial reference frame, 597
- infinite-dimensional vector space, 25
- infinite-range scattering potential, 488
- information, 571–596
- inner product, 26
  - in terms of wavefunctions, 57, 61, 63, 65
  - matrix representation, 39
  - preserved by unitary transformation, 42–43
- instability of charged matter in classical mechanics, 8–9
- intensity and photon occupation numbers, 467
- interaction picture (Dirac picture), 441–445
  - Dyson series solution, 443–444
- interference, 14–15
- interval rule, Landé, 405–406
- intrinsic angular momentum, 16–17, 119, 191
- intrinsic parity, 209–210
- inverse Fourier transform, 31
- inverse metric tensor ( $g^{\mu\nu}$ ), 598
- inverse of an operator, 35
- inverse of matrix, 41
- ionization energy
  - first, multi-electron atoms, 402–403
  - hydrogen (Rydberg), 251
  - total, helium, 334, 363
- irreducible representation of rotation group, 195
- irreducible tensor operator, 291–295, 299–303
- isometric operator, 495
- Jacobi coordinates, 102
- Jacobi identity, 69
- $jj$  coupling approximation, 411–412
- jump (Lindblad) operators for open system, 555
- $K$ -shell electrons, 484
- kernel, 620
- ket, 23
  - generalized (non-normalizable), 29, 33, 57
  - null, 24
- kinetic (mechanical) momentum, 104
  - operator, 106–107, 125–126, 129
  - vs. canonical momentum, 104
- Klein’s inequality, 92
- Klein-Gordon equation, 601–602
- Kossakowski, Andrzej, 555
- Kramers–Pasternack recurrence relation, 257–258
- Kraus operators, 537–538
  - as measurement operators, 545–548
  - non-uniqueness, 540–541
- Kronecker delta symbol, 28

- ladder operator
  - angular momentum, 187
  - harmonic oscillator, 169
- Lagrangian, 98, 625, 641
- Laguerre polynomials, associated, 244–245
  - differential equation, 245
  - explicit form, 245
  - for 3-d harmonic oscillator, 244–245
  - for hydrogen atom, 252–253
  - orthogonality
    - for 3-d harmonic oscillator, 245
    - for hydrogen atom, 253
- Lamb shift, 374–375, 380, 381
- Landé  $g$ -factor
  - hydrogen atom, 377
  - multi-electron atom, 410
- Landé interval rule, 405–406
- Landé projection formula, 297–298, 301, 372
- Landau levels, 218–227
- Laplacian and angular momentum, 199–200, 206
- Larmor formula for radiated power, 9
- Larmor precession frequency, 212
- Legendre functions, associated, 203
- Legendre polynomials, 203, 203
- level repulsion, 349, 350
- Levi-Civita symbol  $\epsilon_{abc}$ , 116
- Lie algebras and groups, 108–109
- lifetime due to spontaneous emission, 474
- light
  - absorption, 464–468
    - electric dipole approximation, 472–473
  - and hydrogen transitions, 260, 261
  - emission, 468–471
    - electric dipole approximation, 473
  - interference, 14–15
  - Planck–Einstein energy-frequency relation, 11, 13
  - quanta (photons), 13
- Lindblad equation, 555
  - damped harmonic oscillator, 561–562
  - simple decoherence example, 560
- line broadening, 467–468
- linear (in)dependence of vectors, 25
- linear operator, 33–34
- linear potential, 153–156
- linear rigid rotor, 429
- linear Stark effect, 346–347
- linearity property of inner product, 26
- Liouville’s Theorem, 90
- Liouvillian superoperator, 553
  - Lindblad form, 555
- Lippmann–Schwinger equation, 494, 497–500
- local conservation of charge, 103
  - Lorentz covariant, 610
- local conservation of probability, 128–131
- local realism, 309–310
  - demise of, 315–319
- logical qubit, 585
- Lorentz force law, 103
- Lorentz tensor, 600–601
- Lorentz transformation, 597–600
  - Dirac spinor, 606–607
  - proper, 600
- Lorentzian (Breit–Wigner) lineshape, 468, 517
- lowering operator
  - angular momentum, 187
  - harmonic oscillator, 169
- $LS$  coupling scheme, 404–409
- Lyman series, 260–261
- Lyman-alpha line, 261
- magnetic dipole (M1) transitions, 476–478
  - selection rules, 477–478
- magnetic field and Landau levels, 218–227
- magnetic flux quantum, 224, 635
- magnetic length, 222
- magnetic moment
  - and spin, 18–19, 104–105
  - deuteron, 524
  - electron, 18–19, 105, 366
  - multi-electron atom, 410
  - neutron, 105–106
  - proton, 105–106, 369–370
- magnetic monopoles, 636–639
- magnetic quantum number  $m$ , 119, 378
- magnetic spin resonance, 215–218
- magneton, Bohr ( $\mu_B$ ) and nuclear ( $\mu_N$ ), 106
- magnitude of a complex number, 23
- many-worlds (Everett) interpretation, 84–85
- Markovian approximation, 553
- matrix element of an operator, 34
- matrix element, reduced, 296, 298, 301–302

- matrix inversion, 41
- matrix representation in an orthobasis, 39–40
- Maxwell's equations, 102
  - Lorentz-covariant form, 609–610
- measurement
  - allowed results, 71
  - and collapse of state, 72–73
    - controversy and unease, 84–85
    - for mixed ensembles, 91–92
  - and decoherence, 564–567
  - and probability, 72
    - for mixed ensembles, 88
  - generalized, 545–551
  - destructive, 549–550
- measurement operators, 545–548
- mechanical (kinetic) momentum, 104
  - operator, 106–107, 125–126, 129
  - vs. canonical momentum, 104
- methane, 428
- metric tensor, 598
- microwave ovens, 437
- Millikan photo-electric effect experiment, 14
- mixed ensemble, 85–95
- mixed state, 89
- modified spherical Bessel functions, 233
- modulus (magnitude) of a complex number, 23
- molecules, 415–437
- moments of inertia, 426–428
- momentum, 59
  - 3-dimensional, 63
  - 4-dimensional, 597
  - as generator of translations, 113, 114
  - canonical, 98–99, 104, 114
  - conservation and translation symmetry, 113–115
  - eigenstates, 60–65
  - Hermiticity, 59
  - kinetic (mechanical), 104, 106–107, 125–126, 129
  - uncertainty in Gaussian wavefunction, 134
- momentum wavefunction and representation, 60–65
- monopole, magnetic, 636–639
- Morse potential, 423–426
  - energy spectrum, 426, 435
  - parameters for common diatomic molecules, 430
  - parameters for H<sub>2</sub>O molecule, 436
- multi-electron atoms, 398–412
- multiplet of angular momentum, 190
- multipole expansion for absorption and emission of light, 475–479
- multipole moment operator, 294–295
- muonic hydrogen, 331
- Moller operators, 493–496
- Neumann functions, spherical, 231–232
- neutron
  - magnetic moment, 105–106
  - mass, 523
- neutron-proton scattering, 523–526
- no-cloning theorem, 584–585
- noble (inert) gases, 402, 407
  - electron scattering, 152–153, 523
- Noether's principle, 110–111
- non-normalizable (generalized) kets, 29, 33, 57
- nonlocality, 309
- norm, 26–27
  - in terms of wavefunction, 57, 65
- normal ordering, 169
- nuclear magnetic resonance, 218
- nuclear magneton ( $\mu_N$ ), 106
- nuclear size, 8, 330–331
- null ket, 24
- number operator (harmonic oscillator), 171
- observable, 53, 71
  - completeness of orthobasis, 53
  - gauge-invariant, 125–127
  - quantum counterpart of classical, 74–75
- occupation numbers, 389
  - of atomic electrons in subshells, 401
  - photons, 467
- on-shell condition, 601
- open systems, 532
  - rules summarized, 551–552
- operator, 33
  - multi-component, 64, 73–74
- operator order ambiguities, 74–75
- operator-sum form of dynamical map, 537–538, 541–543
- Optical Theorem, 501–502
- orbital angular momentum, 115



- orbital states, 400
- Orion nebula, 261
- ortho- and para-
  - diatomic molecules, 432–433
  - helium, 396–397
- orthobasis (orthonormal basis), 28–29
  - corresponding to a CSCO, 55
  - corresponding to Hermitian operator, 47–48
  - Gram–Schmidt construction, 31–32
- orthogonal kets, 28
- orthonormality, 28–29
  - angular momentum eigenstates, 186, 190
  - Dirac, 28–29, 33, 53, 71
    - momentum eigenstates, 60, 64
    - position eigenstates, 57, 64
    - spherical coordinate eigenstates, 200
  - Hermite polynomials, 166
  - Legendre polynomials, 203
  - spherical Bessel functions, 233
  - spherical harmonics, 201
- outer product, 35
- pair-exchange operators for identical particles, 384–385
- para- and ortho-
  - diatomic molecules, 432–433
  - helium, 396–397
- paradox, instantaneous communication, 308
- parallelism, 574–575
- parity, 121–123
  - Dirac spinors, 605
  - eigenvalues, 121
    - angular momentum eigenstates, 208–210
    - composite particle, 209–210
    - intrinsic, 209–210
    - multi-electron atom, 209, 406
  - of operators, 122
    - angular momentum, 123, 478
    - position and momentum, 122
  - violation in weak interactions, 210, 600
- parity selection rule, 123
- partial trace, 533
- partial wave expansion, 507–511
- particle-wave duality, 13–15
- partition function, 94–95
- Paschen series, 261
- Paschen–Back effect
  - hydrogen atom, 378–381
  - multi-electron atoms, 410–411
- path integral, 623–628, 642
  - for harmonic oscillator, 628–633
- Pauli exclusion principle for identical fermions, 272, 386–388, 391, 401
- Pauli matrices, 191–192
- periodic boundary condition, 10, 60, 226
- periodic table of elements, 402, 406–409
- permutation operators for identical particles, 386–387
- perpendicular kets, 28
- perturbation theory
  - stationary-state (time-independent), 321–352
    - almost-degenerate, 348–352
    - Brillouin–Wigner, 334–335
    - Dalgarno–Lewis method, 335–337
    - degenerate, 337–342
    - non-degenerate, 321–334
  - time-dependent, 448–457
    - harmonic, 457–461
- Pfund series, 261
- phase
  - ambiguity, 28, 45, 111
  - global and relative, 28, 80
- phase damping of a spin, 564
- phase shift (scattering), 510–511
- phase velocity vs. group velocity, 134–135
- phase-space coordinates, 98
- photo-electric effect, 12–14
  - for atoms, 479–484
- photon, 13
  - generalized measurement example, 549
- pictures of quantum mechanics
  - Heisenberg, 438–441
  - interaction (Dirac), 441–445
  - Schrödinger, 438
- Planck’s constant ( $h$ ), 11
  - reduced ( $\hbar$ ), 13
- Planck’s formula for black-body radiation, 12
- plane wave
  - in 1 dimension, 61–62
  - in 3 dimensions, 64

- relation to spherical wave, 234–236
- Poincaré recurrence, 569
- pointer states, 558
- Poisson bracket, 90, 99
- Poisson distribution (coherent state), 175–176
- polarizability of hydrogen atom, 346
- polarization vector for light, 463–464
- pole in scattering amplitude
  - bound state, 514–516, 518, 519
  - resonance, 516–519
- populations, 558
- position operator
  - in 1 dimension, 56
  - in 3 dimensions, 63–64
- position wavefunction and representation, 57, 65
- positive map, 541–545
  - not completely positive example, 544–545
- positive operator, 534
- positive operator-valued measure (POVM), 548–549
- positivity of inner product, 26
- positron, 608
- positronium, 290
- postulates of quantum mechanics, 71–73
  - collapse (controversy and unease), 84–85
  - Feynman’s alternative for time evolution, 626
  - for open systems, 551–552
  - in terms of density matrix operator, 85–92
  - spin-statistics for identical particles, 385
- potentials, electromagnetic, 103
- Poynting vector, 103
- precession of spin in magnetic field, 212–215
- principal ( $l = 1$  in spectroscopic notation), 252
- principal axes of rigid body, 426–428
- principal quantum number  $n$  (hydrogen), 250
- private keys, 586–588
- probability
  - current, 128–130
  - density, 58, 65
    - in momentum space, 61, 65
  - local conservation, 128–131
  - of a measurement result, 72
  - for mixed ensembles, 88
- product basis, addition of angular momenta, 266
- projection formula, Landé, 297–298, 301, 372
- projection operator, 35–36, 72
  - and collapse of state, 72–73
  - and measurement probability, 72
  - as observable, 75
  - eigenvalues, 45
- projective measurement, 545
  - as special case of generalized measurement, 548
- propagator, 620–624
  - composition rule, 623–624
  - free particle, 621
  - harmonic oscillator, 621–622
- proper distance, 598
- proper Lorentz transformation, 600
- proton
  - charge, 8, 103
  - magnetic moment, 105–106, 369–370
  - mass, 369, 523
  - size, 8
    - impact on hydrogen energy, 330–331
  - spin, 16, 271
- proton-neutron scattering, 523–526
- pure ensemble, 76
- quadratic Stark effect, 343–346
- quadrupole moment operator, 294–295, 477
- quantum field theory, 389, 608, 642
- quantum tunneling, 151–152
- quark, 8, 103, 105
- quasi-bound state (resonance), 516–519
- qubit
  - logical (synthetic), 585
- qubits, 571–573
- Rabi oscillations, 215–218
- radial wavefunction, 228–229
  - 3-d isotropic oscillator, 244–245
  - finiteness at  $r = 0$ , 228–229
  - hydrogen atom, 252–254
- radius of proton, 330–331
- raising operator
  - angular momentum, 187
  - harmonic oscillator, 169
- Raman effect, 433–434
- Ramsauer–Townsend effect, 152–153, 522–523

- Rayleigh–Jeans formula for black-body radiation, 11–12
- Rayleigh–Ritz variational method, 355–363
  - examples, 358–363
  - for excited states, 356–358
    - example, 360
  - forgiving nature, 355–356
  - helium atom ground state, 361–363
- reduced Compton wavelength, 259
- reduced density (matrix) operator, 533
- reduced mass (in 2-body problem), 101
  - hydrogen-like atoms, 248
  - neutron-proton scattering, 523
- reduced matrix element, 296, 298, 301–302
- reduced Planck’s constant  $\hbar$ , 13
- reducible tensor operator, 294
- redundant encoding, 585
- reflection and transmission ratios  $R$ ,  $T$ , 148
- register of qubits, 573–574
- relabeling summation index trick, 164, 242, 286, 300
- relative phase, 28
- relativity, 73, 259, 597–618
  - Dirac equation, 603–618
  - Klein-Gordon equation, 601–602
- repeated index summation convention, 116, 160
- representation
  - angular momentum, coordinate, 199–208
  - angular momentum, matrix, 190–194
  - momentum, 60–62, 65
  - of operator in an orthobasis, 39–40
  - of rotation operators, 195–198
  - position, 57, 65
  - spin, 190–192
- resonance
  - 1-d scattering, 151–152
  - 3-d scattering, 516–519
  - magnetic spin, 215–218
- rigid bodies, 426–434
  - asymmetric top, 428, 436–437
  - diatomic molecule, 429
  - Hamiltonian, 427
  - linear rotor, 429
  - spherical top, 428
  - symmetric top, 428–429
- Robertson–Schrödinger uncertainty relation, 77–78
- robust states, 558
- rotation operator, unitary, 117–118, 195
  - matrix representations, 195–198
- rotation symmetry and angular momentum conservation, 120–121
- rotation transformations, 115–121
- rotational excitations of molecules, 426–431
- Russell–Saunders coupling approximation, 404–409
- Rutherford scattering, 488
- Rutherford–Geiger–Marsden experiment, 8
- Rydberg (unit of energy), 251, 259
- Rydberg formula for hydrogen spectrum, 260
- $S$ -matrix, 491–493, 496–497
- scalar operator, 292
  - selection rules, 296
- scalar product, 26
- scattering, 146, 486
  - Born approximation, 502–507
  - hard-sphere
    - classical, 487–488
    - quantum, 519–521
  - identical particles, 526–528
  - in 1 dimension, 146–153
    - rectangular barrier potential, 149–151
    - rectangular well potential, 151–153
    - step-function potential, 148–149
  - neutron-proton, 523–526
  - Rutherford, 488
  - spherical potential well, 521–523
  - Yukawa (screened Coulomb) potential, 504–505
- scattering amplitude, 489–490
- scattering cross-section, 486
- scattering length, 513–514
  - for neutron-proton scattering, 524–525
  - for spherical well, 523
- scattering resonance
  - in 1 dimension, 151–152
  - in 3 dimensions, 516–519
- scattering states, 60, 146
- Schmidt decomposition of entangled state, 534–535

- Schrödinger equation, 73, 78–79
  - time-independent, 79
    - differential equation in 1 dimension, 138
    - differential equation in 3 dimensions, 228
- Schrödinger picture, 438
- Schrödinger’s cat, 1, 557–559
- screened Coulomb (Yukawa) potential, 364, 504–505
- selection rules, 47
  - $J = 0$  to  $J = 0$  forbidden, 475
  - Clebsch–Gordan coefficients, 276, 288, 302–303
  - dipole, 298–299
  - electric dipole transitions, 474–475
  - electric quadrupole transitions, 478–479
  - general multipole transitions, 479
  - inner product, 47
  - magnetic dipole transitions, 477–478
  - matrix element, 47
  - parity, 123
  - tensor operator, 302–303
  - time-dependent perturbation theory, 453–454
  - vector operator, 299
- self-adjoint operator, 37
- separable (not entangled) states, 67
- Shannon entropy, 89
- sharp ( $l = 0$  in spectroscopic notation), 252
- shell model for multi-electron atom, 400–404
- Shor, Peter
  - 9-qubit encoding, 593–594, 596
  - factoring algorithm, 578
- short-time approximation, 446–447
- simultaneous measurement of compatible observables, 74
- singlet state of two spins, 269–271
- singular value decomposition, 534–535
- Slater determinant, 387–389
- solid angle differential,  $d\Omega$ , 200, 486
- space inversion (parity), 121–123
- special relativity, 73, 259, 597–618
  - Dirac equation, 603–618
  - Klein-Gordon equation, 601–602
- spectral decomposition, 53–54
  - of propagator, 621
  - of time-evolution operator, 79–80
- spectroscopic notation, 251–252, 284–285, 406–409
- speed of light in vacuum ( $c$ ), 9
- spherical Bessel functions, 231–232
  - modified, 233
- spherical Hankel functions, 232–233, 512
- spherical harmonics, 201–205
  - addition formula, 206–208
  - and parity, 208–209
  - combinations and products, 285–288
  - non-existence for half-integer angular momenta, 197
- spherical Neumann functions, 231–232
- spherical potential well
  - bound states, 239–240
  - scattering, 521–523
- spherical tensor operator, 291–295, 299–303
- spherical top, 428
- spin, 16–17, 119, 191
  - combination of two, 268–270
  - combination with orbital angular momentum, 272–275
  - matrix representation, 190–192
  - precession in magnetic field, 212–215
- spin resonance, magnetic, 215–218
- spin-orbit interaction, 365–366, 369, 399
  - from Dirac equation, 613
- spin-statistics principle, 385–386
  - identical nuclei in diatomic molecules, 431–433
- spinor, 192
  - Dirac, 603
- spinor harmonics, 613–615
- spinor operator, 292
- spontaneous emission of light, 469–471
  - electric dipole approximation, 473
- spooky action at a distance, 307
- square-well potential (1-d)
  - bound states, 142–146
  - scattering, 151–153
- standard deviation and uncertainty, 77
- Stark effect
  - linear, 346–347
  - quadratic, 343–346
- state vector (or state ket), 24, 71
- stationary paths, 640–641

- stationary state, 80
- stationary-state perturbation theory, 321–352
  - almost-degenerate, 348–352
  - Brillouin–Wigner, 334–335
  - Dalgarno–Lewis method, 335–337
  - degenerate, 337–342
  - non-degenerate, 321–326
    - examples, 326–334
- Stern–Gerlach experiment, 17–21
  - analogy with photon polarization, 21
  - sequential, 19–21
- stimulated (induced) emission, 468–469
  - electric dipole approximation, 473
- structure constants of Lie algebra, 109
- subspace of a vector space, 26
- Sudarshan, E.C. George, 555
- sudden approximation, 447–448
- summation convention for indices, 116, 160
- superdense coding, 582–583
- superdeterminism, 319
- superoperator, 537
- superposition, 23, 28
  - principle, 46
- symmetric (triplet) spin combination, 269–271
- symmetries and conserved quantities, 108–111
  - general (Noether’s principle), 110
  - rotations and angular momentum, 120–121
  - time translation and energy, 111
  - translations and momentum, 113–115
- synthetic qubit, 585
- teleportation, 579–582, 596
- temperature, 94
- tensor operator
  - Cartesian, 294
  - irreducible, 291–295, 299–303
  - reducible, 294
- tensor product of Hilbert spaces, 66–68
  - and entanglement, 67
- term symbol (spectroscopic notation), 406–409
- thermal de Broglie wavelength, 570
- Thomas precession, 366
  - from Dirac equation, 613
- Thomson, Joseph J., 8
- time evolution of states, 73, 78–79, 82–84
  - in terms of density matrix operator, 90–91
- time ordering of operators, 83, 444
- time reversal (Lorentz transformation), 600
- time translation symmetry and energy conservation, 111
- time-dependent perturbation theory, 448–457
  - harmonic, 457–461
- time-evolution operator, 78–79, 83–84
  - spectral decomposition, 79
- time-independent Schrödinger equation, 79
- time-independent perturbation theory, 321–352
  - almost-degenerate, 348–352
  - Brillouin–Wigner, 334–335
  - Dalgarno–Lewis method, 335–337
  - degenerate, 337–342
  - non-degenerate, 321–326
    - examples, 326–334
- total angular momentum, 120
- total angular momentum basis, 266–267
- trace of operator, 52
- transformation, 108
  - as unitary operator, 108
  - generator, 108–109
  - rotation, 115–121
  - space inversion (parity), 121–123
  - time translation, 111
  - translation, 111–115
- transition amplitude, 444, 620–624
  - diagrammatic representation, 451–452
  - perturbative expansion, 449–452
  - propagator, 620–624
- transition operator (scattering), 492, 497–500
- translation operator, 111–115
  - 3-dimensional, 114–115
- translation symmetry and momentum conservation, 113–115
- transmission and reflection ratios  $T$ ,  $R$ , 148
- transparency in potential scattering, 151–153, 522–523
- transverse (Coulomb) gauge, 107, 463
- trial state for the variational method, 355
- triangle inequality, 27
- trick
  - completing the square, 132–133, 136
  - dot product of angular momenta, 267–268, 270–271, 367, 377
  - dots and lines, 182

- evaluation of  $1/R$  and  $1/R^2$  expectation values of hydrogen atom, 341–342
- evaluation of matrix element using Hamiltonian commutator, 472, 476
- hole as absence of electron, 409, 608
- periodic boundary conditions, 226
- relabeling of summation index, 164, 242, 286, 300
- spectral decomposition of operator, 53–54
- triplet state of two spins, 269–271
- tritium decay, 448
- tunneling, 151–152
- 21 centimeter line (from hyperfine splitting of hydrogen), 271, 372, 478
- two-body problem, 100–101
  - angular momentum, 120
- ultraviolet catastrophe, 11–12
- unbound states, 60, 146
- uncertainty, 77
- uncertainty relation, 77–78
  - for Gaussian wavefunction, 134
  - position-momentum, 78
  - Robertson–Schrödinger, 77–78
- unitary operator, 37, 41
  - and time evolution, 78–79, 83–84
  - as a change of basis, 42–43
  - as a transformation, 108
  - construction from action on subspace, 543–544
  - eigenvalues, 52
  - matrix, construction from orthobasis components, 41, 49
- units conventions, 8, 11, 13, 102–103
- unstable states, 130–131
- valence electrons, 406
- variational method, 355–363
  - examples, 358–363
  - for excited states, 356–358
  - example, 360
  - forgiving nature, 355–356
  - $H_2^+$  ion, 419–422
  - helium atom ground state, 361–363
- vector operator, 292–293
  - projection formula, 296–298
  - selection rules, 299
- vector space, complex linear, 23–24
  - basis and dimension, 25
  - dual, 32–33
  - examples, 24
  - subspace, 26
- velocity
  - group vs. phase, 134–135
  - of electrons in hydrogen, 259
  - of light in vacuum ( $c$ ), 9
- vibrational excitations of molecules, 422–426, 429–431, 435–436
- Virial Theorem
  - in 1-d, 82
  - in 3-d, 96–97
- virtual bound state, 526
- von Neumann entropy, 89
  - for a qubit, 573
- water molecule, 435–437
- wave-particle duality, 13–15
- wavefunction, 14–15, 57
  - continuity, 139
  - momentum, 1-dimensional, 60–61
  - momentum, 3-dimensional, 65
  - position, 1-dimensional, 57
  - position, 3-dimensional, 65
- width ( $\Gamma$ ) in resonant scattering, 516–517
- Wigner functions (for rotations), 195–198
  - general formula, 198
- Wigner–Eckart theorem, 299–303
  - and selection rules, 302–303
  - for spherical harmonics, 288
- Yukawa (screened Coulomb) potential, 364, 504–505
- $Z_2$  discrete group, 121
- Zeeman effect
  - hydrogen atom, 375–381
  - multi-electron atoms, 409–411
- Zeh, H. Dieter, 558
- zero-point energy (harmonic oscillator), 168
- Zurek, Wojciech, 558